

Title:

The impact of a csDMARD in combination with a TNF inhibitor on drug retention and clinical remission in axial spondylarthritis.

Corresponding author:**Michael Nissen,**

Division of Rheumatology,
Geneva University Hospital
26 Avenue Beau-Séjour
CH-1211, Geneva 14
Switzerland

Phone: + 41 79 55 33 003

Fax: + 41 22 372 35 35

michael.j.nissen@hcuge.ch

ORCID iD: 0000-0002-6326-1764

Co-authors:

Bénédicte Delcoigne[#], Clinical Epidemiology Division, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden

Daniela Di Giuseppe, Clinical Epidemiology Division, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden

Lennart Jacobsson, Department of Rheumatology and Inflammation Research, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden.

Merete Lund Hetland, 1) Copenhagen Center for Arthritis Research (COPECARE), Center for Rheumatology and Spine Diseases, Centre for Head and Orthopaedics, Copenhagen University Hospital, Rigshospitalet, Glostrup, Denmark and 2) Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

Adrian Ciurea, Department of Rheumatology, University Hospital Zurich, University of Zurich, Zurich, Switzerland

Lucie Nekvindova, 1) Faculty of Medicine, Charles University, Prague, Czech Republic 2) Institute of Biostatistics and Analyses, Ltd., Brno, Czech Republic

Florenzo Iannone, Rheumatology Unit, DETO, University of Bari, Italy

Nurullah Akkoc, Division of Rheumatology, Department of Medicine, Celal Bayar University, Manisa Turkey

Tuulikki Sokka-Isler, University of Eastern Finland, Faculty of Health Sciences and Jyväskylä Central Hospital, Jyväskylä, Finland

Karen Minde Fagerli, Division of Rheumatology and Research, Diakonhjemmet Hospital, Oslo, Norway

Maria Jose Santos, 1) Department of Rheumatology, Hospital Garcia de Orta, Almada, Portugal. 2) Reuma.pt

Catalin Codreanu, Center of Rheumatic Diseases, University of Medicine and Pharmacy, Bucharest, Romania

Manuel Pombo-Suarez, Rheumatology Service, Hospital Clinico Universitario, Santiago de Compostela, Spain

Ziga Rotar, 1) Department of Rheumatology, University Medical Centre Ljubljana, Ljubljana, Slovenia; 2)

Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia ORCID: 0000-0002-9323-9189

Bjorn Gudbjornsson, 1) Centre for Rheumatology Research (ICEBIO), University Hospital, Reykjavik, Iceland. 2)

Faculty of Medicine, University of Iceland, Reykjavik, Iceland

Irene van der Horst-Bruinsma, Department of Rheumatology, Amsterdam UMC, location VUmc, Amsterdam, Netherlands

Anne Gitte Loft, 1) Department of Rheumatology, Aarhus University Hospital, Aarhus, Denmark and 2)

Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

Burkhard Möller, Department for Rheumatology and Immunology, Inselspital – University Hospital Bern, Bern, Switzerland

Herman Mann, Institute of Rheumatology and Department of Rheumatology, Faculty of Medicine, Charles University, Prague, Czech Republic

Fabrizio Conti, Rheumatology Unit, Department of Clinical, Internal, Anesthesiological and Cardiovascular Sciences, Sapienza University of Rome, Rome, Italy

1
2
3 **Gozde Yildirim Cetin**, Division of Rheumatology, Department of Medicine, Kahramanmaras Sutcu Imam
4 University, Kahramanmaras, Turkey

5 **Heikki Relas**, Rheumatology, Inflammation Center, Helsinki University Hospital, Helsinki, Finland

6 **Brigitte Michelsen**, 1) Division of Rheumatology and Research, Diakonhjemmet Hospital, Oslo, Norway, 2)
7 Division of Rheumatology, Department of Medicine, Hospital of Southern Norway Trust, Kristiansand, Norway

8 **Pedro Avila Ribeiro**, Rheumatology Department, Hospital de Santa Maria, Centro Hospitalar Universitario
9 Lisboa Norte EPE, Lisboa, Portugal; Rheumatology Research Unit, Instituto de Medicina Molecular, Faculdade
10 de Medicina, Universidade de Lisboa, Lisboa, Portugal.

11 **Ruxandra Ionescu**, Sfanta Maria Hospital, University of Medicine and Pharmacy, Bucharest, Romania

12 **Carlos Sanchez-Piedra**, Health Technology Assessment Agency of Carlos III Institute of Health (AETS), Madrid,
13 Spain.

14 **Matija Tomsic**, 1) Department of Rheumatology, University Medical Centre Ljubljana, Ljubljana, Slovenia; 2)
15 Faculty of Medicine, Ljubljana, University of Ljubljana, Slovenia

16 **Árni Jón Geirsson**, Department for Rheumatology, University Hospital, Reykjavik, Iceland.

17 **Johan Askling**, 1) Clinical Epidemiology Division, Department of Medicine Solna, Karolinska Institutet,
18 Stockholm, Sweden; 2) Rheumatology, Theme Inflammation and Ageing, Karolinska University Hospital,
19 Stockholm, Sweden

20 **Bente Grintborg**, 1) DANBIO and Copenhagen Center for Arthritis Research (COPECARE), Center for
21 Rheumatology and Spine Diseases, Centre of Head and Orthopedics, Copenhagen University Hospital,
22 Rigshospitalet, Glostrup, Denmark; 2) Department of Clinical Medicine, Faculty of Health and Medical Sciences,
23 University of Copenhagen, Copenhagen, Denmark

24 **Ulf Lindström**, Department of Rheumatology and Inflammation Research, Sahlgrenska Academy, University of
25 Gothenburg, Gothenburg, Sweden

26
27
28 #Michael Nissen and Bénédicte Delcoigne are joint first authors.
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rheumatology key messages:

Amongst 24,171 axial spondylarthritis patients treated with a first TNF inhibitor, 32% received csDMARD co-therapy.

AxSpA patients treated with co-therapy had more baseline peripheral arthritis and higher CRP levels.

Co-therapy patients demonstrated higher TNFi retention and remission rates, although the clinical relevance is doubtful.

Key words: Spondylitis, Ankylosing, Methotrexate, Sulfasalazine, Tumour Necrosis Factor Inhibitors, Epidemiology

Abstract**Objectives**

Many axial spondylarthritis (axSpA) patients receive a conventional synthetic disease-modifying anti-rheumatic drug (csDMARD) in combination with a tumour necrosis factor inhibitor (TNFi). However, the value of this co-therapy remains unclear. The objectives were to describe the characteristics of axSpA patients initiating a first TNFi as monotherapy compared to co-therapy with csDMARD, to compare one-year TNFi retention and remission rates, and to explore the impact of peripheral arthritis.

Methods

Data was collected from 13 European registries. One-year outcomes included TNFi retention and hazard ratios (HR) for discontinuation with 95% confidence intervals (95%CI). Logistic regression was performed with adjusted odds ratios (OR) of achieving remission (ASDAS-CRP<1.3 and/or BASDAI<2) and stratified by treatment. Inter-registry heterogeneity was assessed using random-effect meta-analyses, combined results were presented when heterogeneity was not significant. Peripheral arthritis was defined as ≥ 1 swollen joint at baseline (=TNFi start).

Results

Amongst 24,171 axSpA patients, 32% received csDMARD co-therapy (range across countries: 13.5% to 71.2%). The co-therapy group had more baseline peripheral arthritis and higher C-reactive-protein than the monotherapy group. One-year TNFi-retention rates (95%CI): 79% (78-79%) for TNFi monotherapy versus 82% (81-83%) with co-therapy ($p<0.001$). Remission was obtained in 20% on monotherapy and 22% on co-therapy ($p<0.001$); adjusted OR of 1.16 (1.07-1.25). Remission rates at 12 months were similar in patients with/without peripheral arthritis.

Conclusion

1
2
3 This large European study of axial SpA patients showed similar one-year treatment outcomes for TNFi
4 monotherapy and csDMARD co-therapy, although considerable heterogeneity across countries limited the
5 identification of certain subgroups (e.g. peripheral arthritis) that may benefit from co-therapy.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Tumour necrosis factor inhibitors (TNFi) are the mainstay biologic for axial spondylarthritis (axSpA), but the value of combination therapy (co-therapy) with conventional synthetic disease-modifying drugs (csDMARD) remains unclear. According to current international recommendations for the management of axSpA, csDMARDs are not indicated for axial disease, although the potential benefit of combining a csDMARD with a TNFi is on the research agenda.¹⁻³ Recent results from the EuroSpA collaboration, including 22,196 axSpA patients across 13 European countries, found that 31% of patients were treated with a csDMARD in combination with a TNFi at the time of biologic treatment start.⁴

In psoriatic arthritis (PsA), the use of a csDMARDs with TNFi are associated with improved remission rates, specifically with methotrexate in combination with either adalimumab or infliximab.⁵ In rheumatoid arthritis (RA), a TNFi in combination with a csDMARD, such as methotrexate, is consistently associated with improved efficacy and a lower risk of TNFi discontinuation compared to TNFi monotherapy.⁶ In contrast, some observational and pharmacokinetic studies have not demonstrated any added benefit of csDMARD combination therapy in axSpA,⁷⁻¹¹ although other cohort studies reported improved drug retention for TNFi when given in combination with a csDMARD, particularly methotrexate.¹²⁻¹⁵

Considering the relatively frequent use of csDMARD co-therapy in axSpA, an improved understanding of the role of csDMARDs in combination with TNFi in axSpA is a key aspect of management, potentially leading to improved patient outcomes. We therefore aimed to explore whether the co-administration of a csDMARD with a TNFi improved treatment outcomes compared to TNFi monotherapy, for either TNFi retention or the attainment of clinical remission, in a large international cohort of axSpA patients.

METHODS

An observational study based on routine care axSpA registries from 13 European countries, with data aggregated through the EuroSpA collaboration (www.eurospa.eu) as previously described.⁴

Data sources

AxSpA patients ≥ 18 years of age, initiating a first TNFi between 2006 and 2017, and registered in one of the following countries (registries): Czech Republic (ATTRA), Denmark (DANBIO), Finland (ROB-FIN), Iceland (ICEBIO), Italy (GISEA), Norway (NOR-DMARD), Portugal (Reuma.pt), Romania (RRBR), Slovenia (biorx.si), Spain (BIOBADASER), Sweden (SRQ), Switzerland (SCQM) or Turkey (TURKBIO).

Follow-up, time-points and treatment group definitions

Patients were followed from the date of their first registered TNFi start (baseline) for 12 months or until treatment discontinuation, loss to follow-up, end of participation in the registry or death, whichever occurred first. Biosimilars were not distinguished from their originator products, and switches between the two were disregarded.

The methods of registration of csDMARD use (methotrexate, sulfasalazine, leflunomide or other) varied across the registries, with some recording start/stop dates, while others recorded treatment status (use/no use) at specific time points (clinical visits). Exposure to co-therapy at baseline (date of TNFi start) was based on start/stop dates for the csDMARDs when available, or otherwise on csDMARD status (use/no use) at the visit closest to baseline (time-window of -100 to +30 days). The 6- and 12-month follow-up visits were defined as the date of the visit closest to the time-point, within the ranges: days 151-270 and days 271-545, respectively.

The *monotherapy group* included patients starting a first TNFi without any concurrent use of a csDMARD at baseline. The *co-therapy group* included patients starting a first TNFi and either: (i) beginning csDMARD at the TNFi start date, (ii) adding TNFi to an ongoing (and continued) csDMARD-treatment, or (iii) csDMARD subsequently added within 30 days after the initiation of TNFi.

Descriptive statistics

Results from descriptive statistical analyses of the two treatment groups, including baseline data are presented both pooled for the whole study population and by individual country, and compared with t-tests for continuous and Chi-square tests for categorical variables.

Comparative statistics, pooling and adjustment

Results from comparative statistical analyses (regression models, as specified below), per country, were combined in a meta-analysis using a random-effects approach, utilizing the Cochran Q-test and the I² statistic to assess the statistical heterogeneity between countries to evaluate the proportion of the total variation that was due to the between-country variation.¹⁶ Pooled comparative estimates for retention or remission were only reported if the heterogeneity was below 50%. Models were adjusted for age, sex, calendar year of TNFi start, baseline Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) (in quartiles), and disease duration (in quartiles). For BASDAI and disease duration, we included a fifth category including missing values in the regression models.

TNFi retention

The overall proportion of study patients remaining on TNFi at 12-months in the monotherapy and co-therapy groups was described with Kaplan-Meier curves. Hazard ratios (HR) with 95% confidence intervals (95%CI) of TNFi discontinuation (with monotherapy as the reference) were calculated using crude and adjusted Cox regression models. We defined discontinuation as cessation of TNFi due to either lack of efficacy or occurrence of an adverse event. Patients discontinuing for other reasons (e.g., pregnancy, remission) were censored.

Clinical remission

Remission at 12 months after TNFi treatment start was defined as either the Ankylosing Spondylitis Disease Activity Score with C-reactive protein (ASDAS-CRP) <1.3 and/or the BASDAI <2 (0-10 scale).^{17,18} Patients discontinuing TNFi treatment before 12 months because of remission were considered to be in remission if they had not initiated a second TNFi treatment within the 12-month period. For patients missing clinical data at 12 months and still on the same TNFi, ASDAS-CRP and BASDAI values at six months were carried forward

(n=15). Patients discontinuing TNFi treatment within the first year with lack of effectiveness or adverse events as the registered reason for discontinuation were considered as not attaining remission. Remission status in patients discontinuing for an unknown reason were considered missing. In sensitivity analyses, we examined BASDAI<2 and ASDAS-CRP<1.3 separately as criteria for achieving remission. Furthermore, we calculated the difference between 12-month and baseline values (deltas) of ASDAS-CRP and BASDAI.

We performed logistic regression to obtain the odds ratios (OR) of achieving remission at 12 months (with monotherapy as reference), stratified by country, with crude and adjusted models. A pooled OR, adjusted for country, in addition to the variables mentioned above, was calculated. To model the 12-month deltas for BASDAI and ASDAS-CRP, we utilized linear regression in each registry and overall, to compare the monotherapy and the co-therapy groups.

Secondary analyses

The overall proportion of patients remaining on TNFi at 5-years in the monotherapy and co-therapy groups was examined. Stratified analyses were performed according to type of TNFi. Furthermore, stratification was performed according to type of csDMARD co-therapy, with the two most common csDMARDs: methotrexate and/or sulfasalazine. This was done either for all TNFi combined or in separate models for the five different TNFi. Further secondary analyses were performed based on stratification by peripheral joint involvement (28 joints) at baseline (swollen joint count (SJC)=0, SJC≥1, or SJC unknown).

Patient and public involvement

Patients were not involved in the study planning.

RESULTS

We included 24,171 axSpA patients from 13 countries. Baseline characteristics of the study cohort are presented in Table 1. The co-therapy group had significantly higher baseline values for CRP, tender/swollen joint counts, and percentages of patients with ≥1 tender/swollen joint, when compared to the monotherapy group, whereas disease activity scores such as BASDAI and ASDAS-CRP were similar. The proportion of patients on co-therapy ranged from 14% (Italy) to 71% (Finland). Percentages of patients on co-therapy: infliximab 38.9%, adalimumab 31.9%, etanercept 29.9%, golimumab 28.0% and certolizumab 26.6%. Baseline characteristics differed markedly between countries, with a mean BASDAI/ASDAS-CRP of 7.4/4.6 in Romania, compared to 3.6/2.7 in Finland (Supplementary Table S1, available at *Rheumatology* online). Similarly, large inter-country differences were observed in the proportions of patients treated with each specific medication: infliximab use varied from 1% (Slovenia) to 84% (Iceland), while amongst patients on co-therapy, methotrexate use varied from 12% (Romania) to 80% (Slovenia). The baseline rate of infliximab use was higher in the co-therapy group versus the TNFi monotherapy group (30% vs. 23%) as was the use of corticosteroids (21% vs. 7%).

TNFi retention

TNFi retention at 12 months differed substantially between countries and varied from 70% (95% CI 68%-71%) in Denmark to 94% (92%-96%) in Romania (Supplementary Figures S1 and S2, available at *Rheumatology* online). Overall, among the 16,359 patients starting their first TNFi as monotherapy, 79% (78%-79%) remained on the same TNFi treatment after 12 months, while the corresponding percentage for the 7,812 patients starting their first TNFi in combination with csDMARD was 82% (81%-83%) (Figure 1). At 12 months, the proportions of patients who had discontinued their TNFi in the monotherapy/co-therapy groups due to an adverse event were 7%/6% and for lack of effectiveness were 12%/10%, respectively. Amongst patients in the co-therapy group who remained on a TNFi at 12 months and had known csDMARD status at 12 months, 97% (5070/5232) remained on a csDMARD. Reasons for TNFi discontinuation in each registry are displayed in Supplementary Table S2, available at *Rheumatology* online.

In analyses of TNFi retention stratified by country, the HR favoured co-therapy in Denmark, Norway, Switzerland and Sweden, while the HR favoured monotherapy in Italy and Turkey (Figure 2). Due to marked heterogeneity ($I^2=79.4\%$, $p < 0.001$), a pooled HR was not estimated.

Clinical remission

Overall numbers and percentages of patients in remission in the treatment groups are displayed in Table 2, demonstrating significantly higher proportions of remission in patients on co-therapy compared to TNFi monotherapy for each type of remission, although the numerical differences were small. Despite variations in the proportions of remission across countries (Supplementary Table S3, available at *Rheumatology* online), there was no statistically significant heterogeneity in the ORs estimates ($I^2=0\%$, $p=0.658$). In Figure 3, the adjusted ORs comparing the probability of attaining remission at 12 months, with TNFi monotherapy as reference, are presented for each country, as well as the meta-analysis. More than three-quarters of the registries demonstrated an OR higher than 1 (favouring co-therapy, although the majority were not statistically significant), which translated to a pooled adjusted OR of obtaining remission for the co-therapy group of 1.16 (1.07-1.25). Using only BASDAI <2 to define remission status provided a pooled OR which was slightly larger (OR=1.29 (1.12-1.48)).

The improvements in ASDAS-CRP and BASDAI over 12 months are displayed in Supplementary Figures S3A and S3B, respectively, for both the individual registries and the overall pooled data (available at *Rheumatology* online). For the pooled data, crude analyses showed a statistically significant difference between the two treatment groups in favour of co-therapy (p -values < 0.001). Following adjustment, the overall delta-ASDAS-CRP remained highly significant ($p=0.008$), although this was not the case for the overall delta-BASDAI ($p = 0.48$).

Secondary analyses

1
2
3 Overall, among the patients starting their first TNFi as monotherapy, 60% (59%-61%) remained on the same
4 TNFi treatment after 5 years, while the corresponding percentage for co-therapy was 65% (63%-66%)
5 (Supplementary Figure S4, available at *Rheumatology* online). Retention rates for TNFi monotherapy versus co-
6 therapy groups out to 5 years stratified by country are depicted in Supplementary Figure S5, available at
7 *Rheumatology* online. Secondary analyses of pooled data for TNFi retention are not presented due to marked
8 heterogeneity.

9
10
11
12
13 Results of the secondary analyses for remission are displayed in Table 3. In adjusted analyses, co-therapy with
14 any csDMARD compared to TNFi monotherapy was associated with statistically significant increases in the OR
15 of remission for infliximab (1.21 (1.01-1.46)), etanercept (1.28 (1.06-1.56)) and golimumab (1.25 (1.00-1.55)),
16 but not adalimumab (1.08 (0.93-1.27)) or certolizumab (0.96 (0.64-1.44)).

17
18
19
20 Regarding the individual csDMARDs, for all TNFi combined, co-therapy with either methotrexate or
21 sulfasalazine individually improved the rates of remission compared to TNFi monotherapy (OR: 1.24 (1.10-
22 1.38) and 1.26 (1.13-1.41) respectively) in adjusted analyses (Table 3). For the individual TNFi, co-therapy with
23 methotrexate significantly improved the proportions of remission with both infliximab and etanercept, but not
24 the other TNFi; while co-therapy with sulfasalazine significantly improved the proportions of remission with
25 infliximab, etanercept and golimumab, but not adalimumab or certolizumab. Comparing TNFi monotherapy to
26 TNFi combined with both methotrexate and sulfasalazine (N = 868, 11% of co-therapy group), we observed a
27 similar pattern with an overall adjusted OR for attaining remission of 1.67 (1.37-2.04) with the combination of
28 the 3 agents.

29
30
31
32
33
34 In patients with both known baseline peripheral joint status and known 12-month remission status (n=8218),
35 23% had a SJC \geq 1 at baseline. SJC status was unknown for 50% of patients with available remission status. In
36 adjusted analyses, the OR for remission at 12 months was higher for co-therapy in the SJC unknown group (1.3
37 (95% CI: 1.1, 1.5)) but was comparable in the SJC=0 and the SJC \geq 1 subgroups (Supplementary Table S4,
38 available at *Rheumatology* online). Secondary analyses in the subgroups of patients with known classification
39 criteria (ASAS or modified New York) were not possible due to the high proportion of missingness (56% and
40 75% respectively).

41 42 43 44 45 46 47 **DISCUSSION**

48
49 In this study of 24,171 patients with axial SpA treated with a first TNFi, co-therapy with a csDMARD was
50 observed in a third of patients and was associated with the presence of swollen joints and higher CRP levels at
51 baseline, compared to patients on TNFi monotherapy. Although patients on co-therapy demonstrated
52 statistically higher TNFi retention and remission rates compared to monotherapy, the differences were
53 numerically small and potentially influenced by residual confounding, and thus unlikely to be of clinical
54 relevance. This is further demonstrated by a NNT of 43 in order to obtain remission in one additional patient
55 treated with a csDMARD.
56
57
58
59
60

1
2
3 Despite marked variability between countries, overall, a higher proportion of patients on co-therapy
4 (compared to those on TNFi monotherapy) remained on TNFi at 12 months. Our findings suggest that a benefit
5 from concomitant csDMARD therapy on TNFi retention was more apparent in countries in the northern part of
6 Europe (particularly Denmark, Norway, Iceland and Sweden) and Switzerland. Interestingly, these countries
7 had considerably lower 12-month TNFi retention rates of around 75% and thus the potential for an additional
8 beneficial effect from a csDMARD could be greater than in countries with retention rates around 90%.

9
10
11
12
13
14 Our findings regarding the impact of co-medication on TNFi retention are consistent with previous studies of
15 axSpA patients in the southern European countries of Greece and Portugal, which did not demonstrate an
16 advantage of csDMARD co-therapy,^{19 20} and are similarly consistent with studies from Sweden and Switzerland
17 that described a benefit with co-therapy.^{12 13} Conversely, a study of axSpA patients in the DANBIO registry from
18 2010 found that the use of methotrexate had no effect on TNFi retention.⁷ Several differences between the
19 included countries could potentially have an impact on TNFi retention. For example, Switzerland and the
20 Scandinavian countries have approximately three-fold higher gross domestic products (GDP), compared to
21 most of the other countries in the EuroSpA collaboration and this may have an impact on a multitude of
22 factors, including access to healthcare and imaging (such as MRI), as well as the availability of certain
23 medications. GDP has been associated with disease activity and treatment retention in RA.^{21 22} Moreover, the
24 use of a csDMARD prior to starting a TNFi is mandatory in some countries for axSpA, such as Finland, where we
25 observed 71% use of csDMARD at TNFi initiation. Local recommendations/guidelines regarding the use of
26 csDMARD and TNFi vary across countries, as does their reimbursement. For example, Iceland demonstrated
27 the highest proportion of infliximab use (84%), where a tender process exists for the cheapest TNFi.

28
29
30
31
32
33
34
35
36
37 A potential role for csDMARD use with a TNFi in axSpA has been hypothesized due to a reduced clinical
38 response to infliximab being correlated with the formation of anti-infliximab antibodies.²³ Studies in SpA have
39 demonstrated that csDMARDs (particularly methotrexate) improve the retention of infliximab more than other
40 TNFi.^{13 24} Moreover, a study of a US administrative claims database reported that in 3812 AS patients,
41 concomitant use of methotrexate, in contrast to other csDMARDs, was associated with a lower adjusted HR
42 (95% CI) for TNFi discontinuation (0.79 (0.67-0.93)).²⁵ Methotrexate was evaluated in 3 RCTs, with a Cochrane
43 review concluding insufficient evidence to support any benefit in the treatment of AS.²⁶ A Cochrane review of
44 sulfasalazine concluded a statistically significant benefit in reducing the erythrocyte sedimentation rate and
45 easing spinal stiffness, although the effect size was not clinically meaningful and there was no effect on pain or
46 other measures of disease activity.²⁷

47
48
49
50
51
52
53
54 Previous studies in axSpA patients looking at clinical effectiveness, including remission, have not
55 demonstrated an overall benefit for csDMARD co-therapy over TNFi monotherapy.^{7 13 14 28} Although, a Swiss
56 study found a significantly higher proportion of patients with an ASDAS clinically important improvement with
57 co-therapy in the subgroup of patients treated with infliximab and Shimabuco *et al.* found a higher rate of
58 ASDAS remission in patients treated with sulfasalazine (p=0.037), but not methotrexate.^{13 28} Our study found
59
60

1
2
3 significantly higher proportions of clinical remission with co-therapy compared to TNFi monotherapy, both
4 overall and individually with methotrexate and sulfasalazine. This difference could be partly explained by the
5 large sample size. The numerical differences between groups were small (e.g. ASDAS-CRP remission
6 proportions of 22% and 24%, in the 2 treatment groups), thus questioning the clinical relevance at the
7 individual patient level.
8
9

10
11
12 In the current study, co-therapy with a csDMARD significantly improved the remission rates with infliximab,
13 etanercept and golimumab compared to TNFi monotherapy, but not with adalimumab or certolizumab. While
14 it is generally assumed that there are no significant differences between the various TNFi in terms of efficacy in
15 axSpA, no head-to-head trials have compared them directly. A change in prescribing habits over time could
16 also explain this to some extent, with infliximab and etanercept representing the first TNFi on the market,
17 while adalimumab is currently the most prescribed TNFi. Similarly, TNFi retention rates may vary over time,⁴
18 although all analyses were adjusted for calendar year of TNFi start. Despite previous studies describing no
19 benefit of co-therapy with etanercept, our findings in a much larger cohort seem robust as in adjusted analyses
20 similar results were found regardless of the csDMARD utilised. Moreover, given that both BASDAI and ASDAS
21 include patient reported disease outcomes not specific to axial disease (ie “fatigue” and “overall discomfort”
22 with BASDAI and “patient global” with ASDAS), there may be some patients who respond better with
23 etanercept in combination with a csDMARD, compared to etanercept monotherapy which is not so efficient on
24 several extra-musculoskeletal manifestations. Finally, it is difficult to make firm conclusions with regards to
25 certolizumab given the relatively small number of patients.
26
27
28
29
30
31
32
33
34

35 Regarding the individual csDMARDs, we demonstrated that both methotrexate and sulfasalazine had a
36 beneficial role on TNFi remission rates. Several studies have suggested that sulfasalazine might be superior to
37 methotrexate in terms of TNFi retention.^{12 14} Based on the literature in other indications, particularly PsA, one
38 could hypothesize that methotrexate and sulfasalazine may be useful in axSpA patients with peripheral
39 synovitis or extra-articular manifestations such as psoriasis or uveitis.^{5 29 30}
40
41
42
43

44 Our study has several limitations to consider. Similar to many prospective cohort studies, we observed high
45 proportions of missingness for many variables, although information on concomitant csDMARDs was almost
46 complete. The missingness of variables was comparable in both exposure groups and therefore is unlikely to
47 have influenced the results. The high missingness of variables such as peripheral joint involvement, psoriasis,
48 uveitis and BMI prevented their inclusion as potential confounders in adjusted analyses. Much of this
49 missingness was structural, as this data is not explicitly collected in several of the registries. We chose a
50 combined remission score due to the relatively high proportion of missingness in ASDAS-CRP remission data.
51 When comparing patients missing the remission outcome to patients having the outcome, there were no
52 clinically relevant differences (data not shown).
53
54
55
56
57
58
59
60

1
2
3 Another potential limitation is residual confounding by factors influencing the use of a csDMARD, such as local
4 regulations, alcohol intake, desire for pregnancy and the presence of comorbidities such as hepatic
5 impairment. Most registries did not include details that allowed the distinction between patients with and
6 without peripheral joint involvement. We attempted to analyse the effect of peripheral joint involvement with
7 examination of the SJC at baseline, however, channelling bias towards concomitant csDMARD use in certain
8 patient groups and the risk of residual confounding could have influenced the results. In addition, the
9 relatively high rates of missingness for peripheral arthritis of around 50% may have prevented more conclusive
10 results. For a limited number of patients from Finland ($n < 50$) in 2017 a systematic recording error may have
11 occurred for BASDAI scores, which would not have the power to effect either the direction or the magnitude of
12 the overall results.
13
14
15
16
17
18
19

20 The use of alternative statistical methods, such propensity scoring, were considered, but ultimately we
21 concluded that it was not feasible considering the high rates of missing data and risk of residual confounding.
22 Furthermore, multiple imputation was considered, but since this would need to be performed separately for
23 each country, and would thus be impacted by the pattern of missingness which was very closely related to the
24 country variable (very high missingness in some countries and almost zero missingness in others), it was
25 deemed inappropriate. Finally, we were not able to explore drug-related adverse events, which may be more
26 common in the co-therapy group.
27
28
29
30
31

32 In conclusion, this large study of 24,171 axSpA patients demonstrated that csDMARD co-therapy was unlikely
33 to have a clinically meaningful effect on neither TNFi retention nor treatment response. Our results do not
34 support the routine use of csDMARD co-therapy with a TNFi. However, we cannot rule out an additional
35 beneficial effect of co-therapy in certain subgroups, such as those with peripheral joint synovitis or extra-
36 articular manifestations, where further studies may be warranted.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1:

Baseline characteristics of patients initiating their first TNFi as monotherapy or in combination with csDMARD.

	Overall population	Missingness (%)	Co-therapy	TNFi monotherapy
Total number (%)	24171	0	7812 (32.3)	16359 (67.7)
Age (years)	42.5 (12.4)	0	43.3 (12.5)	42.2 (12.4)
Gender, n (% male)	14284 (58)	0	4484 (57)	9800 (60)
Disease duration (years)	5.9 (8.1)	22.6	6.4 (7.9)	5.6 (8.2)
Body mass index	26.5 (6.3)	53.8	26.8 (5.3)	26.4 (6.8)
Year of TNFi start	2012 (3)	0	2012 (3)	2012 (3)
BASDAI (/10)	5.6 (2.2)	38.3	5.7 (2.3)	5.6 (2.1)
ASDAS-CRP	3.5 (1.1)	57.1	3.6 (1.1)	3.4 (1.1)
BASFI (/10)	4.5 (2.5)	51.2	4.5 (2.5)	4.4 (2.5)
Tender joints (n)	2.0 (3.9)	55.6	2.6 (4.2)	1.7 (3.8)
At least 1 tender joint, n (%)	4569 (42.6)	55.6	2059 (56.3)	2510 (35.5)
Swollen joints (n)	0.8 (2.0)	49.6	1.3 (2.6)	0.5 (1.7)
At least 1 swollen joint*, n (%)	3092 (25.4)	49.6	1665 (41.2)	1427 (17.5)
VAS global health (/10)	6.1 (2.5)	34.8	6.0 (2.5)	6.1 (2.5)
VAS pain (/10)	6.0 (2.5)	38.6	6.0 (2.5)	6.1 (2.5)
CRP (mg/L)	16.0 (22.9)	23.9	20.5 (26.6)	13.9 (20.9)
Corticosteroid use, n (%)	2513 (12)	15.6	1558 (21)	955 (7)
csDMARD before TNFi, n (%)	11130 (61)	24.2	5042 (77)	6088 (52)
Type of TNFi				
Adalimumab, n (%)	7553 (31)	0	2408 (31)	5145 (31)
Etanercept, n (%)	5883 (24)	0	1757 (22)	4126 (25)
Infliximab, n (%)	6061 (25)	0	2356 (30)	3705 (23)
Golimumab, n (%)	3515 (15)	0	983 (13)	2532 (16)
Certolizumab pegol, n (%)	1159 (5)	0	308 (4)	851 (5)
Type of csDMARD				
Methotrexate, n (%)	-	0.9	4371 (56)	-
Sulphasalazine, n (%)	-	0.5	3538 (45)	-
Leflunomide, n (%)	-	4.3	260 (3)	-
Other, n (%)	-	2.3	344 (4)	-

Values represent the mean (standard deviation) for continuous variables. Otherwise, numbers and percentages are indicated for categorical variables. Percentages were calculated from the total number of patients with available values. Co-therapy = TNFi in combination with a csDMARD, n = number, VAS = Visual Analogue Score, BASDAI = Bath Ankylosing Spondylitis Disease Activity Index, BASFI = Bath Ankylosing Spondylitis Functional Index, ASDAS-CRP = Ankylosing Spondylitis Disease Activity Score using CRP. *Peripheral arthritis was defined as ≥ 1 swollen joint at baseline.

Table 2.

Percentages and numbers of patients achieving clinical remission at 12 months.

		Remission [#] , % (number)	P-value (mono- vs. co-therapy)	No remission [#] , % (number)	Unknown ^{##} , % (number)	NNT
Any remission	TNFi monotherapy	19.5 (2265)		80.5 (9371)	28.9 (4723)	
	Co-therapy	21.8 (1188)	p<0.0001	78.2 (4255)	30.3 (2369)	43
ASDAS remission (ASDAS-CRP<1.3)	TNFi monotherapy	21.8 (2091)		78.2 (7502)	41.3 (6766)	
	Co-therapy	23.7 (1063)	p = 0.011	76.3 (3418)	42.6 (3331)	53
BASDAI remission (BASDAI<2)	TNFi monotherapy	5.9 (679)		94.1 (10902)	29.2 (4778)	
	Co-therapy	7.7 (418)	p<0.0001	92.2 (4993)	30.7 (2401)	56

Any remission was defined as either BASDAI<2 and / or ASDAS-CRP<1.3. BASDAI = Bath Ankylosing Spondylitis Disease Activity Index, ASDAS-CRP = Ankylosing Spondylitis Disease Activity Score using CRP, NNT=number need to treat (for one additional patient obtaining remission with co-therapy over treatment with monotherapy). [#] Percentages are calculated disregarding patients with unknown remission status. ^{##} Percentages are calculated on the total number of patients, including those with unknown remission status. Chi-square tests were used to compare percentages of patients achieving clinical remission at 12 months between groups.

Table 3:

Secondary analyses of the medication-specific differences for TNFi and csDMARD regarding clinical remission.

Clinical remission at 12 months				
	Total number (mono-/co-therapy)	Missingness [#] (%) (mono-/co-therapy)	Adjusted OR ^{##} (95% CI)	% achieving remission ^{###} (monotherapy vs co-therapy)
Any csDMARD				
Infliximab	3705 / 2356	27 / 33	1.21 (1.01-1.46)*	17 vs 18
Adalimumab	5145 / 2408	30 / 29	1.08 (0.93-1.27)	20 vs 23
Etanercept	4126 / 1757	35 / 35	1.28 (1.06-1.56)*	18 vs 22
Golimumab	2532 / 983	22/21	1.25 (1.00-1.55)*	22 vs 27
Certolizumab	851 / 308	18/25	0.96 (0.64-1.44)	24 vs 23
Methotrexate				
Infliximab	3705 / 1605	27 / 36	1.37 (1.11-1.70)**	17 vs 19
Adalimumab	5145 / 1204	30 / 34	1.19 (0.96-1.47)	20 vs 22
Etanercept	4126 / 922	35 / 40	1.35 (1.04-1.75)*	18 vs 20
Golimumab	2532/478	22 / 24	1.25 (0.93-1.67)	22 vs 26
Certolizumab	851/162	18 / 26	1.24 (0.73-2.07)	24 vs 25
Any TNFi	16359 / 4371	29 / 34	1.24 (1.10-1.38)***	20 vs 21
Sulfasalazine				
Infliximab	3705 / 851	27 / 28	1.32 (1.03-1.69)*	17 vs 22
Adalimumab	5145 / 1210	30 / 24	1.18 (0.97-1.43) ^t	20 vs 28
Etanercept	4126 / 848	35 / 28	1.38 (1.08-1.74)**	18 vs 25
Golimumab	2532/493	22 / 17	1.45 (1.10-1.91)***	22 vs 32
Certolizumab	851/136	18 / 23	0.75 (0.41-1.31)	24 vs 21
Any TNFi	16359 / 3538	29 / 25	1.26 (1.13-1.41)***	20 vs 26
Methotrexate + Sulfasalazine				
Infliximab	3705 / 276	27 / 33	1.95 (1.32-2.85)***	17 vs 30
Adalimumab	5145 / 260	30 / 30	1.85 (1.27-2.67)**	20 vs 37
Etanercept	4126 / 197	35 / 28	1.45 (0.92-2.24) ^t	18 vs 26
Golimumab	2532/112	22 / 19	1.69 (1.01-2.78)*	22 vs 36
Certolizumab	851/23	18 / 22	1.01 (0.28-3.15)	24 vs 28
Any TNFi	16359 / 868	29 / 29	1.67 (1.37-2.04)***	20 vs 32

Analyses based on the presence or absence of co-therapy (reference) with a csDMARD. OR = odds ratio, HR = hazard ratio. Odds ratios (OR) are provided with the 95% confidence intervals (95% CI) in brackets.

[#]Proportions of patients missing data on remission status at 12 months. ^{##}Analyses adjusted for age, gender, calendar year, BASDAI (in quartiles) and disease duration (in quartiles). ^{###}Percentages calculated from the

1
2
3 group of patients with available remission status. $t = \text{trend} = 0.05 \leq p < 0.10$, * $p < 0.05$, ** $p < 0.01$, ***
4 $p < 0.001$.
5
6
7

8 **ACKNOWLEDGMENTS AND AFFILIATIONS**

9
10 The authors would like to acknowledge the following: Prof. Gary Macfarlane for his participation; the medical,
11 nursing and technical staff, as well as patients participating in each national registry; all staff at the EuroSpA
12 coordinating centre; and Novartis Pharma AG and IQVIA for supporting the EuroSpA collaboration.
13
14
15

16 **COMPETING INTERESTS**

17
18 MN reports honoraria and/or consulting fees from Abbvie, Celgene, Janssen, Eli Lilly, Novartis and Pfizer. LJ
19 reports lecture and consulting fees from Pfizer, Abbvie, Novartis, Eli Lilly and Janssen. MLH reports research
20 grants from Abbvie, Biogen, BMS, Celltrion, Eli Lilly, Janssen Biologics B.V, Lundbeck Fonden, MSD, Pfizer,
21 Roche, Samsung Biopics, Sandoz, Novartis; MH chairs the steering committee of the Danish Rheumatology
22 Quality Registry (DANBIO), which receives public funding from the hospital owners and funding from
23 pharmaceutical companies; and MH also co-chairs EuroSpA, which generates real-world evidence of treatment
24 of psoriatic arthritis and axial spondylarthritis based on secondary data and is partly funded by Novartis. AC
25 reports consulting and/or speaking fees from AbbVie, Eli Lilly, Merck Sharp & Dohme, Novartis and Pfizer. FI
26 reports speaking and consulting fees from Abbvie, BMS, Galapagos, Janssen, Lilly, Novartis, Pfizer, Roche,
27 Sanofi, UCB. MJS reports speaker fees from Abbvie, AstraZeneca, Novartis and Pfizer. CC reports speaker and
28 consulting fees from AbbVie, Amgen, Boehringer Ingelheim, Eli Lilly, Ewopharma, Novartis, Pfizer. MPS reports
29 speaker and consulting fees from Janssen and MSD. ZR reports speaker and consultancy fees from Abbvie,
30 Amgen, Biogen, Eli Lilly, Janssen, Medis, MSD, Novartis, Pfizer, Roche, Sanofi. BGu reports consulting fees from
31 Amgen and Novartis. IvdH reports research grants from AbbVie, Pfizer, MSD and UCB, and honoraria/speakers
32 fee from Novartis, BMS, Lilly, AbbVie, MSD, Pfizer and UCB. AGL reports speaking and consulting fees from
33 AbbVie, Janssen, Lilly, MSD, Novartis, Pfizer, UCB. HM reports speaking and consulting fees from Abbvie, Eli
34 Lilly, Janssen, Novartis, Pfizer, UCB. GYC reports speaking and consulting fees from Abbvie, Novartis, Pfizer,
35 Roche, MSD, UCB. HR reports consulting fees from Abbvie, Celgene, Pfizer. BMi reports consulting fees and a
36 research grant from Novartis. JA reports research and/or consulting fees from Abbvie, Astra-Zeneca, BMS, Eli
37 Lilly, MSD, Pfizer, Roche, Samsung Bioepis, Sanofi, and UCB. BGI: research grants from Pfizer, BMS, Abbvie.
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **FUNDING**

52 This work was supported by Novartis. Novartis had no influence on the data collection, statistical analyses,
53 manuscript preparation or decision to submit.
54
55
56

57 **ETHICS APPROVAL**

58 The study was approved by the respective national data protection agencies and the local research ethical
59 committees according to legal regulatory requirements in the participating countries.
60

CONTRIBUTORS

All coauthors have contributed significantly in accordance with contributorship guidelines.

DATA AVAILABILITY STATEMENT

The data underlying this article were accessed from the EuroSpA Research Collaboration Network [<https://eurospa.eu/#registries>] The derived data generated in this research could be shared on reasonable request to the corresponding author, but would require approval from all contributing registries.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Wendling D, Lukas C, Prati C, et al. 2018 update of French Society for Rheumatology (SFR) recommendations about the everyday management of patients with spondyloarthritis. *Joint Bone Spine* 2018;85(3):275-84. doi: 10.1016/j.jbspin.2018.01.006
2. van der Heijde D, Ramiro S, Landewe R, et al. 2016 update of the ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis* 2017;76(6):978-91. doi: 10.1136/annrheumdis-2016-210770
3. Ward MM, Deodhar A, Gensler LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol* 2019;71(10):1599-613. doi: 10.1002/art.41042
4. Ornbjerg LM, Brahe CH, Askling J, et al. Treatment response and drug retention rates in 24 195 biologic-naive patients with axial spondyloarthritis initiating TNFi treatment: routine care data from 12 registries in the EuroSpA collaboration. *Ann Rheum Dis* 2019;78(11):1536-44. doi: 10.1136/annrheumdis-2019-215427
5. Lindstrom U, Di Giuseppe D, Delcoigne B, et al. Effectiveness and treatment retention of TNF inhibitors when used as monotherapy versus comedication with csDMARDs in 15 332 patients with psoriatic arthritis. Data from the EuroSpA collaboration. *Ann Rheum Dis* 2021;80(11):1410-18. doi: 10.1136/annrheumdis-2021-220097
6. Nam JL, Winthrop KL, van Vollenhoven RF, et al. Current evidence for the management of rheumatoid arthritis with biological disease-modifying antirheumatic drugs: a systematic literature review informing the EULAR recommendations for the management of RA. *Ann Rheum Dis* 2010;69(6):976-86. [published Online First: 2010/05/08]
7. Grintborg B, Ostergaard M, Krogh NS, et al. Predictors of treatment response and drug continuation in 842 patients with ankylosing spondylitis treated with anti-tumour necrosis factor: results from 8 years' surveillance in the Danish nationwide DANBIO registry. *Ann Rheum Dis* 2010;69(11):2002-8. doi: 10.1136/ard.2009.124446 [published Online First: 2010/06/01]
8. Heiberg MS, Koldingsnes W, Mikkelsen K, et al. The comparative one-year performance of anti-tumor necrosis factor alpha drugs in patients with rheumatoid arthritis, psoriatic arthritis, and ankylosing spondylitis: results from a longitudinal, observational, multicenter study. *Arthritis Rheum* 2008;59(2):234-40. doi: 10.1002/art.23333 [published Online First: 2008/02/02]
9. Breban M, Ravaud P, Claudepierre P, et al. Maintenance of infliximab treatment in ankylosing spondylitis: results of a one-year randomized controlled trial comparing systematic versus on-demand treatment. *Arthritis Rheum* 2008;58(1):88-97. doi: 10.1002/art.23167 [published Online First: 2008/01/01]
10. Mulleman D, Lauferon F, Wendling D, et al. Infliximab in ankylosing spondylitis: alone or in combination with methotrexate? A pharmacokinetic comparative study. *Arthritis Res Ther* 2011;13(3):R82. doi: 10.1186/ar3350 [published Online First: 2011/06/07]
11. Ternant D, Mulleman D, Lauferon F, et al. Influence of methotrexate on infliximab pharmacokinetics and pharmacodynamics in ankylosing spondylitis. *Br J Clin Pharmacol* 2012;73(1):55-65. doi: 10.1111/j.1365-2125.2011.04050.x [published Online First: 2011/06/23]
12. Lie E, Kristensen LE, Forsblad-d'Elia H, et al. The effect of comedication with conventional synthetic disease modifying antirheumatic drugs on TNF inhibitor drug survival in patients with ankylosing spondylitis and undifferentiated spondyloarthritis: results from a nationwide prospective study. *Ann Rheum Dis* 2015;74(6):970-8. doi: 10.1136/annrheumdis-2014-206616 [published Online First: 2015/02/25]

13. Nissen MJ, Ciurea A, Bernhard J, et al. The Effect of Comedication With a Conventional Synthetic Disease-Modifying Antirheumatic Drug on Drug Retention and Clinical Effectiveness of Anti-Tumor Necrosis Factor Therapy in Patients With Axial Spondyloarthritis. *Arthritis Rheumatol* 2016;68(9):2141-50. doi: 10.1002/art.39691
14. Heinonen AV, Aaltonen KJ, Joensuu JT, et al. Effectiveness and Drug Survival of TNF Inhibitors in the Treatment of Ankylosing Spondylitis: A Prospective Cohort Study. *J Rheumatol* 2015;42(12):2339-46. doi: 10.3899/jrheum.150389
15. Hunter T, Schroeder K, Sandoval D, et al. Persistence, Discontinuation, and Switching Patterns of Newly Initiated TNF Inhibitor Therapy in Ankylosing Spondylitis Patients in the United States. *Rheumatol Ther* 2019;6(2):207-15. doi: 10.1007/s40744-019-0148-4
16. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21(11):1539-58. doi: 10.1002/sim.1186
17. Machado P, Landewe R, Lie E, et al. Ankylosing Spondylitis Disease Activity Score (ASDAS): defining cut-off values for disease activity states and improvement scores. *Ann Rheum Dis* 2011;70(1):47-53. doi: 10.1136/ard.2010.138594
18. van der Heijde D, Dougados M, Landewe R, et al. Sustained efficacy, safety and patient-reported outcomes of certolizumab pegol in axial spondyloarthritis: 4-year outcomes from RAPID-axSpA. *Rheumatology (Oxford)* 2017;56(9):1498-509. doi: 10.1093/rheumatology/kex174
19. Flouri ID, Markatseli TE, Boki KA, et al. Comparative Analysis and Predictors of 10-year Tumor Necrosis Factor Inhibitors Drug Survival in Patients with Spondyloarthritis: First-year Response Predicts Longterm Drug Persistence. *J Rheumatol* 2018;45(6):785-94. doi: 10.3899/jrheum.170477
20. Sepriano A, Ramiro S, van der Heijde D, et al. Effect of Comedication With Conventional Synthetic Disease-Modifying Antirheumatic Drugs on Retention of Tumor Necrosis Factor Inhibitors in Patients With Spondyloarthritis: A Prospective Cohort Study. *Arthritis Rheumatol* 2016;68(11):2671-79. doi: 10.1002/art.39772
21. Finckh A, Neto D, Iannone F, et al. The impact of patient heterogeneity and socioeconomic factors on abatacept retention in rheumatoid arthritis across nine European countries. *RMD Open* 2015;1(1):e000040. doi: 10.1136/rmdopen-2014-000040
22. Sokka T, Kautiainen H, Pincus T, et al. Disparities in rheumatoid arthritis disease activity according to gross domestic product in 25 countries in the QUEST-RA database. *Ann Rheum Dis* 2009;68(11):1666-72. doi: 10.1136/ard.2009.109983
23. de Vries MK, Wolbink GJ, Stapel SO, et al. Decreased clinical response to infliximab in ankylosing spondylitis is correlated with anti-infliximab formation. *Ann Rheum Dis* 2007;66(9):1252-4. doi: ard.2007.072397 [published Online First: 2007/05/03]
24. Fagerli KM, Lie E, van der Heijde D, et al. The role of methotrexate co-medication in TNF-inhibitor treatment in patients with psoriatic arthritis: results from 440 patients included in the NOR-DMARD study. *Ann Rheum Dis* 2014;73(1):132-7. doi: annrheumdis-2012-202347 [published Online First: 2013/01/08]
25. George MD, Baker JF, Ogdie A. Comparative Persistence of Methotrexate and Tumor Necrosis Factor Inhibitors in Rheumatoid Arthritis, Psoriatic Arthritis, and Ankylosing Spondylitis. *J Rheumatol* 2020;47(6):826-34. doi: 10.3899/jrheum.190299
26. Chen J, Veras MM, Liu C, et al. Methotrexate for ankylosing spondylitis. *Cochrane Database Syst Rev* 2013;2:CD004524. doi: 10.1002/14651858.CD004524.pub4 [published Online First: 2013/03/02]
27. Chen J, Lin S, Liu C. Sulfasalazine for ankylosing spondylitis. *Cochrane Database Syst Rev* 2014;11:CD004800. doi: 10.1002/14651858.CD004800.pub3 [published Online First: 2014/11/28]
28. Shimabuco AY, Goncalves CR, Moraes JCB, et al. Factors associated with ASDAS remission in a long-term study of ankylosing spondylitis patients under tumor necrosis factor inhibitors. *Adv Rheumatol* 2018;58(1):40. doi: 10.1186/s42358-018-0040-x

- 1
2
3 29. Wilsdon TD, Whittle SL, Thynne TR, et al. Methotrexate for psoriatic arthritis. *Cochrane Database*
4 *Syst Rev* 2019;1:CD012722. doi: 10.1002/14651858.CD012722.pub2
5
6 30. Gangaputra S, Newcomb CW, Liesegang TL, et al. Methotrexate for Ocular Inflammatory
7 Diseases. *Ophthalmology* 2009;116(11):2188-98. doi: 10.1016/j.opthta.2009.04.020
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1




Figure 1. TNFi retention rates to 12 months in TNFi monotherapy versus TNFi + csDMARD co-therapy groups.

The shaded zone represents 95% confidence intervals for pooled data from 13 European registries.

Figure 2




Figure 2. Meta-analysis of TNFi retention (HR)

Meta-analysis of adjusted hazard ratios (HR) and 95% confidence intervals (95% CI) for discontinuation of TNFi based on the presence or absence of co-therapy with a csDMARD by country. Weights are from a random effects analysis. Overall estimate was not presented due to statistically significant heterogeneity ($I^2=79.4\%$, $p < 0.001$).

Figure 3




Figure 3. Meta-analysis of remission (OR)

Meta-analysis of remission (defined as either an ASDAS-CRP <1.3 and/or a BASDAI <2) at 12 months after TNFi treatment start by country, demonstrating the adjusted odds ratios (OR) and 95% confidence intervals (95% CI) based on the presence or absence of co-therapy (reference) with a csDMARD.