



# Facemasks do not lead to abnormal gas exchange during treadmill exercise testing in children

Copyright ©The authors 2022

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact [permissions@ersnet.org](mailto:permissions@ersnet.org)

Received: 29 Oct 2021  
Accepted: 29 Nov 2021

To the Editor:

Facemasks help reducing the spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) during the COVID-19 pandemic. The United Nations Children's Fund and World Health Organization recommend their use in children from 6 years of age [1]. However, many parents worry about health effects of facemasks in children, especially during exercise. They fear that facemasks might cause breathing difficulties, impair oxygen uptake and cause carbon dioxide retention [2]. Parents of children with underlying respiratory problems are particularly concerned and some request medical certificates to exempt their children from wearing facemasks during sports at school.

Studies in adults have shown that it is safe to wear facemasks even during intense physical exercise [3–6] but data for children are lacking. The few paediatric studies included healthy children observed at rest or during mild physical exercise [7, 8]. We thus aimed to assess whether wearing a surgical facemask during intense physical exercise leads to abnormal gas exchange in children with underlying respiratory problems.

We conducted a cross-sectional study nested in the Swiss Paediatric Airway Cohort (SPAC), a multicentre observational cohort study of children referred to paediatric respiratory outpatient clinics for investigation of respiratory problems including wheeze, cough and exercise-induced symptoms [9]. The study is hosted in Bern and has ethical approval (KEB 2016-02176).

Between December 2020 and October 2021, children with exercise-induced symptoms (EIS) performing an exercise-induced asthma (EIA) test on a treadmill were recruited at the Lucerne Children's Hospital, one of the SPAC centres. We obtained informed consent from caregivers. All children had to wear a surgical mask during the EIA test, which was performed according to the American Thoracic Society guidelines [10]. Briefly, the children exercised during 6 (if aged <12 years) to 8 (if aged ≥12 years) min on a treadmill at a heart rate of 80–90% of the predicted maximum (220 – age in years). The forced expiratory volume in 1 s (FEV<sub>1</sub>) was measured before and at 2, 5, 10 and 15 min after exercise. Oxygen saturation (S<sub>pO<sub>2</sub></sub>) was measured before and after exercise with a fingertip oximeter. Capillary blood gas analysis was performed immediately after exercise to assess gas exchange. Measurements were performed in all children, including those who prematurely interrupted the exercise test because of breathing difficulties or exhaustion. We summarised categorical data using counts and percentages, and continuous data using medians, interquartile range and range. Differences between medians were assessed with the Wilcoxon signed rank test.

We included 25 participants. Median age was 14 years (range 9–16 years) and 76% (19) were girls. Median heart rate attained during the exercise test was 185 beats per min (range 175–200 beats per min). Four participants terminated exercise prematurely because of shortness of breath, burning sensation in the throat and chest or dizziness and nausea. One participant reported shortness of breath and throat/chest tightness, and one reported cough during the exercise, but completed it. Two participants had chest tightness, one had shortness of breath and throat tightness, one dry cough, one dizziness and one tachypnoea, at the end of the exercise.

15 children (60%) had a drop in FEV<sub>1</sub> of ≥10%, suggestive of an exercise-induced airway obstruction. Median FEV<sub>1</sub> was thus lower after exercise (2.27 L) than before (2.83 L, *p*<0.001) (table 1). The clinicians



Shareable abstract (@ERSpublications)

No episodes of oxygen desaturation or carbon dioxide retention were observed in this cross-sectional study assessing children with exercise-induced symptoms wearing a surgical facemask during a submaximal treadmill exercise test <https://bit.ly/3GuxhvO>

Cite this article as: Mallet MC, Hitzler M, Lurà M, *et al.* Facemasks do not lead to abnormal gas exchange during treadmill exercise testing in children. *ERJ Open Res* 2022; 8: 00613-2021 [DOI: 10.1183/23120541.00613-2021].



**TABLE 1** Physiological parameters in children wearing a surgical facemask during exercise-induced asthma (EIA) test on a treadmill (N=25)

	Before EIA test			After EIA test			p-value
	Median	IQR	Range	Median	IQR	Range	
<b>Lung function</b>							
FEV <sub>1</sub> <sup>#</sup> (L)	2.83	2.07–3.01	1.53–4.10	2.27	1.92–2.65	1.14–3.39	<0.001
Drop in FEV <sub>1</sub> <sup>¶</sup> (%)				11	7–18	0–43	
<b>Oxygen saturation (%)</b>	98	97–99	95–99	97	96–98	93–99	0.096
<b>Capillary blood gas</b>							
pH				7.30	7.25–7.34	7.15–7.40	
P <sub>CO<sub>2</sub></sub> (kPa)				4.54	4.22–4.99	3.90–5.67	
Bicarbonate (mmol·L <sup>-1</sup> )				16.1	14.7–18.8	12.7–21.3	
Lactate (mmol·L <sup>-1</sup> )				7.9	5.9–10.6	3.0–13.6	

IQR: interquartile range; FEV<sub>1</sub>: forced expiratory volume in 1 s; P<sub>CO<sub>2</sub></sub>: carbon dioxide tension. <sup>#</sup>: after EIA test, FEV<sub>1</sub> was the lowest value recorded within 15 min after exercise <sup>¶</sup>: difference between the pre-exercise FEV<sub>1</sub> value and the lowest FEV<sub>1</sub> value recorded within 15 min after exercise expressed as a percentage of the pre-exercise value. p-values compare the median values before and after the EIA test.

who interpreted the exercise test and considered the full clinical picture diagnosed EIA in 11 children (44%). Other diagnoses included induced laryngeal obstruction, dysfunctional breathing and insufficient fitness level. We observed no episodes of oxygen desaturation after exercise. The lowest S<sub>pO<sub>2</sub></sub> was 93%. We also found no difference in S<sub>pO<sub>2</sub></sub> before and after exercise (p=0.096). Capillary blood gas analysis revealed lactic acidosis in most participants (n=19, 76%) but carbon dioxide retention was not observed in any child. The highest carbon dioxide tension (5.67 kPa) was well below the upper limit of normal (6.0 kPa).

This cross-sectional study does not provide any evidence that wearing surgical facemasks during submaximal treadmill exercise testing leads to oxygen desaturation or carbon dioxide retention in children with EIS. Although exercise-induced airway obstruction occurred in around two-thirds of the children, none had abnormal gas exchange.

As no similar studies have been performed in children, we cannot compare our data to others. LUBRANO *et al.* [8] conducted a study among 47 0–12-year-old healthy children wearing surgical masks during play activities and a 12-min walk test (in those >2 years old). During the first 60 min of evaluation, there was no significant change in S<sub>pO<sub>2</sub></sub> or end-tidal carbon dioxide tension. Studies conducted in adults found that wearing masks could reduce performance but did not affect gas exchange in healthy individuals undergoing moderate to vigorous exercise [3, 4, 6] or in patients with severe lung impairment after a 6-min walk test [5].

Our data do not allow us to state whether the wearing of facemasks can lead to breathing difficulties or exacerbate breathing problems. It is thought that masks can increase airflow resistance with subsequent increased breathing effort, resulting in a feeling of discomfort frequently associated with their use [4, 6, 7]. Further limitations of our study include the small sample size, the short period of exercise, the assessment of surgical masks only and the absence of a control group without facemasks. However, it was not feasible to conduct a controlled study as it was recommended at that time to perform EIA tests with a mask to minimise the risk of SARS-CoV-2 transmission.

In conclusion, this first study assessing the impact of surgical facemasks on oxygen saturation and carbon dioxide retention in children with EIS undergoing submaximal treadmill exercise testing did not reveal evidence of abnormal gas exchange. Even if these findings are reassuring and may help promote the use of surgical masks among children, controlled studies with a larger number of children are needed to confirm our preliminary findings.

Maria Christina Mallet<sup>1,2</sup>, Michael Hitzler<sup>3</sup>, Marco Lurà<sup>3</sup>, Claudia E. Kuehni<sup>1,4</sup> and Nicolas Regamey<sup>3</sup>

<sup>1</sup>Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland. <sup>2</sup>Graduate School for Health Sciences, University of Bern, Bern, Switzerland. <sup>3</sup>Division of Paediatric Pulmonology,

Children's Hospital, Cantonal Hospital Lucerne, Lucerne, Switzerland. <sup>4</sup>Division of Paediatric Respiratory Medicine and Allergology, Dept of Paediatrics, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland.

Corresponding author: Nicolas Regamey ([Nicolas.regamey@luks.ch](mailto:Nicolas.regamey@luks.ch))

Provenance: Submitted article, peer reviewed.

Acknowledgements: We thank the participants, their families and the technicians who performed the EIA treadmill tests.

Conflicts of interest: All authors have nothing to disclose

Support statement: This study was funded by the Swiss National Science Foundation (SNF grant 320030\_182628). Funding information for this article has been deposited with the Crossref Funder Registry.

### References

- 1 World Health Organisation, United Nations Children's Fund. Advice on the use of masks for children in the community in the context of COVID-19: annex to the advice on the use of masks in the context of COVID-19, 21 August 2020. <https://apps.who.int/iris/handle/10665/333919>. Date last accessed: 8 May 2021. Date last updated: 21 August 2020.
- 2 Villani A, Bozzola E, Staiano A, *et al.* Facial masks in children: the position statement of the Italian Pediatric Society. *Ital J Pediatr* 2020; 46: 132.
- 3 Epstein D, Korytny A, Isenberg Y, *et al.* Return to training in the COVID19 era: the physiological effects of face masks during exercise. *Scand J Med Sci Sports* 2021; 31: 70–75.
- 4 Shaw K, Butcher S, Ko J, *et al.* Wearing of cloth or disposable surgical face masks has no effect on vigorous exercise performance in healthy individuals. *Int J Environ Res Public Health* 2020; 17: 8110.
- 5 Samannan R, Holt G, Calderon-Candelario R, *et al.* Effect of face masks on gas exchange in healthy persons and patients with chronic obstructive pulmonary disease. *Ann Am Thorac Soc* 2021; 18: 541–544.
- 6 Mapelli M, Salvioni E, De Martino F, *et al.* "You can leave your mask on": effects on cardiopulmonary parameters of different airway protection masks at rest and during maximal exercise. *Eur Respir J* 2021; 58: 2004473.
- 7 Goh DYT, Mun MW, Lee WLJ, *et al.* A randomised clinical trial to evaluate the safety, fit, comfort of a novel N95 mask in children. *Sci Rep* 2019; 9: 18952.
- 8 Lubrano R, Bloise S, Testa A, *et al.* Assessment of respiratory function in infants and young children wearing face masks during the COVID-19 pandemic. *JAMA Netw Open* 2021; 4: e210414.
- 9 Pedersen ESL, de Jong CCM, Ardura-Garcia C, *et al.* The Swiss Paediatric Airway Cohort (SPAC). *ERJ Open Res* 2018; 4: 00050–2018.
- 10 American Thoracic Society. Guidelines for methacholine and exercise challenge testing – 1999. *Am J Respir Crit Care Med* 2000; 161: 309–329.