

Implementation of an electronic, secure web-based application to support routine hand hygiene observation with immediate direct feedback and anonymised benchmarking

S. Simonet , J. Marschall , R. Kuhn , M. Schlegel , CR. Kahlert

PII: S0196-6553(22)00356-X
DOI: <https://doi.org/10.1016/j.ajic.2022.04.006>
Reference: YMIC 6221



To appear in: *AJIC: American Journal of Infection Control*

Please cite this article as: S. Simonet , J. Marschall , R. Kuhn , M. Schlegel , CR. Kahlert , Implementation of an electronic, secure web-based application to support routine hand hygiene observation with immediate direct feedback and anonymised benchmarking, *AJIC: American Journal of Infection Control* (2022), doi: <https://doi.org/10.1016/j.ajic.2022.04.006>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Inc. on behalf of Association for Professionals in Infection Control and Epidemiology, Inc.

Implementation of an electronic, secure web-based application to support routine hand hygiene observation with immediate direct feedback and anonymised benchmarking

S. Simonet^{a,*}, J. Marschall^b, R. Kuhn^a, M. Schlegel^a, CR. Kahlert^{a,c}

^aInfectious diseases and Hospital Epidemiology, Cantonal Hospital of St. Gallen, St. Gallen, Switzerland

^bInfectious Diseases, Bern University Hospital, Bern, Switzerland, Inselspital Bern, Bern, Switzerland; Division of Infectious Diseases, Washington University School of Medicine, St. Louis, MO, United States

^cInfectious diseases and Hospital Epidemiology, Children's Hospital of Eastern Switzerland, St. Gallen, Switzerland

*Corresponding author: Simon Simonet. Address: Infectious diseases and Hospital Epidemiology, Cantonal Hospital of St. Gallen, Rorschacher Str. 95, 9007 St. Gallen, Switzerland. Tel.: +41 (0)41 494 10 28; E-mail address: simon.simonet@hin.ch

Declaration of Competing Interest

There are no conflicts of interest to declare.

Abstract

A secure web-based electronic tool was developed and implemented to record adherence to hand hygiene in the routine and to provide direct feedback including anonymised benchmarking. It was found suitable for documenting hand hygiene improvements in a local campaign and following rollout to other institutions in 2013, the tool is currently used in >100 hospitals in Switzerland and will play a major part in upcoming national hand hygiene campaigns.

Introduction

An ongoing challenge in modern medicine is the prevention of healthcare-associated infections.[1] Hygienic hand disinfection (or hand hygiene, HH) is considered the simplest and single most effective measure to prevent the transmission of multidrug-resistant microorganisms.[2]

Improving HH adherence by means of continuous education, monitoring using direct observation as gold standard [3] and direct feedback [4] is therefore a key role of any infection control team. In clinical practice, 2/5 of the required hand disinfection are actually omitted. [5]

In 2005/06, as part of a national campaign in Switzerland, measurements of HH adherence were conducted through direct observations with data collection on paper forms.

The main disadvantage of that method was the time-consuming and costly manual processing of the data with a long delay before the results were reported back to the institutions.

Our aim was to develop a standardised electronic tool that eliminates these disadvantages and allows easy digital recording of adherence to HH. The tool should be easy to introduce in everyday clinical routine and then be evaluated for its suitability in the context of a local campaign. In addition to data collection with a uniform methodology for epidemiological

studies, immediate analysis should be feasible to allow for direct feedback with anonymised benchmarking. We hypothesized that this facilitates both the use in training and for disease outbreak interventions.

Methods

'CleanHands' was developed as a platform-independent web application that allows data entry during direct observation of HH adherence via a touch-enabled device and by using a pictogram-based input mask. After data collection via an internet-enabled mobile device, the user interface immediately displays automated graphical results with anonymised benchmarking and allows extensive grouping and filtering to further specify the analysis and make stratifications. A technical description can be found in the supplement [6].

The following independent predictors were considered in the analysis: profession group (nurses, physicians, others), indication (all 5 moments of HH according to WHO [7] as well as "between patients" until 2014 as employed during the national campaign), departments and the temporal relation to an intervention.

Initially, 'CleanHands' was tested at a 700-bed tertiary care hospital when conducting a direct observational study without intervention on the state of HH adherence, and to evaluate 'CleanHands' for everyday clinical routine and as part of a local HH campaign. The multi-modal HH campaign was conducted from January to July 2011. In addition to adherence measurement with instant analysis and direct feedback, the campaign consisted of training and educational sessions and the placement of campaign reminders on the doors of patients' rooms.

HH measures were assessed independently of the wearing of gloves and since hand washing is comparatively rarely done, no distinction between hand disinfection with an alcohol-based preparation and hand washing with soap and water was made in the data collection.

Results

From January 2009 to December 2016, 15,487 indications were recorded by 19 different observers across 46 inpatient wards. The average adherence was 79% (Table 1).

Small but significant differences in HH adherence between the professional groups were detected. The mean HH adherence of nurses was 79%, higher than that of physicians (76%) and paramedical hospital staff (70%).

Mean HH adherence to WHO indications after patient contact (touching the patient 86%, contact with body fluids 86%, patient environment 80% and additionally "between patients" 86%) were significantly higher than before patient contact (touching the patient 67%, aseptic activity 75%).

In addition, 1,380 (9%) non-coded actions (hand disinfection without any indication) were recorded during the study period.

In comparison, a significantly lower mean HH adherence of 71% was observed on the intensive care units compared to normal wards, with values around 80%.

HH adherence increased from 78% before the multi-modal HH campaign to 85% during the campaign and dropped back to 78% later during campaign. Among physicians, there was a significant increase in HH adherence from 69.5% to 87.8% during the campaign, while HH adherence among nurses increased only slightly from 80.5% to 83.4% (Table 2).

Discussion

Here, we demonstrate the successful implementation of a newly developed secure web-based electronic tool to assess HH adherence with instant analysis for direct feedback in clinical routine. As the tool has been adopted and implemented by the Swiss National Centre for Infection Prevention (Swissnoso) in >100 hospitals in Switzerland since August 2014 [8], it does not only support routine HH surveillance and training but also mono- or multicentre HH campaigns and research.

Our results confirm findings from traditional assessments by paper questionnaires.[9] Numerous studies investigated the difference between occupational groups in terms of HH adherence.[5] We found that disparities have decreased overall as a result of the campaign, primarily due to better adherence among physicians.

Significantly lower adherence was observed for pre-patient versus post-patient contact indications. In previous studies, a greater concern regarding self-protection over potential contamination was mentioned as a possible explanation. [5] Targeted training concerning this indication led to an improvement in adherence 'before patient contact' from 59% in 2009 to 71% in 2016.

The intervention as part of a multi-modal campaign showed that further improvement in adherence is possible even where HH adherence is already quite high.[10] However, without regularly raising awareness of the topic and repeated training, the achievements do not seem sustainable.

The study has some limitations. Although the observations were made as inconspicuously as possible, it is likely that the people being observed changed their behaviour during the ongoing data collection. Therefore, adherence tends to be somewhat overestimated (i.e. Hawthorne effect). On the other hand, observations were optimized through structured training of observers and supervision of observations by experienced trainers with the goal of reducing inter-observer variability. Another limitation is that no observations were made during night shifts and weekends when staffing is reduced, which could lead to a slight overestimation of adherence.

In conclusion, "CleanHands" has replaced the tedious traditional hygiene adherence assessment using paper questionnaires at our centre. It has also proven to be a suitable tool in a local HH campaign and beyond that, on a national level for easy benchmarking of this important performance indicator.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

Funding

This study was not funded.

Authors' contributions

Conceptualization and Methodology: SS, MS and CRK; Software: SS; SS, JM, RK, MS and CRK contributed to acquisition, analysis and interpretation of data. SS: wrote the first draft of the manuscript. JM, RK, MS, and CRK critically revised the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank the health care workers participating in this study.

References

- [1] World Health Organization (Who). Report on the Burden of Endemic Health Care-Associated Infection Worldwide. WHO Libr Cat Data 2011:40. http://www.who.int/gpsc/country_work/burden_hcai/en/.
- [2] Sickbert-Bennett EE, DiBiase LM, Schade Willis TM, Wolak ES, Weber DJ, Rutala WA. Reducing health care-associated infections by implementing a novel all hands on deck approach for hand hygiene compliance. *Am J Infect Control* 2016;44. doi:10.1016/j.ajic.2015.11.016.
- [3] Stewardson AJ, Pittet D. Hand hygiene. *Infect. Control Hosp.*, 2014, p. 22.
- [4] Dubbert PM, Dolce J, Richter W, Miller M, Chapman SW. Increasing ICU Staff Handwashing: Effects of Education and Group Feedback. *Infect Control Hosp Epidemiol* 1990. doi:10.1086/646148.
- [5] Erasmus V, Daha TJ, Brug H, Richardus JH, Behrendt MD, Vos MC, et al. Systematic review of studies on compliance with hand hygiene guidelines in hospital care. *Infect Control Hosp Epidemiol* 2010;31:283-94.
- [6] Simonet S, Kahlert C, Schlegel M, Marshall J. "CleanHands" Development - Technical Description 2022. doi:10.5281/ZENODO.6399317.
- [7] Safety WHOP, Organization WH, others. WHO guidelines on hand hygiene in health care. 2009.
- [8] Swissnos National Center for Infection Control website. CleanHands das Modul. 2019. <https://www.swissnos.ch/module/ccm-cleanhands/ccm-cleanhands/das-modul> (Only in German or French available.) (accessed July 17, 2021).
- [9] Allegranzi B, Gayet-Ageron A, Damani N, Bengaly L, McLaws M-L, Moro M-L, et al. Global implementation of WHO's multimodal strategy for improvement of hand hygiene: a quasi-experimental study. *Lancet Infect Dis* 2013;13:843-51.
- [10] Kohler P, Kahlert C, Simonet S, Rettenmund G, Schöbi B, Rafeiner P, et al. improvement of hand hygiene adherence in physicians after a hospital-wide campaign: p1978. *Clin Microbiol Infect* 2012;18:567.

Table 1
Hand hygiene adherence as a function of the different study variables

Variable	Number of indications		Adherence %	Univariate analysis		Multivariate analysis	
	n	(%)		Odds ratio† (95% CI)	P values	Odds ratio† (95% CI)	P values
Occupation							
Nursing‡	12460	(80.5)	79.4	1.00	-	1.00	-
Physicians	2833	(18.3)	76.5	0.84 (0.77 - 0.93)	<.001	0.81 (0.73 - 0.90)	<.001
Other	194	(1.2)	69.6	0.59 (0.44 - 0.81)	<.001	0.73 (0.53 - 1.01)	.059
Indication							
Before patient‡	3985	(25.7)	66.8	1.00	-	1.00	-
Between patients*	1049	(6.8)	85.9	3.03 (2.52 - 3.65)	<.001	3.05 (2.52 - 3.68)	<.001
After patient	5076	(32.8)	85.9	3.02 (2.72 - 3.35)	<.001	3.00 (2.70 - 3.32)	<.001
After body fluid	1300	(8.4)	86.0	3.06 (2.58 - 3.62)	<.001	3.09 (2.60 - 3.68)	<.001
Before invasive procedure	2106	(13.6)	75.0	1.49 (1.33 - 1.68)	<.001	1.43 (1.27 - 1.62)	<.001
After environment	1971	(12.7)	80.0	1.99 (1.75 - 2.26)	<.001	2.07 (1.82 - 2.37)	<.001
Unit							
Medicine‡	4848	(31.3)	83.0	1.00	-	1.00	-
Surgery	4847	(31.3)	79.1	0.77 (0.70 - 0.86)	<.001	0.80 (0.72 - 0.89)	<.001
Gynaecology/Obstetric	166	(1.1)	80.7	0.85 (0.58 - 1.27)	.434	0.84 (0.57 - 1.26)	.411
Intensive care/monitoring	3675	(23.7)	71.5	0.51 (0.46 - 0.57)	<.001	0.53 (0.47 - 0.59)	<.001
Mixed	1686	(10.9)	80.7	0.86 (0.74 - 0.99)	.031	0.91 (0.79 - 1.06)	.235
Neonatology	265	(1.7)	80.0	0.82 (0.60 - 1.11)	.201	0.94 (0.69 - 1.29)	.715
HH campaign (01-06/2011)							
Before‡	3224	(20.8)	78.5	1.00	-	1.00	-
During	1459	(9.4)	85.1	1.56 (1.32 - 1.84)	<.001	1.64 (1.38 - 1.95)	<.001
After	10804	(69.8)	78.0	0.97 (0.88 - 1.07)	.513	1.11 (1.00 - 1.23)	.041
Total	15487		78.7				

A P<0.05 was considered significant. Multivariate analysis using logistic regression models.

† Odds ratio and 95% confidence interval given for adherence to hygiene rules

‡ Reference category for odds ratio

* Recorded until 08/2014

HH = hand hygiene

Table 2

Hand hygiene adherence among occupational groups before, during (01-06/2011) and after local HH campaign

Variable	Number of indications		Adherence %	Univariate analysis	
	n	(%)		Odds ratio† (95% CI)	P values
Nursing‡	12460				-
Before	2602	(20.9)	80.5	1.00	
During	872	(7.0)	83.4	1.21 (0.99 - 1.49)	.062
After	8986	(72.1)	78.7	0.89 (0.80 - 1.00)	.044
Physicians	2833				
Before	560	(19.8)	69.5	1.00	
During	583	(20.6)	87.8	3.17 (2.33 - 4.31)	<.001
After	1690	(59.6)	74.9	1.31 (1.06 - 1.62)	.011

A P<0.05 was considered significant.

‡ Reference category for odds ratio

HH = hand hygiene