

Preoperative application of the Enzian classification for endometriosis (The cEnzian Study): a prospective international multicenter study

Short title: cEnzian Study

Authors:

Simon-Hermann Enzelsberger¹, Peter Oppelt¹, Konstantinos Nirgianakis², Beata Seeber³, Jan Drahoňovský⁴, Leopold Wanderer⁵, Bernhard Krämer⁶, Kristin Nannette Grübling⁷, Sudip Kundu⁸, Darius Salehin⁹, Maciej Mierzwiński¹⁰, Harald Krentel¹¹, Philipp Hermann¹², Helga Wagner^{12,13}, Omar Shebl¹, Sebastian Schäfer¹⁴

Affiliations:

- 1: Department of Gynecology, Obstetrics and Gynecological Endocrinology, Kepler University Hospital, Johannes Kepler University Linz, Altenberger Strasse 69, 4040 Linz, Austria
- 2: Department of Obstetrics and Gynecology, University Hospital and University of Bern, Friedbühlstraße 19, Bern 3010, Switzerland
- 3: Department of Gynecologic Endocrinology and Reproductive Medicine, Medical University of Innsbruck, Anichstraße 35, 6020 Innsbruck, Austria
- 4: Institute for the Care of Mother and Child, Podolské nábřeží 157, 147 00 Prague 4, Czech Republic
- 5: Department of Gynecology and Obstetrics, Landesklinikum Melk, Krankenhausstraße 11, 3390 Melk, Austria
- 6: Department of Gynecology, University Hospital Tübingen, Calwer Straße 7, 72076 Tübingen, Germany
- 7: Department of Gynecology and Obstetrics, University Hospital Carl Gustav Carus, Fetscherstraße 74, 01307 Dresden, Germany
- 8: Department Gynaecology and Obstetrics, Hanover Medical School, Carl-Neuberg-Str. 1, 30625 Hannover, Germany
- 9: Department of Gynecology and Obstetrics, Johanniter Krankenhaus Mönchengladbach, Ludwig-Weber-Straße 15, 41061 Mönchengladbach, Germany
- 10: Department of Gynecology and Obstetrics, St.-Johannes-Hospital, Johannesstraße 9-13, 44137 Dortmund, Germany
- 11: Department of Gynecology, Obstetrics and Gynecological Oncology, Bethesda Hospital, Academic Teaching Hospital, Heerstraße 219, 47053 Duisburg, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](https://doi.org/10.1111/1471-0528.17235). Please cite this article as doi: [10.1111/1471-0528.17235](https://doi.org/10.1111/1471-0528.17235)

12: Center for Clinical Studies, Johannes Kepler University Linz, Altenberger Strasse 69, 4040 Linz, Austria

13: Institute of Applied Statistics, Johannes Kepler University Linz, Altenberger Strasse 69, 4040 Linz, Austria.

14: Department of Gynecology and Obstetrics, University Hospital Münster, Albert-Schweitzer Campus 1, Building A1, 48149 Münster, Germany

Corresponding Author:

Dr. Simon-Hermann Enzelsberger, MSc

Department of Gynecology, Obstetrics and Gynecological Endocrinology, Kepler University Hospital, Johannes Kepler University Linz, Altenberger Strasse 69, 4040 Linz, Austria

- E-Mail: simon-hermann.enzelsberger@kepleruniklinikum.at

- Tel: +43 (0)5 7680 84-0

- ORCID: 0000-0002-3487-4561

Abstract:

Objective: To assess the diagnostic performance of preoperative application of the Enzian classification (cEnzian) using surgical findings as reference standard

Design: A prospective international non-interventional study

Setting: 12 endometriosis centers in four European countries (Austria, Germany, Switzerland, and Czech Republic)

Population: 1062 women with endometriosis surgery

Methods: Extent of endometriosis was preoperatively classified using the cEnzian classification based on gynecological examination and/or transvaginal ultrasound (TVS) and/or magnetic resonance imaging (MRI). After subsequent surgery, the surgeon classified the intraoperative findings using the Enzian classification.

Main Outcome Measures: Sensitivity, specificity, PPV, NPV, LR+, LR- and accuracy were calculated. Conditional frequencies of intraoperative Enzian codings and the corresponding 95 % confidence intervals were computed for each preoperative coding and visualised in plots.

Results: Although overall consistency of cEnzian and Enzian was poor (35.14 %, 95%-CI 32.26-38.03), high specificities and negative predictive values (NPVs) of the cEnzian compartments could be demonstrated. Looking at the individual parts of the Enzian classification, the poorest diagnostic performance was detected for compartment B and the highest PPVs were found for category 3-lesions (> 3 cm), independently of the compartment.

Conclusions: Using the Enzian classification in a non-invasive setting is a useful tool providing us with a 'at a glance' summary of the diagnostic workup regarding deep endometriosis with high specificities and NPVs. An attempt to merge the two new endometriosis classification systems (#Enzian and AAGL 2021) seems reasonable - with consideration of the respective advantages of each other.

Tweetable Abstract: The Enzian classification is a useful tool for a 'at a glance' description of preoperative workup regarding DE

Keywords: endometriosis; classification; Enzian

1. INTRODUCTION/BACKGROUND:

Finding an adequate classification system for endometriosis is a long-lasting and ongoing process. In 2017, a consensus statement of the World Endometriosis Society (WES) was the first global attempt to guide the usage of existing endometriosis classification systems (1). The recommendation was that surgeons should use a uniform classification toolbox including the revised American Society for Reproductive Medicine (r-ASRM) classification (2) and, where appropriate, the Enzian classification (3) for deep endometriosis (DE) and the Endometriosis Fertility Index (EFI) (4) for fertility prediction.

All these above-mentioned classification systems are based on findings during a surgical procedure – usually laparoscopy. This is plausible, since a laparoscopic workup with histological verification was considered the gold standard of endometriosis diagnostics for a long time (5, 6). However, the most accurate preoperative assessment of the extent of the disease is crucial for successful surgical therapy. Furthermore, the rise of new alternative medical therapies and advances in diagnostic imaging techniques have raised the importance of using non-invasive diagnostic tools prior to proceeding with surgery of potentially high morbidity.

The question arises as how accurate current preoperative diagnostics in endometriosis can assess the extent of the disease by using an existing classification system. For the Enzian classification, a few small single-center diagnostic studies have been published so far, based either on sonography (7, 8) or magnetic resonance imaging (9-11). All of these studies showed promising results for a widespread usage in a non-invasive setting.

The objective of this large multicenter study was to understand the diagnostic performance of the preoperative application of the Enzian classification as a summary of non-invasive workup regarding deep endometriosis (using surgical findings as a reference standard).

2. METHODS:

The cEnzian study is an international prospective non-interventional study, which investigates whether or in which quality the findings during subsequent surgery can be predicted by the preoperative application of the Enzian classification (version 2012) based on findings in imaging techniques and gynecological clinical examination. The main research aim was not to perform a diagnostic accuracy study for the single diagnostic modalities, but to assess the diagnostic performance of the complete preoperative evaluation in a large prospective study. The preoperative application of the classification has been named *cEnzian* (for *clinical Enzian*) and is mostly identical to the known Enzian 2012 classification (Figure S1 in the Supporting Information) (3, 12). Only the parameter 'FU' – representing intrinsic ureteric endometriosis in the Enzian version of 2012 – was omitted in the cEnzian, since the preoperative differentiation between intrinsic and extrinsic ureteral endometriosis is usually not possible. For sizing, the largest diameter of the endometriotic lesion was used. If multiple compartments (A, B, C) were involved in one lesion, the size in each compartment was evaluated and coded individually. Precise instructions for using the cEnzian and Enzian classification including the complete definitions of the compartments (Table S5 in the Supporting Information) were printed on each case report form. In 2017, the time of study planning, gathering information on writing these instructions was an enormous effort, because detailed instructions on usage of the latest Enzian classification (version 2012) have never been published and could only be achieved in close cooperation with the creators of the classification. To maybe provide a better reading flow for readers unexperienced in the Enzian classification, we put short definitions of the compartments at the beginning of the respective parts in the result section.

Women aged 18 years or older with a planned endometriosis surgery were recruited from 12 centers in four European countries (Austria, Germany, Switzerland, and Czech Republic). All of the participating centers are specialised endometriosis clinics certified by *EuroEndoCert*. The surgery itself was not a study procedure, as it was planned independently of the study depending on the patient's complaints. The cancellation of the planned surgery or a time interval of more than six months between preoperative evaluation and surgery were defined as dropout criteria. Women without preoperative signs for deep endometriosis could be included in the study as well. A recruitment period of 2 years with at least 1000 included women was planned – with the expectation of a 10 % drop-out rate.

The actual recruitment phase was from January 2018 to October 2019. Following the informed consent and study inclusion, common diagnostic measures were carried out routinely. Depending on the use or need, 'gynecological examination with palpation' and/or 'transvaginal ultrasound (TVS)' and/or 'magnetic resonance imaging (MRI)' were performed. After each examination modality the corresponding section on the CRF was filled in. The final cEnzian was the summary of the results of all

the diagnostic measures preoperatively performed. The examiners for palpation and sonography were gynecologists / gynecologic surgeons. MRI-ratings were assigned by a gynecologist together with a specialised radiologist of the respective center. For imaging evaluation (both sonography and MRI), anatomical structures were defined according to the IDEA consensus of 2016 (13). In a study of Indrielle-Kelly et al (14), it was shown that the IDEA consensus can be properly used for MRI as well. In case of discrepancies between diagnostic modalities, the responsible gynecologist in the study center determined the final cEnzian coding in the relevant compartment taking into account the respective experience in the different diagnostic modalities. An intraoperative examination under anesthesia directly before the start of surgery was not considered in the cENZIAN assignment. Directly after surgery, the surgeon recorded his or her intraoperative impressions in the postoperative section of the questionnaire (Enzian 2012 classification). The histological workup had no influence on the determination of postoperative codings. Due to the local requirements of national societies (recommendation that examiner and surgeon should be the same person if possible) and ethical concerns, the blinding of surgeons was not possible. The completed data sheets were returned pseudonymised to the leading study center in Linz, Austria.

The study was approved by the institutional ethics committee of the Johannes Kepler University (Nr. 1002/2017) and registered in the German Clinical Trials Register (ID: DRKS00013614). The STROBE checklist was used for reporting (15). There was no patient or public involvement (PPI) in this research. Terminology used in this publication is as advised by an international working group of AAGL, ESGE, ESHRE, and WES (16).

Statistical Analysis:

The entire data set was analysed using descriptive statistics. The compartments A/B/C were analysed based on the ratings from 0 to 3 as well as the binary categorization 'affected' / 'not affected' (presence or absence of a DE lesion in a certain compartment). Ratings of the different locations in the F-compartment (FA, FB, FI, FO) are binary ('affected' or 'not affected', no classification by size). The surgery-based Enzian classification was used as gold standard for the evaluation of the preoperative classification. For each compartment sensitivity (SENS), specificity (SPEC), positive predictive value (PPV), negative predictive value (NPV), likelihood ratio of a positive test result (LR+), likelihood ratio of a negative test result (LR-) and accuracy (ACC) were calculated. Taking compartment A as an example, sensitivity describes the correct detection of DE by the cEnzian in compartment A whereas specificity states the correct exclusion of DE by the cEnzian in this compartment. Furthermore, PPV and NPV were calculated for each numeric coding (0-3) in the compartments A, B and C. Normal approximation was used to compute 95 %-confidence intervals. Receiver operating curves (ROCs) could not be determined due to the binary outcomes with no underlying continuous variable.

Accepted Article

Regarding the exact severity grades (0-3) in the compartments A, B and C, conditional frequencies of intraoperative Enzian codings and the corresponding 95% confidence intervals were computed for each preoperative coding and visualised in plots (e.g. frequency of an intraoperative A0, A1, A2, and A3 in the case of a preoperatively coded A0). Heatmaps are provided to visualise the association between the cEnzian and Enzian classification.

Associations between nominal variables are tested using Fisher's exact test. As mentioned above, all the preoperative examinations were performed in specialised endometriosis centers, however by clinicians with different experience in endometriosis diagnostics. To estimate effects of potential covariates (e.g. BMI) while taking into account the potential heterogeneity in endometriosis diagnosis experience of raters or centers, two mixed effects logistic regression models for overall exact consistency with random effects for raters and centers, respectively, were fitted.

Missing values were not replaced. 'Not assessable'-codings in single compartments of the postoperative Enzian classification were transformed to missing values, whereas a 'not assessable'-coding in a cEnzian compartment was not transformed to a missing value but scored as a mismatch (worst-case scenario for the preoperative "not assessable"-codings in the diagnostic performance analysis). The term 'valid ratings' refers to the number of ratings without missing values. Data analysis was performed using the statistics software R (17). Level of significance was set to 5 %.

3. RESULTS:

1144 women were enrolled in the study. After taking into account all of the drop-out criteria, the number of cases was reduced to 1062. The reasons for the drop-outs are listed in Figure S2. The median age in the final dataset was 32 years (IQR 27-37) showing a median BMI of 23.03 (IQR 20.70-26.43).

Of all the women included, 1048 (98.68 %), 1057 (99.53 %), and 168 (15.82 %) were examined with a gynecological examination (including palpation), ultrasound, and MRI, respectively. Except for 1 case, every woman included in the study had at least one imaging modality (TVS or MRI) and not only a gynecological examination with palpation. Preoperatively, deep endometriosis was suspected in 41.34 % (n=439) of cases. After surgery, 53.11 % (n=564) were classified as having deep endometriosis. The distribution of all coded cEnzian and Enzian values is provided in Table S2 in the Supporting Information. In 539 cases (50.75 % of all cases), the cEnzian rater and the Enzian rater were the same person.

Overall consistency of cEnzian and Enzian:

In total, cEnzian and Enzian ratings were exactly equal for 369 women (35.14 % of 1050 valid ratings, 95 % CI 32.26-38.03) – meaning that in these patients all compartments were coded with the exact same values in the preoperative and postoperative classification. The number of matches increased to 423 women (40.29 %, 95 % CI 37.32-43.25) after categorizing the numerical ratings in compartments A/B/C into ‘affected’ (combining the values 1, 2, and 3) and ‘not affected’ (0 coded).

A further analysis with non-consideration of the FA compartment (corresponding to adenomyosis) was performed as well: Then, the number of matches were 486 and 548, respectively for exact and categorized comparison. Focusing only at valid cases with intraoperatively confirmed deep endometriosis (n=553), exact consistency was only seen in 59 women (10.67 %, 95% CI 8.10-13.24) and categorized consistency in 113 patients (20.43 %, 95% CI 17.07-23.79).

Compartment A:

Compartment A describes DE of the rectovaginal septum and/or the vagina and/or generally DE medial of the uterosacral ligaments retrocervically (without rectal wall). 824 of 1058 valid numerical cEnzian ratings (77.88 %) concerning compartment A were exactly equal to the postoperative allocated Enzian A rating. When excluding the cases without intraoperatively confirmed DE in compartment A (corresponding to the exclusion of Enzian A0 ratings), only 115 of 277 ratings were exactly equal (41.52 %). Figure 1 shows the conditional frequencies of intraoperative Enzian codings with their 95 %

confidence intervals for each preoperative coding (cEnzian A0, A1, A2, and A3) and the corresponding heat map.

Categorizing the numerical ratings of compartment A into 'A affected' (A1, A2 or A3 coded) and 'A not affected' (A0 coded) increases the number of matches to 883 of 1058 ratings (83.46 %). Presence of DE in compartment A could be preoperatively detected with a sensitivity of 62.82 % (95% CI 59.90-65.73) and its non-occurrence with a specificity of 90.78 % (95% CI 89.04-92.52). The corresponding PPV and NPV were 71.90 % and 87.64 %, respectively. The further calculated PPVs for the exact cEnzian-codings A1, A2 and A3 were 29.67 %, 44.12 % and 87.76 %, respectively. These numbers for compartment A are also displayed in Table 1. The low PPVs of cEnzian A1 and A2 with a wide spread of corresponding Enzian codings are also clearly visible in Figure 1.

Compartment B:

Compartment B includes DE of the uterosacral ligaments (from the uterine base to the pelvic sidewall), the cardinal ligament and the pelvic sidewall. The calculated values for compartment B can be found in Table 1. In analogy to Figure 1, Figure 2 shows the conditional frequencies of intraoperative Enzian codings with their 95 % confidence intervals for each preoperative B coding and the corresponding heat map.

Looking only at the cases with intraoperatively-coded deep endometriosis in compartment B, the number of exact numerical matches was very low (27.03 %). Using all cases, categorizing into 'B affected' and 'B not affected' improved consistency to 69.22 %. The low number of exact matches is also reflected in the poor PPVs for B1, B2, and B3 (34.38 %, 45.10 % and 68.75 % respectively). In comparison to compartments A and C, compartment B shows the lowest sensitivity (46.59 %, 95% CI 43.58-49.60) and specificity (86.36 %, 95% CI 84.29-88.43).

Compartment C:

Respective numbers for compartment C (DE of the rectum) are again listed in Table 1 and conditional frequencies of intraoperative Enzian codings including the heat map are shown in Figure 3. Specificity was the highest (96.25 %, 95% CI 95.10-97.40) of the three major compartments, crowning the generally high specificity of the cEnzian compartments and resulting in a good NPV of 91.17 % for the C-compartment. On the other hand, poor exact PPVs for C-lesions < 1 cm (C1, 39.29 %) and 1-3 cm (C2, 30.43 %) were observed.

Compartment F:

Codings regarding adenomyosis (FA) had the highest (but still poor) sensitivity of all compartments (71.25 %). The other F-variables (FB, FI, FO) showed a low sensitivity with high specificity (>99 %). Exact values are again listed in Table 1.

Influence of diagnostic modalities used:

The measures of diagnostic accuracy were also computed for each diagnostic method (clinical examination, sonography, MRI) and reported in Table S3. As expected, no postoperative 'FI'-coding could be detected preoperatively via gynecological examination. Sensitivity for the B-compartment was remarkably poor for 'sonography' (20.14 %) and not much better for 'gynecological examination' (38.32 %). Lowest sensitivity overall (besides FI by gynecological examination) was seen for FB-coding by gynecological examination (18.97 %). Preoperative codings by MRI showed the highest sensitivities in comparison to gynecological examination and sonography (except for FA and FO). Overall, cases with MRI showed better overall consistencies than cases without MRI (Fisher's exact test, $p=0.02$). The distributions of the different examination modalities for each numerical Enzian coding are displayed in the Supporting Information (Figures S3, S4, and S5).

Inter-Rater and Inter-Center Variability:

Two separate mixed effects logistic regression models were fitted to estimate effects of predictive factors on the probability of overall exact consistency with random effects for study centers and cEnzian raters, respectively, to take heterogeneity with respect to study centers or cEnzian raters into account. A negative effect of the presence of deep endometriosis and a positive effect of rater relationship (whether the cEnzian and Enzian raters were the same person or different ones) was significant in both regression models (decreasing and increasing the odds ratio of overall consistency, respectively). Overall consistency was not significantly different for patients with preoperative evaluation and surgery more than 90 days apart compared to those with both within 90 days. Details on the mixed effects logistic regression models are provided in the Supporting Information (Table S4).

4. DISCUSSION:

Main Findings:

To our knowledge, the cEnzian study is by far the largest prospective clinical study on the preoperative application of the Enzian classification for endometriosis. Although overall consistency of cEnzian and Enzian was poor, high specificities and good negative predictive values of the cEnzian compartments could be observed in a real-world clinical setting. Looking at the individual parts of the Enzian classification, the poorest diagnostic performance was detected for compartment B and the highest PPVs were found for category 3-lesions (> 3 cm), independently of the compartment.

Interpretation:

The presented poor exact overall consistency between the preoperative and postoperative classifications (35.14% in the whole study population and 10.67% in women with DE) can probably be explained to a large extent by the fact that the Enzian classification is composed of multiple compartments with multiple numerical codings, resulting in over 2000 possible combinations for the Enzian version of 2012. Given this high number of possible combinations for the whole classification, it seems to make more sense to assess the Enzian compartments separately. This is reflected by the fact that none of the previous studies, conducted on the preoperative prediction of the Enzian classification (based on sonography or MRI), evaluated the overall agreement but only the different compartments (7-11).

In 2021, a single-center study by Hudelist et al (7) compared preoperatively-assigned Enzian codings using transvaginal sonography (TVS) with the intraoperatively-assigned ones. The preoperative estimation of deep endometriosis in 195 women using the Enzian classification was satisfactory for the authors, only assessment of deep endometriotic lesions in compartment B was less precise. When looking at the performance of sonography in detail, the published data of Hudelist et al on sensitivities and specificities of TVS regarding Enzian compartments A (SENS 84 %, SPEC 85 %), B (SENS 91 %, SPEC 73%) and C (SENS 92 %, SPEC 95 %) showed clearly better sensitivities in comparison to the cEnzian study. This difference could be explained by the fact that the mentioned study was a single-rater study with an expert in TVS for mapping endometriosis. It is known that the operator's experience in performing TVS is crucial for detecting deep endometriosis via sonography (18). Another prospective study by Goncalves et al (8) on the replacement of diagnostic laparoscopy by transvaginal ultrasound showed a substantial correlation between the compartments A (weighted Kappa = 0.827), B (weighted Kappa = 0.670) and C (weighted Kappa = 0.670) either obtained by TVS with bowel preparation or diagnostic laparoscopy.

Di Paola et al in 2013 (9), Burla et al in 2019 (10), and Thomassin-Naggara et al in 2020 (11) used MRI to assign a preoperative Enzian coding. In the retrospective study of Di Paola (9) including 82 women with histopathologically confirmed deep endometriosis, MRI-based Enzian codings showed a good accuracy in the detection of lesions in the different Enzian compartments (81 % for compartment A, 89 % for B, 82 % for C, 100 % for FA and 37 % for FB). The study of Burla et al (10) was also retrospective containing 63 patients. Sensitivities and negative predictive values for compartment A, B, C, FA, FB, and FI were 95.2 % and 91.7 %, 78.4 % and 56 %, 91.4 % and 89.7 %, 57.1 % and 94.1 %, 85.7 % and 98.3 %, and 73.3 % and 92.2 %, respectively. The authors concluded that the Enzian classification may be used as an anatomical land map and valuable communication tool between radiologists and gynecologists. The third retrospective MRI-based prediction study by Thomassin-Naggara et al (11) included 150 women showing concordance with the surgical Enzian codings for compartment A in 78.7 %, for compartment B in 34.7 % and for compartment C in 82.7 %. Similar to our study, MRI seems to have the worst diagnostic performance in compartment B as well (except the study results of Di Paola 2013).

Beside the study of Hudelist et al (7), none of the listed studies investigated the exact numerical ratings. Comparing the results of the cEnzian study and the data of Hudelist et al, the numeric values of 0 (no DE) and 3 (> 3 cm) in the different compartments could be predicted reliably, while the codings of 1 (< 1cm) and 2 (1-3 cm) showed poor positive predictive values. This might open the discussion that in a future revision of the Enzian classification a fusion of these two size groups should be taken into consideration. A classification that already uses this three centimeter cut-off is the new AAGL 2021 endometriosis classification (19).

Strength and Limitations:

The strengths of the cEnzian study are the prospectively collected data, the large study population recruited by multiple competence centers for endometriosis and the acquisition of real-world data. Nevertheless, some limitations of the study should be considered as well:

- For statistical analysis, the surgery-based Enzian classification was used as a reference. However, since validation studies on the Enzian classification regarding its clinical validity, accuracy and reproducibility are lacking, this should be considered a weakness of the study. Furthermore, in a study from Goncalves et al (8) the accuracy to detect rectosigmoid endometriosis was shown to be better for preoperative TVS than for diagnostic laparoscopy.
- The cEnzian raters are all clinicians part of competence centers for endometriosis, nevertheless the expertise of each rater is different. As mentioned above, especially in the field of mapping endometriosis by transvaginal ultrasound, a certain expertise is necessary. As

a regional feature, it needs to be mentioned that in German-speaking countries TVS for mapping and surgical planning is usually performed by the responsible gynecologists/surgeons themselves and not by a professional subgroup of 'gynecologic sonographers'.

- As rater blinding was not possible for ethical reasons (provision of all available relevant preoperative information to the surgeon), an influence of the Enzian codings by the present cEnzian counterpart could not be excluded (bias towards increasing the ENZIAN prediction quality). A hint towards this effect is reflected in the positive effect of rater agreement in the mixed effects logistic regression model.

A new revised version of the Enzian classification (the "#Enzian"-classification) was released in 2021 (20) – providing a more comprehensive classification for endometriosis with compartments for peritoneal endometriosis (compartment P), ovarian endometriosis (O), and adhesions of the tubo-ovarian unit (T). Some minor changes to the ABCF-compartments were done as well (e.g. separate listing of the B-compartment by left and right side) without relevant impact on the study results presented. Definition for "FU"-coding was changed to "*intrinsic or extrinsic involvement of the ureters with hydroureteric changes or hydronephrosis*" and can now be used in a preoperative setting as well (in contrast to the 2012 version).

5. CONCLUSION:

Using the Enzian classification in a preoperative or non-invasive setting is a useful tool providing us with a short, 'at a glance' summary of the diagnostic workup regarding deep endometriosis. Due to the high specificities and NPVs shown for the different compartments, adequate planning of a surgery seems to be possible as well as invasive procedures may be avoided in case of negative results. Preoperative cEnzian ratings for compartment B are of limited value and the clinical relevance of a subdivision into endometriotic nodules <1 cm and 1-3 cm should be evaluated to consider a possible fusion into a single group. In general, an attempt to merge the two new endometriosis classification systems (#Enzian and AAGL 2021) seems reasonable - with consideration of the respective advantages of each other. If the availability of an expert sonographer is limited, preoperative MRI should be considered.

DECLARATIONS:

Disclosure of interests:

All authors (SE, PO, KN, BS, JD, LW, BK, KNG, SK, DS, MM, HK, PH, HW, OS, and SS) declare that they have no conflict of interest.

Contribution to authorship:

- SE: project development, data collection, data analysis, manuscript writing
- PO: project development, manuscript editing, supervision
- KN: data collection, manuscript editing
- BS: data collection, manuscript editing
- JD: data collection, manuscript editing
- LW: data collection, manuscript editing
- BK: data collection, manuscript editing
- KNG: data collection, manuscript editing
- SK: data collection, manuscript editing
- DS: data collection, manuscript editing
- MM: data collection, manuscript editing
- HK: data collection, manuscript editing
- PH: data analysis, manuscript editing
- HW: data analysis, manuscript editing
- OS: manuscript editing, supervision
- SS: project development, data collection, data analysis, manuscript editing

Funding:

The study was supported by a research grant from ‘Stiftung Endometriose Forschung’. This financial support had no influence on study design, data collection, analysis, interpretation, writing or decision to submit for publication.

Data sharing and data availability:

The data that support the findings of this study are available from the corresponding author upon reasonable request. The data are not publicly available due to privacy or ethical restrictions.

Details of Ethics Approval:

The study was approved by the institutional ethics committee of the Johannes Kepler University (submission number 1002/2017, date of approval 20.12.2017).

Clinical Trial Registration:

DRKS - German Clinical Trials Register, ID: DRKS00013614, <http://www.drks.de/DRKS00013614>

REFERENCES:

1. Johnson NP, Hummelshoj L, Adamson GD, Keckstein J, Taylor HS, Abrao MS, et al. World Endometriosis Society consensus on the classification of endometriosis. *Hum Reprod*. 2017;32(2):315-24.
2. Revised American Society for Reproductive Medicine classification of endometriosis: 1996. *Fertil Steril*. 1997;67(5):817-21.
3. Stiftung-Endometriose-Forschung. ENZIAN 2012. Consensus meeting, 7th Conference of the Stiftung Endometriose Forschung (Foundation for Endometriosis Research); February 2011; Weissensee, Austria.
4. Adamson GD, Pasta DJ. Endometriosis fertility index: the new, validated endometriosis staging system. *Fertil Steril*. 2010;94(5):1609-15.
5. Dunselman GA, Vermeulen N, Becker C, Calhaz-Jorge C, D'Hooghe T, De Bie B, et al. ESHRE guideline: management of women with endometriosis. *Hum Reprod*. 2014;29(3):400-12.
6. Hirsch M, Begum MR, Paniz E, Barker C, Davis CJ, Duffy J. Diagnosis and management of endometriosis: a systematic review of international and national guidelines. *BJOG*. 2018;125(5):556-64.
7. Hudelist G, Montanari E, Salama M, Dauser B, Nemeth Z, Keckstein J. Comparison between Sonography-based and Surgical Extent of Deep Endometriosis Using the Enzian Classification - A Prospective Diagnostic Accuracy Study. *J Minim Invasive Gynecol*. 2021.
8. Goncalves MO, Siufi Neto J, Andres MP, Siufi D, de Mattos LA, Abrao MS. Systematic evaluation of endometriosis by transvaginal ultrasound can accurately replace diagnostic laparoscopy, mainly for deep and ovarian endometriosis. *Hum Reprod*. 2021;36(6):1492-500.
9. Di Paola V, Manfredi R, Castelli F, Negrelli R, Mehrabi S, Pozzi Mucelli R. Detection and localization of deep endometriosis by means of MRI and correlation with the ENZIAN score. *Eur J Radiol*. 2015;84(4):568-74.
10. Burla L, Scheiner D, Samartzis EP, Seidel S, Eberhard M, Fink D, et al. The ENZIAN score as a preoperative MRI-based classification instrument for deep infiltrating endometriosis. *Arch Gynecol Obstet*. 2019;300(1):109-16.
11. Thomassin-Naggara I, Lamrabet S, Crestani A, Bekhouche A, Wahab CA, Kermarrec E, et al. Magnetic resonance imaging classification of deep pelvic endometriosis: description and impact on surgical management. *Hum Reprod*. 2020;35(7):1589-600.
12. Enzian 2012 classification [Available from: <https://www.endometriose-sef.de/aktivitaeten/klassifikation-enzian/>].
13. Guerriero S, Condous G, van den Bosch T, Valentin L, Leone FP, Van Schoubroeck D, et al. Systematic approach to sonographic evaluation of the pelvis in women with suspected endometriosis, including terms, definitions and measurements: a consensus opinion from the International Deep Endometriosis Analysis (IDEA) group. *Ultrasound Obstet Gynecol*. 2016;48(3):318-32.
14. Indrielle-Kelly T, Fruhauf F, Fanta M, Burgetova A, Lavu D, Dundr P, et al. Diagnostic Accuracy of Ultrasound and MRI in the Mapping of Deep Pelvic Endometriosis Using the International Deep Endometriosis Analysis (IDEA) Consensus. *Biomed Res Int*. 2020;2020:3583989.
15. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet*. 2007;370(9596):1453-7.
16. International Working Group of Aagl EE, Wes, Tomassetti C, Johnson NP, Petrozza J, Abrao MS, et al. An International Terminology for Endometriosis, 2021. *Facts Views Vis Obgyn*. 2021;13(4).
17. R-Core-Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2021.

18. Tammaa A, Fritzer N, Strunk G, Krell A, Salzer H, Hudelist G. Learning curve for the detection of pouch of Douglas obliteration and deep infiltrating endometriosis of the rectum. *Hum Reprod.* 2014;29(6):1199-204.
19. Abrao MS, Andres MP, Miller CE, Gingold JA, Rius M, Neto JS, et al. AAGL 2021 Endometriosis Classification: An Anatomy-Based Surgical Complexity Score. *Journal of Minimally Invasive Gynecology.* 2021.
20. Keckstein J, Saridogan E, Ulrich UA, Sillem M, Oppelt P, Schweppe KW, et al. The #Enzian classification: A comprehensive non-invasive and surgical description system for endometriosis. *Acta Obstet Gynecol Scand.* 2021.


Figure 1: Concordance in Compartment A

Figure 2: Concordance in Compartment B


Figure 3: Concordance in Compartment C

		Compartment A	Compartment B	Compartment C	FA	FB	FI	FO / F() *
Exact matches - total		824/1058 (77.88 %)	642/1056 (60.80 %)	899/1054 (85.29 %)	809/1062 (76.18 %)	1015/1062 (95.57 %)	1001/1062 (94.26 %)	1025/1062 (96.52 %)
Exact matches - only DE		115/277 (41.52 %)	123/455 (27.03 %)	52/174 (29.89 %)	389/546 (71.25 %) #	19/59 (32.20 %)	23/76 (30.26 %)	23/53 (43.40 %)
Categorized matches		883/1058 (83.46 %)	731/1056 (69.22 %)	938/1054 (88.99 %)	NA	NA	NA	NA
Sensitivity		62.82 % (59.90-65.73)	46.59 % (43.58-49.60)	52.30 % (49.28-55.31)	71.25 % (68.52-73.97)	32.20 % (29.39-35.01)	30.26 % (27.50-33.03)	43.40 % (40.42-46.38)
Specificity		90.78 % (89.04-92.52)	86.36 % (84.29-88.43)	96.25 % (95.10-97.40)	81.40 % (79.05-83.74)	99.30 % (98.80-99.80)	99.19 % (98.65-99.73)	99.31 % (98.81-99.81)
PPV	categorized	71.90 %	72.11 %	75.83 %	80.21 %	73.08 %	74.19 %	76.67 %
	1: DE < 1cm	29.67 %	34.38 %	39.29 %	NA	NA	NA	NA
	2: DE 1-3 cm	44.12 %	45.10 %	30.43 %	NA	NA	NA	NA
	3: DE > 3cm	87.76 %	68.75 %	58.70 %	NA	NA	NA	NA
NPV		87.64 %	68.47 %	91.17 %	72.79 %	96.14 %	94.86 %	97.09 %
LR+		6.81	3.42	13.95	3.83	46.14	37.30	62.55
LR-		0.41	0.62	0.50	0.35	0.68	0.70	0.57


NA = not applicable



BJO_17235_Figure 1 - cEnzian Study.jpg



BJO_17235_Figure 2 - cEnzian Study.jpg



BJO_17235_Figure 3 - cEnzian Study.jpg