




ELSEVIER

Contents lists available at ScienceDirect

Best Practice & Research Clinical Endocrinology & Metabolism

journal homepage: www.elsevier.com/locate/beem

Basics of androgen synthesis and action

Rawda Naamneh Elzenaty, MSc, PhD Candidate ^{a, b, c},
 Therina du Toit, PhD, Postdoc ^b,
 Christa E. Flück, PhD, MD, Professor ^{a, b, *}

^a Division of Pediatric Endocrinology, Diabetology and Metabolism, Department of Pediatrics, Bern University Hospital, University of Bern, Switzerland

^b Department of Biomedical Research, University of Bern, Switzerland

^c Graduate School of Cellular and Biomedical Sciences, University of Bern, Switzerland

ARTICLE INFO

Article history:

Available online xxx

Keywords:

testosterone
 androgen receptor
 11-oxygenated steroids
 PCOS
 prostate cancer
 androgen deficiency

Androgens are essential sex steroid hormones for both sexes. Testosterone (T) is the predominant androgen in males, while in adult females, T concentrations are about 15-fold lower and androgen precursors are converted to estrogens. T is produced primarily in testicular Leydig cells in men, while in women precursors are biosynthesised in the adrenal cortex and ovaries and converted into T in the periphery. The biosynthesis of T occurs via a series of enzymatic reactions in steroidogenic organs. Notably, the more potent androgen, dihydrotestosterone, may be synthesized from T in the classic pathway, however, alternate metabolic pathways also exist. The classic action of androgens on target organs is mediated through the androgen receptor, which regulates nuclear receptor gene transcription. However, the androgen–androgen receptor complex may also interact directly with membrane proteins or signaling molecules to exert more rapid effects. This review summarizes the current knowledge of androgen biosynthesis, mechanisms of action and endocrine effects in human biology, and relates these effects to respective human congenital and acquired disorders.

© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Pediatric Endocrinology, Diabetology and Metabolism, University Children's Hospital, Freiburgstrasse 15 / C845, 3010 Bern, Switzerland.

E-mail addresses: Rawda.naamneh@students.unibe.ch (R. Naamneh Elzenaty), Therina.dutoit@dbmr.unibe.ch (T. du Toit), christa.flueck@dbmr.unibe.ch (C.E. Flück).

<https://doi.org/10.1016/j.beem.2022.101665>

1521-690X/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article as: R. Naamneh Elzenaty, T. du Toit and C.E. Flück Basics of androgen synthesis and action, Best Practice & Research Clinical Endocrinology & Metabolism, <https://doi.org/10.1016/j.beem.2022.101665>

List of abbreviations

T	testosterone/4-androsten-17 β -ol-3-one
DHT	dihydrotestosterone/5 α -androstan-17 β -ol-3-one
AR	androgen receptor
DHEA-S	dehydroepiandrosterone-sulfate/5-androsten-3 β -ol-17-one-sulfate
DHEA	dehydroepiandrosterone/5-androsten-3 β -ol-17-one
A4	androstenedione/4-androsten-3,17-dione
HPG	hypothalamus-pituitary-gonadal
GnRH	gonadotropin-releasing-hormone
LH	luteinizing hormone
FSH	follicular-stimulating hormone
11OHA4	11 β -hydroxyandrostenedione/4-androsten-11 β -ol-3,17-dione
ACTH	adrenocorticotrophic hormone
MC	mineralocorticoids
GC	glucocorticoids
PREG	pregnenolone/5-pregnen-3 β -ol-20-one
17OHPREG	17 α -hydroxypregnenolone/5-pregnen-3 β ,17 α -diol-20-one
androstenediol	5-androsten-3 β ,17 β -diol
PROG/P4	progesterone/4-pregnen-3,20-dione
17OHPROG/17OHP4	17 α -hydroxyprogesterone/4-pregnen-17 α -ol-3,20-dione
17OH-ALLO	17 α -hydroxy-allopregnanolone/5 α -pregnan-3 α ,17 α -diol-20-one
androsterone	5 α -androstan-3 α -ol-17-one
DHPROG	dihydroprogesterone/5 α -pregnan-3,20-dione
11OHT	11 β -hydroxy-T/4-androsten-11 β ,17 β -diol-3-one
11KT	11-keto-T/4-androsten-17 β -ol-3,11-dione
11KDHT	11-keto-DHT/5 α -androstan-17 β -ol-3,11-dione
11KA4	11-keto-A4/4-androsten-3,11,17-trione
17OH-DHP	17 α -hydroxy-dihydroprogesterone/5 α -pregnan-17 α -ol-3,20-dione
11OHDHT	11 β -hydroxydihydrotestosterone/5 α -androstan-11 β ,17 β -diol-3-one
11KAST	11-ketoandrosterone/5 α -androstan-3 α -ol-11,17-dione
11OHAST	11 β -hydroxyandrosterone/5 α -androstan-3 α ,11 β -diol-17-one
11KP4	11-ketoprogesterone/4-pregnan-3,11,20-trione
11KDHP4	5 α -pregnanetrione/5 α -pregnan-3,11,20-trione
11OHDHP4	11OH-dihydroprogesterone/5 α -pregnan-11 β -ol-3,20-dione
Alfaxalone	5 α -pregnan-3 α -ol-11,20-dione
3 α ,11 β -diOH-DHP4	3 α , 11 β -dihydroxy-dihydroprogesterone
21dF	21-deoxycortisol/11 β ,17 α -dihydroxypregnan-4-ene-3,20-dione
21dE	21-deoxycortisone/17 α -hydroxypregnan-4-ene-3,11,20-trione
11OHPdione	5 α -pregnan-11 β , 17 α -diol-3,20-dione
11KPDione	5 α -pregnan-17 α -ol-3,11,20-trione
zR	zona reticularis
StAR	steroidogenic acute regulatory protein
TSPO	translocator proteins
CYP11A1	cytochrome P450 side chain cleavage P450 _{sc}
FDX1	ferrodoxin
FDXR	ferrodoxin-reductase
CYP17A1	cytochrome P450 17 α -hydroxylase/17,20-lyase
POR	cytochrome P450 oxidoreductase
CYB5	cytochrome <i>b</i> ₅
HSD3B2/3 β HSDII	3 β -hydroxysteroid dehydrogenase type II
HSD17B3	17 β -hydroxysteroid dehydrogenase type 3
AKR1C3/HSD17B5	17 β -hydroxysteroid dehydrogenase type 5
SRD5A2/5 α Red2	steroid-5 α -reductase type II
SRD5A1/5 α Red1	steroid-5 α -reductase type I

HSD11B	11 β -hydroxysteroid dehydrogenase
AKR1C2/4	aldo-keto reductase family 1 member C2/C4
CAH	congenital adrenal hyperplasia
PCOS	polycystic ovary syndrome
CRPC	castration-resistant prostate cancer
CYP11B1	11 β -hydroxylase
NR3C4	nuclear receptor subfamily 3, group C, member 4
NTD	N-terminal domain
AF1	activation functional motif
DBD	DNA binding domain
LBD	ligand binding domain
AF2	activation functional motif
AIS	androgen insensitivity syndrome
APOD	apolipoprotein D
HSPs	heat shock proteins
ARE	androgen response elements
MAPK	mitogen-activated protein kinases
ERK	extracellular signal-regulated kinases
PC	prostate cancer
PI3K	phosphoinositide 3-kinases
Akt	protein kinase B
ARKO	AR knockout mouse
LC-ARKO	Leydig cell-specific ARKO
GU-ARKO	gubernaculum-specific ARKO
DSD	disorders/differences of sex development
CYP21A2	cytochrome P450 21-hydroxylase
CYP19A1	cytochrome P450 aromatase
SF1/NR5A1	steroidogenic factor 1/nuclear receptor subfamily 5 group A member 1
CAIS	complete androgen insensitivity
PAIS	partial androgen insensitivity

Introduction

Androgens are steroid hormones that are essential for human sexual development and reproduction, but they also modulate other organs including bone, muscle, adipose tissue, skin, hair, the brain and the cardiovascular system, thereby effecting growth, body shape and human behavior.

Androgens are produced in the adult female ovaries and male testes and in the adrenal glands, from where they are secreted into circulation to exert their biological effects on target organs. In addition, secreted androgens also serve as substrates for peripheral organs for the intermediate steroid metabolism and its action. Androgens also serve as precursors for estrogen biosynthesis in the gonads and peripherally. Ultimately, the androgens biosynthesised in the adrenals and gonads are metabolized in the liver, and then excreted in urine.

In male adults, testosterone (T, 4-androsten-17 β -ol-3-one) is the most abundant androgen produced in the testes that is present in circulation. Total T concentrations are about 10–30 nmol/L at age 30 years in men and decline at an average rate of 1–2% per year with aging (Fig. 1) [1–6]. T can be converted to the most potent endogenous androgen, dihydrotestosterone (DHT, 5 α -androstan-17 β -ol-3-one) [7,8], as DHT has about 5-10-fold greater affinity for the androgen receptor (AR) compared to T [8]. In a 30 year old woman, the most abundant androgens in circulation are dehydroepiandrosterone-sulfate (DHEA-S, 5-androsten-3 β -ol-17-one-sulfate; 1.2–10 μ mol/L), dehydroepiandrosterone (DHEA, 5-androsten-3 β -ol-17-one; 0.1–23 nmol/L) and androstenedione (A4, 4-androsten-3,17-dione; 0.5–7.9 nmol/L) [9], which are all considered weak androgens according to their low affinity towards the AR [10]. However, these weak androgens can be metabolised to more potent androgens (such as T or DHT) in peripheral tissues through multiple pathways. In menstruating women, circulating androgens originate in part from the adrenal cortex (mainly DHEA(-S)) and more so from the ovaries (A4)




Fig. 1. Lifetime curves of circulating testosterone (T) concentrations in males and females. **A.** Prenatal period. Androgens are produced by the fetal testis and adrenals, while T is predominantly biosynthesised in the testes. In the fetal testis, the Leydig cells start producing T for sexual differentiation of male sex organs at 6 weeks of gestation. T levels keep rising until the end of the second trimester, peaking with values similar to levels measured in adult men; thereafter T's concentration decrease until the end of the third trimester. T levels in a female fetus remain low during pregnancy. **B.** Neonatal Period. At birth (0–3 days of life), T levels are increased in males due to a surge of luteinizing hormone (LH)/follicular-stimulating hormone (FSH), and again rises with minipuberty between 2 and 3 months of life stimulated once again by LH. **C.** Pubertal and adult period. During adrenarache around the age of 6–8 years, T levels increase slightly in both sexes. Finally, activation of the hypothalamic–pituitary–gonadal axis at the age of 12 years (puberty in boys) results in a significant increase in T levels peaking between the second and fourth decade of life in men. With aging, however, T production slightly decreases in men. Although T levels also rise during puberty in females, after about 10 years of age, T remains markedly lower in middle-aged women and decreases further with menopause.

[11]. Total plasma T levels of a 30-year-old woman are about 10 to 15-fold lower than in a same-aged male (0.4–2.1 nmol/L) (Fig. 1) [2,3]. After menopause, when ovarian steroidogenesis has ceased, circulating A4 levels are cut in half and total T levels decrease by about 25% [12].

Production of androgens in the adult gonads is controlled by the hypothalamus-pituitary-gonadal (HPG) axis, which involves the gonadotropin-releasing-hormone (GnRH), luteinizing hormone (LH) and follicular-stimulating hormone (FSH), and comprises of balanced feed forward and feedback loops (Fig. 2). This system is sex specific and thus characterized by the testis or ovary as target organs. Within the testis, Leydig cells produce and secrete T abundantly. By contrast in the ovary, theca cells produce A4 that is mostly transferred to granulosa cells as precursor for the production of estrogens, while only small amounts are secreted into circulation (Fig. 2). The adult adrenal cortex produces adrenal androgens in its zona reticularis (zR) (predominantly DHEA/-S, A4 and 11 β -hydroxyandrostenedione [11OHA4, 4-androsten-11 β -ol-3,17-dione]) [13]. However, unlike androgen production in the HPG axis, the exact regulation of adrenal androgen production is still not fully elucidated, although adrenocorticotrophic hormone (ACTH) is an important co-regulator [11].

The human gonads and adrenals contribute to androgen production pre- and postnatally in a developmental specific fashion [14]. As both the adrenals and the gonads originate from a common embryologic anlage, the urogenital ridge, they share many genetic and steroidogenic characteristics. Prenatally, the fetal adrenal cortex consists of a large fetal zone that resembles the adult zR and produces predominantly DHEA-S which, together with the conversion to its hepatic 16 α -hydroxylated metabolite, 16 α -DHEA-S (5-androsten-3 β ,16 α -diol-17-one-sulfate), contribute to the androgen pool of the fetal-placental unit [11,14]. These androgens are efficiently converted to estradiol and estriol in the placenta due to its aromatase activity, thereby regulating the androgen exposure to the fetus and the mother.

After birth the fetal adrenals involute and give rise to the adult adrenals which start to produce mineralocorticoids (MC) and glucocorticoids (GC) immediately after birth, while production of androgens in the zR occurs only around age 6–8 years at adrenarche. Similarly, the fetal testis is formed early and produces high amounts of T throughout the first and second trimester of pregnancy to masculinize the male fetus (Fig. 1) [7,15]. After birth T levels fall dramatically but raise again for a short time in the event of minipuberty (days 30–100 of life) predominantly in males [16,17]. After that T levels stay low until puberty (Fig. 1). In contrast to the testis, the fetal ovary seems steroidogenic inactive during the prenatal and neonatal period and starts its steroid production only with puberty (Fig. 2).

While the biochemistry of the classic androgen biosynthesis pathway is long known and the action of its conventional products on the AR well described, alternative pathways and novel active androgens have recently been reported (Fig. 3) [11]. In addition, newer studies on the regulation of the AR itself and its stimulated signaling have enhanced our knowledge on androgen action in health and disease [18]. The purpose of this review is to give an update on recent advances in androgen synthesis and action. In our literature search we therefore focused on the literature of the past decade whenever possible.

Androgen synthesis and metabolism

Androgen biosynthesis by the classic pathway

All steroid hormones are produced from cholesterol through a cascade of enzymes which are encoded by genes that are common to all steroid producing organs [11,19]. In most cases the organ specific gene expression determines the steroid profile of each specialized organ. For instance, the testis is determined to produce T from cholesterol while the theca cells in the ovary produce A4 (Figs. 2 and 3A). Cholesterol is transported to the inner mitochondrial membrane by the steroidogenic acute regulatory protein (StAR). Its cooperation with translocator proteins (TSPO) for this transport has been suggested but remains controversial [20]. Inside the mitochondria, cholesterol is the substrate for the first step of steroidogenesis. In the mitochondria, cholesterol is converted to pregnenolone (PREG, 5-pregnen-3 β -ol-20-one) by the side chain cleavage system comprised of CYP11A1 (cytochrome P450 side chain cleavage, P450sc), ferredoxin (FDX1) and ferredoxin-reductase (FDXR). In the classic

Hypothalamic-Pituitary-Gonadal Axis




Fig. 2. Schematic diagram of the hypothalamic-pituitary-gonadal (HPG) axis, resulting in androgen biosynthesis and their biological actions. Androgens are produced upon central gonadotropin-releasing-hormone (GnRH)-luteinizing hormone (LH)/follicular-stimulating hormone (FSH) stimulation of the gonads. In the testicular Leydig cells, the most abundant androgen testosterone (T) is produced at a daily rate of 5–7 mg. About 90–95% of the circulating T reaches the target organs and exerts its effect via the androgen receptor (AR) (left thick green arrow), while 5–10% of T is converted to dihydrotestosterone (DHT) by 5 α -reductase activity in the prostate or skin. Only a small amount of T (0.1%) is converted to estrogens by aromatase (CYP19A1) activity in peripheral tissues (e.g., bone, fat or the brain), where these hormones exert their effect via the estrogen receptor (ER). In the female ovarian theca cells, androstenedione (A4) is produced, where most of it is converted to estradiol by aromatase in the ovarian granulosa cells. Only a small amount of A4 is converted peripherally to T and DHT in hair follicles or in sweat glands. Eventually all steroids, including T, DHT and estrogens, exert negative feedback on the HPG axis.

pathway, PREG is then converted through the delta-5 pathway by cytochrome P450 17 α -hydroxylase/17,20-lyase (CYP17A1, P450c17) to 17 α -hydroxypregnenolone (17OHPREG, 5-pregnen-3 β ,17 α -diol-20-one) and DHEA with the first reaction catalyzed by its 17 α -hydroxylation activity supported by cytochrome P450 oxidoreductase (POR) and the second reaction by the 17,20-lyase activity supported by POR and additionally also cytochrome *b*₅ (CYB5). DHEA is then turned over to A4 or androstenediol (5-androsten-3 β ,17 β -diol) by the enzymes 3 β -hydroxysteroid dehydrogenase type II (HSD3B2/3 β HSDII) and 17 β -hydroxysteroid dehydrogenase type 3 or type 5 (HSD17B3 or AKR1C3 also known as HSD17B5) (Fig. 3A), or to T through the action of both enzymes. Thereafter, T is converted to DHT by steroid-5 α -reductase activities; either type II (SRD5A2/5 α Red2), which is specifically expressed in skin and the prostate, or type I (SRD5A1/5 α Red1), which is expressed more abundantly in the liver, cerebellum and hypothalamus [21]. In humans, only little conversion to A4 occurs through the delta-4 pathway following the conversion of PREG to progesterone (PROG, 4-pregnen-3,20-dione) to 17 α -hydroxyprogesterone (17OHPROG, 4-pregnen-17 α -ol-3,20-dione) and then to A4, as 17OHP4 is a poor substrate of CYP17A1 compared to 17OHPREG [22].

In the ovary, androgen production leads to A4 in theca cells (Fig. 2), which is either further converted to estrogens in the granulosa cells or secreted into the periphery where it serves as a substrate for the intermediate androgen metabolism. Likewise, adrenal androgens such as DHEA, DHEA-S and 11OHA4 are secreted into the peripheral androgen pool and further converted to active and inactive (androgen) products by different pathways that have been explored in greater detail recently, but are still not fully understood [23–25].

Novel androgens and biosynthesis pathways

In addition to the well-known classic pathway of T and DHT biosynthesis, alternative biochemical pathways and novel bioactive androgens have been revealed more recently (Fig. 3B–D) [26,27]. It has been recognized that the intermediate metabolism and peripheral androgen pool play an important role for the formation of active androgen metabolites that has thus far mostly been regarded as 'inactive' products on their way out of the body. This has been illustrated for several disorders which divert normal steroidogenesis and produce alternative steroids, and these disorders therefore serve as models to fully understand the role of novel steroids and metabolic pathways.

Androgen production by the so-called **backdoor, alternative pathway** (Fig. 3B) was first described while studying sexual development of the tamar wallaby pouch young [28,29]. The same pathway has also been described in mice [30], before its metabolites were identified in the human steroid disorders, POR deficiency [31,32] and 21-hydroxylase deficiency [33]. Soon after, the first human mutations in genes (aldo-keto reductase family member C2/C4, AKR1C2/4) specific to this pathway were reported, thereby establishing a role for this pathway in humans [34]. In this pathway, mainly 17OHPROG is 5 α -reduced (by SRD5A1) and then 3 α -reduced (by AKR1C2/4), producing 17 α -hydroxyallopregnanolone (5 α -pregnan-3 α ,17 α -diol-20-one), a perfect substrate for CYP17A1, which is then converted to androsterone (5 α -androstan-3 α -ol-17-one) (Fig. 3B). Androsterone is thereafter converted to androstenediol (5 α -androstan-3 α ,17 β -diol) and finally to DHT. Alternatively, the backdoor pathway may start with the 5 α -reduction of PROG, producing dihydroprogesterone (5 α -pregnan-3,20-dione) (Fig. 3B). Of note, androsterone and androstenediol are androgenic, however, androsterone has only an AR affinity of one-seventh compared to T and androstenediol has only very low affinity and thus weak androgenic activity [35].

The existence of 11-oxy androgens has long been known, but that the human adrenal cortex produces abundant precursors (predominantly 11OHA4, less 11 β -hydroxy-T [11OHT, 4-androsten-11 β ,17 β -diol-3-one]) [10] to feed this **C11-oxy pathway** and provide an important source for the production of active androgens in the periphery (11-keto-T [11KT, 4-androsten-17 β -ol-3,11-dione], 11-keto-DHT [11KDHT, 5 α -androstan-17 β -ol-3,11-dione]), has only recently been recognized (Fig. 3C) [13,36]. In this C11-oxy pathway, the 11 β -hydroxylation (by 11 β -hydroxylase, CYP11B1) of A4 in the adrenals leads to 11OHA4 [37], which can be converted to 11-keto-A4 (11KA4, 4-androsten-3,11,17-trione) (by 11 β -hydroxy steroid dehydrogenase type II, HSD11B2), then 11KT (by HSD17B3/5) and finally 11KDHT (by SRD5A1/2) in peripheral organs [38]. Other parallel metabolic paths may also be used, but they do not lead to other active androgens (Fig. 3C) [27,35]. Overall, 11-oxy androgens are associated with androgen




Fig. 3. Classic and alternative androgen biosynthesis pathways. A. The classic pathway of testosterone (T) biosynthesis, metabolically active in endocrine organs such as the human testis Leydig cells. Cholesterol is transported to the inner mitochondrial membrane by steroidogenic acute regulatory protein (StAR). At the inner mitochondrial membrane cholesterol is then converted to pregnenolone (PREG) by the side-chain cleavage catalytic unit consisting of the enzyme cytochrome P450 side chain cleavage (CYP11A1), ferredoxin (FDX1) and ferredoxin-reductase (FDXR). PREG is then sequentially converted to 17 α -hydroxypregnenolone (17OHPREG) and dehydroepiandrosterone (DHEA) by cytochrome P450 17 α -hydroxylase/17,20-lyase (CYP17A1) supported by cytochrome P450 oxidoreductase (POR) in the first hydroxylase step and cytochrome b₅ (CYB5) in the second lyase step. Following the biosynthesis of DHEA, both androstenediol and androstenedione (A4) are biosynthesized by the enzyme 17 β -hydroxysteroid dehydrogenase (AKR1C3/HSD17B3) and 3 β -hydroxysteroid dehydrogenase (HSD3B2), respectively. Thereafter, both metabolites are

excess disorders such as congenital adrenal hyperplasia (CAH), premature adrenarche and polycystic ovary syndrome (PCOS) [13,24,39–46], and diseases such as castration-resistant prostate cancer (CRPC) [47–50] and (most recently) breast cancer [51].

Most recently, a possible connection between the C11-oxy and the backdoor pathways has been suggested through *in vitro* studies (Fig. 3D) [52–55]. The entry point to this **C11-oxy backdoor pathway** could be either through PROG or 17OHPROG, which are 11 β -hydroxylated by CYP11B in the first step, and then turned over to 11KDHT (a very potent androgen) by multiple conversions involving enzymes common to the backdoor pathway (Fig. 3B-D). Whether this pathway plays a role in human biology, to what degree and under which circumstances remains to be studied.

Androgen action and the androgen receptor

Androgens exert their effect typically through the AR and modulate a vast range of different biological processes (Table 1) [56–58]. Accordingly, the AR is differentially expressed in many tissues including female and male reproductive organs, bones, muscles, the brain, the cardiovascular system, neural tissues, as well as the immune and haematopoietic systems (Fig. 4). Once androgens are bound to the AR, their effect may be at the genomic or at the non-genomic level (Fig. 4) [7,57,59].

Characteristics and regulation of the androgen receptor

The AR is a member of the class I nuclear receptor transcription factor family, also known as NR3C4 (nuclear receptor subfamily 3, group C, member 4). It is a ligand-activated transcription factor. In humans, the AR gene is located on the X chromosome at position Xq11–12. Hence, 46,XX females have two copies of the AR and 46,XY males have only one copy. The AR consists of eight exons which are separated by relatively large introns (Fig. 4A). The AR expressed region encodes a 110-kDa protein of

converted to T via the same enzymes. Small amounts of T or A4 may also be converted to estrogens (estradiol and estrone, respectively) in the testis by aromatase (CYP19A1). Secreted T is then partially converted to dihydrotestosterone (DHT) in peripheral tissues by steroid-5 α -reductases (SRD5A1/2). **B. The alternative backdoor pathway.** In this pathway 17 α -hydroxyprogesterone (17OHPROG/17OHP4) or progesterone (PROG/P4) serve as substrate for 5 α -reduction (by SRD5A1) to enter the backdoor pathway to DHT synthesis. SRD5A1 converts 17OHPROG to 17 α -hydroxy-dihydroprogesterone (17OH-DHP, 5 α -pregnan-17 α -ol-3,20-dione) and the 3 α -hydroxyandrogen dehydrogenase activity of AKR1C2/4 will yield 17 α -hydroxy-allopregnanolone (17OH-ALLO); this metabolite is then converted to androsterone by CYP17A1. Further conversion by AKR1C3/HSD17B3 leads to androstenediol and by AKR1C2/4 or retinol-like dehydrogenase (RoDH) to DHT. **C. The alternative C₁₁-oxy pathway.** The starting point of this pathway is the 11 β -hydroxylation of A4 and T (by 11 β -hydroxylase (CYP11B1)) yielding 11 β -hydroxyandrostenedione (11OHA4) and 11 β -hydroxytestosterone (11OHT), which are then metabolized by SRD5As to 11 β -hydroxy-5 α -androstane-3,17-dione (11OH5 α DIONE) and 11 β -hydroxydihydrotestosterone (11OHDHT, 5 α -androstan-11 β ,17 β -diol-3-one), respectively. Alternatively, 11OHA4 can also be metabolized to 11-ketoandrostenedione (11KA4) by 11 β -hydroxysteroid dehydrogenase (HSD11B). 11KA4 is reversibly metabolized to 11-ketotestosterone (11KT) by HSD17Bs, and the latter to the 11-ketoandrostanolone (11KDHT) by SRD5A. Furthermore, 11OH5 α DIONE can be converted to 11K5 α DIONE, and 11OHDHT to 11KDHT by HSD11B. 11K5 α DIONE and 11KDHT are both reversibly metabolized to 11-ketoandrosterone (11KAST, 5 α -androstan-3 α -ol-11,17-dione) and 11keto-5 α -androstane-3 α ,17 β -diol (11K3 α DIOL) by AKR1C2/4, while 11OH5 α DIONE is also reversibly metabolized to 11 β -hydroxyandrostosterone (11OHAST, 5 α -androstan-3 α ,11 β -diol-17-one), and 11OHDHT to 11OH-3 α -androstane-diol (11OH3 α DIOL) by AKR1C2/4. Finally, 11OHAST and 11OH3 α DIOL are metabolized to 11KAST and 11K3 α DIOL by HSD11B. **D. The alternative C₁₁-oxy backdoor pathway.** In this pathway the 11 β -hydroxylation (CYP11B) of PROG/P4 and 17OHPROG/17OHP4 is the starting point. From P4, 11OHP4 is reversibly metabolized to 11-ketoprogesterone (11KP4, 4-pregnen-3,11,20-trione) by HSD11B, whereafter both 11KP4 and 11OHP4 are metabolized by SRD5A to either 5 α -Pregnanetriolone (11KDHP4, 5 α -pregnan-3,11,20-trione) or 11OH-dihydroprogesterone (11OHDHP4, 5 α -pregnan-11 β -ol-3,20-dione), respectively, with these metabolites also interconverted by HSD11B. Next, 11KDHP4 and 11OHDHP4 are reversibly metabolized to alfaxalone (5 α -pregnan-3 α -ol-11,20-dione) and 3 α ,11 β -dihydroxy-dihydroprogesterone (3 α ,11 β -diOH-DHP4), respectively, by AKR1C2/4, and these metabolites may be converted to 11OHAST and 11KAST by CYP17A1. On the other hand, from 17OHP4, 21-deoxycortisol (21dF, 11 β ,17 α -dihydroxypregn-4-ene-3,20-dione) is reversibly metabolized to 21-deoxycortisone (21dE, 17 α -hydroxypregn-4-ene-3,11,20-trione) by HSD11B. Then, 21dF and 21dE are converted by SRD5A to 5 α -pregnan-11 β ,17 α -diol-3,20-dione (11OHPdione, 5 α -pregnan-11 β ,17 α -diol-3,20-dione) and 5 α -pregnan-17 α -ol-3,11,20-trione (11Kpdione, 5 α -pregnan-17 α -ol-3,11,20-trione), respectively, with these metabolites also interconverted by HSD11B. Thereafter, 11OHPdione and 11Kpdione can be metabolized by AKR1C2/4 to 5 α -pregnan-3 α ,11 β ,17 α -triol-20-one (11OHPdiol) and 5 α -pregnan-3 α ,17 α -diol-11,20-dione (11Kpdil), respectively (with these metabolites also metabolized by HSD11B). Finally, 11OHAST and 11KAST is biosynthesised by CYP17A1 from the latter metabolites, and in the final steps of this pathway, 11KAST is converted to 11K3 α DIOL (by HSD17Bs), followed by the production of 11KDHT (by AKR1C2/4 or RoDH).

919 amino acids [56,57,60,61]. The AR protein is comprised of three functional domains: the N-terminal domain (NTD) which harbours the first activation functional motif (AF1), the DNA binding domain (DBD) which is responsible for the AR interactions and dimerization with specific DNA sequences, and the ligand binding domain (LBD) which contains the second activation functional motif (AF2) which is responsible for most AR activities [56,57,60,61].

Activity of the AR is regulated at different levels. At the genomic level, the AR can comprise inactivating variants itself, or its expression can be epigenetically regulated by methylation or cofactors. Studies in patients with a profile of androgen resistance/insensitivity syndrome (AIS type I) revealed many pathogenic variants in the AR gene and are listed in the AR mutation database [62–65]. In many other patients without an AR gene mutation, low expression of the AR and low activity on DHT stimulated AR-dependent apolipoprotein D (APOD) expression in genital skin fibroblasts was found and consequently classified as AIS type II [18,66]. In some of these patients, further studies showed a direct correlation between the AR mRNA expression and the methylation of CpG regions within the proximal AR promoter [18,67]. In addition, a highly conserved region at the 3'-UTR of the AR gene has been identified through which RNA stability can be regulated by RNA binding proteins in multiple cell lines and tissues [68–71].

At the protein level, the expressed AR can be regulated by posttranslational modifications, which may affect its transcription activity, cellular localization, and stability (Fig. 4A) [72]. Posttranslational modifications that have thus far been identified include phosphorylation, methylation, acetylation, ubiquitination and SUMOylation. These modifications were discovered by performing overexpression experiments in androgen dependent cell lines [72]. Identified phosphorylation modifications were mostly mapped to the NTD, and several studies indicated that differentially phosphorylated residues regulate AR stability, cellular localization, and transcription [72,73]. Moreover, acetylation or methylation of residues in the AR hinge region regulate the ligand-dependent activation and nuclear localization, respectively [72,73]. Ubiquitination of residues in the AR LBD region and SUMOylation in the NTD region also alter AR transcriptional activity and stability and provide another level of regulation of the AR protein [72,73].

Androgen receptor signaling













Androgens are typically secreted from the female and male reproductive systems (ovaries and testes) and the cortex of the adrenal glands to exert their effect on AR in various tissues [57]. After reaching the targeted AR expressing cells through the blood stream (Fig. 4B), androgens bind to the AR. ARs are located in the cytoplasm bound to a complex of different heat shock proteins (HSPs), chaperones and co-chaperones (such as HSP40, HSP70, HSP90 and p23) in a conformation that is competent for ligand-binding [59,74]. Androgen-AR (A-AR) binding leads to signaling activation through a change in AR conformation and dissociation of the chaperones from the AR-complex [59,74]. Once the AR is activated, it can act either at the genomic level or at the non-genomic level.

In the **classical genomic action pathway**, the A-AR complex is transited through the nuclear pore complex into the nucleus. In the nucleus, the A-AR binds as a dimer through its DBD to androgen response elements (ARE) comprised in promoters of target genes [56]. Upon binding of an AR dimer, several co-regulators join the A-AR complex to promote the transcription of target genes [75]. Co-regulators may affect the complex assembly, formation and stability. Moreover, they can affect the chromatin occupancy, looping and remodelling, which can all modulate AR action [75].

Apart from the genomic action, AR signaling in the cell can also occur through a rapid **non-genomic action**. This signaling path of the AR may be activated within seconds or minutes and usually targets plasma membrane proteins or receptors, which activate intracellular kinase cascades that enhance cell proliferation and survival (Fig. 4B) [76,77]. One of the well-studied non-genomic pathways activated by the AR is mitogen-activated protein kinases (MAPK)/extracellular signal-regulated kinases (ERK) signaling. In this signaling pathway, the activated A-AR binds to the tyrosine kinase Src, which then becomes unfolded and auto-phosphorylated resulting in modulation of the MAPK/ERK cascade (Fig. 4B) [76,77]. MAPK/ERK signaling then targets AR-independent transcription, regulating immediate early genes in the cell nucleus through activated phospho-ERK [76]. Interestingly, this modulation seems dose-dependent in prostate cancer (PC) cell lines where low concentrations of androgens




Table 1

Circulatory levels (reference ranges) of bioavailable androgens in females and males (group age: \pm 18–49 [F; follicular phase or entire cycle]/18–62 [M] years) and their potency towards the mammalian AR in comparison to T. (Not available, N/a; these steroids are not commercially available for bioassays; not established, reference ranges for these hormones have not been confirmed *in vivo*). Abbreviations can be found in the list of abbreviations. *Only in tissue [10,35,115–122].

Androgen metabolites	Circulatory levels (ng/dL)		Potency toward the AR
	F	M	
T	1.1–14.3	95–430	
11K3 α DIOL	<i>Not established</i>		N/a
11K5 α DIONE	<i>Not established</i>		
11KA4	17.4–21	8.4–9.5	
11KAST	<1	<1	
11KDHT	<1	<1	
11 KT	5.0–60.6	9.5–70.8	
11OH3 α DIOL	<i>Not established</i>		N/a
11OH5 α DIONE	<i>Not established</i>		Inactive
11OHA4	19.2–233	36.4–313	Inactive
11OHA5T	3.9–4.7	4.4–5.3	<
			
11OHDHT	<i>Not established</i>		
11OHT	8.2–10.8	8.5–9	
A4	28–230	44–186	
Androstenediol-glucuronide	11–249	112–1046	Inactive
Androstenediol*	<i>Not established</i>		
Androstenedione	<i>Not established</i>	5.8–20.2	

(continued on next page)

Table 1 (continued)

Androgen metabolites	Circulatory levels (ng/dL)		Potency toward the AR
	F	M	
Androstenediol	38–45	116–125.5	
Androsterone	31.3–35.2	17.1–18.6	
DHEA	31–701	31–701	Inactive
DHEA-S	17,000–372,000	16,000–523,000	Inactive
DHT	4–22	30–85	

activate the cascade, while high concentrations inhibit it [76,77]. As shown in non-tumoral cells and PC cell lines, activated A-AR may also directly interact with other signaling molecules like phosphoinositide 3-kinases (PI3K), thereby activating the protein kinase B (Akt) kinase pathway [76].

At the systemic level, much of the current knowledge on the role of the AR comes from mouse knockout models [78,79]. The global male AR knockout mouse (ARKO) revealed its important role for male fertility and the development of the testes, while the global female ARKO mouse showed an important role in normal breast and ovary development [78,79]. To study the exact role of the AR for specific cells, tissues and developmental processes, cell-specific ARKO mice were investigated [78,79]. Thus Leydig cell-specific ARKO (LC-AKRO) mice present with infertility, absence of sperm, arrested spermatogenesis and low serum T levels, indicating that the AR is critical for maintaining normal T production and spermatogenesis [78]. In gubernaculum-specific ARKO (GU-ARKO) mice, the crucial role of the AR for regulating testicular descent was illustrated [78]. In ovarian cell-specific ARKO mice, AR-mediated action for normal folliculogenesis and fertility was shown, for example its role to optimize follicular development, maximize ovulation rates, and maintain follicle health. Likewise, its role for cellular proliferation and growth of the uterus has also been shown [78,79].

Implications of lack or excess of androgen action

Males and females need androgens and its action to develop normally and stay healthy, but the dosage of androgens is highly sex specific (Fig. 1). Both androgen deficiency and excess may lead to endocrine disorders that may manifest with a phenotype of disorders/differences of sex development (DSD) at birth, disturb pubertal development, and sexual functioning and fertility. Both lack and excess of androgen action may also manifest at any time of life with adverse effects on other organ systems including overall metabolism, the cardiovascular system, muscles, bones, brain and psychological system. Non-endocrine and non-reproductive effects are described in more details elsewhere [44,80,81].

The effects of androgen deficiency and excess on sexual development and function are best studied in patients carrying rare disease-causing monogenetic variants in genes involved in androgen biosynthesis and action. By contrast, acquired disorders of androgen deficiency and excess will not cause a DSD, but may affect pubertal development, fertility and overall health, nevertheless. Thus, androgen excess in adults is a common problem in women suffering from PCOS and in men with prostate hyperplasia/tumors [82–88]. Furthermore, androgen decline/deficiency is a controversial topic in healthy aging males and females [89,90].

To understand the clinical manifestations of genetic disorders of gonadal and adrenal steroidogenesis at birth it is important to keep in mind that adrenal and gonadal androgens are part of a concerted steroid network *in utero*. During fetal development, the liver and the placenta play major




Fig. 4. Androgen action through the androgen receptor (AR). **A.** The human AR gene is located on the X chromosome at position Xq11–12. The AR consists of eight exons which are separated by long introns and the AR gene can be epigenetically regulated by methylation. Notably, the AR protein is comprised of three functional domains, the N-terminal domain (NTD), the DNA binding domain (DBD) and the ligand binding domain (LBD) and can be regulated by posttranslational modifications. **B. The macroscopic view:** androgens are secreted from the adrenal gland and the gonads. The secreted androgens exert their effect in AR-expressing tissues, including the kidney, liver, adrenal gland and gonads. **The microscopic view:** once the androgens (A) are released into the blood stream, they enter the AR-expressing cell and activate the AR. The A-AR complex can have a genomic action where it binds androgen receptor elements (ARE) in the target gene and initiates gene expression. While in the non-genomic action, the A-AR can bind membrane-bound receptors and initiate signaling cascades, which increases the cell proliferation and survival.

roles in this network [91]. The fetal liver functions to metabolise and inactivate androgens, and while the placenta is not considered an androgen producing organ, it does express all the necessary steroidogenic enzymes to convert maternal PROG to androgens, including those that can biosynthesise T [91–96]. Additionally, a role of the placenta in the fetal backdoor pathway in the production of androgens has also been suggested [91]. Moreover, predominant androgen levels have recently been measured in placental tissue [97], highlighting the production of placental androgens and that androgens are exchanged in the fetoplacental unit. Notably, placental aromatase would convert A4 and T

to estrogens, while other androgens might not be biotransformed to estrogen derivatives *in vivo* [98,99]. As the development of the typical female and male external genitalia relies largely on the absence or presence of T and DHT, any disturbance in androgen production may result in apparent virilization of a 46,XX fetus or undermasculinization of a 46,XY fetus. After birth and minipuberty, gonadal steroidogenesis is quiescent until puberty (Fig. 1), when activation of the HPG axis commands to resume sex steroid production for normal sexual maturation, fertility and reproduction (Fig. 2). Also, in postnatal/adult life, steroids secreted by the adrenals and gonads are converted to active and inactive metabolites by peripheral organs, and this complex peripheral steroid metabolism may then be responsible for the formation of unusual steroids through alternate pathways in genetic disorders of steroidogenesis (Fig. 3).

Specific monogenetic defects of androgen biosynthesis

Pathogenic variants in all genes involved in human androgen biosynthesis and metabolism may cause androgen deficiency and/or excess. Table 2 gives an overview of all genes, for which variants have been reported with a human phenotype associated with abnormal androgen production. For details we refer the reader to [11,57,100]. In the following section we provide a summary of possible genetic defects underlying specific phenotypes.

Genetic defects affecting early steps of steroid biosynthesis and cortisol production in particular are known as **CAH** [11,100]. These disorders may be grouped in defects causing 46,XY DSD CAH, 46,XX DSD CAH or DSD CAH in both chromosomal sexes according to the current DSD classification [101].

To the **first group of 46,XY DSD CAH** belong mutations in the genes for *StAR*, *CYP11A1*, *CYP17A1* and lack of androgens. Patients with autosomal recessive variants in these genes with loss of function are not able to synthesize cortisol and androgens. Thus, affected chromosomal male fetuses are born with typical female external genitalia and raised as girls. However, genetic mutations in these genes with partial loss of function may not affect prenatal sex development and only manifest with primary adrenal insufficiency in early life, while sex hormone failure may develop later. Affected 46,XX individuals with severe mutations in the *StAR*, *CYP11A1*, *CYP17A1* genes manifest at birth with adrenal insufficiency only and show typically lack of pubertal development later.

The **second group of 46,XX DSD CAH** with androgen excess comprises autosomal recessive defects of 21-hydroxylase (*CYP21A2*) and *CYP11B1* necessary for GC and MC synthesis. With these defects that affect GC and MC synthesis in the adrenal cortex, the lack of cortisol leads to negative feedback stimulation of the HPA axis, elevated ACTH and thus increased adrenal androgen production. Affected girls show variable degrees of external genital virilization at birth as the androgen excess already occurs very early *in utero*, when the female fetus needs to be safeguarded from high androgens to develop normally [102]. Affected 46,XY boys have no DSD phenotype but still suffer from neonatal onset adrenal insufficiency with severe variants.

Genetic mutations in *HSD3B2* and *POR* belong to the **third group** that can manifest with cortisol deficiency and variable severity of a **DSD at birth in both chromosomal sexes** depending on the specific variants.

Genetic defects that only affect sex hormone biosynthesis also display some sex specific characteristics. 46,XY DSD individuals with an isolated moderate to severe undervirilization at birth may harbour a defect in the *CYP17A1* (isolated 17,20-lyase deficiency), *CYP5A1*, *POR*, *AKR1C2/4*, *HSD17B3* or *SRD5A2* genes (Fig. 3). Regarding these genetic variants, only 46,XX females with variants in *CYP17A1* and *POR* have been found with a DSD phenotype (*POR*), with primary or secondary failure of pubertal development (*POR*, *CYP17A1*), or with a PCOS-like phenotype (*POR*). Moreover, aromatase (*CYP19A1*) mutations inhibiting the aromatase activity and the conversion of androgens to estrogens, cause fetal and maternal androgen excess during pregnancy and beyond. Affected girls are virilized at birth and show abnormal pubertal development and fertility. Affected boys are asymptomatic during childhood years, but then come to medical attention for tall stature and failure to stop growing. Bone age is typically delayed because estrogens are crucial for epiphyseal maturation and closure, and decreased bone mineral density can be observed later in life. A negative impact on glucose homeostasis and lipid profile has also been described in both adult males and females with aromatase deficiency.

Finally, genetic variants in the steroidogenic factor 1/nuclear receptor subfamily 5 group A member 1 (*SF1/NR5A1*) may cause a broad phenotypical range of 46,XY DSD, 46,XX DSD and failure of pubertal development and fertility [103,104]. In rare cases, *NR5A1* mutations also cause adrenal insufficiency. *SF1* is an essential transcription regulator for many genes involved in sex determination, sex differentiation as well as (sex) steroid biosynthesis [103–107].

Genetic defects of androgen action – AIS

Complete or partial androgen insensitivity syndromes (CAIS/PAIS) are caused by genetic mutations affecting AR function [18,108]. In these syndromes of 46,XY phenotypic undermasculinization, androgen concentrations are typically elevated [62]. AIS incidence is reported internationally in 1 in 20,400 live born 46,XY infants, with CAIS occurring at a higher rate than PAIS [109]. This high incidence might be explained by low mortality and morbidity of AIS. So far about five hundred *AR* mutations have been described in individuals with AIS, classified as **AIS type I**. In CAIS, severe hemizygous *AR* mutations cause the loss of AR signaling in more than 95% of cases. Affected 46,XY individuals show male typical inner genital organs and undescended gonads with the prostate, vas deferens and seminal vesicles missing. External genitalia are typical female with a vaginal pouch. Female carriers are phenotypically normal. In 46,XY PAIS, the phenotypic variability in undermasculinization is large and depends on the residual activity of the AR. In the mildest form of PAIS, gynecomastia and male infertility may be the only clinical signs. Unlike in CAIS, the underlying genetic defect in PAIS is only found in the *AR* gene in about 40% of cases [64].

By contrast, individuals with AIS without *AR* mutations are classified as **AIS type II**. In these individuals largely unidentified regulators or cofactors of the AR are responsible for the impaired AR signaling as revealed by an AR-dependent bioassay using genital skin fibroblasts and the targeted *APOD* as a biomarker [66]. As AR activity can be regulated at various levels, the possible mechanisms of AIS type II are manifold. Thus far, altered DNA methylation of the *AR* promoter has been found in some individuals with PAIS [67].

Common acquired disorders of androgen excess

Androgen levels and AR activity control development, proliferation and growth of a variety of cells in many organs and regulate metabolic processes in both males and females (Table 1) [57].

Unbalanced levels of androgens are involved in the development of several types of cancers including PC and CRPC [57,82–84]. In men, PC is the second leading cause of cancer death worldwide. PC growth and development critically dependent on androgens and AR signaling [60,82,83]. Androgens promote prostate cell proliferation resulting in continuous growth of malignant cells. But it is important to note that there is no evidence of an association between the lifetime T exposure and the appearance of PC [110]. Therefore, local prostate tumors are cared for by watchful waiting or treated with surgery or radiation therapy, while high-grade and metastatic PCs are treated with androgen-deprivation therapies, lowering T levels. However, over time many PCs escape this treatment and progress to CRPC [83]. CRPC might be driven by potent androgens synthesized from alternate, non-gonadal steroid metabolites (for example metabolites originating from the adrenals) [47,49,50] that stimulate AR signaling [83]. However, the specific mechanisms associated are still unclear [82–84].

In women, PCOS is the most common endocrine condition affecting about 10% of reproductive-age women [85]. PCOS is characterized by hyperandrogenism, menstrual disturbances and polycystic ovaries [86,111]. Studies performed in animals and humans support the hypothesis that excess androgen and AR action are key players of PCOS pathomechanism as they are clearly associated with adverse reproductive and metabolic outcome [86], but the exact interplay remains a conundrum. Numerous studies have shown that androgens of the classic pathway (such as T and A4; Fig. 3A) are elevated in circulation of most women with PCOS when compared to healthy controls [85,87]. In line with these findings are *in vitro* studies showing ovarian theca cells isolated from PCOS women with steroidogenic hyperfunction resulting in androgen excess [88].

Interestingly, comprehensive steroid profiling in more recent studies showed that C11-oxy androgens of alternate pathways (such as 11OHA4, 11KA4, 11OHT and 11KT; Fig. 3C and D) are even more

Table 2

Characteristics of monogenetic disorders of steroidogenesis affecting androgen production and the gonads in 46,XX and 46,XY individuals [11,57,100].

Disorder	Gene/ OMIM	46,XY Gonadal Phenotype (T Deficiency)	46,XX Gonadal Phenotype (E2 Deficiency)	Fertility	Adrenal Insufficiency	Other Features
Lipoid congenital adrenal hyperplasia (LCAH)	<i>StAR</i> 201710	<i>Classic form:</i> 46,XY DSD, gonadal insufficiency <i>Non-classic form:</i> normal or NK	<i>Classic:</i> primary or secondary ovarian insufficiency (POI) <i>Non-classic:</i> NK or normal	<i>Classic:</i> Absent in 46,XY; variable in 46,XX	YES	
P450 side chain cleavage syndrome (CAH)	<i>CYP11A1</i> 118485	<i>Classic form:</i> 46,XY DSD, gonadal insufficiency <i>Non-classic form:</i> normal	<i>Classic:</i> primary or secondary ovarian insufficiency (POI) <i>Non-classic:</i> NK or normal	Reported in 46,XX	YES	
3β-Hydroxysteroid dehydrogenase II deficiency (CAH)	<i>HSD3B2</i> 201810	46,XY DSD, gonadal insufficiency <i>Non-classic form:</i> normal, but premature adrenarche	46,XX DSD with atypical genital development; gonadal insufficiency <i>Non-classic form:</i> normal, but premature adrenarche	Absent in 46,XY; reported in 46,XX	YES	
21-Hydroxylase deficiency (CAH)	<i>CYP21A2</i> 201910	<i>Classic form:</i> normal <i>Non-classic form:</i> normal	46,XX DSD with atypical genital development through androgen excess; <i>Non-classic form:</i> premature adrenarche, virilization, PCO	Normal in both 46,XX and 46,XY, if treated	YES	Cave: Testicular adrenal rest tumor (m >> f) CAH-X (when combined with Ehlers-Danlos syndrome with contiguous gene variants) Hypertension
11β-hydroxylase deficiency (CAH)	<i>CYP11B1</i> 202010	<i>Classic form:</i> normal <i>Non-classic form:</i> normal	46,XX DSD with atypical genital development through androgen excess; <i>Non-classic form:</i> premature adrenarche, virilization, PCO	Normal in both 46,XX and 46,XY, if treated	YES	
Combined 17α-hydroxylase, 17,20 lyase deficiency (CAH)	<i>CYP17A1</i> 202110	46,XY DSD, gonadal insufficiency	Lack of pubertal development, POI	Possible in 46,XX with assisted fertility measures	Rare	Hypertension and hypokalemic alkalosis (not seen with isolated lyase deficiency)
P450 oxidoreductase deficiency (CAH)	<i>POR</i> 124015 201750	Mild to severe 46,XY DSD, gonadal insufficiency	46,XX DSD with atypical genital development or premature adrenarche (androgen excess), virilisation, POI, PCO	Reported	Variable	Maternal virilization during pregnancy; Antley-Bixler skeletal malformation syndrome; changes in drug metabolism
Cytochrome <i>b</i>₅ deficiency	<i>CYB5A</i> 613218	46,XY DSD	NK	NK	NO	Methemoglobinemia

Table 2 (continued)

Disorder	Gene/ OMIM	46,XY Gonadal Phenotype (T Deficiency)	46,XX Gonadal Phenotype (E2 Deficiency)	Fertility	Adrenal Insufficiency	Other Features
17β-Hydroxysteroid dehydrogenase III deficiency/17- ketosteroid reductase deficiency	<i>HSD17B3</i> 264300	46,XY DSD; progressive virilisation and gynecomastia at puberty	Normal	Decreased or absent in 46,XY	NO	
5α-Reductase II deficiency	<i>SRD5A2</i> 607306	46,XY DSD; progressive virilisation and gynecomastia at puberty	Normal	Impaired in 46,XY	NO	
3α-Hydroxysteroid dehydrogenase deficiency	<i>AKR1C2/4</i> 600450 600451	46,XY DSD; gonadal insufficiency	Normal	NK	NO	
Aromatase deficiency	<i>CYP19A1</i> 107910	Normal	46,XX DSD with variable degree of virilisation at birth, gonadal insufficiency, POI	Impaired in 46,XX	NO	Overgrowth and metabolic anomalies in males
Steroidogenic factor 1	<i>NR5A1/SF1</i> 184757	Mild to severe 46,XY DSD; gonadal insufficiency – very variable	Mostly POI or normal 46,XX ovotesticular DSD	Mostly impaired in 46,XY; variable in 46,XX	Rare	

Abbr.: DSD – Disorder/Difference of Sexual Development; E2 – estradiol; NK – not known; PCO – polycystic ovaries; POI – primary ovarian insufficiency; T – testosterone.

elevated than classic androgens in PCOS patients [24,112]. This indicates that androgen excess in PCOS originates from both the ovaries and non-gonadal tissues including the adrenals and peripheral tissues as well as the intermediary metabolism. This also suggests that the underlying defect of PCOS leads to a dysregulation of androgen production through different steroid producing pathways. In addition, hyperandrogenism in PCOS is very often associated with insulin resistance [43,86,111], and both may lead to metabolic dysfunction in the reproductive tissues and beyond [88]. However, although the metabolic consequences of PCOS have been well described, the underlying cause and detailed molecular mechanisms are still unsolved.

Paradoxically, there are no clear pathophysiological conditions in men where T is too high, and no such conditions in women where it is too low [113]. However, there are some discussions about a male PCOS phenotype with androgen excess and a female lack of libido with deficiency.

Acquired disorders of androgen deficiency

In principle, the loss of circulating androgens in adults is caused by gonadal failure. This might be due to any disorders affecting the HPG axis (Fig. 2). In men, hypogonadism causes impaired sexual functioning, infertility, alterations in body composition and osteoporosis [89]. Moreover, it has adverse effects on metabolic and cardiovascular health as well as a negative impact on brain health [90]. When in women low androgen levels are due to hypogonadism, estrogen levels are also decreased, and the consequences of low estrogens are the same as with hypogonadism in men [89,113]. By contrast, no adverse consequences are seen when androgen levels are decreased because of a loss of adrenal androgens with primary or secondary adrenal disorders [6]; although this has been challenged in some studies for women without convincing evidence [114].

Summary

Clinical, genetic and functional characterization of human mutations in genes involved in androgen biosynthesis and action have taught us plenty about the crucial role of androgens for sexual development, reproduction and beyond. All the same, many patients manifesting with rare congenital or common acquired disorders of androgen deficiency or excess remain unexplained indicating that androgen biology is still not fully understood. Only in the last two decades, two novel alternate pathways of androgen biosynthesis have been recognized to play important roles in normal biology and disease states. But the exact interplay and specific contribution of the different biosynthesis pathways to the active androgen pool need further investigation. Likewise, the regulation of the AR is more complex than initially assumed. In theory, multiple regulators and co-regulators may modulate AR signaling at the transcriptional and posttranslational level. Studies of patients with PAIS found differential AR promoter methylation in some, but other regulatory factors have yet to be identified.

Understanding the complex network regulating androgen biosynthesis, metabolism and action is essential for finding new targets for better diagnostic and therapeutic opportunities.

Funding

Work related to this review is supported by the Swiss National Science Foundation (320020_197725), Switzerland. TDT was supported by a Marie Skłodowska-Curie Individual Fellowship (# 101023999), European Commission.

Declaration of competing interest

The authors declare no competing interest.

Practice points

- Androgens are essential sex steroid hormones for both sexes.
- Testosterone is the predominant circulating androgen in males, while in females the major circulating classic androgens are DHEA-S, DHEA and androstenedione.
- Androgens exert their effect through binding to the androgen receptor to modulate many different biological processes.
- Androgen receptor signaling occurs at the genomic and non-genomic level.
- Rare human genetic disorders have been described for all genes involved in androgen biosynthesis and for the androgen receptor.
- Common disorders associated with androgen imbalance are castration resistant prostate cancer in men and polycystic ovary syndrome in women.

Research agenda

- To better understand the role of newly discovered, alternate androgen biosynthesis pathways in health and disease.
- To study the impact of alternate androgens on the androgen receptor in various tissues.
- To identify regulators and co-regulators of the androgen receptor affecting its expression, stability and activity.
- To understand the role of androgens' action in common disorders like prostate hyperplasia and cancers, and in polycystic ovary syndrome in order to identify targets for better treatments.
- To investigate the treatment potential of candidate drugs for disorders and diseases marked by androgen imbalances, such as selective androgen receptor modulators (SARMs).

References

- [1] Kelsey TW, Li LQ, Mitchell RT, et al. A validated age-related normative model for male total testosterone shows increasing variance but no decline after age 40 years. *PLoS One* 2014;9(10):e109346.
- [2] Fanelli F, Baronio F, Ortolano R, et al. Normative basal values of hormones and proteins of gonadal and adrenal functions from birth to adulthood. *Sex Dev* 2018;12(1–3):50–94.
- [3] Kulle AE, Riepe FG, Melchior D, et al. A novel ultrahigh-pressure liquid chromatography tandem mass spectrometry method for the simultaneous determination of androstenedione, testosterone, and dihydrotestosterone in pediatric blood samples: age- and sex-specific reference data. *J Clin Endocrinol Metab* 2010;95(5):2399–409.
- [4] Harman SM, Metter EJ, Tobin JD, et al. Baltimore Longitudinal Study of A. Longitudinal effects of aging on serum total and free testosterone levels in healthy men. Baltimore Longitudinal Study of Aging. *J Clin Endocrinol Metab* 2001;86(2):724–31.
- [5] Feldman HA, Longcope C, Derby CA, et al. Age trends in the level of serum testosterone and other hormones in middle-aged men: longitudinal results from the Massachusetts male aging study. *J Clin Endocrinol Metab* 2002;87(2):589–98.
- [6] Bhasin S, Brito JP, Cunningham GR, et al. Testosterone therapy in men with hypogonadism: an endocrine society clinical practice guideline. *J Clin Endocrinol Metab* 2018;103(5):1715–44.
- [7] Fluck CE, Pandey AV. Steroidogenesis of the testis – new genes and pathways. *Ann Endocrinol* 2014;75(2):40–7.
- [8] Wilson JD. The role of 5 α -reduction in steroid hormone physiology. *Reprod Fertil Dev* 2001;13(7–8):673–8.
- [9] Skiba MA, Bell RJ, Islam RM, et al. Androgens during the reproductive years: what is normal for women? *J Clin Endocrinol Metab* 2019;104(11):5382–92.
- [10] Rege J, Nakamura Y, Satoh F, et al. Liquid chromatography–tandem mass spectrometry analysis of human adrenal vein 19-carbon steroids before and after ACTH stimulation. *J Clin Endocrinol Metab* 2013;98(3):1182–8.
- [11] Miller WL, Flück CE, Breault DT, et al. 14 - the adrenal cortex and its disorders. In: Sperling MA, editor. *Sperling pediatric Endocrinology*. 5th ed. Philadelphia: Elsevier; 2021. p. 425–90.
- [12] Haring R, Hannemann A, John U, et al. Age-specific reference ranges for serum testosterone and androstenedione concentrations in women measured by liquid chromatography–tandem mass spectrometry. *J Clin Endocrinol Metab* 2012;97(2):408–15.
- *[13] Turcu AF, Rege J, Auchus RJ, et al. 11-Oxygenated androgens in health and disease. *Nat Rev Endocrinol* 2020;16(5):284–96.
- [14] Pignatti E, Fluck CE. Adrenal cortex development and related disorders leading to adrenal insufficiency. *Mol Cell Endocrinol* 2021;527:111206.
- [15] Becker M, Hesse V. Minipuberty: why does it happen? *Horm Res Paediatr* 2020;93(2):76–84.
- *[16] Dhayat NA, Dick B, Frey BM, et al. Androgen biosynthesis during minipuberty favors the backdoor pathway over the classic pathway: insights into enzyme activities and steroid fluxes in healthy infants during the first year of life from the urinary steroid metabolome. *J Steroid Biochem Mol Biol* 2017;165(Pt B):312–22.
- [17] von Schnakenburg K, Bidlingmaier F, Knorr D. 17-hydroxyprogesterone, androstenedione, and testosterone in normal children and in prepubertal patients with congenital adrenal hyperplasia. *Eur J Pediatr* 1980;133(3):259–67.
- *[18] Hornig NC, Holterhus PM. Molecular basis of androgen insensitivity syndromes. *Mol Cell Endocrinol* 2021;523:111146.
- [19] Miller WL, Auchus RJ. The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr Rev* 2011;32(1):81–151.
- [20] Costa B, Da Pozzo E, Martini C. Translocator protein and steroidogenesis. *Biochem J* 2018;475(5):901–4.
- [21] Russell DW, Wilson JD. Steroid 5 α -reductase: two genes/two enzymes. *Annu Rev Biochem* 1994;63:25–61.
- [22] Fluck CE, Miller WL, Auchus RJ. The 17, 20-lyase activity of cytochrome P450c17 from human fetal testis favors the delta5 steroidogenic pathway. *J Clin Endocrinol Metab* 2003;88(8):3762–6.
- [23] Barnard L, du Toit T, Swart AC. Back where it belongs: 11beta-hydroxyandrostenedione compels the re-assessment of C11-oxy androgens in steroidogenesis. *Mol Cell Endocrinol* 2021;525:111189.
- [24] Swart AC, du Toit T, Gourgari E, et al. Steroid hormone analysis of adolescents and young women with polycystic ovarian syndrome and adrenocortical dysfunction using UPC(2)-MS/MS. *Pediatr Res* 2021;89(1):118–26.
- [25] Barnard M, Mostaghel EA, Auchus RJ, et al. The role of adrenal derived androgens in castration resistant prostate cancer. *J Steroid Biochem Mol Biol* 2020;197:105506.
- [26] Pretorius E, Arlt W, Storbek KH. A new dawn for androgens: novel lessons from 11-oxygenated C19 steroids. *Mol Cell Endocrinol* 2017;441:76–85.
- *[27] Swart AC, Storbek KH. 11beta-Hydroxyandrostenedione: downstream metabolism by 11betaHSD, 17betaHSD and SRD5A produces novel substrates in familial pathways. *Mol Cell Endocrinol* 2015;408:114–23.
- [28] Wilson JD, Auchus RJ, Leihy MW, et al. 5 α -androstane-3 α ,17 β -diol is formed in tammar wallaby pouch young testes by a pathway involving 5 α -pregnane-3 α ,17 α -diol-20-one as a key intermediate. *Endocrinology* 2003;144(2):575–80.
- *[29] Auchus RJ. The backdoor pathway to dihydrotestosterone. *Trends Endocrinol Metabol* 2004;15(9):432–8.
- [30] Mahendroo M, Wilson JD, Richardson JA, et al. Steroid 5 α -reductase 1 promotes 5 α -androstane-3 α ,17 β -diol synthesis in immature mouse testes by two pathways. *Mol Cell Endocrinol* 2004;222(1–2):113–20.
- [31] Arlt W, Walker EA, Draper N, et al. Congenital adrenal hyperplasia caused by mutant P450 oxidoreductase and human androgen synthesis: analytical study. *Lancet* 2004;363(9427):2128–35.
- [32] Homma K, Hasegawa T, Nagai T, et al. Urine steroid hormone profile analysis in cytochrome P450 oxidoreductase deficiency: implication for the backdoor pathway to dihydrotestosterone. *J Clin Endocrinol Metab* 2006;91(7):2643–9.
- [33] Kamrath C, Hochberg Z, Hartmann MF, et al. Increased activation of the alternative "backdoor" pathway in patients with 21-hydroxylase deficiency: evidence from urinary steroid hormone analysis. *J Clin Endocrinol Metab* 2012;97(3):E367–75.
- *[34] Fluck CE, Meyer-Boni M, Pandey AV, et al. Why boys will be boys: two pathways of fetal testicular androgen biosynthesis are needed for male sexual differentiation. *Am J Hum Genet* 2011;89(2):201–18.

- [35] Handelsman DJ, Cooper ER, Heather AK. Bioactivity of 11 keto and hydroxy androgens in yeast and mammalian host cells. *J Steroid Biochem Mol Biol* 2022;218:106049.
- [36] Storbeck KH, Bloem LM, Africander D, et al. 11beta-Hydroxydihydrotestosterone and 11-ketodihydrotestosterone, novel C19 steroids with androgenic activity: a putative role in castration resistant prostate cancer? *Mol Cell Endocrinol* 2013;377(1–2):135–46.
- [37] Schloms L, Storbeck KH, Swart P, et al. The influence of *Aspalathus linearis* (Rooibos) and dihydrochalcones on adrenal steroidogenesis: quantification of steroid intermediates and end products in H295R cells. *J Steroid Biochem Mol Biol* 2012;128(3–5):128–38.
- [38] Swart AC, Schloms L, Storbeck KH, et al. 11beta-hydroxyandrostenedione, the product of androstenedione metabolism in the adrenal, is metabolized in LNCaP cells by 5alpha-reductase yielding 11beta-hydroxy-5alpha-androstenedione. *J Steroid Biochem Mol Biol* 2013;138:132–42.
- [39] Turcu AF, Nanba AT, Chomic R, et al. Adrenal-derived 11-oxygenated 19-carbon steroids are the dominant androgens in classic 21-hydroxylase deficiency. *Eur J Endocrinol* 2016;174(5):601–9.
- [40] Bacila I, Adaway J, Hawley J, et al. Measurement of salivary adrenal-specific androgens as biomarkers of therapy control in 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 2019;104(12):6417–29.
- [41] Jha S, Turcu AF, Sinaii N, et al. 11-Oxygenated androgens useful in the setting of discrepant conventional biomarkers in 21-hydroxylase deficiency. *J Endocr Soc* 2021;5(2):bvaa192.
- [42] Schröder MAM, Turcu AF, O'Day P, et al. Production of 11-oxygenated androgens by testicular adrenal rest tumors. *J Clin Endocrinol Metab* 2021;107(1):e272–80.
- [43] Kempegowda P, Melson E, Manolopoulos KN, et al. Implicating androgen excess in propagating metabolic disease in polycystic ovary syndrome. *Ther Adv Endocrinol Metab* 2020;11. 2042018820934319.
- [44] Yoshida T, Saito K, Kawamura T, et al. Circulating steroids and mood disorders in patients with polycystic ovary syndrome. *Steroids* 2021;165:108748.
- [45] Tosi F, Villani M, Garofalo S, et al. Clinical value of serum levels of 11-oxygenated metabolites of testosterone in women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2022;107(5):e2047–55. <https://doi.org/10.1210/clinem/dgab920>.
- [46] Rege J, Turcu AF, Kasa-Vubu JZ, et al. 11-Ketotestosterone is the dominant circulating bioactive androgen during normal and premature adrenarche. *J Clin Endocrinol Metab* 2018;103(12):4589–98.
- [47] du Toit T, Bloem LM, Quanson JL, et al. Profiling adrenal 11beta-hydroxyandrostenedione metabolites in prostate cancer cells, tissue and plasma: UPC(2)-MS/MS quantification of 11beta-hydroxytestosterone, 11keto-testosterone and 11keto-dihydrotestosterone. *J Steroid Biochem Mol Biol* 2017;166:54–67.
- [48] du Toit T, Swart AC. Inefficient UGT-conjugation of adrenal 11beta-hydroxyandrostenedione metabolites highlights C11-oxy C19 steroids as the predominant androgens in prostate cancer. *Mol Cell Endocrinol* 2018;461:265–76.
- [49] Snatser G, van Dessel LF, van Riet J, et al. 11-Ketotestosterone is the predominant active androgen in prostate cancer patients after castration. *JCI Insight* 2021;6(11).
- [50] Wright C, O'Day P, Alyamani M, et al. Abiraterone acetate treatment lowers 11-oxygenated androgens. *Eur J Endocrinol* 2020;182(4):413–21.
- [51] Houghton LC, Howland RE, Wei Y, et al. The steroid metabolome and breast cancer risk in women with a family history of breast cancer: the novel role of adrenal androgens and glucocorticoids. *Cancer Epidemiol Biomarkers Prev* 2021;30(1):89–96.
- [52] du Toit T, Swart AC. The 11beta-hydroxyandrostenedione pathway and C11-oxy C21 backdoor pathway are active in benign prostatic hyperplasia yielding 11keto-testosterone and 11keto-progesterone. *J Steroid Biochem Mol Biol* 2020;196:105497.
- [53] Barnard L, Gent R, van Rooyen D, et al. Adrenal C11-oxy C21 steroids contribute to the C11-oxy C19 steroid pool via the backdoor pathway in the biosynthesis and metabolism of 21-deoxycortisol and 21-deoxycortisone. *J Steroid Biochem Mol Biol* 2017;174:86–95.
- [54] van Rooyen D, Gent R, Barnard L, et al. The in vitro metabolism of 11beta-hydroxyprogesterone and 11-ketoprogesterone to 11-ketodihydrotestosterone in the backdoor pathway. *J Steroid Biochem Mol Biol* 2018;178:203–12.
- [55] van Rooyen D, Yadav R, Scott EE, et al. CYP17A1 exhibits 17alpha-hydroxylase/17,20-lyase activity towards 11beta-hydroxyprogesterone and 11-ketoprogesterone metabolites in the C11-oxy backdoor pathway. *J Steroid Biochem Mol Biol* 2020;199:105614.
- [56] Sutinen P, Malinen M, Palmvo JJ. Androgen receptor. In: Simoni M, Huhtaniemi I, editors. *Endocrinology of the testis and male reproduction*. Cham: Springer International Publishing; 2016. p. 1–22.
- [57] Flück CE, Pandey AV. Testicular steroidogenesis. In: Simoni M, Huhtaniemi IT, editors. *Endocrinology of the testis and male reproduction*. Cham: Springer International Publishing; 2017. p. 343–71.
- [58] Chang C, Yeh S, Lee SO, et al. Androgen receptor (AR) pathophysiological roles in androgen-related diseases in skin, bone/muscle, metabolic syndrome and neuron/immune systems: lessons learned from mice lacking AR in specific cells. *Nucl Recept Signal* 2013;11:e001.
- *[59] Chaturvedi AP, Dehm SM. Androgen receptor dependence. *Adv Exp Med Biol* 2019;1210:333–50.
- [60] Tan MH, Li J, Xu HE, et al. Androgen receptor: structure, role in prostate cancer and drug discovery. *Acta Pharmacol Sin* 2015;36(1):3–23.
- *[61] Hunter I, Hay CW, Esswein B, et al. Tissue control of androgen action: the ups and downs of androgen receptor expression. *Mol Cell Endocrinol* 2018;465:27–35.
- [62] Mongan NP, Tadokoro-Cuccaro R, Bunch T, et al. Androgen insensitivity syndrome. *Best Pract Res Clin Endocrinol Metab* 2015;29(4):569–80.
- [63] Ahmed SF, Cheng A, Dovey L, et al. Phenotypic features, androgen receptor binding, and mutational analysis in 278 clinical cases reported as androgen insensitivity syndrome. *J Clin Endocrinol Metab* 2000;85(2):658–65.
- [64] Ahmed SF, Bashamboo A, Lucas-Herald A, et al. Understanding the genetic aetiology in patients with XY DSD. *Br Med Bull* 2013;106:67–89.

- [65] Audi L, Fernandez-Cancio M, Carrascosa A, et al. Novel (60%) and recurrent (40%) androgen receptor gene mutations in a series of 59 patients with a 46,XY disorder of sex development. *J Clin Endocrinol Metab* 2010;95(4):1876–88.
- [66] Hornig NC, Ukat M, Schweikert HU, et al. Identification of an AR mutation-negative class of androgen insensitivity by determining endogenous AR activity. *J Clin Endocrinol Metab* 2016;101(11):4468–77.
- *[67] Hornig NC, Rodens P, Dorr H, et al. Epigenetic repression of androgen receptor transcription in mutation-negative androgen insensitivity syndrome (AIS type II). *J Clin Endocrinol Metab* 2018;103(12):4617–27.
- [68] Zhou H, Mazan-Mamczarz K, Martindale JL, et al. Post-transcriptional regulation of androgen receptor mRNA by an ErbB3 binding protein 1 in prostate cancer. *Nucleic Acids Res* 2010;38(11):3619–31.
- [69] Barker A, Epis MR, Porter CJ, et al. Sequence requirements for RNA binding by HuR and AUF1. *J Biochem* 2012;151(4):423–37.
- [70] Zhao J, Zhang Y, Liu XS, et al. RNA-binding protein Musashi2 stabilizing androgen receptor drives prostate cancer progression. *Cancer Sci* 2020;111(2):369–82.
- [71] Yeap BB, Wilce JA, Leedman PJ. The androgen receptor mRNA. *Bioessays* 2004;26(6):672–82.
- [72] van der Steen T, Tindall DJ, Huang H. Posttranslational modification of the androgen receptor in prostate cancer. *Int J Mol Sci* 2013;14(7):14833–59.
- [73] Coffey K, Robson CN. Regulation of the androgen receptor by post-translational modifications. *J Endocrinol* 2012;215(2):221–37.
- [74] Anestis A, Zoi I, Papavassiliou AG, et al. Androgen receptor in breast cancer-clinical and preclinical Research insights. *Molecules* 2020;25(2).
- [75] Heemers HV, Tindall DJ. Androgen receptor (AR) coregulators: a diversity of functions converging on and regulating the AR transcriptional complex. *Endocr Rev* 2007;28(7):778–808.
- [76] Leung JK, Sadar MD. Non-genomic actions of the androgen receptor in prostate cancer. *Front Endocrinol* 2017;8:2.
- [77] Liao RS, Ma S, Miao L, et al. Androgen receptor-mediated non-genomic regulation of prostate cancer cell proliferation. *Transl Androl Urol* 2013;2(3):187–96.
- *[78] Chang C, Lee SO, Wang RS, et al. Androgen receptor (AR) physiological roles in male and female reproductive systems: lessons learned from AR-knockout mice lacking AR in selective cells. *Biol Reprod* 2013;89(1):21.
- [79] Walters KA, Simanainen U, Gibson DA. Androgen action in female reproductive physiology. *Curr Opin Endocrinol Diabetes Obes* 2016;23(3):291–6.
- [80] Bulant J, Hill M, Velikova M, et al. Changes of BMI, steroid metabolome and psychopathology in patients with anorexia nervosa during hospitalization. *Steroids* 2020;153:108523.
- [81] Kelava I, Chiaradia I, Pellegrini L, et al. Androgens increase excitatory neurogenic potential in human brain organoids. *Nature* 2022;602(7895):112–6.
- [82] Zhou Y, Bolton EC, Jones JO. Androgens and androgen receptor signaling in prostate tumorigenesis. *J Mol Endocrinol* 2015;54(1):R15–29.
- [83] Auchus RJ, Sharifi N. Sex hormones and prostate cancer. *Annu Rev Med* 2020;71:33–45.
- [84] Storbek KH, Mostaghel EA. Canonical and noncanonical androgen metabolism and activity. *Adv Exp Med Biol* 2019;1210:239–77.
- [85] Rodríguez Paris V, Bertoldo MJ. The mechanism of androgen actions in PCOS etiology. *Med Sci* 2019;7(9).
- [86] Lizneva D, Gavrilova-Jordan L, Walker W, et al. Androgen excess: investigations and management. *Best Pract Res Clin Obstet Gynaecol* 2016;37:98–118.
- [87] Abdelazim IA, Alanwar A, AbuFaza M, et al. Elevated and diagnostic androgens of polycystic ovary syndrome. *Prz Menopauzalny* 2020;19(1):1–5.
- [88] Sanchez-Garrido MA, Tena-Sempere M. Metabolic dysfunction in polycystic ovary syndrome: pathogenic role of androgen excess and potential therapeutic strategies. *Mol Metabol* 2020;35:100937.
- [89] Richard-Eaglin A. Male and female hypogonadism. *Nurs Clin* 2018;53(3):395–405.
- [90] Hermoso DAM, Bizerra PFV, Constantin RP, et al. Association between metabolic syndrome, hepatic steatosis, and testosterone deficiency: evidences from studies with men and rodents. *Aging Male* 2020;23(5):1296–315.
- [91] O'Shaughnessy PJ, Antignac JP, Le Bizec B, et al. Alternative (backdoor) androgen production and masculinization in the human fetus. *PLoS Biol* 2019;17(2):e3000002.
- [92] Pezzi V, Mathis JM, Rainey WE, et al. Profiling transcript levels for steroidogenic enzymes in fetal tissues. *J Steroid Biochem Mol Biol* 2003;87(2–3):181–9.
- [93] Li Y, Isomaa V, Pulkka A, et al. Expression of 3beta-hydroxysteroid dehydrogenase type 1, P450 aromatase, and 17beta-hydroxysteroid dehydrogenase types 1, 2, 5 and 7 mRNAs in human early and mid-gestation placentas. *Placenta* 2005;26(5):387–92.
- [94] Hill M, Paskova A, Kanceva R, et al. Steroid profiling in pregnancy: a focus on the human fetus. *J Steroid Biochem Mol Biol* 2014;139:201–22.
- [95] Karahoda R, Kallol S, Groessl M, et al. Revisiting steroidogenic pathways in the human placenta and primary human trophoblast cells. *Int J Mol Sci* 2021;22(4).
- [96] Pasqualini JR. Enzymes involved in the formation and transformation of steroid hormones in the fetal and placental compartments. *J Steroid Biochem Mol Biol* 2005;97(5):401–15.
- [97] Yoshida T, Matsumoto K, Miyado M, et al. Quantification of androgens and their precursors in full-term human placenta. *Eur J Endocrinol* 2021;185(5):K7–11.
- [98] Imamichi Y, Yuhki KI, Orisaka M, et al. 11-Ketotestosterone is a major androgen produced in human gonads. *J Clin Endocrinol Metab* 2016;101(10):3582–91.
- [99] Barnard L, Schiffer L, Louw du-Toit R, et al. 11-Oxygenated estrogens are a novel class of human estrogens but do not contribute to the circulating estrogen pool. *Endocrinology* 2021;162(3).
- [100] Boettcher C, Fluck CE. Rare forms of genetic steroidogenic defects affecting the gonads and adrenals. *Best Pract Res Clin Endocrinol Metabol* 2021:101593.
- [101] Hughes IA, Houk C, Ahmed SF, et al. Consensus statement on management of intersex disorders. *Arch Dis Child* 2006;91(7):554–63.

- [102] Goto M, Piper Hanley K, Marcos J, et al. In humans, early cortisol biosynthesis provides a mechanism to safeguard female sexual development. *J Clin Invest* 2006;116(4):953–60.
- [103] Kouri C, Sommer G, Fluck CE. Oligogenic causes of human differences of sex development: facing the challenge of genetic complexity. *Horm Res Paediatr* 2021:279–89.
- [104] Fabbri-Scallet H, de Sousa LM, Maciel-Guerra AT, et al. Mutation update for the NR5A1 gene involved in DSD and infertility. *Hum Mutat* 2020;41(1):58–68.
- [105] Camats N, Fluck CE, Audi L. Oligogenic origin of differences of sex development in humans. *Int J Mol Sci* 2020;21(5).
- [106] Mazen I, Abdel-Hamid M, Mekawy M, et al. Identification of NR5A1 mutations and possible digenic inheritance in 46,XY gonadal dysgenesis. *Sex Dev* 2016;10(3):147–51.
- [107] El-Khairi R, Achermann JC. Steroidogenic factor-1 and human disease. *Semin Reprod Med* 2012;30(5):374–81.
- [108] Batista RL, Costa EMF, Rodrigues AS, et al. Androgen insensitivity syndrome: a review. *Arch Endocrinol Metab* 2018;62(2):227–35.
- [109] Shukla GC, Plaga AR, Shankar E, et al. Androgen receptor-related diseases: what do we know? *Andrology* 2016;4(3):366–81.
- [110] Baillargeon J, Kuo YF, Fang X, et al. Long-term exposure to testosterone therapy and the risk of high grade prostate cancer. *J Urol* 2015;194(6):1612–6.
- [111] Azziz R. Polycystic ovary syndrome. *Obstet Gynecol* 2018;132(2):321–36.
- [112] O'Reilly MW, Kempegowda P, Jenkinson C, et al. 11-Oxygenated C19 steroids are the predominant androgens in polycystic ovary syndrome. *J Clin Endocrinol Metab* 2017;102(3):840–8.
- [113] Hammes SR, Levin ER. Impact of estrogens in males and androgens in females. *J Clin Invest* 2019;129(5):1818–26.
- [114] Davis SR, Baber R, Panay N, et al. Global consensus position statement on the use of testosterone therapy for women. *J Sex Med* 2019;16(9):1331–7.
- [115] Labcorp. Expected values & S.I. unit conversion tables.
- [116] Caron P, Turcotte V, Guillemette C. A quantitative analysis of total and free 11-oxygenated androgens and its application to human serum and plasma specimens using liquid-chromatography tandem mass spectrometry. *J Chromatogr A* 2021;1650:462228.
- [117] Kancheva L, Hill M, Vcelakova H, et al. The identification and simultaneous quantification of neuroactive androstane steroids and their polar conjugates in the serum of adult men, using gas chromatography-mass spectrometry. *Steroids* 2007;72(11–12):792–801.
- [118] Yazawa T, Uesaka M, Inaoka Y, et al. Cyp11b1 is induced in the murine gonad by luteinizing hormone/human chorionic gonadotropin and involved in the production of 11-ketotestosterone, a major fish androgen: conservation and evolution of the androgen metabolic pathway. *Endocrinology* 2008;149(4):1786–92.
- [119] Bloem LM, Storbeck KH, Swart P, et al. Advances in the analytical methodologies: profiling steroids in familiar pathways-challenging dogmas. *J Steroid Biochem Mol Biol* 2015;153:80–92.
- [120] Campana C, Rege J, Turcu AF, et al. Development of a novel cell based androgen screening model. *J Steroid Biochem Mol Biol* 2016;156:17–22.
- [121] Yazawa T, Sato T, Nemoto T, et al. 11-Ketotestosterone is a major androgen produced in porcine adrenal glands and testes. *J Steroid Biochem Mol Biol* 2021;210:105847.
- [122] Miyamoto H, Yeh S, Lardy H, et al. Delta5-androstenediol is a natural hormone with androgenic activity in human prostate cancer cells. *Proc Natl Acad Sci U S A* 1998;95(19):11083–8.