

Journal Pre-proof

Responses of reconstituted human bronchial epithelia from normal and health-compromised donors to non-volatile particulate matter emissions from an aircraft turbofan engine

Mathilde N. Delaval, Hulda R. Jonsdottir, Zaira Leni, Alejandro Keller, Benjamin T. Brem, Frithjof Siegerist, David Schönenberger, Lukas Durdina, Miriam Elser, Matthias Salathe, Nathalie Baumlin, Prem Lobo, Heinz Burtscher, Anthi Liati, Marianne Geiser

PII: S0269-7491(22)00735-7

DOI: <https://doi.org/10.1016/j.envpol.2022.119521>

Reference: ENPO 119521

To appear in: *Environmental Pollution*

Received Date: 6 October 2021

Revised Date: 19 May 2022


Accepted Date: 20 May 2022

Please cite this article as: Delaval, M.N., Jonsdottir, H.R., Leni, Z., Keller, A., Brem, B.T., Siegerist, F., Schönenberger, D., Durdina, L., Elser, M., Salathe, M., Baumlin, N., Lobo, P., Burtscher, H., Liati, A., Geiser, M., Responses of reconstituted human bronchial epithelia from normal and health-compromised donors to non-volatile particulate matter emissions from an aircraft turbofan engine, *Environmental Pollution* (2022), doi: <https://doi.org/10.1016/j.envpol.2022.119521>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Ltd.





Responses of reconstituted human bronchial epithelia from normal and health-compromised donors to non-volatile particulate matter emissions from an aircraft turbofan engine

Mathilde N. Delaval^{1,8*}, Hulda R. Jonsdottir^{1,9*}, Zaira Leni¹, Alejandro Keller², Benjamin T. Brem^{3,10}, Frithjof Siegerist⁴, David Schönenberger^{3,11}, Lukas Durdina^{3,12}, Miriam Elser^{3,13}, Matthias Salathe⁵, Nathalie Baumlin⁵, Prem Lobo⁶, Heinz Burtcher², Anthi Liati^{7#} & Marianne Geiser^{1§}

¹Institute of Anatomy, University of Bern, 3012 Bern, Switzerland. ²Institute for Sensors and Electronics, University of Applied Sciences and Arts Northwestern Switzerland, 5210 Windisch, Switzerland. ³Empa, Swiss Federal Laboratories for Materials Science and Technology, Laboratory for Advanced Analytical Technologies, 8600 Dübendorf, Switzerland. ⁴SR Technics, 8302 Kloten, Switzerland. ⁵Department of Internal Medicine, University of Kansas Medical Center, Kansas City, Kansas, USA. ⁶Metrology Research Centre, National Research Council Canada, Ottawa, Ontario K1A 0R6, Canada. ⁷Empa, Swiss Federal Laboratories for Materials Science and Technology, Automotive Powertrain Technologies Laboratory, 8600 Dübendorf, Switzerland.

⁸Present address: Joint Mass Spectrometry Center (JMSC) at Comprehensive Molecular Analytics (CMA), Helmholtz Zentrum München, D-85764 Neuherberg, Germany.

⁹Present address: Department of Rheumatology, Immunology, and Allergology, Inselspital University Hospital, Bern, Switzerland. Department of BioMedical Research, University of Bern, Bern, Switzerland. Spiez Laboratory, Federal Office for Civil Protection, Spiez, Switzerland

¹⁰Present address: Laboratory of Atmospheric Chemistry, Paul Scherrer Institute, 5232 Villigen, Switzerland

¹¹Present address: Empa, Laboratory for Air Pollution/Environmental Technology, 8600 Dübendorf, Switzerland

¹²Present address: Centre for Aviation, School of Engineering, Zurich University of Applied Sciences, 8401 Winterthur, Switzerland

¹³Present address: Empa, Swiss Federal Laboratories for Materials Science and Technology, Automotive Powertrain Technologies Laboratory, 8600 Dübendorf, Switzerland

Author Information

* Equal contribution

Date of death: April 1, 2021

§Corresponding author: Marianne Geiser, Institute of Anatomy, Baltzerstrasse 2, University of Bern, 3012 Bern, Switzerland. Email: marianne.geiser@unibe.ch

Abstract

Health effects of particulate matter (PM) from aircraft engines have not been adequately studied since controlled laboratory studies reflecting realistic conditions regarding aerosols, target tissue, particle exposure and deposited particle dose are logistically challenging. Due to the important contributions of aircraft engine emissions to air pollution, we employed a unique experimental setup to deposit exhaust particles directly from an aircraft engine onto re-differentiated human bronchial epithelia (HBE) at air-liquid interface under conditions similar to in vivo airways to mimic realistic human exposure. The toxicity of non-volatile PM (nvPM) from a CFM56-7B26 aircraft engine by sampling was evaluated under realistic engine conditions and exposing HBE derived from donors of normal and compromised health status to exhaust for one hour followed by biomarker analysis 24hours post exposure. Particle deposition varied depending on the engine thrust levels with 85% thrust producing the highest nvPM mass and number emissions with estimated surface deposition of 3.17×10^9 particles cm^{-2} or 337.1 ng cm^{-2} . Transient increase in cytotoxicity was observed after exposure to nvPM in epithelia derived from a normal donor as well as a decrease in the secretion of interleukin 6 and monocyte chemotactic protein 1. Non-replicated multiple exposures of epithelia derived from a normal donor to nvPM primarily led to a pro-inflammatory response, while both cytotoxicity and oxidative stress induction remained unaffected. This raises concerns for the long-term implications of aircraft nvPM for human pulmonary health, especially in occupational settings.

Keywords: Aerosol, aircraft engine exhaust, bronchial epithelial cell culture, cellular response, non-volatile particulate matter.

1. Introduction

Aircraft engine emissions contribute significantly to both global and local air pollution (Lee et al., 2020). Detailed characterization of these emissions and their adverse health effects is essential for the health and safety of airport workers as well as communities living in proximity of large airports. Due to the continuous growth of commercial air travel, emissions from aircraft engines have contributed to an increase in global air pollution in the last decades and are predicted to keep doing so in the future despite the reduced air-traffic during the COVID-19 pandemic (Manisalidis et al., 2020; Masiol and Harrison, 2014; Mazareanu, 2021; Peeters, 1998; Price and Probert, 1995). Non-volatile particulate matter (nvPM) from aircraft engines is very small, with mean mobility diameters typically smaller than 100 nm, also known as ultrafine particles (UFP), and can traverse the entire human respiratory tract upon inhalation (Lobo et al., 2015; Sturm, 2016a, b). These non-volatile particles are mainly comprised of soot and contain a minor fraction of inorganic, non-combustible ash PM (also referred to as metal-PM). Studies on ash PM, mainly from the automotive sector, have shown that ash consists of metal compounds formed by the combustion of fuel impurities, additives in lubricating oil, as well as metals (*e.g.*, iron, chromium, nickel, copper, tin) from the corrosion and mechanical wear of engine components (Gagné et al., 2021; Liati et al., 2013; Sappok and Wong, 2006; Vaarasmahti et al., 2005). We have previously shown that nvPM from a CFM56-7B26 aircraft engine induces oxidative stress in bronchial epithelial cells at ground-idle (GI) thrust and that the observed biological effects are not determined by deposited particle mass or number alone but rather influenced by particle morphology (Jonsdottir et al., 2019). Furthermore, variability between particle emissions generated by the combustion of fuels with different compositions has been reported, which adds complexity to determining the contribution of aircraft engine emissions to air pollution and its resulting health effects (Jonsdottir et al., 2019; Liati et al., 2019; Saffaripour et al., 2020). Overall, little is known about the general health hazards associated with PM from aircraft engine emissions. However, in recent years, several studies on the composition of aircraft engine exhaust and their effects on pulmonary health have been conducted (Cavallo et al., 2006; Gawron et al., 2020; Habre et al., 2018; He et al., 2020; He et al., 2018; Jonsdottir et al., 2019; Møller et al., 2014; Wing et al., 2020). Thus far, reported results suggest that since PM from aircraft engine emissions shares structural similarities with PM emitted from other combustion sources, particularly diesel exhaust, the resulting health effects would be similar (Bendtsen et al., 2019; He et al., 2020). Furthermore, repeated short (5h) exposures to UFP near a major airport have been associated with decreased lung function in healthy volunteers, further demonstrating the need for comprehensive characterization of aircraft engine emissions and their effects on human health (Lammers et al., 2020). Such studies are especially relevant to airport workers who spend extended periods of time on the tarmac, *e.g.*, baggage handlers, mechanics. Frequent

travelers, especially those with underlying respiratory diseases, might also be disproportionately affected.

As previously stated, PM emitted by aircraft engines is small in size, even smaller than that observed in road traffic pollution (Bendtsen et al., 2021; Harris and Maricq, 2001; Stacey, 2019). It is, therefore, able to deposit with high efficiency in the respiratory tract of humans and animals. However, as previously mentioned, absolute quantities of nvPM number and mass from aircraft engines are not the only metrics important to human health. We, and others, have observed a correlation between the physicochemical properties of these combustion-generated particles, both from aircraft and other combustion engines, and adverse health effects (Bendtsen et al., 2020; Jonsdottir et al., 2019). Moreover, the adequate evaluation of the adverse effects of aircraft engines on the human respiratory tract requires the use of a representative model system. In the current study, we exposed re-constituted human bronchial epithelia (HBE) to aircraft engine nvPM at the air-liquid interface, mimicking *in vivo* exposure. These epithelial cell cultures are three-dimensional, pseudostratified, and contain the characteristic epithelial cell types of this lung compartment, such as basal, goblet, and ciliated cells. They produce mucus and exhibit coordinated ciliary beating similar to *in vivo* respiratory epithelium (de Jong et al., 1994; de Jong et al., 1993). In the present study, we evaluated the health effects of nvPM from a CFM56-7B26 aircraft turbofan engine burning standard Jet A-1 fuel at different thrust levels in HBE derived from three individual donors of different background. We deposited nvPM from aircraft engine exhaust directly onto the apical surface of the epithelium under physiological conditions using the portable Nano-Aerosol Chamber for In-Vitro Toxicity (NACIVT) (Jeannet et al., 2015). Additionally, we studied the morphology of the deposited particles by Transmission Electron Microscopy (TEM). The combination of a realistic particle source and a physiological three-dimensional cell culture model provides a unique platform to study the effects of aviation emissions on the human respiratory tract in a controlled experimental set-up.

2. Materials & Methods

2.1 Summary of experimental design

Combustion aerosol was emitted by a CFM56-7B26 turbofan engine and sampled with a standardized sampling system as previously described (Jonsdottir et al., 2019; Liati et al., 2019). To investigate the correlation between nvPM generated at high and low engine thrust levels and cellular effects in pulmonary cells, we sampled aerosols from three distinct thrust levels, 85% (climb-out), 7% (taxi), and ground idle (GI, 3%), along with filtered aerosol (at 65% thrust) as particle-free (P-free) control. The exposure campaign took place in July 2018 at Zürich Airport, Zürich, Switzerland. Aerosol was sampled on 4 consecutive days. We exposed re-differentiated primary human bronchial epithelium from three donors of varying background to exhaust at physiological conditions for one hour. Each individual exposure was repeated two or three times, depending on the availability of cellular material. Overview over number of exposures, replicates, and donors can be found in Table S1. At 24-hours post exposure, we assessed distinct biomarkers of pulmonary injury. In addition to physicochemical characterization of the exhaust, we measured the morphology and composition of nvPM by analytical microscopy.

2.2 Aerosol generation, sampling, and characterization

An airworthy CFM56-7B26 turbofan engine, running in a test cell at SR Technics at Zürich Airport in Switzerland was used as aerosol source. The engine was fueled with Jet A-1 with fuel properties well within the allowable range for commercial jet fuel. The engine operating conditions were determined using the engine combustor inlet temperature (T3), which correlated to sea-level static thrust levels corrected to international standard atmospheric conditions (15 °C, 1013.25 hPa). This approach has been used for emissions certification of aircraft engines and research experiments (Durdina et al., 2017; ICAO, 2018; Lobo et al., 2020). The aircraft engine nvPM emissions were collected using a single point sampling probe and a standardized sampling system, compliant with the specifications listed in the standards and recommended practices (ICAO, 2017; SAE International, 2018), also used in our previous study (Jonsdottir et al., 2019). Briefly, the extracted nvPM sample is diluted with dry synthetic air by a factor of 8–14 and then transferred to the diagnostic instruments using a 25 m long carbon-loaded, electrically grounded polytetrafluoroethylene line. The nvPM mass concentration was measured using a Micro Soot Sensor (MSS, Model 483, AVL List GmbH, Austria) (Schindler et al., 2004), the nvPM number concentration was determined with an AVL particle counter (APC, Model 489, AVL List GmbH, Austria) (Lobo et al., 2020; Lobo et al., 2015), and particle size distributions were measured with a scanning mobility particle sizer (SMPS, Model 3938, consisting of a long differential mobility analyzer Model 3081A, a soft X-ray aerosol neutralizer Model 3088, and a condensation particle

counter Model 3776, TSI Inc., USA). The lower particle size cut-off was 10 nm for the APC (50% of particles counted) and 6 nm for the SMPS measurements. Particle size distribution data from the SMPS were analyzed using Aerosol Instrument Manager (AIM 10.2, TSI Inc.). The SMPS instrument made one scan every 30 seconds. The reported distributions are averages of multiple scans during the 60-minute sampling window. The NACIVT chamber was connected to the diluted PM sampling line in parallel to particle instrumentation. Volatile organic compounds were removed upstream of the NACIVT chamber with a customized low flow thermodenuder (Fierz et al., 2007) operated at 200 °C on the preconditioning and first absorption sections, and 100 °C on the second absorption section. All PM data were plotted using Igor Pro 7.0 (Wavemetrics Inc.) and Origin 2019 (Originlab Inc). Estimated deposition of nvPM onto the apical surface of the re-differentiated epithelia was calculated as previously described (Jeannet et al., 2015). General chemical characteristics of Jet-A1 aviation fuel were described previously (Jonsdottir et al., 2019).

2.3 Cell Cultures

Human bronchial epithelial cells (HBEC) were isolated from human lungs unsuitable for transplantation appropriately consented for donation and recovered by the Life Alliance Organ Recovery Agency (LAORA) Miami (Miami, Florida, USA). The cells were collected from the proximal conducting airways of three individual donors with different background: one normal healthy donor with no smoking history (donor 1) and two asthmatic donors with varying smoking history (donors 2 and 3) (Table S1). Air-Liquid Interface (ALI) cultures of re-differentiated HBE were generated as previously described (Jonsdottir and Dijkman, 2015; Künzi et al., 2015; Künzi et al., 2013; Schmid et al., 2010). Briefly, bronchial epithelial cells were maintained in submerged two-dimensional culture in Bronchial Epithelial Cell Growth Medium (BEGM - LHC base media with supplements, Gibco, Fisher Scientific, Reinach, Switzerland). Cells were thereafter seeded onto porous 0.33 cm² Transwell® inserts (Corning International, Fisher Scientific, Reinach, Switzerland) in chemically defined medium that induces terminal differentiation. Once confluent, apical medium was removed establishing ALI and the epithelium allowed to differentiate over a period of 4 weeks, with apical washes and basal media change three times a week (Jonsdottir and Dijkman, 2015). Terminal differentiation results in a pseudostratified ciliated epithelium with established air-liquid interface. Mucus secretion and ciliary beating were routinely checked visually and by light microscopy, respectively. Differentiated epithelia were washed with Dulbecco's phosphate-buffered saline (DPBS with Ca²⁺ and Mg²⁺, Invitrogen, Lucerne, Switzerland) 1 to 2 h before aerosol exposure.

2.4 Aerosol exposure

Bronchial epithelia derived from three separate human donors were divided into two or three sets of 6 inserts per donor, and each series of inserts was exposed to combustion-generated aerosol from a CFM56-7B26 turbofan engine at different thrust levels (85%, 7% and GI) for 60 min within the NACIVT at physiological conditions, *i.e.*, 37 °C, 5% CO₂, and > 85% relative humidity on separate days Table S1). For P-free air exposure, we mounted a Balston DFU Model 9933-11, grade BQ filter (Parker Hannifin Corporation, New York, USA) between the aerosol exhaust line and the thermodenuder. Particle deposition was observed in real-time with Lab View 9.0.1. After exposure, cell cultures were incubated at the same conditions for 1 h before collecting apical wash samples and subsequently incubated for additional 23 hours followed by final sampling.

2.5 Analyses of cellular responses

To evaluate the cellular effects of the exposure to combustion aerosol, we analyzed several biomarkers at 1-h and/or 24-h post exposure. To assess cytotoxicity, we quantified the release of Adenylate Kinase (AK) by damaged cells into the apical compartment using the commercial ToxiLight kit (LONZA, Visp, Switzerland) according to the manufacturer's instructions. Cytotoxicity is reported as fold AK release over P-free (particle-filtered exhaust) controls. To assess oxidative stress, we analyzed the gene expression of Heme Oxygenase 1 (HMOX-1) by quantitative real-time polymerase chain reaction (qRT-PCR) by extracting total cellular RNA from exposed epithelia with Trizol (Sigma Aldrich, Buchs, Switzerland) and ZymoResearch DirectZol Mini Prep Plus columns (LucernaChem, Luzern, Switzerland) according to the manufacturer's protocol. Complementary DNA (cDNA) was prepared using the QuantiTect® reverse transcription kit (Qiagen, Hombrechtikon, Switzerland) according to the manufacturer's instructions. Briefly, genomic DNA (gDNA) was removed from all samples by incubating extracted RNA in gDNA Wipeout buffer for 2 min at 42 °C and immediately transferring samples to ice. Reverse transcription (RT) of 150 ng total RNA was performed with the provided RT mixes for 15 min at 42 °C and inactivated for 3 min at 95 °C. For gene expression analysis, 0.2 µL of total cDNA was amplified using the Applied Biosystems 7900HT system (Thermo Fisher Scientific) using the following cycling parameters: 15 min at 95 °C, 45 cycles of 15 s at 94 °C, 30 s at 55 °C, and 30 s at 72 °C, followed by a dissociation step to confirm product specificity. Data were normalized to Hypoxanthine-Guanine Phosphoribosyl Transferase (HPRT) using the $\Delta\Delta C_t$ method (Livak and Schmittgen, 2001). Biological replicates (n = 3–9 cultures) were analyzed three times using Applied Biosystems SDS v2.4. Data are presented as fold change over P-free controls. The release of the inflammatory mediators, Interleukin (IL)-6, IL-8, and Monocyte Chemoattractant Protein (MCP)-1 into basal media was assessed 24 h after

aerosol exposure. Cytokine release was measured using a sandwich Enzyme Linked Immunosorbent Assay (ELISA) according to the manufacturer's instructions (DuoSet ELISA; R&D Systems Europe, Abingdon, United Kingdom). All samples were stored at -20 °C from sampling to analysis.

2.6 Transmission Electron Microscopy (TEM)

During experiments, nvPM was collected directly on TEM grids, in parallel with cell exposures. Two types of TEM grids were used in the current study: i) Cu-supported holey carbon film grids for detailed study of soot and ii) ultra-thin carbon film grids for better optical contrast facilitating observation of ash particles. TEM imaging was performed with a JEOL 2200FS microscope fitted with an Omega filter, a Schottky field emission gun at 200 kV, and a 0.23 nm point-to-point resolution (Electron Microscopy Center of Empa, Dübendorf, Switzerland). The TEM instrument is equipped with an EDX detector (JEOL EDX detector: EX-24065JGT), which was used for the elemental analysis of the inorganic, ash particles. Gatan DigitalMicrograph® was used for image analysis.

2.7 Statistical analysis

Re-differentiated epithelia ($n = 12-18$, *i.e.* 6 replicates per exposure) were exposed to either aircraft engine exhaust aerosol or to particle-filtered air. The number of biological replicates depended primarily on the availability of cellular material from the different donors and space in the deposition chamber (24 slots). Furthermore, specific focus was put on GI thrust levels based on our previous publication (Jonsdottir et al., 2019), which was analyzed in all donors. Biological data are presented as fold change over P-free control or absolute quantification (pg mL^{-1}). Data are presented as mean \pm standard deviation (SD). Statistical significance was determined with the commercial software GraphPad Prism 8.0 for Windows (GraphPad Software Inc., La Jolla, California, USA). Non-matching one-way analysis of variance (ANOVA) with Tukey's or Sidak's multiple comparison tests were used for statistical comparison of cytotoxicity and biological responses to negative control (particle-free air exposure). Parameters and applied statistical tests can be found in the corresponding figure legends. p values < 0.05 were considered statistically significant.

3. Results

3.1 Successful deposition of nvPM from aircraft engine exhaust on re-differentiated human bronchial epithelia

Electrometer data collected within the NACIVT deposition chamber show distinct diffusion charging currents for the three thrust levels sampled (Figure 1a). Currents of -3995 ± 340 fA obtained for 85% thrust indicate the highest particle deposition, while the -225 ± 24 and -83 ± 9 fA registered for GI and 7% thrust, respectively, point to low deposition. Precipitation voltages for P-free exhaust (-43 ± 3 fA) were identical to those observed for synthetic air (SA, dashed line), demonstrating the successful removal of nvPM by the particle filters. The resulting particle size distributions (Figure 1b) were unimodal and lognormal. Geometric median diameters (GMDs) of particles from GI and 7% thrust are similar at 16.3–16.4 nm. Particles sampled at 85% thrust displayed the largest median diameters at 45.8 nm. Estimated deposition of both mass and number of nvPM from the exhaust onto bronchial epithelia (Figures 1c and 1d) was calculated based on Jeannet et al. (Jeannet et al., 2015) and correlates with the observed precipitation voltages from within the chamber, indicating successful deposition of nvPM. The average deposition per surface area of cell culture was 3.17×10^9 particles cm^{-2} or 337.1 ng cm^{-2} for the 85% thrust condition, 0.22×10^9 particles cm^{-2} or 0.9 ng cm^{-2} for 7% thrust, and 1.04×10^9 particles cm^{-2} or 2.3 ng cm^{-2} for GI.

3.2 TEM, HRTEM and TEM-EDX analyses of soot and ash collected on TEM grids during exposure

TEM analysis of particles collected at high (85%) and low (GI) thrust level conditions revealed a clear correlation between engine thrust level and the amount of collected soot. It is widely accepted that due to higher surface-to-volume ratio, soot with small particle sizes is more reactive with atmospheric components than that with larger sizes (Al-Qurashi and Boehman, 2008; Harris, 1990; Pahalagedara et al., 2012; Vander Wal et al., 2010; Yehliu et al., 2011). Based on this fact, we determined the size of the soot agglomerates (maximum length) and of their primary particle constituents (diameter of their circular projection, Figure 2). The results revealed soot agglomerates sampled from 85% thrust conditions are significantly larger (size mode: 60–160 nm, more rarely 200 nm) than those sampled from GI thrust conditions (< 40 nm). The size of the primary soot particles within the agglomerates also correlated with thrust levels. The primary soot particles from GI were 5–10 nm, while those from 85% thrust were 10–25 nm, more rarely 30 nm. These results are in line with previous observations (Jonsdottir et al., 2019; Liati et al., 2019; Saffaripour et al., 2020). Ash was present in minor yet detectable amounts among collected particles on examined TEM grids. As characterization of particles by electron microscopy is generally not a quantitative technique, it cannot be implied whether ash can

be considered to occur in negligible amounts among nvPM (Gagné et al., 2021). The TEM/EDX analyses of ash under GI conditions revealed that the most prominent element is iron (Fe), often together with Chromium (Cr) \pm Nickel (Ni) and/or Titanium (Ti), in form of single particles, fragments or aggregates, a few hundreds of nanometers large, rarely down to tens of nanometers. The aggregates themselves consist of tens of nm-large single particles with the same or different chemical composition. Another frequently found element is Calcium (Ca), in the form of an oxide but also together with Sulfur (S) \pm Phosphor (P) \pm Chloride (Cl). Ca-bearing particles occur as single fragments or as aggregates with sizes of a few hundreds of nanometers. Iron (Fe)- and, more often, Ca-particles were also found inside soot. Other elements detected in ash particles include Aluminum (Al), Magnesium (Mg), Tin (Sn), Silicon (Si), and more rarely Potassium (K), Manganese (Mn), Zinc (Zn) and Bismuth (Bi), the latter a few tens of nanometers large. Analysis of ash from high engine thrust conditions (85%) revealed the same chemical composition as that from low (GI) thrust. However, Silver (Ag) was identified repeatedly in high thrust ash samples with sizes of a few tens of nm. Ash particle sizes, in general, were similar among both examined thrust conditions.

3.3 Biological responses in epithelia from a normal donor after 1-hour exposure to nvPM

Exposures to both GI and 7% thrust led to a statistically significant, immediate (1 hour), two-fold increase of AK release in normal cells (donor 1) compared to control cells exposed to P-free air ($p < 0.0001$ and $p = 0.0001$, respectively). This increase of AK release is also observed at 24-h post exposure to 7% thrust, although at lower level ($p = 0.0144$). There was no significant increase in cytotoxicity over P-free controls after exposure to nvPM at 85% thrust. There was no correlation between deposited particle mass or number and the observed cytotoxicity. Exposure to nvPM from 7% thrust led to a statistically significant decrease in the release of both the pro-inflammatory cytokine IL-6 ($p = 0.0092$) and the chemokines MCP-1 ($p = 0.253$) and IL-8 ($p = 0.0299$) compared to P-free controls. MCP-1 levels also decreased after exposure to GI thrust ($p = 0.0024$).

3.4 Comparison of biological responses after 1-hour of exposure to ground idle thrust in epithelia from three individual donors

Comparison of the same biological parameters in epithelia derived from three separate donors and exposed to GI thrust revealed significant differences in the release of adenylylate kinase (AK) at 1 h and 24-h post exposure (Figure 4a) and the secretion of MCP-1 at 24-h post exposure (Figure 4c) for donor 1. The epithelia from the two additional donors did not exhibit any alterations in the tested parameters at 24-h post exposure. Additionally, the overall biological responses of epithelia derived from donors

2 and 3 were similar apart from baseline secretions of IL-8, which were higher in epithelia of donor 3 (Figure 4c).

3.5 Non-replicated analysis of biological responses after multiple exposures to nvPM emissions in a normal donor

To determine the adverse effects of multiple exposures to nvPM from distinct thrust levels, we exposed N-HBE to high (85%) and low (7%) engine thrust levels for one hour, once a day for up to 3 days (Figure S1). Multiple exposures to nvPM at 7% thrust did not lead to an increase in cytotoxicity, while a single exposure caused a significant, immediate increase of AK release (average 1.6-fold, $p < 0.05$) that had resolved 24-h post exposure. Furthermore, nvPM from 7% thrust induced a significant, non-dose dependent decrease of IL-6 ($p < 0.01$ – $p < 0.05$) and a significant, dose-dependent decrease of MCP-1 ($p < 0.001$ – $p < 0.0001$), similar to 85% thrust. However, multiple exposures to 7% thrust did not affect IL-8 levels, unlike 85% thrust, which caused an almost two-fold increase of IL-8 after the second exposure ($p < 0.0001$) compared to controls. A second exposure to 85% thrust induced a decrease of MCP-1 ($p < 0.001$) compared to P-free controls. Multiple exposures to nvPM of low and high engine thrust levels did not induce significant changes in the expression of the HMOX-1 gene, indicating low oxidative stress.

4. Discussion

In the present study, we evaluated the respiratory health effects of nvPM from a CFM56-7B26 turbofan, one of the most commonly used aircraft engines in the world, by using a unique experimental setup for particle sampling and cellular deposition. We sampled nvPM directly from the engine exhaust under representative operating conditions and deposited the particles onto the apical surface of re-differentiated bronchial epithelia (HBE) to evaluate their effects on acute toxicity and inflammation. This unique combination of experimental systems is able to realistically represent *in vivo* particle deposition in the human airways. Furthermore, we assessed these effects in HBE from single donors of normal and compromised health status, providing a relevant overview of the potential adverse respiratory effects of nvPM from aircraft engine exhaust.

The nvPM emission profiles of the CFM56-7B26 turbofan engine have been extensively characterized in previous studies (Brem et al., 2015; Durdina et al., 2021; Lobo et al., 2020). The nvPM mass-based emissions have been found to be higher at engine levels corresponding to idle (3–7% thrust), decreasing to a minimum at 15–30% engine thrust levels, and then increasing to maximum rated thrust. Similarly, the nvPM number-based emissions are higher at engine idle and decrease at low engine thrust levels, however, they increase up to 60% thrust and then decrease again to maximum rated thrust. The size distributions vary with engine thrust levels, with geometric mean diameter of particles ranging from ca. 10 nm to 40 nm (Durdina et al., 2021; Lobo et al., 2020). Furthermore, the particle effective density has been observed to increase with engine thrust levels and decrease with particle size (Abegglen et al., 2015; Durdina et al., 2014). The chemical composition and radiative properties of nvPM emissions from the CFM56-7B26 have also been reported (Elser et al., 2019).

Upon exposure to nvPM from three distinct engine thrust levels, we observed immediate damage in epithelia derived from a normal donor (donor 1), especially after exposure to lower thrust levels (7% and GI), indicating that this observed effect is not related to either nvPM mass or number, since the highest nvPM deposition was observed after exposure to 85% thrust. This is in line with other published studies as well as our previous observations (BéruBé et al., 2007; Jaramillo et al., 2018; Jonsdottir et al., 2019; Schmid and Stoeger, 2016). Exposure to PM emissions is known to interfere with inflammatory cytokine homeostasis, where both up- and down-regulation of modulators have been reported in both experimental and human studies (Bendtsen et al., 2019; Habre et al., 2018; He et al., 2020; He et al., 2018). However, most current studies on the biological effects of aircraft emissions cover total PM and do not focus on non-volatile PM specifically. For instance, mice exposed to total PM from aircraft exhaust for up to 90 days displayed an acute inflammation with immune cell influx to the lungs, which persisted for up to 28 days (Bendtsen et al., 2019). Additionally, Habre et al. detected acute systemic inflammation, characterized by increased circulation of the pro-inflammatory

cytokine IL-6 in asthmatic, non-smoking adults exposed to total UFP downwind of Los Angeles International Airport (LAX) for 2 hours (Habre et al., 2018). In the present study, we did not observe an increase in IL-6 release in any donor after single 1-hour exposures. Increased expression of IL-6 has also been observed in bronchial epithelial cell lines exposed to both low and high doses of UFP from aircraft exhaust (He et al., 2020; He et al., 2018) and those results are in line with our previous observations with the undifferentiated human bronchial cell line, BEAS-2B (Jonsdottir et al., 2019). In contrast, here we observed a decrease in the secretion of IL-6, MCP-1, and IL-8 in epithelia from a normal donor after exposure to 7% thrust. Secretion of MCP-1 also decreased after exposure to GI thrust. Non-repeated analysis of multiple exposures of the normal donor to nvPM appeared to down-regulate the inflammatory response even further. No significant changes in cytokine secretions were observed in the other two donors.

We previously observed an increase in acute cellular toxicity and oxidative stress in BEAS-2B cells after exposure to ground idle (GI) thrust using the same experimental setup (Jonsdottir et al., 2019). Similarly, in the present study, exposure to GI and 7% thrust conditions resulted in increased cytotoxicity in epithelia derived from a normal donor (donor 1) 1 h post exposure. After 24 hours, only epithelia exposed to 7% thrust exhibit minor increased cytotoxicity. In contrast to our previous study with BEAS-2B cells, we found no increase in the expression of HMOX-1 in any donor after exposure to any thrust condition, indicating that re-differentiated human bronchial cells are more robust, when it comes to the induction of the oxidative stress response compared to two-dimensional BEAS-2B cells. Additionally, this could indicate that one-hour exposures are simply not long enough to induce drastic changes in re-constituted airway epithelia. The airways are in constant contact with various types of aerosols throughout our lives and having a high tolerance for aerosol exposures would be pertinent to their function. Furthermore, we did not observe any effect of GI aerosol exposure in HBE cultures of the health-compromised donors (donors 2 and 3). We consider it very likely that, in general, diseased epithelia are more susceptible to damage from nvPM exposure and therefore, we hypothesize that this could be related to the asthmatic background and smoking history of these donors, since their derived bronchial epithelia show characteristics of diseased epithelium. In the present study, we combined a state-of-the art sampling system coupled to a realistic aerosol source with complex three-dimensional cell cultures and conducted aerosol exposures at physiological conditions. This experimental system is unique but not without its limitations. While this study was conducted with more physiologically relevant cell culture models, we could only analyze one donor per health status. Although this limits the generality of our results, we still consider these data a strong basis for further investigation into the likely adverse health effects from exposure to aircraft exhaust. Additionally, although the re-constituted epithelia used here is comprised of many different cell types, we are only able to elucidate the epithelial response to aerosol exposure due to the absence of an immune system

or cells of the adjoining connective tissue (*i.e.*, fibroblasts, endothelial cells). However, given that the respiratory epithelium is the entry point of emission aerosols, the epithelial response is likely to represent a sizable portion of the resulting effects making it a strongly relevant experimental parameter. Though exposure times were realistic, they were also limited due to the operational schedule of the engine test cell and fuel costs. Longer exposures may have potentially resulted in an increased epithelial response.

Despite these limitations, we attempted to cover a broad variety of parameters and found that adverse effects do not seem to be primarily related to deposited nvPM mass and number, since very low doses of nvPM (7%, GI) did cause significant cytotoxicity in normal HBE. The size of nvPM (below 100 nm) is inversely proportional to its surface reactivity, which could explain the higher effect of nvPM from the two lowest thrust levels, in comparison to those from 85% thrust. Generally, physicochemical characterization of the particles is necessary to fully understand the potential adverse effects caused by nvPM. Here, we observed a clear correlation between engine thrust levels and the number, as well as the size of soot agglomerates and their primary particle constituents. GI thrust produced substantially fewer and smaller soot particles than those from 85% thrust, correlating with electrometer data obtained from within the deposition chamber. Based on the difference in size of the soot particles collected under different engine thrust levels, soot from GI thrust is more reactive than that collected from 85% thrust, which may explain the toxicological effects observed in response to nvPM from low engine thrust levels. There was no difference in ash particle size observed among the different thrust level conditions. Compared to soot, ash production occurs in minute amounts and the most frequent element in ash is iron (Fe), either alone or in combination with Chromium (Cr), Nickel (Ni), and/or Titanium (Ti), (mainly derived from corrosion and the mechanical wear of engine components. Other observed metallic components from combustion of fuel and/or lube oil additives were primarily Ca \pm S \pm P \pm Cl, less frequently Al, Mg, Sn, Si, more rarely K, Mn, Zn, Bi, and, at high thrust levels, Ag. Based on these physicochemical properties, aircraft exhaust particles are similar to those in diesel exhaust (Bendtsen et al., 2021).

Realistic *in vitro* exposures are crucial to comprehensively decipher the effects of UFP on human health; this includes using representative doses for human exposure. In this study, we exposed pulmonary epithelia to realistic doses of NvPM from three separate thrust levels, *i.e.*, climb-out (85%), taxi (7%) and ground idle (3%). Here, the deposited mass of particles ranged from 0.9 ng cm⁻² to 337.1 ng cm⁻² after 1 hour of exposure, covering the range of ambient exposures to high daily occupational exposures (Paur et al., 2011). Although airport workers are mainly exposed to particles generated at low thrust levels, particles emitted at high levels are also relevant to human exposure and health, since the air mass in and around the airport is constantly being mixed and influenced by both airport

schedules and prevailing winds. Indeed, Buonanno et al. found that particle number concentrations measured in the vicinity of the runway present several main short-term peaks during the workday, related to take-off and landing of aircraft and pre-flight operations. They measured an average total concentration of 6.5×10^3 particles cm^{-3} at a static receptor site. However, when measuring the occupational exposure to airborne particles and other pollutants at the airport, it could be as high as 2.5×10^4 particles cm^{-3} and these exposure levels vary depending on the role of the worker at the airport (Buonanno et al., 2012). Similarly, Møller et al. found that baggage handlers are exposed to more ultrafine particles than employees working inside the airport. Levels of exposure were evaluated from 5×10^3 UFP cm^{-3} for inside workers up to 3.7×10^4 UFP cm^{-3} for baggage handlers (Møller et al., 2014). However, these studies seem to report particle number concentrations on the lower end (Masiol and Harrison, 2014). Furthermore, the study of a cohort of male workers at the Copenhagen airport may shed much needed light on the incidence of respiratory diseases developed and/or exacerbated in relation to aircraft exhaust exposure by estimating the contribution of aviation related UFP to total personal exposure (Møller et al., 2017). In this context, our study highlights the importance of evaluating the effects of aircraft exhaust nvPM, even at a low dose, since surface exposure as low as 1×10^9 particles cm^{-2} and 1 ng cm^{-2} can cause a biological response, especially in health-compromised epithelia.

5. Conclusions

Toxicological data on the adverse health effects of particles emitted by aircraft engines are still scarce. This is partly due to general experimental difficulty, *i.e.*, particle collection at the airport and performing realistic *in vitro* and *in vivo* toxicity studies relevant to the human health hazard and risk assessments. Here, using a unique experimental setup for particle sampling directly from the engine running under representative thrust conditions and an instrument specifically developed to mimic inhalation exposure, we demonstrated that exposure to non-volatile particle matter causes biological responses in epithelia derived from three single donors of different background. Our results support the need to further characterize the pulmonary toxicity of aircraft engine exhaust and reinforce the need for establishing Occupational Exposure Limits (OELs) for such particles to protect airport workers, especially those with underlying respiratory conditions. Based on the current data, clarifications of the adverse effects of multiple exposures to nvPM from aircraft emissions should be of specific interest to both airport employees and neighboring communities.

Acknowledgements

We thank Y. Arroyo, Electron Microscopy Center, Empa, for assistance during TEM imaging.

Funding sources

This research was supported by the Swiss Federal Office for Civil Aviation (FOCA), projects 2016-037 (granted to M.G.) and 2015-113 (granted to B.T.B.). M.S. and N.B. acknowledge funding by the National Institutes of Health (NIH) grants NHLBI (R01 HL133240 and R01 HL157942).

Competing interests

The authors declare no competing interests.

Data availability

All relevant raw data are available from the authors upon reasonable request.

Figure Legends:

Fig. 1. Deposition of non-volatile particulate matter (nvPM) from a CFM56-7B26 turbofan onto re-differentiated human bronchial epithelia (HBE). **a)** Recorded diffusion charging current for each thrust level (0.32mv/fA). **b)** Size distributions of deposited particles. **c)** Mass of nvPM deposited per surface area of cell culture after one hour of exposure. **d)** Number of nvPM deposited per surface area of cell culture after one hour of exposure. Highest deposition is observed for 85% thrust and lowest for the second lowest thrust level (7%), which is less deposition than after exposure to ground idle (GI, 3%). Data for diffusion current, particle mass and number are reported as mean and SD of four (85%) or three (7%, GI) independent aerosol generations for cell exposures.

Fig. 2. Morphology of non-volatile particulate matter (nvPM) soot emissions. Transmission electron microscopy (TEM) images showing the larger size and higher abundance of soot particles, **a)** at 85% thrust compared to **b)** ground idle (GI) condition. Scale bar: 200 nm.

Fig. 3. Biological responses in bronchial epithelia of a normal donor after 1 hour of exposure to non-volatile particulate matter (nvPM) from different thrust levels. **a)** Adenylate kinase (AK) released from damaged cells into the apical compartment, presented as fold change over particle-filtered (P-free) controls, at 1-h and 24-h post exposure (hpe). **b)** HMOX-1 gene expression 24 hpe as determined by qPCR, presented as fold change over P-free controls. **c)** IL-6, **d)** MCP-1, and **e)** IL-8 secretions into the basal compartment 24 hpe as determined by ELISA, presented as pg mL⁻¹. Data is presented as mean \pm SD and $n = 9-18$ cultures from three independent exposures. Statistical significance was assessed using a non-matching two-way analysis of variance (ANOVA) with Sidak's multiple comparison test (AK release) or a non-matching one-way analysis of variance (ANOVA) with Tukey's multiple comparison test (HMOX-1, IL-6, MCP-1, IL-8): * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, and **** $p < 0.0001$.

Fig. 4. Comparison of biological responses after 1-hour of exposure to ground idle thrust conditions in three individual donors. **a)** Adenylate kinase (AK) released from damaged cells into the apical compartment, presented as fold change over particle-free (P-free) controls, at 1 and 24-hours post exposure (hpe). **b)** HMOX-1 gene expression 24 hpe as determined by qPCR, presented as fold change over P-free controls. **c)** IL-6, MCP-1, and IL-8 secretions into the basal compartment 24 hpe as determined by ELISA, presented as pg mL⁻¹. Data is presented as mean \pm SD and $n = 9-18$ cultures from three independent exposures. Statistical significance was assessed using a non-matching one-way analysis of variance (ANOVA) with Tukey's multiple comparison test (HMOX-1, IL-6, MCP-1, IL-8): *** $p < 0.001$, and **** $p < 0.0001$.

References


- Abegglen, M., Durdina, L., Brem, B.T., Wang, J., Rindlisbacher, T., Corbin, J.C., Lohmann, U., Sierau, B., 2015. Effective density and mass–mobility exponents of particulate matter in aircraft turbine exhaust: Dependence on engine thrust and particle size. *Journal of Aerosol Science* 88, 135-147.
- Al-Qurashi, K., Boehman, A.L., 2008. Impact of exhaust gas recirculation (EGR) on the oxidative reactivity of diesel engine soot. *Combustion and Flame* 155, 675-695.
- Bendtsen, K.M., Bengtsen, E., Saber, A.T., Vogel, U., 2021. A review of health effects associated with exposure to jet engine emissions in and around airports. *Environmental Health* 20, 10.
- Bendtsen, K.M., Brostrøm, A., Koivisto, A.J., Koponen, I., Berthing, T., Bertram, N., Kling, K.I., Dal Maso, M., Kangasniemi, O., Poikkimäki, M., Loeschner, K., Clausen, P.A., Wolff, H., Jensen, K.A., Saber, A.T., Vogel, U., 2019. Airport emission particles: exposure characterization and toxicity following intratracheal instillation in mice. *Particle and Fibre Toxicology* 16, 23.
- Bendtsen, K.M., Gren, L., Malmborg, V.B., Shukla, P.C., Tunér, M., Essig, Y.J., Kraus, A.M., Clausen, P.A., Berthing, T., Loeschner, K., Jacobsen, N.R., Wolff, H., Pagels, J., Vogel, U.B., 2020. Particle characterization and toxicity in C57BL/6 mice following instillation of five different diesel exhaust particles designed to differ in physicochemical properties. *Particle and Fibre Toxicology* 17, 38.
- Brem, B.T., Durdina, L., Siegerist, F., Beyerle, P., Bruderer, K., Rindlisbacher, T., Rocci-Denis, S., Andac, M.G., Zelina, J., Penanhoat, O., Wang, J., 2015. Effects of Fuel Aromatic Content on Nonvolatile Particulate Emissions of an In-Production Aircraft Gas Turbine. *Environmental Science & Technology* 49, 13149-13157.
- Buonanno, G., Bernabei, M., Avino, P., Stabile, L., 2012. Occupational exposure to airborne particles and other pollutants in an aviation base. *Environ Pollut* 170, 78-87.
- BéruBé, K., Balharry, D., Sexton, K., Koshy, L., Jones, T., 2007. Combustion-derived nanoparticles: mechanisms of pulmonary toxicity. *Clin Exp Pharmacol Physiol* 34, 1044-1050.
- Cavallo, D., Ursini, C.L., Carelli, G., Iavicoli, I., Ciervo, A., Perniconi, B., Rondinone, B., Gismondi, M., Iavicoli, S., 2006. Occupational exposure in airport personnel: Characterization and evaluation of genotoxic and oxidative effects. *Toxicology* 223, 26-35.
- de Jong, P.M., van Sterkenburg, M.A., Hesseling, S.C., Kempenaar, J.A., Mulder, A.A., Mommaas, A.M., Dijkman, J.H., Ponc, M., 1994. Ciliogenesis in human bronchial epithelial cells cultured at the air-liquid interface. *Am J Respir Cell Mol Biol* 10, 271-277.
- de Jong, P.M., van Sterkenburg, M.A., Kempenaar, J.A., Dijkman, J.H., Ponc, M., 1993. Serial culturing of human bronchial epithelial cells derived from biopsies. *In Vitro Cell Dev Biol Anim* 29a, 379-387.
- Durdina, L., Brem, B.T., Abegglen, M., Lobo, P., Rindlisbacher, T., Thomson, K.A., Smallwood, G.J., Hagen, D.E., Sierau, B., Wang, J., 2014. Determination of PM mass emissions from an aircraft turbine engine using particle effective density. *Atmospheric Environment* 99, 500-507.
- Durdina, L., Brem, B.T., Elser, M., Schönenberger, D., Siegerist, F., Anet, J.G., 2021. Reduction of Nonvolatile Particulate Matter Emissions of a Commercial Turbofan Engine at the Ground Level from the Use of a Sustainable Aviation Fuel Blend. *Environmental Science & Technology* 55, 14576-14585.


- 541 Durdina, L., Brem, B.T., Setyan, A., Siegerist, F., Rindlisbacher, T., Wang, J., 2017. Assessment of Particle
542 Pollution from Jetliners: from Smoke Visibility to Nanoparticle Counting. *Environmental Science &*
543 *Technology* 51, 3534-3541.
- 544 Elser, M., Brem, B.T., Durdina, L., Schönenberger, D., Siegerist, F., Fischer, A., Wang, J., 2019. Chemical
545 composition and radiative properties of nascent particulate matter emitted by an aircraft turbofan
546 burning conventional and alternative fuels. *Atmos. Chem. Phys.* 19, 6809-6820.
- 547 Fierz, M., Vernooij, M.G.C., Burtscher, H., 2007. An improved low-flow thermodenuder. *Journal of*
548 *Aerosol Science* 38, 1163-1168.
- 549 Gagné, S., Couillard, M., Gajdosechova, Z., Momenimovahed, A., Smallwood, G., Mester, Z., Thomson,
550 K., Lobo, P., Corbin, J.C., 2021. Ash-Decorated and Ash-Painted Soot from Residual and Distillate-Fuel
551 Combustion in Four Marine Engines and One Aviation Engine. *Environmental Science & Technology* 55,
552 6584-6593.
- 553 Gawron, B., Białocki, T., Janicka, A., Zawisłak, M., Górniak, A., 2020. Exhaust toxicity evaluation in a gas
554 turbine engine fueled by aviation fuel containing synthesized hydrocarbons. *Aircraft Engineering and*
555 *Aerospace Technology* 92, 60-66.
- 556 Habre, R., Zhou, H., Eckel, S.P., Enebish, T., Fruin, S., Bastain, T., Rappaport, E., Gilliland, F., 2018. Short-
557 term effects of airport-associated ultrafine particle exposure on lung function and inflammation in
558 adults with asthma. *Environment International* 118, 48-59.
- 559 Harris, S., Maricq, M.M., 2001. Signature size distributions for diesel and gasoline engine exhaust
560 particulate matter. *Journal of Aerosol Science* 32, 749-764.
- 561 Harris, S.J., 1990. Surface Growth and Soot Particle Reactivity. *Combustion Science and Technology* 72,
562 67-77.
- 563 He, R.-W., Gerlofs-Nijland, M.E., Boere, J., Fokkens, P., Leseman, D., Janssen, N.A.H., Cassee, F.R., 2020.
564 Comparative toxicity of ultrafine particles around a major airport in human bronchial epithelial (Calu-
565 3) cell model at the air-liquid interface. *Toxicology in Vitro* 68, 104950.
- 566 He, R.-W., Shirmohammadi, F., Gerlofs-Nijland, M.E., Sioutas, C., Cassee, F.R., 2018. Pro-inflammatory
567 responses to PM_{0.25} from airport and urban traffic emissions. *The Science of the total environment*
568 640-641, 997-1003.
- 569 ICAO, 2017. Environmental Protection: Vol. II Aircraft Engine Emissions, 4th edn, Annex 16 to the
570 Convention on International Civil Aviation. ICAO, Montréal.
- 571 ICAO, 2018. Environmental Protection: Vol. II Aircraft Engine Emissions, 4th edn, Annex 16 to the
572 Convention on International Civil Aviation. ICAO, Montréal.
- 573 Jaramillo, I.C., Sturrock, A., Ghiassi, H., Woller, D.J., Deering-Rice, C.E., Lighty, J.S., Paine, R., Reilly, C.,
574 Kelly, K.E., 2018. Effects of fuel components and combustion particle physicochemical properties on
575 toxicological responses of lung cells. *Journal of Environmental Science and Health, Part A* 53, 295-309.
- 576 Jeannet, N., Fierz, M., Kalberer, M., Burtscher, H., Geiser, M., 2015. Nano Aerosol Chamber for In-Vitro
577 Toxicity (NACIVT) studies. *Nanotoxicology* 9, 34-42.
- 578 Jonsdottir, H.R., Delaval, M., Leni, Z., Keller, A., Brem, B.T., Siegerist, F., Schönenberger, D., Durdina, L.,
579 Elser, M., Burtscher, H., Liati, A., Geiser, M., 2019. Non-volatile particle emissions from aircraft turbine
580 engines at ground-idle induce oxidative stress in bronchial cells. *Communications Biology* 2, 90.


- 581 Jonsdottir, H.R., Dijkman, R., 2015. Characterization of human coronaviruses on well-differentiated
582 human airway epithelial cell cultures. *Methods Mol Biol* 1282, 73-87.
- 583 Künzi, L., Krapf, M., Daher, N., Dommen, J., Jeannet, N., Schneider, S., Platt, S., Slowik, J.G., Baumlin,
584 N., Salathe, M., Prévôt, A.S.H., Kalberer, M., Strähli, C., Dübgen, L., Sioutas, C., Baltensperger, U.,
585 Geiser, M., 2015. Toxicity of aged gasoline exhaust particles to normal and diseased airway epithelia.
586 *Scientific Reports* 5, 11801.
- 587 Künzi, L., Mertes, P., Schneider, S., Jeannet, N., Menzi, C., Dommen, J., Baltensperger, U., Prévôt, A.S.H.,
588 Salathe, M., Kalberer, M., Geiser, M., 2013. Responses of lung cells to realistic exposure of primary and
589 aged carbonaceous aerosols. *Atmospheric Environment* 68, 143-150.
- 590 Lammers, A., Janssen, N.A.H., Boere, A.J.F., Berger, M., Longo, C., Vijverberg, S.J.H., Neerincx, A.H.,
591 Maitland - van der Zee, A.H., Cassee, F.R., 2020. Effects of short-term exposures to ultrafine particles
592 near an airport in healthy subjects. *Environment International* 141, 105779.
- 593 Lee, D.S., Fahey, D.W., Skowron, A., Allen, M.R., Burkhardt, U., Chen, Q., Doherty, S.J., Freeman, S.,
594 Forster, P.M., Fuglestad, J., Gettelman, A., De León, R.R., Lim, L.L., Lund, M.T., Millar, R.J., Owen, B.,
595 Penner, J.E., Pitari, G., Prather, M.J., Sausen, R., Wilcox, L.J., 2020. The contribution of global aviation
596 to anthropogenic climate forcing for 2000 to 2018. *Atmospheric Environment*, 117834.
- 597 Liati, A., Schreiber, D., Alpert, P.A., Liao, Y., Brem, B.T., Corral Arroyo, P., Hu, J., Jonsdottir, H.R.,
598 Ammann, M., Dimopoulos Eggenschwiler, P., 2019. Aircraft soot from conventional fuels and biofuels
599 during ground idle and climb-out conditions: Electron microscopy and X-ray micro-spectroscopy.
600 *Environmental Pollution* 247, 658-667.
- 601 Liati, A., Schreiber, D., Dimopoulos Eggenschwiler, P., Arroyo Rojas Dasilva, Y., 2013. Metal Particle
602 Emissions in the Exhaust Stream of Diesel Engines: An Electron Microscope Study. *Environmental*
603 *Science & Technology* 47, 14495-14501.
- 604 Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative
605 PCR and the 2⁻($\Delta\Delta C_T$) Method. *Methods* 25, 402-408.
- 606 Lobo, P., Durdina, L., Brem, B.T., Crayford, A.P., Johnson, M.P., Smallwood, G.J., Siegerist, F., Williams,
607 P.I., Black, E.A., Llamado, A., Thomson, K.A., Trueblood, M.B., Yu, Z., Hagen, D.E., Whitefield, P.D.,
608 Miake-Lye, R.C., Rindlisbacher, T., 2020. Comparison of standardized sampling and measurement
609 reference systems for aircraft engine non-volatile particulate matter emissions. *Journal of Aerosol*
610 *Science* 145, 105557.
- 611 Lobo, P., Durdina, L., Smallwood, G.J., Rindlisbacher, T., Siegerist, F., Black, E.A., Yu, Z., Mensah, A.A.,
612 Hagen, D.E., Miake-Lye, R.C., Thomson, K.A., Brem, B.T., Corbin, J.C., Abegglen, M., Sierau, B.,
613 Whitefield, P.D., Wang, J., 2015. Measurement of Aircraft Engine Non-Volatile PM Emissions: Results
614 of the Aviation-Particle Regulatory Instrumentation Demonstration Experiment (A-PRIDE) 4 Campaign.
615 *Aerosol Science and Technology* 49, 472-484.
- 616 Manisalidis, I., Stavropoulou, E., Stavropoulos, A., Bezirtzoglou, E., 2020. Environmental and Health
617 Impacts of Air Pollution: A Review. *Frontiers in Public Health* 8.
- 618 Masiol, M., Harrison, R.M., 2014. Aircraft engine exhaust emissions and other airport-related
619 contributions to ambient air pollution: A review. *Atmospheric Environment* 95, 409-455.
- 620 Mazareanu, E., 2021. Air traffic - passenger growth rates forecast 2019-2040.
621 <https://www.statista.com/statistics/269919/growth-rates-for-passenger-and-cargo-air-traffic/>.


- 622 Møller, K.L., Brauer, C., Mikkelsen, S., Loft, S., Simonsen, E.B., Koblauch, H., Bern, S.H., Alkjær, T.,
 623 Hertel, O., Becker, T., Larsen, K.H., Bonde, J.P., Thygesen, L.C., 2017. Copenhagen Airport Cohort: air
 624 pollution, manual baggage handling and health. *BMJ Open* 7, e012651.
- 625 Møller, K.L., Thygesen, L.C., Schipperijn, J., Loft, S., Bonde, J.P., Mikkelsen, S., Brauer, C., 2014.
 626 Occupational Exposure to Ultrafine Particles among Airport Employees - Combining Personal
 627 Monitoring and Global Positioning System. *PLOS ONE* 9, e106671.
- 628 Pahalagedara, L., Sharma, H., Kuo, C.-H., Dharmarathna, S., Joshi, A., Suib, S.L., Mhadeshwar, A.B.,
 629 2012. Structure and Oxidation Activity Correlations for Carbon Blacks and Diesel Soot. *Energy & Fuels*
 630 26, 6757-6764.
- 631 Paur, H.-R., Cassee, F.R., Teeguarden, J., Fissan, H., Diabate, S., Aufderheide, M., Kreyling, W.G.,
 632 Hänninen, O., Kasper, G., Riediker, M., Rothen-Rutishauser, B., Schmid, O., 2011. In-vitro cell exposure
 633 studies for the assessment of nanoparticle toxicity in the lung—A dialog between aerosol science and
 634 biology. *Journal of Aerosol Science* 42, 668-692.
- 635 Peeters, J.H.A.M., 1998. Aviation and air pollution, in: Schneider, T. (Ed.), *Studies in Environmental*
 636 *Science*. Elsevier, pp. 601-613.
- 637 Price, T., Probert, D., 1995. Environmental impacts of air traffic. *Applied Energy* 50, 133-162.
- 638 SAE International, 2018. ARP 6320 Procedure for the Continuous Sampling and Measurement of Non-
 639 Volatile Particulate Matter Emissions from Aircraft Turbine Engines. SAE International, Warrendale.
- 640 Saffaripour, M., Thomson, K.A., Smallwood, G.J., Lobo, P., 2020. A review on the morphological
 641 properties of non-volatile particulate matter emissions from aircraft turbine engines. *Journal of*
 642 *Aerosol Science* 139, 105467.
- 643 Sappok, A.G., Wong, V.W., 2006. Comparative Particulate Trap Performance Using Fischer-Tropsch and
 644 Conventional Diesel Fuels in a Modern CI Engine, ASME 2006 Internal Combustion Engine Division
 645 Spring Technical Conference, pp. 93-107.
- 646 Schindler, W., Haisch, C., Beck, H.A., Niessner, R., Jacob, E., Rothe, D., 2004. A Photoacoustic Sensor
 647 System for Time Resolved Quantification of Diesel Soot Emissions. SAE International.
- 648 Schmid, A., Sutto, Z., Schmid, N., Novak, L., Ivonnet, P., Horvath, G., Conner, G., Fregien, N., Salathe,
 649 M., 2010. Decreased soluble adenylyl cyclase activity in cystic fibrosis is related to defective apical
 650 bicarbonate exchange and affects ciliary beat frequency regulation*. *The Journal of Biological*
 651 *Chemistry* 285, 29998-30007.
- 652 Schmid, O., Stoeger, T., 2016. Surface area is the biologically most effective dose metric for acute
 653 nanoparticle toxicity in the lung. *Journal of Aerosol Science* 99, 133-143.
- 654 Stacey, B., 2019. Measurement of ultrafine particles at airports: A review. *Atmospheric Environment*
 655 198, 463-477.
- 656 Sturm, R., 2016a. Local lung deposition of ultrafine particles in healthy adults: experimental results and
 657 theoretical predictions. *Annals of translational medicine* 4, 420-420.
- 658 Sturm, R., 2016b. Total deposition of ultrafine particles in the lungs of healthy men and women:
 659 experimental and theoretical results. *Annals of translational medicine* 4, 234-234.

- 660 Vaaraslahti, K., Keskinen, J., Giechaskiel, B., Solla, A., Murtonen, T., Vesala, H., 2005. Effect of Lubricant
661 on the Formation of Heavy-Duty Diesel Exhaust Nanoparticles. *Environmental Science & Technology*
662 39, 8497-8504.
- 663 Vander Wal, R.L., Bryg, V.M., Hays, M.D., 2010. Fingerprinting soot (towards source identification):
664 Physical structure and chemical composition. *Journal of Aerosol Science* 41, 108-117.
- 665 Wing, S.E., Larson, T.V., Hudda, N., Boonyarattaphan, S., Fruin, S., Ritz, B., 2020. Preterm Birth among
666 Infants Exposed to in Utero Ultrafine Particles from Aircraft Emissions. *Environmental Health*
667 *Perspectives* 128, 047002.
- 668 Yehliu, K., Vander Wal, R.L., Boehman, A.L., 2011. A comparison of soot nanostructure obtained using
669 two high resolution transmission electron microscopy image analysis algorithms. *Carbon* 49, 4256-
670 4268.
- 671









- Exposure to aircraft nvPM causes transient toxicity in airway epithelia in-vitro
- Lower thrust levels elicit more severe biological responses.
- Differential responses of airway epithelia from donors of normal and compromised health status.

1 **Mathilde N. Delaval**: Investigation, Data curation, Formal analysis, Methodology, Validation,
2 Visualization, Writing - original draft, Writing - review & editing. **Hulda R. Jonsdottir**:
3 Conceptualization, Investigation, Data curation, Formal analysis, Methodology, Supervision,
4 Validation, Visualization, Writing - original draft, Writing - review & editing. **Zaira Leni**: Methodology,
5 Writing - review & editing. **Alejandro Keller**: Data curation, Formal analysis, Methodology, Validation,
6 Visualization, Writing - review & editing. **Benjamin T. Brem**: Data curation, Methodology, Validation,
7 Funding acquisition, Writing - review & editing. **Frithjof Siegerist**: Resources. **David Schönenberger**:
8 Resources, Methodology. **Lukas Durdina**: Investigation, Data curation, Formal analysis, Methodology,
9 Visualization, Writing - review & editing. **Miriam Elser**: Investigation, Data curation, Formal analysis,
10 Methodology, Writing - review & editing. **Matthias Salathe**: Methodology, Validation, Writing review
11 & editing. **Nathalie Baumlin**: Methodology, Validation. **Prem Lobo**: Investigation, Data curation,
12 Formal analysis, Methodology, Writing review & editing. **Heinz Burtscher**: Investigation,
13 Methodology, Writing - review & editing. **Anthi Liati**: Formal analysis, Data curation, Methodology,
14 Validation, Visualization, Writing - original draft. **Marianne Geiser**: Conceptualization, Data curation,
15 Funding acquisition, Methodology, Project administration, Supervision, Validation, Writing - original
16 draft, Writing - review & editing.

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

--