Functional design of bacterial superoxide:Quinone oxidoreductase



Abbas Abou-Hamdan, Roman Mahler, Philipp Grossenbacher, Olivier Biner, Dan Sjöstrand, Martin Lochner, Martin Högbom, Christoph von Ballmoos

PII:	S0005-2728(22)00052-4	
DOI:	https://doi.org/10.1016/j.bbabio.2022.148583	
Reference:	BBABIO 148583	
To appear in:	BBA - Bioenergetics	
Received date:	30 January 2022	
Revised date:	10 May 2022	
Accepted date:	31 May 2022	

Please cite this article as: A. Abou-Hamdan, R. Mahler, P. Grossenbacher, et al., Functional design of bacterial superoxide:Quinone oxidoreductase, *BBA - Bioenergetics* (2021), https://doi.org/10.1016/j.bbabio.2022.148583

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier B.V.

# Functional design of bacterial superoxide:quinone oxidoreductase

Abbas Abou-Hamdan<sup>+1</sup>, Roman Mahler<sup>+1</sup>, Philipp Grossenbacher<sup>2</sup>, Olivier Biner<sup>1,3</sup>, Dan Sjöstrand<sup>4</sup>, Martin Lochner<sup>2</sup>, Martin Högbom<sup>4</sup> and Christoph von Ballmoos<sup>1,\*</sup>

#### Affiliations:

<sup>1</sup> Department of Chemistry, Biochemistry and Pharmaceutical Sciences, University of Bern, 3012 Bern, Switzerland.

<sup>2</sup> Institute of Biochemistry and Molecular Medicine, University Cilleru, 3012 Bern, Switzerland

<sup>3</sup> Department of Plant and Microbial Biology, University of Zürich, 8008 Zürich, Switzerland

<sup>4</sup> Stockholm center for Biomembrane Research, Department of Biochemistry and Biophysics, Stockholm University, 10691 Stockholm, Swedon.

\*Correspondence to: christoph.vonbalin o/,s@unibe.ch

+ Contributed equally

## Abstract

The superoxide anion - molecular oxygen reduced by a single electron - is produced in large amounts by enzymatic and adventitious reactions. It can perform a range of cellular functions, including bacterial warfare and iron uptake, signalling and host immune response in eukaryotes. However, it also serves as precursor for more deleterious species such as the hydroxyl anion or peroxynitrite and defense mechanisms to neutralize superoxide are important for cellular health. In addition to the soluble proteins superoxide dismutase and superoxide reductase, recently the membrane embedded diheme cytochrome  $b_{561}$  (CybB) from *E. coli* has been proposed to act as a superoxide:quinone oxidoreductase. Here, we confirm superoxide and cellular ubiquinones or menaquine ner as natural substrates and show that quinone binding to the enzyme accelerater the eaction with superoxide. The reactivity of the substrates is in accordance with the line or determined midpoint potentials of the two *b* hemes (+48 and -23 mV / NHE). Our data subcreat that the enzyme can work near the diffusion limit in the forward direction and can also catalyse the reverse reaction efficiently under physiological conditionr. The clata is discussed in the context of described cytochrome  $b_{561}$  proteins and potential physiological roles of CybB.

## Introduction

In all aerobic organisms, oxygen plays a vital role in the conversion of reducing equivalents into the universal cellular energy currency ATP. During the last step of respiration, terminal oxidases catalyze the highly exergonic reduction of molecular oxygen to water. This reaction is coupled to the formation of a transmembrane electrochemical proton gradient (or proton motive force, *pmf*), which is harnessed by the ATP synthase to recycle ATP from ADP and inorganic phosphate. In contrast to this well controlled reaction, adventitious single electron transfer to molecular oxygen results in formation of reactive oxygen species (ROS), representing incompletely reduced oxygen variants such as superoxide  $(O_2^{-})$ , hydrogen peroxide  $(H_2O_2)$  and hydroxyl radical  $(OH^-)$ . Depending of their concentration, ROS can either be toxic or function as signalling molecules, p ayir g important roles in normal physiological and pathophysiological processes<sup>1</sup>. High and sustained concentrations of ROS are known to have deleterious effects on cellular health by oxidizing the three major biomolecule classes DNA, proteins, and lipids<sup>2</sup>. *Is* molecular oxygen preferably partitions into the lipophilic environment of the monicrane, the electron transfer reactions of the membrane embedded respiratory chain crzymes are a prominent source of ROS, e.g. at the flavin sites of complex  $I^3$  and  $II^4$  and the ubiquinol oxidation center of complex  $III^5$ . Single electron transfer to molecular o.y get yields superoxide, which shows relatively mild deleterious cellular effects, but has been shown to directly damage the metal centers of enzymes<sup>6</sup>. In eukaryotes, superovide is actively produced by the enzyme NADPH oxidase as a second messenger, but also to serve as a first line of defense against bacterial infections<sup>7-9</sup>. In prokaryotes, extrace.'un superoxide production has been observed in marine bacteria to support metal uptake<sup>1°</sup>, as well as in *Enterococcus faecalis, E. coli* and *Vibrio cholerae*<sup>11-13</sup>. The main defense mechanism is production of superoxide dismutase, first described in 1969, which greatly accelerates the disproportionation of two superoxide molecules to one molecule of hydrogen peroxide and one molecule of oxygen<sup>14,15</sup>. Some twenty years later, a second enzyme, superoxide reductase, was described in anaerobic bacteria to react specifically with superoxide, forming hydrogen peroxide using reduced rubredoxin as electron donor<sup>16</sup>.

Yet another 20 years later, in 2018, our groups described a third, membrane-embedded enzyme, encoded by the *Escherichia coli* gene *cybB* with the gene product known as

cytochrome  $b_{561}$  (CybB). In contrast to the two other enzymes, CybB catalyzes the oxidation of superoxide to oxygen and transfers the extracted electron to the oxidized quinone pool in the membrane<sup>17</sup> (Fig. 1A). Therefore, CybB is a superoxide:ubiquinone oxidoreductase or, in short, superoxide oxidase (SOO). The crystal structure and functional data suggested that in the forward reaction, superoxide binds to a patch of positively charged amino acid residues at the periplasmic surface. This triggers a transmembrane electron transfer reaction via two *b*-type hemes, and the reduction of ubiquinone bound near the cytoplasmic surface of the protein (Fig. 1A). The two hemes show a close edge-to-edge distance of 11 Å allowing a fast electron transfer<sup>18,19</sup>. When CybB was mixed with reduced ubiquinol, the reverse reaction, i.e., reduction of CybB hemes and production of superoxide was observed. The direction of catalysis under physiological conditions and the meta'sonthological conditions and the meta'sonthological the However, *mRNA* levels of *cybB* were found to be the more than 10-times higher in the exponential than in the stationary phase whilst their gunaffected by the presence of oxygen<sup>17</sup>.

Here, we thoroughly characterize detergent-solubilised superoxide:quinone oxidoreductase from *E. coli* using a range of biochemical and biophysical techniques. We provide evidence that superoxide and cellular quinone are indeed its natural substrate and measure the pH dependence of the reverse reaction. The data are then interpreted in connection to the heme mid-point potential (also determined in this study) in the absence and presence of quinone. Using a coupled ecogmatic assay, we determine that the Michaelis-Menten constant  $K_m$  for quinones in the forward direction is in the submicromolar range.

## Materials and methods

#### **Expression of CybB**

Expression of CybB was done as published previously<sup>17</sup>. Briefly, the plasmid pcybB-His encoding for the *E. coli cybB* gene with an 8xHis-tag fusion at its C-terminus was used to transform chemically competent *E. coli* BL21(DE3) pLysS cells. Transformed bacteria were grown on LB-agar plates containing 30  $\mu$ g ml<sup>-1</sup> kanamycin overnight at 37 °C. One colony was picked to inoculate a culture in 100ml LB media containing kanamycin (30  $\mu$ g ml<sup>-1</sup>). Large culture growth was conducted at 37 °C using a LeX48 system (Epiphyte 3, Canada) where conventional shaking was replaced by continuous air bubble  $\alpha$ . At a culture absorbance (OD<sub>600</sub>) between 0.9 and 1.1, expression of CybB was induced by the addition of 0.2 mM IPTG. Cells were harvested 4 hours after induction by centr fugation at 7500 x g and the pellet was resuspended in buffer (50 mM Hepes pH 7 5, 200 mM NaCl and 7.5 % glycerol).

#### Membrane preparation and protein purification

To prepare membranes, the resuspended  $pe^{-1}$  et was mixed with 1 mM PMSF, 2 mM MgCl<sub>2</sub>, some crystals of PEFA-bloc (Biomol, Germany), lysozyme and DNase I (Merck, Germany). The mixture was stirred for 30 min at RT and the homogenized cells were broken at least twice at 2000 Bar utilizing a Maxir 1a'.o High Pressure Homogenizer HPL6 (Maximator AG, Switzerland). Unbroken cells we'e removed by centrifugation at 7500 x g for 15 min and the supernatant transferred to a Berkman Ti45 ultracentrifuge rotor and centrifuged at 40000 rpm for 45 min. The memory is pellet was resuspended in the resuspension buffer and diluted up to 10 mg ml ciluted membranes were solubilized by the addition of 1 % of OGNG (or 1 % DDM) (/ matrace) in the presence of 1 mM PMSF and some crystals of PEFA-Bloc for 1 h at 4 °C. Insoluble material was cleared by ultracentrifugation for 45 min at 40000 rpm in a Beckman Ti70 rotor. The resulting supernatant was loaded onto the nickelcharged Profinity IMAC Resin (Bio-Rad, USA) and washed with 15 column volumes (CV) of wash buffer (10 mM Hepes pH 7.5, 200 mM NaCl, 7.5 % glycerol and 0.1 % OGNG) and 15 CV of wash buffer 2 (10 mM Hepes pH 7.5, 200 mM NaCl, 7.5 % glycerol, 0.1 % OGNG and 5 mM Histidine). CybB was eluted with 10CV of elution buffer (10 mM Hepes pH 7.5, 200 mM NaCl, 7.5 % glycerol, 0.1 % OGNG containing either 200 mM histidine, 100 mM EDTA or 200 mM Imidazole) and concentrated up to 0.5 ml using Amicon Ultra-15 centrifugal filters with MWCO of 50 kDa (Merck, Germany). The sample was then applied to a size exclusion

chromatography on a Superdex 200 increase 10 /300GL column equilibrated with wash buffer. Fractions containing CybB were pooled together, concentrated with a MWCO 50 kDa concentrator, and flash-frozen in liquid nitrogen.

#### **Reduction of CybB**

Reduction of CybB was monitored using a Cary 60 UV-Vis Spectrophotometer (Agilent Technologies, USA) by measuring heme absorbance changes at either 428 nm or 561 nm upon addition of various substrates in aerobic conditions. All measurements were done in 100 mM sodium phosphate pH 8, 0.1 mM diethylenetriaminepentaacetic acid (DTPA) and 0.05 % DDM. In those measurements, 0.25 to 1  $\mu$ M of CybB was used and the reaction was started by various strategies to produce superoxide. I.) Environmentally using 0.01 U xanthine oxidase (bovine milk (Sigma, USA)) in the presence of (.1,..., hypoxanthine. II.) Chemically by dissolving 7.1 mg KO<sub>2</sub> in 1 ml of anhydrous DMSC in the presence of 80mg of 18-crown-6 ether to stabilize produced superoxide under stea n of  $N_2$  as described.  $^{20}$  III.) By mixing 1 mM NADH with 10  $\mu$ M PMS<sup>21</sup>, or other reducta to such as 1 mM dithiothreitol (DTT) and 1 mM ascorbate. For control measuremer  $c_{s_1}$   $\lambda_{s_2}$   $\mu$ g ml<sup>-1</sup> of superoxide dismutase (Sigma) was used to disproportionate superoxide anion  $\mu$  the presence of 20 µg ml<sup>-1</sup> of catalase (Sigma). Full reduction of CybB was achieved and addition of sodium dithionite and used hence to calculate the percentage of relative tieme reduction. Ubiquinone analogues were either synthetized (Q<sub>1</sub>, Q<sub>2</sub> and decy biquinone) as described in the supplementary material or purchased from Sigma. Ubiquinone analogues  $(Q_1, Q_2 \text{ and decylubiquinone})$  and naphthalene analogue. (1,4-naphthoquinone and menadione, a soluble form of menaquinone) were ad led from stock solution of 50 mM quinone and 1 M DTT. Alternatively, quinones were pre-reduced with sodium borohydride NaBH<sub>4</sub>.

For the pH dependency measurements, we used the same protocol as before and measured either superoxide or  $Q_1H_2$  induced heme reduction in the presence of different buffered solutions. In the first set of experiment, we used 100 mM sodium phosphate (pH 7 or 8) or 100 mM CHES and 100 mM sodium phosphate dibasic (pH 9 or 10) containing 20 mM KCl, 200 mM NaCl, 0.1 mM DPTA, 0.05 % DDM. In the second set, we used 100 mM sodium phosphate (for pH ranging from 6 to 8.5) or 20 mM Tris-HCl (for pH ranging from 9 to 10) containing each 20 mM KCl, 200mM NaCl, 0.1 mM DPTA, 0.05 % DDM. In both experiments, the percentage of relative heme reduction compared to a full dithionite reduction of CybB

6

was calculated. Anaerobic measurements were performed after removal of dioxygen using a combined system of argon blowing in the presence of 53  $\mu$ g ml<sup>-1</sup> catalase, 26  $\mu$ g ml<sup>-1</sup> glucose oxidase and 5 mM glucose mixture (all purchased from Sigma).

#### **Oxidation of reduced CybB**

Oxidation of reduced CybB was monitored under anaerobic conditions. Anaerobicity was achieved as described above (section: *Reduction of CybB*). 15 min after argon blowing, 0.25  $\mu$ M of CybB was fully pre-reduced using a tiny amount of dithionite (6.6  $\mu$ M in 1.5 ml of 100 mM sodium phosphate pH 8, 20 mM KCl, 200 mM NaCl, 0.05 % DDM) before addition of 10  $\mu$ M of diverse quinone (ubiquinone and naphthalene analogue) to oxidize it. A full oxidation of reduced CybB was achieved by adding 10  $\mu$ M of potentiation ferricyanide and used to calculate the percentage of relative heme oxidation.

#### Superoxide assay and determination of Michaelis-N. ant an constant

We used the previously established WST-1 (V:ter soluble Tetrazolium dye) assay<sup>22</sup> and adapted it<sup>17</sup> to determine either the Mic<sup>1</sup> is Nenten constant for ubiquinone of CybB or to measure superoxide production by the chzyme. In both assays, formazan formation upon reduction of WST-1 by superoxide wa: monitored at 438 nm in assay buffer (100 mM sodium phosphate pH 8, 0.1 mM DT<sup>2</sup>*L*, 0.1 mM HPX, and 0.05 % DDM) containing 50  $\mu$ M WST-1, 20  $\mu$ g ml<sup>-1</sup> catalase and 60 nM *bo3* oxidase. Measurements were done either individually in a cuvette using Cary 60 UV-Vis spectrophotometer or in a 96-well plate in SpectraMax ABS Plus Microplate Reader (Molecular Devices, USA). Ubiquinone Q<sub>1</sub> or Q<sub>2</sub> were added to the previous mixture to measure the overall activity of CybB. The Michaelis-Menten constant (*K*<sub>m</sub>) for quinone was calculated using a constant amount of CybB (50 nM) while varying the quinone concentration between 0 and 100  $\mu$ M. To measure superoxide production and determine the apparent constant affinity for quinol, the reaction was initiated by addition of 50 nM of CybB to the assay buffer containing different concentration of quinol (0 - 100  $\mu$ M) in the absence of *bo*<sub>3</sub> oxidase and catalase.

The second order rate (catalytic efficiency) of CybB was calculated based on the known value for SOD ( $2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ )<sup>23</sup>. Briefly, the concentration for both enzymes SOD and CybB was titrated using the former assay to reduce the formazan formation rate by 50 %. The determined concentrations were used to calculate the catalytic efficiency of CybB.

#### **Potentiometric redox titrations**

Redox titrations were performed anaerobically at room temperature in a Spectroelectrochemical cell (Prosense, Netherlands), by following heme absorption changes upon a gradual addition of dithionite solution as electron source, instead of electrochemistry as in<sup>24</sup> using a UV-Vis spectrophotometer.

Throughout the measurement, anaerobicity was maintained by a continuous flow of argon. Prior to measuring, 10  $\mu$ M CybB, in the absence or presence of 20  $\mu$ M Q<sub>1</sub>, and 5  $\mu$ M mediators were incubated together in 100 mM sodium phosphate pH 7, 0.1 mM DTPA, 0.1 mM HPX, and 0.05 % DDM and placed in a Thunberg currette and made anaerobic by alternative cycles of  $N_2$  vacuum. The following mediators we clusted: benzyl viologen (-350 mV), sodium anthraquinone 2-sulfonate (-225 mV), 2-h, droxy 1,4-naphtoquinone (-145 mV), resorufin (-51 mV), methylene blue (+11 r IV), phenazine ethosulfate (+55 mV), phenazine methosulfate (+80 mV), 2,6-Dichlc ophenol-indophenol (+217 mV) and tetramethyl-p-phenylendediamine (+276 mV). K dc x potentials were measured using an Ag / AgCl 3 M KCl reference electrode associated to a platinum counter electrode (Pt) (Prosense, Netherlands). The redox potential was noted after equilibration, and an optical spectrum was recorded between 405 500 nm. The measured absorbance of heme reduction at 561nm was plotted against the measured redox potentials and curves were fitted using the Nernst equation with n=2 4ue to two successive one electron reductions corresponding to the reduction of both home. All midpoint potentials referenced here were adjusted by the addition of +210 mV rela ive to the normal hydrogen electrode (vs NHE) at 25 °C.

## Results

#### Enzyme production and purification

The gene sequence of *E. coli cybB*, equipped with a C-terminal octahistidine tag was cloned into a pET-28b vector and expressed in BL21(DE3) pLysS. The protein is well expressed yielding cells and inverted membranes with a distinct pink color. Here, we further optimized the purification of CybB based on our published protocol<sup>17</sup>. The membrane fraction was solubilized with either 1 % OGNG or 1 % DDM and purified via Ni-IDA chromatography. The protein was eluted using either imidazole, histidine or EDTA, concentrated and assessed using size-exclusion chromatography (Superdex 200 Increase, 10 /300GL, 500 µl injected), using either 0.1 % OGNG or 0.1 % DDM. The results displated in supplementary Figure S1 show a major impact of both the detergent and the Ni-DA eluant used. While both detergents extracted the protein similarly from the memorane, only extractions with OGNG produced a monomeric peak (~14.7 ml), as judged them the size exclusion chromatograms (Fig. S1 A and E). In addition to this peak (Fig S1A), two further discrete elution peaks were observed, indicating aggregate forms of the protein. If the protein was eluted with either 100 mM EDTA or 200 mM histidine (Fig. 1 A and E) instead of imidazole, a homogenous preparation with a single monomeric peak was obtained. In the following, 1 % OGNG and 0.1% OGNG/ 200 mM histidine was a for solubilization and elution, respectively.

#### Spectroscopic properties and reaction with cellular reductants

The crystal structure of CyhB (F.?B ID 5OC0) shows a highly distorted 4-helix bundle with the *b*-hemes fully coordinated by histidines (Fig. 1B)<sup>17</sup>. The protein is almost entirely in contact with the lipid bilayer with no peripheral domains. Heme 2 is partly exposed towards the periplasm by a narrow cavity that contains a variety of positively charged residues that are proposed to guide the negatively charged superoxide anion in proximity of the heme to allow electron transfer. Heme 1 is located towards the cytoplasm and buried in the membrane with no polar access route, supporting the idea of a binding site for a hydrophobic compound such as ubiquinone. The two hemes are rapidly reduced by dithionite and the reduced spectrum shows a characteristic Soret peak at 428 nm as well as  $\alpha$  and  $\beta$  peaks at 530 and 561 nm as described earlier. Lundgren et al<sup>17</sup> found that detergent solubilized CybB is rapidly, but only partly reduced (~50 %) by the DTT-reduced short-chain ubiquinone analogue Q<sub>1</sub>. Only partial reduction by superoxide was also observed if xanthine

oxidase was used as a superoxide producing system. No or only very slow reduction is observed with reducing agents such as ascorbate, DTT and NADH (Fig. 1C). We therefore aimed to better understand the reduction behaviour of detergent solubilized CybB in a series of kinetic and steady state experiments.



*Figure 1: CybB structure and the corresponding substrates.* A) Proposed reaction mechanism of CybB. Upon oxidation of superoxide anion at the heme 2, electrons are transferred through the two hemes to reduce the membrane quinone pool with protons being sequestrated from the cytoplasm B) Distorted helical arrangement of CybB (produced by ECSF Chimera X using PDB entry 50C0). The 4 alpha helices are shown in grey, heme 2 and heme 1 in dark red and ubiquinone UQ<sub>8</sub> in cyan. Visualization of ubiquinone Q<sub>8</sub> (PubChem CID 5283546) as suggested by molecular docking to the crystal structure (pdb 50C0) using AutoDock Vina<sup>25</sup>. C) Kinetic traces monitoring the CybB reduction at 428 nm upon addition of different substrates: 1 mM sodium dithionite (blue), 1 mM dithiothreitol (DTT) (red), 1 mM DTT / 1 uM ubiquinone Q<sub>2</sub> (orange), 0.1 mM Hypoxanthine / 7.6 mU Xanthine oxidase (HPX / XO) (black), 1 mM Ascorbate (green) and 1 mM NADH (purple). Measurements were done in 100 mM sodium phosphate pH 8, 0.1 mM DTPA, 0.05 % DDM. When xanthine oxidase is used, 0.1 mM of hypoxanthine was added to the previous buffer. 8.8  $\mu$ g ml<sup>-1</sup> of SOD was present before addition of substrates (brown).

#### Reaction of CybB with superoxide produced by KO2 and NADH / PMS

Superoxide in aqueous solution has a half-life that is strongly pH dependent, as the spontaneous disproportionation to hydrogen peroxide and oxygen occurs via the protonated O<sub>2</sub>H form. Consequently, superoxide is more stable at higher pH values, i.e., t½~1 s at 10  $\mu$ M and pH 8<sup>26</sup>. If superoxide was indeed the reducing agent, the reaction of superoxide with CybB should be more complete at higher pH values. To test this, KO<sub>2</sub> dissolved in anhydrous DMSO stabilized with 18-crown-6 ether was used to produce a stable superoxide stock solution. From this solution, a small amount was added to an aqueous solution containing detergent solubilized CybB (1 µM) and the reduction of CybB was followed spectrophotometrically (mixing time ~2 s) at 428 in at different pH values (Fig. 2A). Both the rate and the level of reduction increased with a preasing pH correlating with a prolonged presence of superoxide at higher pH values (Fig. 2 A, B). Essentially full reduction was observed at pH values  $\geq 10$ . If the reaction at pH 10 vas supplemented with SOD (8.8 µg ml<sup>-1</sup>), reduction of CybB was suppressed, indicating the superoxide specificity of the system. The enzyme showed an unaltered UV-VIS sp<sup>2</sup> ctrum at pH 10 for prolonged time suggesting that the protein is stable also at this pl. (S) pplementary. Fig. S2). A further system that has been described to produce superoxide is NADH mixed with the electron mediator phenazine methosulfate  $(PMS)^{21}$ . Indeed, when into mixture was incubated with detergent solubilized CybB, rapid reduction of the nen.2s was observed and the reaction was completely suppressed in the presence of SOD (Fig. 2C).



**Figure 2:** CybB reduction by varioe. sources of superoxide. A) Kinetic traces monitoring CybB reduction at 428 nm upon addition of 5  $\mu$ M superoxide from a 2 solution at different pH values. Full reduction of CybB was achieved by addition of sodium dithionite. B) percentage of the relative level of reduction after 20 s incubation with KO<sub>2</sub> (Data from Fig. 2A). C) Kinetic traces monitoring CybC reduction in presence of various amounts of NADH and PMS at pH 8. A baseline was measured in the presence of NAD I (and SOD (8.8  $\mu$ g ml<sup>-1</sup>) for the control) and the reaction was started by addition of PMS after ~2 s.

#### Reaction of CybB with reduced ubiquinol

To investigate the reverse reaction, we mixed detergent solubilized CybB (1  $\mu$ M) with ubiquinol QH<sub>2</sub>, either prereduced or reduced *in situ* by 1 mM DTT 1 min prior to addition of the enzyme. CybB reduction by DTT alone is very slow and can be ignored (Fig. 3A). As depicted in Figure 3A, rapid reduction is observed if solubilized CybB is mixed with 1 mM DTT and 10  $\mu$ M ubiquinone Q<sub>2</sub> (purple trace). If 1  $\mu$ M Q<sub>2</sub> is used, the reaction is slightly slower, but the same level of reduction is reached. At 0.1  $\mu$ M Q<sub>2</sub>, the reaction was slowed

down >10 times, and the maximum reduction was not reached within the measuring period. Next, we varied the pH of the reaction mixture, keeping DTT /  $Q_1$  constant at 1 mM / 10  $\mu$ M. As depicted in Fig. 3B, reduction levels of CybB were higher at alkaline pH values, while essentially no heme reduction was observed at pH 6 or below. We had previously observed that upon the reaction with quinol, the enzyme consumes oxygen and produces superoxide, as detected with WST-1<sup>17</sup> (see also Fig. 6E). Consequently, under aerobic conditions, the enzyme is turning over, and the observed reduction level reflects steady-state catalysis. Therefore, the experiment was repeated under anaerobic conditions and while a similar trend was found, reduction levels were increased at all physicalues compared to aerobic conditions (Fig. 3B). The difference between aerobic and anarobic reduction was most pronounced at pH values <7.5. A plausible explanation fc, this behaviour is that the level of reduction correlates with the midpoint potential of the 'wo reactants, the b hemes of CybB and ubiquinone. However, only the latter is strongly, H cependent (~-60 mV per pH unit), as two protons participate in the reaction<sup>27</sup>. A higher pH value thus corresponds to a lower midpoint potential, thus increasing the reducing power, while the midpoint potentials of protein embedded hemes are expected or leless sensitive to pH changes<sup>28</sup>.



**Figure 3: Reduction of CybB with short chain ubiquinone A)** Kinetic traces monitoring CybB reduction at 428 nm upon addition of DTT alone or premixed with different amounts of ubiquinone  $Q_2$  at pH 8. **B)** percentage of heme reduction measured at 561 nm after 20 s incubation of CybB with 1 mM DTT / 10  $\mu$ M  $Q_1$  at different pH values under aerobic and anaerobic conditions.

#### Redox titration of CybB

In 1986<sup>29</sup>, the midpoint potential of the two hemes was determined with the purified enzyme and a single value of ~20 mV was obtained at pH 7.4. However, it is unlikely that both hemes have an identical mid-point potential. If the forward reaction was favourable, a lower midpoint potential is expected for the periplasmic heme 2 compared to the cytoplasmic heme 1 to support electron transfer from superoxide via heme 2 and heme 1 to ubiquinone (Fig. 1A). We therefore performed a redox titration from the oxidized to the reduced state with dithionite under anaerobic conditions in the presence of a set of electron mediators and followed the potential with a voltme er and the reduction state spectrophotometrically at 561 nm. Several titrations were done in the absence and presence of ubiquinone Q<sub>1</sub>, and the obtained traces, acpicted in Figure 4A, were fitted for two hemes. The two waves have similar amplitudes, erch of them reflecting ~50 % of the relative reduction signal as expected for two hences with similar absorption properties. In the absence of ubiquinone, midpoint potent. Is or  $-23 \pm 12$  mV and  $+ 48.5 \pm 12.5$  mV NHE were determined, close to the reported values, and we assigned them to heme 2 and heme 1, respectively (Figure 1B). If ubiquinone  $Q_1$  is present, the two potentials are shifted, from -23 mV to - 8 mV for heme 2 ard num 48.5 to 100 mV for heme 1. Importantly, the difference in midpoint potent als is increased (108 mV), suggesting that binding of ubiquinone thermodynamically incilitates electron transfer from heme 2 to heme 1 (Fig. 4B). In addition, the increase on the potential at heme 2 facilitates reduction by superoxide, while the potential at ten - 1 (100 mV) is still fully compatible with quinone reduction. From these measurements, it can be speculated that binding of quinone to the oxidized enzyme could influence the reaction of the enzyme with superoxide. We therefore incubated CybB with stoichiometric amounts of ubiquinone Q<sub>1</sub> before superoxide production was initiated by the addition of xanthine oxidase. As depicted in Figure 4C, the observed reaction rate of superoxide with the quinone-preincubated enzyme increased dramatically (red and blue trace). If guinone was added in a 20-fold excess the effect was even more pronounced, indicating saturation of the binding site (green trace). The reduction kinetics in presence of ubiquinone was biphasic, with a fast first phase and a slower second phase. We speculated that the slow phase originates from reaction of ubiquinol that has been formed either by the adventitious reaction of superoxide with guinone or by the reaction of superoxide with

CybB. To test this hypothesis, purified ubiquinol  $bo_3$  oxidase was added to keep the ubiquinone pool oxidised and thus prevent reduction of CybB by ubiquinol. As observed in Fig. 4C (purple trace), the addition of  $bo_3$  oxidase resulted in a monophasic reduction of CybB with an identical first phase as observed in the absence of  $bo_3$  oxidase (red trace), but lacking the slow phase, indicating that reduction occurred via superoxide only.



**Figure 4: Heme redox potentials. A)** Midpoint redox potential titrations curves of 10  $\mu$ M CybB in the absence (black) and in the presence (red) of 20  $\mu$ M Q<sub>1</sub>. Redox potential measurements were performed from the oxidized to the reduced state anaerobically in 100 mM sodium phosphate pH 7, 0.1 mM DTPA, 0.1 mM HPX and 0.05 % DDM and redox potentials are expressed in mV against NHE. B) Energy diagram of potentials calculated from fits of Fig. 4A. Shown are the values for the low (E<sub>2</sub>) and high potential heme (E<sub>1</sub>) and their energy difference( $\Delta E_1 E_2$ ) in the presence and absence of quinone. **C)** Kinetic traces showing reduction of 0.5  $\mu$ M of CybB by superoxide anion produced by xanthine oxidase in the absence and in presence of an equimolar (0.5  $\mu$ M) or an excess amount (10  $\mu$ M) of ubiquinone Q<sub>1</sub> and 60 nM of bo<sub>3</sub> oxidase. Measurements were done in 100 mM sodium phosphate pH 8, 0.1 mM DTPA, 0.1 mM HPX and 0.05 % DDM.

#### Both ubiquinone and menaquinone are substrates of CybB

In contrast to the reverse reaction, the forward reaction is more complicated to monitor due to the short half-life of superoxide and its hard-to-control production. We have therefore simplified the system and pre-reduced CybB under anaerobic conditions with stoichiometric amounts of dithionite, until ~90-95 % of the enzyme is reduced.



**Figure 5: Ubiquinone and menaquinone as substrates for CybB. A)** Kinetic traces of CybB oxidation as observed at 428 nm. CybB was reduced, under anaerobic conditions, with small amounts of dithionite (6.6  $\mu$ M) until >90 % reduction was obtained, after which 10  $\mu$ M of different quinones was added. After the reaction was complete, CybB was fully reduced by addition of 3.3 mM dithionite. **B)** Summary of the percentage of reduced heme oxidation, including the data from addition of air-saturated water (not included in Fig. 1A). **C)** Kinetic traces of CybB reduction, under aerobic conditions, as observed at 428 nm. Different quinols were added at 10  $\mu$ M concentration. **D)** Summary of the results obtained from Fig. 5C. (Note that the reference sample

(Dithionite) has no trace in 5C). The bars in figures 5B and 5D have the same color as the respective oxidized (5A) or reduced (5C) quinone traces.

Addition of ubiquinone 10  $\mu$ M Q<sub>1</sub> showed a rapid oxidation of CybB, indicating that ubiquinone is a substrate (Fig. 5A, red trace). A similar behavior was found with other shortchain ubiquinones such as Q<sub>2</sub> and decylubiquinone. Next, we tested the two menaquinone analogues menadione and 1,4-naphtoquinone. Reaction with 1,4-naphtoquinone proceeded also quickly, but enzyme oxidation was less complete, while the reaction with menadione was >10 slower and less enzyme was oxidized. A summary of the different experiments is found in Figure 5B, showing that reaction with ubiquinones reached almost complete CybB oxidation, in reference to the oxidation by ferricyanide. Addition of air saturated water also led to oxidation of CybB, indicating that the reduced enzyme car deliver electrons to oxygen in the absence of quinones.

Next, we compared the ability of the reduced form of these different quinones to reduce CybB under aerobic conditions, i.e., the reverse maction. When using ubiquinol  $Q_1H_2$  at pH 8, ~60 % of reduction of CybB was observed as discussed above (Fig. 5C).  $Q_2H_2$  and decylubiquinol showed similar kinetics t ut an overall lower level of reduction. In contrast to the forward reaction, incubation with the reduced menaquinol analogues led to almost complete reduction of the enzyme ( $t_{12}$  5 C, D). Hence, the observed data correlate well with our hypothesis that the  $t_{12}Wect$  midpoint potential of menaquinones (compared to ubiquinones) lead to a more complete reduction of CybB.

# Measurement of steady and e activity of the forward reaction and determination of quinone affinity

Having the two most likely substrates identified, we aimed to develop a turnover assay that reports the actual protein function, converting superoxide to reduced quinol. However, it is not trivial to directly measure quinol formation (see discussion). We therefore adapted a superoxide dismutase assay using WST-1, a water-soluble indicator for superoxide, which after reaction with superoxide is converted into a stable formazan that strongly absorbs at 438 nm<sup>17</sup>. As a constant superoxide producing system, xanthine oxidase and hypoxanthine was employed, which in phosphate buffer has been reported to produce ~50 % superoxide and 50 % hydrogen peroxide<sup>30,31</sup>. Although neither CybB nor WST-1 has been shown to react with hydrogen peroxide, catalase is included in the assay buffer to avoid accumulation of

hydrogen peroxide. Consequently, the linear increase observed at 438 nm represents the continuous conversion of WST-1 by xanthine oxidase produced superoxide into the stable formazan (Fig. 6B, blue trace). Winterbourn and others have previously used this assay to estimate SOD content of tissue extracts, as any present SOD competes with WST-1 for superoxide leading to a decreased formazan formation<sup>22</sup>. We adapted this idea and added the required components for CybB to compete with WST-1 for superoxide. If CybB (50 nM) and  $Q_2$  (10  $\mu$ M) were present (orange trace on Fig. 6), the initial signal was a bit lower, but quickly regained the initial slope. We interpreted this result as competition between CybB and WST-1 for superoxide that stops after enough ubiquinone has been converted to ubiquinol. We therefore added purified *E. coli bo*<sub>3</sub> quinol oxi  $\mu$  so to regenerate the quinone pool, which led to a continuous suppression of formaza' p. oduction (red trace on Fig. 6). Importantly, if  $bo_3$  and  $Q_2$  were present in the absence of CybB, no suppression was observed (green trace on Fig. 6). This indicates that he pontaneous reaction of superoxide with ubiquinone to semiubiquinone - accompanied by the disproportionation to quinone and quinol (that can be consumed by the vo<sub>3</sub> oxidase) - is not sufficient to explain the observed suppression. The overall scheme of the assay with all components is depicted in Figure 6A.

We used this approach to estimate both the second order rate constant of the reaction and the Michaelis-Menten constant  $K_{-}$  for ubiquinone. First, in the presence of all components including 10  $\mu$ M Q<sub>2</sub> and  $bo_3$  criticase, different amounts of SOD and CybB were added and the decrease in formazar regretation was followed. As depicted in Figure 6C, about 10-times more CybB than SOD wes required to obtain similar quenching values. Taking the previously determined second order rate constant of 2 x 10<sup>9</sup> M<sup>-1</sup> s<sup>-1</sup> for SOD, a rate constant in the range of 1 to 3 x 10<sup>8</sup> M<sup>-1</sup> s<sup>-1</sup> can be estimated for CybB (see table 1). To measure the Michaelis-Menten constant  $K_m$  of the reaction for ubiquinone, all components except ubiquinone were mixed, and a steady formazan formation was monitored for 30 s, before different amount of Q<sub>2</sub> were added from stock solutions, leading to decreased levels of formazan formation (Figure 6D). Using Michaelis-Menten kinetics, a  $K_m$  of 390 ± 40 nM was obtained. We determined very similar  $K_m$  values by either subsequent addition of increasing amounts of quinone as well as with different quinone concentrations measured in parallel in a 96-well plate.

Finally, we also determined a Michaelis-Menten constant  $K_m$  for the reverse reaction. To this end, superoxide formation by CybB was monitored following formazan formation from WST-1 (Fig. 6E and supplementary Fig. S3) after addition of NaBH<sub>4</sub>-prereduced ubiquinol. As depicted in Figure 6F, if initial rates were plotted against quinol concentrations, an apparent  $K_m$  of ~10 µM was obtained, thus being ~25 higher than for the forward reaction (see table 1).



**Figure 6:** Superoxide assay and Michaelis-Menten constant  $K_m$  determination. A) Cartoon depicting the coupled assay to follow competition (COMP) between CybB and WST-1 for superoxide. B) Kinetic traces monitoring formazan formation at 438 nm. The different conditions are indicated in the legend. C) Titration of different amounts of purified CybB and purified SOD under otherwise identical experimental conditions (0.01 U xanthine oxidase, 10  $\mu$ M Q<sub>2</sub>, 60 nM bo<sub>3</sub> oxidase). Relative decrease in formazan rate formation (inhibition of WST-1 reduction %) upon addition of either SOD (red trace) or CybB

(black trace). **D)** Determination of the apparent  $K_m$  of ubiquinone  $Q_2$  in the forward reaction. Relative decrease in formazan rate formation upon addition of different amounts of  $Q_2$ . **E)** Cartoon depicting the assay scheme to determine reverse activity, i.e., formation of superoxide by quinol. **F)** Determination of apparent  $K_m$  of ubiquinol  $Q_2H_2$  in the reverse reaction. Initial slopes of formazan formation ( $\Delta A_{438nm} \times s^{-1} (\times 10^3)$ ) upon addition of different amounts of prereduced ubiquinol  $Q_2H_2$  (as depicted in Supplementary Fig. S3).

### Discussion

#### Architecture of a superoxide: quinone oxidoreductase

The enzyme described here, CybB or cytochrome b<sub>561</sub> from *E. coli* was proposed to act as a membrane bound superoxide:ubiquinone oxidoreductase<sup>17</sup>. It is only the third enzyme described, after SOD and SOR, to specifically react with super pxide. However, in contrast to the two other enzymes that contain a catalytic dimetal center, CybB is a hemoprotein containing two *b* hemes. Superoxide produced in biological systems is either the product of an enzymatic or adventitious reaction of an electron with molecular oxygen. It has a standard midpoint potential of -160 mV, which h, we er is expected to be much higher under biological conditions as the  $O_2 / O_2^{-1}$  ratio much larger than 1 (e.g., at a ratio of  $O_2 / O_2^{-1}$  $O_2^{-}$  = 1,000 the potential would increase to ~+42 mV). Superoxide spontaneously disproportionates into oxygen and hyd ogen peroxide, which are more stable products. In principle, superoxide can act as a mild reductant, however, it is better known as an oxidant oxidizing lipids or enzyme metal certer.<sup>6</sup> Superoxide reacts with many organic molecules in solution, including ubiquinone. We molecules of the product, ubisemiquinone, have been shown to disproportionate in ubiquinone and ubiquinol<sup>32</sup>. However, this spontaneous reaction is not expected conficution biological systems, as superoxide is considered membrane impermet ble due its negative charge, while ubiquinones are tightly embedded in the membrane. Supproxide production can also be envisioned to happen at the membrane surface as oxygen preferentially partitions into the membrane, and quinone binding sites of respiratory enzymes have been described as points of superoxide formation<sup>3,33-35</sup>. In a similar scenario, water-soluble ascorbate and membrane-bound ubiquinone are unable to react in cells, even though they have been shown to rapidly react in solution<sup>36</sup>. The proposed design of a membrane embedded superoxide:quinone oxidoreductase therefore makes sense if electron transfer between the two otherwise separated substrates should happen. The orientation of CybB in the membrane predicts that reduction occurs from the periplasm via the hemes to the quinone binding site located on the cytoplasmic side. The idea of a cytoplasmic quinone binding site is consistent with the

glycerol found in the crystal structure and docking studies with ubiquinones<sup>17</sup>. The binding of superoxide from the periplasmic side is supported by an easily accessible heme from the protein surface surrounded by positively charged amino acids. Such a positive patch would attract, and guide, negatively charged superoxide anion radical O2<sup>-</sup>, creating an entry path for superoxide. Similar positive patches have also been found in the human Cu,Zn-SOD<sup>37</sup>, Alvinella pompejana Cu,Zn-SOD<sup>38</sup> or the superoxide reductase (SOR) of Desulfoarculus *baarsii* in complex with Fe(CN)<sub>6</sub> <sup>39,40</sup>. However, the strongest indication for a specific interaction of superoxide with CybB is the recent structure of the transmembrane domain of the NADPH oxidase (Nox5) from Cylindrospermum stagnale<sup>41</sup>. The protein consists of a transmembrane domain formed by six helices, with an overall similar fold as CybB, harboring two *b* hemes in close distance. The protein procrutes into the cytoplasm with an NADPH binding domain containing FAD as cofactor. This topology allows the enzyme to use cellular NADPH to produce superoxide and hydrogen peroxide on the extracellular side to act as chemical weapon in cellular warfare to combat bacterial infections. Notably, the two proteins, Nox5 and CybB, show a very similar electrostatic environment around the periplasmic heme, which is surrounded by positively charged amino acids creating an easy access route for negatively charged species in the periplasmic compartment (see Fig. 7). In particular, it has been proposed t'a'. rg256 present in this cavity could electrostatically promote the catalytic production on superoxide'. The overall reaction of Nox5, oxidation of NADPH and reduction of oxygen to superoxide is thus very similar to the reverse reaction of CybB, ubiquinol oxidation and uperoxide formation, and the structural arrangement of the two enzymes, especially with respect to the exposed heme and the positive patch, is also strikingly similar.



**Figure 7: Comparison of the electrostatic potential map**. The charges distribution around the exposed edge of heme group (green) of CybB (pdb 50C0) and Nox5 from Cylindrospermum stagnale (pdb 50C0),  $\frac{1}{2}$ , w a similar positive surface potential (blue) that is presumed to funnel the superoxide anion to the heme edge. Imc ges were produced in Pymol.

#### **Reaction of CybB with its substrates**

A short description of enzyme purification is included in the results section for a reason. Even though the purified CybB was very pure on SDS-PAGE gel immediately after purification via Ni-IDA resin and the endeed dithionite spectra appeared identical to those after gel filtration, functional characterization revealed different kinetics and levels of heme reduction in different enzyme preparations when reduced with DTT/Q<sub>1</sub> or superoxide. We therefore performed the classcribed experiments by varying detergent and IMAC elution methods, until we found conditions that yielded protein with reproducible properties. Possible reasons for the inconsistent behaviour in the other preparations are different levels of delipidation and protein aggregation.

The experiments presented here provide strong evidence that superoxide and both ubiquinone and menaquinone are natural substrates of CybB. This quinone substrate diversity is not surprising, as all terminal *E. coli* oxidases (*bo*<sub>3</sub>, *bdI* and *bdII*) have been shown to accept either UQ<sub>8</sub>, DMQ<sub>8</sub> or MQ<sub>8</sub><sup>42</sup>, as well as *E. coli* nitrate reductase (NarGHI) during nitrate respiration<sup>43-45</sup>. We show that CybB is readily reduced by short chain ubiquinols in a strongly pH dependent manner. The standard midpoint potential of ubiquinone / ubiquinol is ~70 mV at pH 7<sup>46</sup>, but decreases by – 60 mV per pH unit, rendering it a stronger reductant

at higher pH values in accordance with our data. Menaquinols, that have a lower midpoint potential (~-100 mV) reduce the enzyme more completely than ubiquinol. The opposite picture is observed, if the prereduced enzyme is mixed with the oxidized form of the different quinone analogues under anaerobic conditions. While incubation with oxygen and ubiquinone led to a complete oxidation comparable to ferricyanide, incubation with the lower potential menaquinone led only to a partial oxidation of the enzyme.

The specific reaction of CybB with superoxide is more complex to analyse because of difficulties to produce sufficient concentrations due to the short lifetime of superoxide. Here, we have used three well-established methods to provide superoxide in aqueous solutions, whilst superoxide specificity was tested by signal suppression after the addition of SOD<sup>22,47,48</sup>. First, superoxide was produced enzymatically by xanthine oxidase, which is routinely used to investigate SODs. However, it is nearly impossible to quantify the superoxide directly, as the percentage of superoxide, oduced per hypoxanthine molecule varies strongly and depends on the buffer (or,d'tions and on the enzyme preparation itself<sup>31</sup>. Second, we used inorganic potusium dioxide mixed with a crown-ether that produces a stable superoxide solution in organic solvents, e.g., DMSO or acetonitrile and finally we used NADH and phenazine methosulfate (PMS), which produces superoxide in aqueous solution<sup>21</sup>. All three systems rapidly reduced CybB and, in all cases, addition of SOD suppressed CybB reduction. To estimate the unspecific reaction of superoxide with other heme-containing enzymes,  $v_{1}$  incubated purified  $bo_{3}$  oxidase with xanthine oxidase/ hypoxanthine and found essentially no heme b reduction over the time course of CybB reduction (Supplementary Fig. S4). In the case of KO<sub>2</sub>, we analysed the pH dependency of the reaction and found that CybB can be fully reduced at higher pH values. Since the redox potential of the oxygen / superoxide couple is not proton dependent, we attribute this observation to the increased stability of the superoxide anion (and hence a higher substrate concentration). It is important to point out that all tested low redox potential substrates such as NADH (-340 mV), DTT (-330 mV), glutathione (-240 mV), lactate (-160 mV) or hypoxanthine (-400 mV) were unable to reduce the enzyme at pH 8. Earlier reports that NADH and lactate reduce CybB in membrane vesicles are likely due to adventitious superoxide production during respiration by NADH and lactate dehydrogenases, respectively<sup>49</sup>. A notable exception is ascorbic acid (+60 mV), described as natural substrate

for many other cytochrome  $b_{561}$  proteins, which reduced CybB of *E. coli* slowly and in a pH dependent fashion (no reduction below pH 7), likely due to its decreased potential at higher pH values. Interestingly, in cytochrome  $b_{561}$  of secretory vesicles that regenerate internal ascorbate from semidehydroascorbate via transmembrane electron transfer, the reduction of the enzyme is not pH sensitive between pH 6 and 8, indicating that a different mode of action with ascorbate is utilised in this enzyme compared to CybB<sup>50</sup>.

#### Midpoint potential of CybB hemes

We determined the midpoint potential of the two hemes by titrating dithionite to an anaerobic solution of CybB in the presence of electron mediator. While Murakami et al<sup>29</sup> found a single potential at +20 mV in a similar experiment, we were able to fit the titration to two waves with ~50 % signal amplitude each, indicating that the two hemes have different potentials, i.e. -23 mV and +48.5 mV, assigned to periplasmic side faced heme 2 and cytoplasmic faced heme 1, respectively (Figure 1A). These values are close to the ones reported by Murakami and well within the range found for other diheme b enzymes, i.e. heme b in E. coli bo<sub>3</sub> oxidase and nitrate reductase have been shown to be +50 mV<sup>51</sup> and +20 mV<sup>52</sup>, respectively. For previously described proteins denoted as cytochrome  $b_{561}$ , typically slightly higher potentials  $h_{2}$ , been found. In cytochrome  $b_{561}$  from bovine adrenal chromaffin vesicles, the redox price tials of high and low potential hemes were determined to be +150 and +60 mV, respectively<sup>53</sup>. A value of +80 mV ± 30 mV has been reported for the human duodenal cytochrcme  $\dot{\nu}_{561}$  that catalyse transmembrane electron transfer from external ascorbate to in rav, scular Fe<sup>3+ 54</sup>. In plants, the reported midpoint redox potential cytochrome  $b_{561}$  proteins range between +80 to +190 mV for the high potential heme and between -20 to +60 mV for the low potential heme, in accordance with the potentials of the proposed substrate couple dehydroascorbate / ascorbate (+60 mV)<sup>55</sup>. Potential measurements in the presence of 20  $\mu$ M ubiquinone Q<sub>1</sub> primarily shifted the midpoint potential of heme 1 to higher values, increasing the potential difference. We thus hypothesize that binding of the substrate quinone close to heme 1 increases its potential, facilitating electron transfer from heme 2 to heme 1. We tested this hypothesis by comparing the reaction of CybB with superoxide in the presence and absence of quinone. In the presence of saturating quinone concentration the reaction was faster than the mixing time ( $t_{\chi}$ <1 s). This effect is in accordance with the observation that binding of the

semiquinone analogue inhibitor 'HQNO' decreased the potential of the low potential heme in succinate: menaquinone oxidoreductase from *Bacillus subtilis*<sup>56</sup>. Future experiments using EPR spectroscopy will help to measure and assign heme potentials reliably and pulse radiolysis studies will enable fast reaction kinetics of CybB with superoxide.

Substrate	Vmax (A <sub>438 nm</sub> . s <sup>-1</sup> )	<i>K<sub>m</sub></i> (μM)	Second order rate (M <sup>-1</sup> s <sup>-1</sup> )
Ubiquinone Q <sub>2</sub>	37.49 ± 0.68	0.39 ± 0.04	$2.9 \pm 0.4 \times 10^8$
			$(SOD = 2.0 \times 10^{9})^{23}$
Ubiquinol Q <sub>2</sub> H <sub>2</sub>	31.02 ± 2.79	10.29 ± 2.40	nd

**Table 1:** Summary of kinetics parameters as determined using WST-1 assay.

#### **Enzyme turnover measurements**

Here, we used a coupled enzyme assay that shows that C, bB actively competes with WST-1 for superoxide and that this competition is dependent on the CybB and quinone concentrations, as well as on quinone regeneration by bo<sub>3</sub> oxidase. This assay has been used and validated to detect SOD activity in tiscurs and purified fractions. We confirm that only in the presence of all components, suppression of formazan formation is observed. If all components but CybB was present, WCT-1 reduction by xanthine oxidase was unaffected. This is noteworthy, as it has been discribed that superoxide reacts in solution with short chain quinones spontaneously to form semiquinone and oxygen, and that two molecules of semiquinone can disproportion to fully reduced ubiquinol and fully oxidized ubiquinone 57 <sup>58</sup>. Alternatively, semiquinone can react with oxygen to form superoxide again. However, if this reaction occured a rignificant rate, formazan formation would already be decreased in the presence of quinone and  $bo_3$  oxidase (without CybB), which was not the case. To eliminate potential pitfalls, we have used the exact same conditions, including the presence of quinone and  $bo_3$  oxidase, and performed the experiment with purified SOD. We were surprised to find that only 10x more CybB than SOD was required to suppress WST-1 formation to a similar extent. This suggests that the second order rate constant of CybB is about 10 times lower than the one of SOD, yielding a still impressive value of ~2.9 x  $10^8 \text{ M}^{-1}$  $s^{-1}$  in the presence of 10  $\mu$ M ubiquinone Q<sub>2</sub> (see table 1). Such a rate makes CybB a true competitor for cellular superoxide even in the presence of SOD (see below).

This same assay was used to titrate quinone as a substrate, rendering a  $K_m$  of ~390 nM, which is lower than what has been found for other quinone reducing enzymes in bacteria (typically in the micromolar range)<sup>59-63</sup>. Obviously, such high turnover rates require a fast association of CybB with superoxide and a fast reduction of ubiquinone, and the rates of these have not been measured yet, but might be accessible by pulse radiolysis experiments, where defined concentrations of superoxide can be produced within nanoseconds<sup>64</sup>. The observed prerequisite for quinone binding to CybB to allow rapid electron transfer from superoxide might reflect a strategy to suppress reactive intermediates in the membrane. Mechanistically, it is important to remember that two molecules of superoxide must bind to fully reduce quinone to quinol and that protons must be taken up before reduced quinol can be released to the membrane and be used as a substrate for e.g., respiration. Future mutagenesis studies and experiments in proteolipologic should allow us to identify residues that are critical for these processes.

#### **Concluding remarks**

The presented data strongly support or, in tial hypothesis, which was based on the crystal structure and a set of functional experimence<sup>17</sup> that CybB of *E. coli* can work as a superoxide oxidase *in vitro*. While fast pre-steece state measurements like those performed for SOD and SOR are still outstanding, we find the present set of data corroborates our initials findings. The measured redox notentials of the two hemes are lower than those of other cytochrome  $b_{561}$  proteins, crigge sting a different substrate than ascorbate. In addition to the proposed forward reaction also the reverse reaction (formation of superoxide from ubiquinol) is readily observed, and the similarity of the structural arrangement to the NADPH: superoxide oxidase from *Cylindrospermum stagnale* is intriguing. The Michaelis-Menten constant  $K_m$  for quinone is 25 times lower in the forward direction than for quinol in the reverse direction, making the forward reaction kinetically favourable.

To date, the physiological role of CybB is unknown, as simple growth experiments have not yielded a phenotype in a knockout strain. It seems unlikely, however that the forward reaction takes place under aerobic conditions with a high oxygen / superoxide ratio. Under such conditions, the reverse reaction, the release of superoxide to periplasm is more likely to occur. In this context, it is noteworthy that periplasmic superoxide formation has been described for *E. coli*<sup>12</sup>, and the reaction of mena-semiguinone with oxygen has been

27

suggested as a pathway for this. Based on our results with menaquinol, we suggest that CybB can catalyse such a reaction when the menaquinone pool is reduced during respiration, equipping *E. coli* with a "superoxide gun" to attack competitors. The idea is especially attractive since periplasmic SOD is mainly expressed during stationary phase, while CybB transcription is enhanced during the logarithmic phase<sup>17</sup>. Finally, the forward reaction can be envisioned to happen under microaerobic conditions where the  $O_2$  / superoxide ratio is low; like the environment *E. coli* faces as a commensal bacterium of the human gut.

#### Acknowledgments

Work in the labs of CvB and ML is supported by the Swiss National Science Foundation. Work in the lab of MH is supported by the Swedish research council and the Knut and Alice Wallenberg foundation.

SUILOR

# References

- 1 D'Autréaux, B. & Toledano, M. B. ROS as signalling molecules: mechanisms that generate specificity in ROS homeostasis. *Nature reviews. Molecular cell biology* **8**, 813-824, doi:10.1038/nrm2256 (2007).
- 2 Dröge, W. Free radicals in the physiological control of cell function. *Physiological reviews* **82**, 47-95, doi:10.1152/physrev.00018.2001 (2002).
- 3 Pryde, K. R. & Hirst, J. Superoxide is produced by the reduced flavin in mitochondrial complex I: a single, unified mechanism that applies during both forward and reverse electron transfer. *J Biol Chem* **286**, 18056-18065, doi:10.1074/jbc.M110.186841 (2011).
- 4 Quinlan, C. L. *et al.* Mitochondrial complex II can generate reactive oxygen species at high rates in both the forward and reverse reactions. *J Biol Chem* **287**, 27255-27264, doi:10.1074/jbc.M112.374629 (2012).
- 5 Dröse, S. & Brandt, U. The Mechanism of Mitochondria Superoxide Production by the Cytochrome bc1 Complex\*. *Journal of Biological Chemistry* **283**, 21649-21654, doi:https://doi.org/10.1074/jbc.M803236200 (2008).
- 6 Imlay James, A. & Slauch James, M. Oxidetive Stress. *EcoSal Plus* **3**, doi:10.1128/ecosalplus.5.4.4 (2009).
- 7 Magnani, F. *et al.* Crystal structures and atomic moot of NADPH oxidase. *Proceedings of the National Academy of Sciences* **114**, 6764, doi:10.1073, pnas.1702293114 (2017).
- 8 Vignais, P. V. The superoxide-generating NADPH coidase: structural aspects and activation mechanism. *Cellular and molecular life sciences : CMLS* **59**, 1428-1459, doi:10.1007/s00018-002-8520-9 (2002).
- 9 Nguyen, G. T., Green, E. R. & Mecsas. I. Steutrophils to the ROScue: Mechanisms of NADPH Oxidase Activation and Bacterial Resist; nee. Frontiers in cellular and infection microbiology 7, 373, doi:10.3389/fcimb.2017.0037. 2017).
- 10 Diaz, J. M. *et al.* Widespread <sup>D</sup>roduction of Extracellular Superoxide by Heterotrophic Bacteria. *Science* **340**, 1223-122<sup>C</sup>, du: doi:10.1126/science.1237331 (2013).
- 11 Huycke, M. M. *et al.* Extracell ll r uperoxide production by Enterococcus faecalis requires demethylmenaquinone and is actenuated by functional terminal quinol oxidases. *Mol Microbiol* **42**, 729-740, dc::10106/j.1365-2958.2001.02638.x (2001).
- 12 Korshunov, S. & Imlay, I., Detection and quantification of superoxide formed within the periplasm of Escherichia c.!i. *Journal of bacteriology* **188**, 6326-6334, doi:10.1128/jb.00554-06 (2006).
- 13 Lin, P. C., Tür', K. H.se, C. C., Fritz, G. & Steuber, J. Quinone reduction by the Na+translocating NAL.' dehydrogenase promotes extracellular superoxide production in Vibrio cholerae. *Journal*, *bacteriology* **189**, 3902-3908, doi:10.1128/jb.01651-06 (2007).
- 14 Fridovich, I. Superoxide dismutases. *Annu Rev Biochem* **44**, 147-159, doi:10.1146/annurev.bi.44.070175.001051 (1975).
- 15 McCord, J. M. & Fridovich, I. Superoxide Dismutase: AN ENZYMIC FUNCTION FOR ERYTHROCUPREIN (HEMOCUPREIN). *Journal of Biological Chemistry* **244**, 6049-6055, doi:https://doi.org/10.1016/S0021-9258(18)63504-5 (1969).
- 16 Jenney Francis, E., Verhagen Marc, F. J. M., Cui, X. & Adams Michael, W. W. Anaerobic Microbes: Oxygen Detoxification Without Superoxide Dismutase. *Science* **286**, 306-309, doi:10.1126/science.286.5438.306 (1999).
- 17 Lundgren, C. A. K. *et al.* Scavenging of superoxide by a membrane-bound superoxide oxidase. *Nature Chemical Biology* **14**, 788-793, doi:10.1038/s41589-018-0072-x (2018).
- 18 Winkler, J. R. & Gray, H. B. Electron Flow through Metalloproteins. *Chemical Reviews* **114**, 3369-3380, doi:10.1021/cr4004715 (2014).
- 19 Page, C. C., Moser, C. C., Chen, X. & Dutton, P. L. Natural engineering principles of electron tunnelling in biological oxidation-reduction. *Nature* **402**, 47-52, doi:10.1038/46972 (1999).

- 20 Hayyan, M., Hashim, M. A. & AlNashef, I. M. Superoxide Ion: Generation and Chemical Implications. *Chemical Reviews* **116**, 3029-3085, doi:10.1021/acs.chemrev.5b00407 (2016).
- 21 Nishikimi, M., Appaji Rao, N. & Yagi, K. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochemical and Biophysical Research Communications* **46**, 849-854, doi:https://doi.org/10.1016/S0006-291X(72)80218-3 (1972).
- 22 Peskin, A. V. & Winterbourn, C. C. Assay of superoxide dismutase activity in a plate assay using WST-1. *Free Radical Biology and Medicine* **103**, 188-191, doi:https://doi.org/10.1016/j.freeradbiomed.2016.12.033 (2017).
- 23 Abreu, I. A. & Cabelli, D. E. Superoxide dismutases—a review of the metal-associated mechanistic variations. *Biochimica et Biophysica Acta (BBA) Proteins and Proteomics* **1804**, 263-274, doi:https://doi.org/10.1016/j.bbapap.2009.11.005 (2010).
- 24 Cerda, J. F. *et al.* Spectroelectrochemical measurements of redox proteins by using a simple UV/visible cell. *Electrochemistry Communications* **33**, 76-79, doi:https://doi.org/10.1016/j.elecom.2013.05.004 (2013).
- 25 Trott, O. & Olson, A. J. AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry* **31**, 455-461, doi:https://doi.org/10.1002/jcc 213 }4 (2010).
- 26 Bielski, B. H. J., Cabelli, D. E., Arudi, R. L. & Ross, B. Reactivity of HO2/O-2 Radicals in Aqueous Solution. *Journal of Physical and Cherrical Reference Data* **14**, 1041-1100, doi:10.1063/1.555739 (1985).
- 27 Nicholls, D. G. & Ferguson, S. J. in *Bioenergetics (Fourth Edition)* (eds David G. Nicholls & Stuart J. Ferguson) ix-x (Academic Press, 2C 1.?.).
- 28 Alric, J., Pierre, Y., Picot, D., Lavergne, J. S. Rappaport, F. Spectral and redox characterization of the heme ci of the cytochrome b of complex. *Proc Natl Acad Sci U S A* **102**, 15860-15865, doi:10.1073/pnas.0508102102 (2005)
- 29 Murakami, H., Kita, K. & Anraku, Y. Puritication and properties of a diheme cytochrome b561 of the Escherichia coli respiratory chain. *Journal of Biological Chemistry* **261**, 548-551, doi:https://doi.org/10.1016/SC 071-2258(17)36126-4 (1986).
- 30 Fridovich, I. Quantitative are of the production of superoxide anion radical by milk xanthine oxidase. *J Biol Che.* **245**, 4053-4057 (1970).
- 31 Porras, A. G., Olson, J. S. Palmer, G. The reaction of reduced xanthine oxidase with oxygen. Kinetics of peroxide and uperoxide formation. *Journal of Biological Chemistry* 256, 9096-9103, doi:https://doi.org/10.1016/S0021-9258(19)52513-3 (1981).
- 32 Samoilova, R. J. C ofts A. R. & Dikanov, S. A. Reaction of Superoxide Radical with Quinone Molecules. *The Journal of Physical Chemistry A* **115**, 11589-11593, doi:10.1021/jp204891n (2011).
- 33 Takahashi, M. & Asada, K. Superoxide production in aprotic interior of chloroplast thylakoids. *Archives of Biochemistry and Biophysics* **267**, 714-722, doi:https://doi.org/10.1016/0003-9861(88)90080-X (1988).
- 34 Husen, P. & Solov'yov, I. A. Spontaneous Binding of Molecular Oxygen at the Qo-Site of the bc1 Complex Could Stimulate Superoxide Formation. *Journal of the American Chemical Society* **138**, 12150-12158, doi:10.1021/jacs.6b04849 (2016).
- 35 Guo, J. & Lemire, B. D. The Ubiquinone-binding Site of the Saccharomyces cerevisiae Succinate-Ubiquinone Oxidoreductase Is a Source of Superoxide\*. *Journal of Biological Chemistry* **278**, 47629-47635, doi:https://doi.org/10.1074/jbc.M306312200 (2003).
- 36 Roginsky, V. A., Mohr, D. & Stocker, R. Reduction of ubiquinone-1 by ascorbic acid is a catalytic and reversible process controlled by the concentration of molecular oxygen. *Redox Report* **2**, 55-62, doi:10.1080/13510002.1996.11747027 (1996).
- 37 Getzoff, E. D. *et al.* Faster superoxide dismutase mutants designed by enhancing electrostatic guidance. *Nature* **358**, 347-351, doi:10.1038/358347a0 (1992).

- 38 Shin, D. S. *et al.* Superoxide Dismutase from the Eukaryotic Thermophile Alvinella pompejana: Structures, Stability, Mechanism, and Insights into Amyotrophic Lateral Sclerosis. *Journal of Molecular Biology* **385**, 1534-1555, doi:https://doi.org/10.1016/j.jmb.2008.11.031 (2009).
- 39 Sheng, Y. *et al.* Superoxide Dismutases and Superoxide Reductases. *Chemical Reviews* **114**, 3854-3918, doi:10.1021/cr4005296 (2014).
- 40 Adam, V., Royant, A., Nivière, V., Molina-Heredia, F. P. & Bourgeois, D. Structure of Superoxide Reductase Bound to Ferrocyanide and Active Site Expansion upon X-Ray-Induced Photo-Reduction. *Structure* **12**, 1729-1740, doi:https://doi.org/10.1016/j.str.2004.07.013 (2004).
- 41 Bedard, K. & Krause, K. H. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiological reviews* **87**, 245-313, doi:10.1152/physrev.00044.2005 (2007).
- 42 Sharma, P., Teixeira de Mattos, M. J., Hellingwerf, K. J. & Pekker, M. On the function of the various quinone species in Escherichiacoli. *The FEBJ Journal* **279**, 3364-3373, doi:https://doi.org/10.1111/j.1742-4658.2012.08608.x (2012).
- 43 Wissenbach, U., Kröger, A. & Unden, G. The specific functions of menaquinone and demethylmenaquinone in anaerobic respiration with fumarate, dimethylsulfoxide, trimethylamine N-oxide and nitrate by Escherichia Cli. Archives of Microbiology **154**, 60-66, doi:10.1007/BF00249179 (1990).
- 44 Tyson, K., Metheringham, R., Griffiths, L. & Cole, Characterisation of Escherichia coli K-12 mutants defective in formate-dependent nitrite reduction: essential roles for hemN and the menFDBCE operon. *Archives of Microbiology* **168**, 403-411, doi:10.1007/s002030050515 (1997).
- 45 Wallace, B. J. & Young, I. G. Role of qui iones in electron transport to oxygen and nitrate in Escherichia coli. Studies with a usi menA- double quinone mutant. *Biochimica et Biophysica Acta (BBA) Biochergetics* **461**, 84-100, doi:https://doi.org/10.1016/0005-2728(77)90071-8 (1977).
- 46 Ingledew, W. J. & Poole, R. K. The respiratory chains of Escherichia coli. *Microbiol Rev* **48**, 222-271, doi:10.1128/mr.48.3.222-271.1984 (1984).
- 47 Imlay, J. A. Cellular Defenser against Superoxide and Hydrogen Peroxide. *Annual Review of Biochemistry* **77**, 755-776, doi:10.1146/annurev.biochem.77.061606.161055 (2008).
- 48 Quick, K. L., Hardt, J. I. C Dugan, L. L. Rapid microplate assay for superoxide scavenging efficiency. *Journal of Neuroscience Methods* **97**, 139-144, doi:https://doi.org/10.016/S0165-0270(00)00179-5 (2000).
- 49 Murakami, H., K. 3. K. & Anraku, Y. Cloning of cybB, the gene for cytochrome b561 of Escherichia coli K12. *Molecular and General Genetics MGG* **198**, 1-6, doi:10.1007/BF00328692 (1984).
- 50 Jalukar, V., Kelley, P. M. & Njus, D. Reaction of ascorbic acid with cytochrome b561. Concerted electron and proton transfer. *Journal of Biological Chemistry* **266**, 6878-6882, doi:https://doi.org/10.1016/S0021-9258(20)89583-0 (1991).
- 51 Schultz, B. E., Edmondson, D. E. & Chan, S. I. Reaction of Escherichia coli Cytochrome bo3 with Substoichiometric Ubiquinol-2: A Freeze-Quench Electron Paramagnetic Resonance Investigation. *Biochemistry* **37**, 4160-4168, doi:10.1021/bi971714y (1998).
- 52 Magalon, A. *et al.* Heme Axial Ligation by the Highly Conserved His Residues in Helix II of Cytochrome b (Narl) of Escherichia coliNitrate Reductase A (NarGHI)\*. *Journal of Biological Chemistry* **272**, 25652-25658, doi:https://doi.org/10.1074/jbc.272.41.25652 (1997).
- 53 Takeuchi, F., Hori, H., Obayashi, E., Shiro, Y. & Tsubaki, M. Properties of Two Distinct Heme Centers of Cytochrome b561 from Bovine Chromaffin Vesicles Studied by EPR, Resonance Raman, and Ascorbate Reduction Assay. *The Journal of Biochemistry* **135**, 53-64, doi:10.1093/jb/mvh006 (2004).

- 54 Oakhill, J. S., Marritt, S. J., Gareta, E. G., Cammack, R. & McKie, A. T. Functional characterization of human duodenal cytochrome b (Cybrd1): Redox properties in relation to iron and ascorbate metabolism. *Biochimica et Biophysica Acta (BBA) Bioenergetics* **1777**, 260-268, doi:https://doi.org/10.1016/j.bbabio.2007.12.001 (2008).
- 55 Iyanagi, T., Yamazaki, I. & Anan, K. F. One-electron oxidation-reduction properties of ascorbic acid. *Biochimica et Biophysica Acta (BBA) Bioenergetics* **806**, 255-261, doi:https://doi.org/10.1016/0005-2728(85)90103-3 (1985).
- 56 Smirnova, I. A., Hägerhäll, C., Konstantinov, A. A. & Hederstedt, L. HOQNO interaction with cytochrome b in succinate:menaquinone oxidoreductase from Bacillus subtilis. *FEBS Letters* **359**, 23-26, doi:https://doi.org/10.1016/0014-5793(94)01442-4 (1995).
- 57 Land, E. J. & Swallow, A. J. One-electron reactions in biochemical systems as studied by pulse radiolysis. 3. Ubiquinone. *J Biol Chem* **245**, 1890-1894 (1970).
- 58 Maroz, A., Anderson, R. F., Smith, R. A. & Murphy, M. P. Reactivity of ubiquinone and ubiquinol with superoxide and the hydroperoxyl radical: in plications for in vivo antioxidant activity. *Free Radic Biol Med* **46**, 105-109, doi:10.1016/j.freera. biomed.2008.09.033 (2009).
- 59 Yap, L. L. *et al.* The quinone-binding sites of the cytoch om bo3 ubiquinol oxidase from Escherichia coli. *Biochimica et Biophysica Acta (BBA) 2:Jenergetics* **1797**, 1924-1932, doi:https://doi.org/10.1016/j.bbabio.2010.04.011 (2017).
- 60 Ma, J., Puustinen, A., Wikström, M. & Gennis, B. Tryptophan-136 in Subunit II of Cytochrome bo3 from Escherichia coli May facticipate in the Binding of Ubiquinol. Biochemistry **37**, 11806-11811, doi:10.1021/bi980\_0777 (1998).
- 61 Murai, M. *et al.* Characterization of the Ubiquinone Binding Site in the Alternative NADH-Quinone Oxidoreductase of Saccharor v.e. cerevisiae by Photoaffinity Labeling. *Biochemistry* **49**, 2973-2980, doi:10.1021; bi100005j (2010).
- 62 Blaza, J. N. *et al.* The mechanism of catalysis by type-II NADH:quinone oxidoreductases. *Scientific Reports* **7**, 40165, doi:10.10.75/srep40165 (2017).
- 63 Anderson, R. F. *et al.* Electron-Transfer Fathways in the Heme and Quinone-Binding Domain of Complex II (Succinate Dehydrogenase). *Biochemistry* **53**, 1637-1646, doi:10.1021/bi401630m (2014 .
- 64 Kobayashi, K. Pulse Radio<sup>1</sup>, S. Cludies for Mechanism in Biochemical Redox Reactions. *Chemical Reviews* **119**, 4412, 4462, doi:10.1021/acs.chemrev.8b00405 (2019).

#### **Declaration of interests**

 $\boxtimes$ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

# Highlights

- cybB of E. coli is a true superoxide:quinone oxidoreductase
- both ubiquinone and menaquinone can serve as electron acceptors or donors
- quinone binding to cybB accelerates reaction with superoxide
- apparent  $K_M$  values for both reactions are in the physiological range