

Review Article

Subcutaneous Drugs and Off-label Use in Hospice and Palliative Care: A Scoping Review



Ursina Wernli, MSc, Fabienne Dürr, BSc, Sibylle Jean-Petit-Matile, MD, Andrea Kobleder, PhD, and Carla Meyer-Massetti, PhD

Clinical Pharmacology and Toxicology (U.W., C.M.M.), Inselspital Bern, Switzerland; Graduate School for Health Sciences (U.W.), University of Bern, Switzerland; Institute of Primary Health Care BIHAM (F.D., C.M.M.), University of Bern, Switzerland; Hospice of Central Switzerland (S.J.P.M.), Littau, Switzerland; Institute of Applied Nursing Science (A.K.), Eastern Switzerland University of Applied Sciences OST, St Gallen, Switzerland

Abstract

Background. Subcutaneous drug administration is an interesting approach for symptom control in hospice and palliative care. However, most drugs have no marketing authorization for subcutaneous administration and are therefore used off-label. In order to meet the requirements of a safe and effective drug therapy, especially in highly vulnerable patients, it is essential to investigate the scope of evidence of these common practices.

Objectives. The purpose of this scoping review was to provide an overview of available data on the tolerability and/or effectiveness of subcutaneously administered and off-label used drugs.

Method. We performed a scoping review according to the PRISMA extension to identify data available on the tolerability and/or effectiveness of 17 predefined drugs that are commonly administered subcutaneously in Swiss hospices and hospice-like institutions and that have no marketing authorization (*off-label use*).

Results. The scoping review identified 57 studies with most data available on their tolerability (68% local, 54% systemic), clinical effects (82%), details on dosage (96%) and routes of application (100%). Information on pharmacokinetic properties was mostly missing and only available for fentanyl, levetiracetam, midazolam, and ondansetron. For seven drugs, less than five articles were identified and no studies on codeine or clonazepam were available.

Conclusion. This work provides an overview of current evidence on subcutaneous and off-label used drugs in hospice and palliative care. Although both are common practices, evidence on tolerability and effectiveness, particularly pharmacokinetic data, is limited and the identified information gaps need to be closed. This work establishes a basis for further research in this area. *J Pain Symptom Manage* 2022;64:e250–e259. © 2022 The Authors. Published by Elsevier Inc. on behalf of American Academy of Hospice and Palliative Medicine. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Key Words

Subcutaneous drug administration, off-label use, hospice care, palliative care, medication safety

Key Message

This scoping review summarizes clinical aspects of 17 drugs that are commonly administered subcutaneously in hospice and palliative care despite not holding a marketing authorization for this route of administration (*off-label use*). The identified lack of structured practice guidelines and

pharmacokinetic data indicate a need for further research.

Introduction

Subcutaneous (SC) drug administration offers a minimal invasive alternative to oral drug

All authors contributed to the study conception and design. Preparation, data collection and analysis for the scoping review were performed by F.D., C.M.M. and U.W. They were supported by S.J.P.M., who contributed her specialist knowledge from hospice care to this project. F.D. performed the scoping review (support by U.W. and C.M.M.). Title-abstract screening was performed by F.D. and U.W. The first draft of

the manuscript was written by U.W. and all authors commented on all versions of the manuscript. All authors read and approved the final manuscript.

Address correspondence to: Ursina Wernli, MSc E-mail: ursina.wernli@extern.insel.ch

Accepted for publication: 12 July 2022.

administration for symptom control, preferably when oral intake of drugs is severely limited (e.g., due to dysphagia, vomiting or impaired consciousness).¹ It is less invasive than intravenous administration and less painful than intramuscular injections,² which complies with the comfort-oriented considerations of hospice and palliative care, provided only drugs well-tolerable for SC administration with measurable effect are applied.³⁻⁵ A decline in patients' cognition may lead to a shift in preferences between routes of administration, from mainly oral medication towards parenteral medication. Therefore, the SC administration route becomes increasingly important.^{6,7}

Parenteral drugs can be administered as bolus injection or via infusion. Continuous SC infusions deliver drugs individually or in some cases as a mixture, usually over a period of 24 hours, using a syringe pump.⁸ Many drugs used in palliative care hold no marketing authorization for SC administration and are thus used *off-label*.⁹ Even though off-label use is a common practice in palliative care, precise requirements and structured practice guidelines for SC drug administration are lacking.¹⁰⁻¹²

In Switzerland, the term *off-label use* is not regulated by law. The Swiss Academy of Medical Sciences (SAMW) defines *off-label use* as «the use of readily available drugs with marketing authorization in Switzerland that deviate from the purpose approved by Swissmedic¹ and from characteristics published in the Swiss drug compendium».¹³ The definition thus includes administration of a drug that deviates from its registered and approved indication, dosage, route of application, duration of therapy, and administration in specific patient groups (e.g., children). Off-label use is permissible provided that due diligence is done and there is compliance with established best practice guidelines.¹⁴ The responsibility for the off-label use rests solely with the prescribing physician. To be able to justify off-label uses within the meaning of federal requirements, the physician must demonstrate that the decision is evidence-based or based on solid recommendations (i.e., guidelines of professional associations), and the benefit must clearly exceed any risk. Affected patients must be adequately informed and consent to the treatment must be obtained.^{14,15} As cost coverage of off-label prescriptions is limited, off-label use requires prior approval by the individual's health insurance provider according to the Swiss Health Insurance Ordinance.¹⁶ Off-label use should not be confused with *unlicensed use*, which alludes to drugs for which no marketing authorization for any indication has been granted by the relevant licensing authority (i.e., Swissmedic).

Examples are the import from countries where the drug is licensed by authorities with comparable regulatory drug control or pharmaceutical modifications to registered and approved drugs (e.g., crushing tablets to prepare a solution) and dispensing it in a different form.^{13,17,18}

To meet the requirements of a safe and effective drug therapy, especially in highly vulnerable patients, it is essential to investigate the scope of evidence of these common practices. The purpose of this scoping review was to obtain a scope of evidence from the literature on the tolerability and/or effectiveness of drugs that are administered subcutaneously and off-label in Swiss hospices and hospice-like institutions.

Method

A protocol was used to document the process of the scoping review that was performed and documented according to the PRISMA extension.¹⁹ The protocol was not previously published. Based on a previous survey study performed in Swiss hospices and hospice-like institutions, we identified 14 drugs that are used subcutaneously and off-label.²⁰ Additionally, three representatives of the therapeutic drug group of PPIs (esomeprazole, omeprazole, pantoprazole) were included in the scoping review, as a particular request for information on this drug group emerged from the survey (see Table 1).

Information Sources and Search

We performed a scoping review searching in PubMed, Embase, and CINAHL databases. The search string was designed using two topic blocks "Palliative Care Setting" and "Subcutaneous Drug Administration," using MeSH terms and keywords. It was initially developed in PubMed and then translated for use in Embase (using Emtree) and CINAHL (using Subject Headings). The full electronic search strategy from the search in PubMed is available as a *Supplemental file*. No filters were applied. If a full text article was not available online, employees of the university's library were contacted for procurement.

Table 1
Drugs That Were Identified to Be Used Subcutaneously and Off-label in Swiss Hospices and Hospice-like Institutions

Drugs Used Subcutaneously and Off-Label in Switzerland (n = 17)

• Ceftriaxone	• Levomepromazine
• Codeine	• Metamizole
• Clonazepam	• Metoclopramide
• Esomeprazole ^a	• Midazolam
• Fentanyl	• Olanzapine
• Furosemide	• Omeprazole ^a
• Haloperidol	• Ondansetron
• Ketamine	• Pantoprazole ^a
• Levetiracetam	

¹ Swissmedic is the Swiss national authorization and supervisory authority for drugs and medical products.

^aadditionally added n = 3 representatives of proton pump inhibitors

Table 2
Eligibility Criteria for Inclusion and Exclusion of Studies in Scoping Review

Eligibility Criteria IN	Eligibility Criteria OUT
<ul style="list-style-type: none"> • Primary literature (intervention and observational studies including case reports and case series) • Studies reporting local and/or systemic tolerability and/or effectiveness (clinical effect, blood plasma levels) of $n = 17$ predefined drugs • Drug administration and investigation in patients ≥ 18 years that receive palliative care • Referral of reported tolerability and/or effectiveness (clinical effect, blood plasma levels) to a specific drug or a mixture of active substances that contains at least one of the $n=17$ drugs of interest, must be feasible • Language English, French or German 	<ul style="list-style-type: none"> • Nonprimary literature (e.g., reviews), editorials, conference abstracts, expert opinions • Reports of drugs that are not among the predefined drugs • Drug administration and investigation in patients < 18 years, non-palliative, or healthy study subjects • Hypodermoclysis • Other routes of application • Other languages

Selection Sources of Evidence

The final search was performed on April 6, 2021. After removing duplicates, title-abstract screening was performed according to the eligibility criteria (see *Eligibility criteria*, Table 2) by two independent reviewers (F. D., U. W.); any discrepancies were resolved through discussion until consensus was reached. In reviews that were excluded according to the predefined eligibility criteria, “backward citation chasing” was applied to identify potentially missed studies. Full text screening was performed by one reviewer (F. D.) and discussed with two additional reviewers (U. W., C. M. M.).

Eligibility Criteria

Eligibility criteria are listed in Table 2. Included were all publications reporting tolerability and/or effectiveness (clinical effect, blood plasma or serum levels) of the 17 predefined drugs of interest (SC administration and off-label use). No restrictions for time of publication were defined. Hypodermoclysis (i.e., SC fluid infusions) was excluded as this scoping review aimed at putting an emphasis on the administration of drugs only.

Data Charting Process and Data Items

A detailed table was created for data extraction (i.e., study type, drug, number of patients, tolerability, clinical effect, details on drug administration, and references) with one row for each included publication. The detailed table is provided by the authors upon special request. The charted data was transferred to another, less detailed table (see *Synthesis of results*).

Data was charted independently by one reviewer (F. D.) and independently repeated in $n = 6$ (approximately 10%) randomly selected studies by a second reviewer (U. W.). The charted data was discussed among all three reviewers (F.D., U.W., C.M.M.) to resolve any discrepancies.

Synthesis of Results

To summarize the charted data, a table was created (see *Results of individual sources of evidence*, Table 3). References investigating the same drug were grouped and shaded to make them more discernible in the table.

Results

Selection Sources of Evidence

A total of $n = 58$ identified articles were included for data extraction. Two of the articles were written by the same author^{21,22} and reported the same findings but they were published in different journals at different times. The two articles were combined for data extraction leading to a total of $N = 57$ included articles (see Fig. 1).

Characteristics of Sources of Evidence

Of the included sources, 57.9% ($n = 33/57$) were of European origin. Almost half and thus the majority of the European sources (48.5%, $n = 16/33$) originated from the United Kingdom²¹⁻³⁷ followed by Spain ($n = 7/33$, 21.2%),³⁸⁻⁴⁴ France,⁴⁵⁻⁴⁷ and Germany⁴⁸⁻⁵⁰ (both with $n = 3/33$, 9.1% each). Other represented countries were Denmark,⁵¹ Italy,⁵² Portugal,⁵³ and Norway,⁵⁴ each contributing one source. Australia was the second most represented continent with $n = 12/57$ (21.1%) sources,⁵⁵⁻⁶⁶ followed by North America, with $n = 7/57$ (12.3%) sources. Of those, 57.1% ($n = 4/7$) originated from Canada⁶⁷⁻⁷⁰ and 42.9% ($n = 3/7$) from the United States.⁷¹⁻⁷³ One Canadian article⁶⁹ was a collaboration with a palliative care unit in Switzerland. The remaining articles originated from South America ($n = 3/57$, 5.3%) with one contribution each from Argentina,⁷⁴ Brazil,⁷⁵ and Uruguay.⁷⁶ 3.5% ($n = 2/57$) were of Asian origin, with one article each from China⁷⁷ and Japan.⁷⁸

Of the included articles, 17.5% ($n = 10/57$) were intervention studies.^{29,52,59-63,71,74,76} Eight of them were

Table 3
Overview of Extracted Data From $N = 57$ Articles

References (see below)	total (n=)	total %	Study type	Investigated drug																																																									Tolerability
				observational	intervention	ceftiaxone	clonazepam	codeine	esomeprazole	fentanyl	furosemide	haloperidol	ketamine	levetiracetam	levomepromazine	metamizole	metoclopramide	midazolam	olanzapine	omeprazole	ondansetron	pantoprazole																																							
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	
				Papa et al. 2021 (Uruguay)	Sutherland et al. 2021 (UK)	Furtado et al. 2018 (Portugal)	Sutherland et al. 2018 (UK)	Murray-Brown et al. 2016 (UK)	Wells et al. 2016 (UK)	Rémi et al. 2014 (Germany)	López-Saca et al. 2013 (Spain)	Birch et al. 2021 (UK)	Brown et al. 2020 (UK)	Picazo Sánchez et al. 2018 (Spain)	Farless et al. 2013 (USA)	Zacharias et al. 2011 (UK)	Goenaga et al. 2004 (Spain)	Elisayem et al. 2010 (USA)	Michelson et al. 2019 (France)	Desmids et al. 2009 (France)	Agar et al. 2004 (Australia)	Gremaud et al. 1998 (Australia)	Ramani et al. 1996 (USA)	Wilcock et al. 1996 (UK)	Bleasel et al. 1994 (Australia)	Stiefel et al. 1992 (Canada/Switzerland)	Burke et al. 1991 (Australia)	McNamara et al. 1991 (UK)	Bottomley et al. 1990 and 1992 (UK)	Armsbury et al. 1989 (UK)	D Sousa et al. 1988 (UK)	Barbosa et al. 2020 (Brazil)	Hardy et al. 2012 (Australia)	Jackson et al. 2010 (Australia)	Jackson et al. 2001 (Australia)	Bell 1999 (Norway)	Oshima et al. 1990 (Japan)	Hunt et al. 1999 (Australia)	Watanabe et al. 1998 (Canada)	Miller et al. 1995 (Australia)	Paix et al. 1995 (Australia)	Schlunk et al. 1994 (Germany)	Eisenschlas et al. 2005 (Argentina)	Kennett et al. 2005 (UK)	Hutton et al. 1995 (UK)	Bruera et al. 1987 (Canada)	Bruera et al. 1986 (Canada)	Porcel et al. 1998 (Spain)	Philpot 1993 (Australia)	Mulvenna et al. 1992 (UK)	Hardy et al. 2010 (Australia)	Negro et al. 2010 (Spain)	Back et al. 2007 (UK)	Negro et al. 2006 (Spain)	Ventafredda et al. 1990 (Italy)	Roubaud-Baudron et al. 2017 (France)	Jensen et al. 2020 (Denmark)	Bartz et al. 2014 (Germany)	Negro et al. 2002 (Spain)	Thorsen et al. 1994 (China)	

(Continued)




Fig. 1. Flowchart of articles identified in scoping review. Adapted from: Page MJ et al. (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews¹⁹

reactions usually occurred only in a few individuals, they were primarily described in studies on furosemide^{25,26,37} and ketamine.^{54,62,63} Clinical effects were reported in $n = 47/57$ (82.5%) articles. Most of these effects were reported for midazolam ($n = 8$), levetiracetam ($n = 7$), and furosemide and ketamine, both with six studies reporting clinical effects (see Table 3).

Information on PK properties (i.e., plasma levels) was available for fentanyl, levetiracetam, midazolam, and ondansetron. Three articles reported plasma concentrations of SC levetiracetam,^{39,49,76} but only one study performed a full PK analysis for SC levetiracetam.⁷⁶ This was the only PK analysis identified among the totally $N = 57$ included articles.

Discussion

This scoping review provides the first overview on important clinical aspects (i.e., tolerability and effectiveness) of SC drug administration in hospice and palliative care. These clinical aspects need to be considered in order to meet the requirements of safe and effective symptom management, especially if drugs are used off-label. The identified scope of evidence reflects that SC administration is essential for symptom control in hospice and palliative care. If in compliance with best practice guidelines,¹⁴ off-label use offers treatment options for patients with special symptom control needs for whom conventionally approved routes of administration are inadequate. Hence, in hospice and palliative care, SC drug administration is often associated with off-label use.

Summary of Evidence

Information gaps on the tolerability and/or effectiveness (clinical effect, blood plasma or serum levels) of the drugs of interest became evident in the scoping review. Considering the substantial number of drugs of interest, only a rather small number of sources qualified for inclusion and data extraction. For seven drugs (ceftriaxone, esomeprazole, metamizole, olanzapine, omeprazole, ondansetron, pantoprazole), less than five studies were included. Interestingly, all three representatives of the therapeutic group of proton pump inhibitors, that were included in the scoping review upon request, were among these seven drugs. For two drugs (clonazepam, codeine), no studies were identified at all. However, both substances seem to have become less important in hospice and palliative care. The most recent “Model List of Essential Medicines,”⁷⁹ published by the WHO in 2021, lists only codeine tablets for oral administration among the medicines considered essential for hospice and palliative care, clonazepam is not listed at all. Although the evidence remains low, both drugs are still used in Swiss hospices and hospice-like institutions. The lack of identified information in the literature on the SC administration of most included drugs of interest reflects the need for more evidence to support clinical decision-making by hospice and palliative care physicians, as the responsibility for off-label use rests solely with them. Decisions need to be evidence-based or based on solid recommendations, and the benefits must clearly exceed any risk.

It is particularly important to increase medication safety in highly vulnerable patients. Evidence-based structured guidelines can help to improve medication safety in clinical settings. A lack of structured guidelines on SC drug administration and off-label use pertaining to hospice and palliative care was identified in this scoping review. Structured guidelines are desirable to support clinical decision-making, especially when drugs are used off-label. Guidelines are preferably based on evidence from studies that have investigated the safety and/or effectiveness of drug administration, particularly when administered to highly vulnerable patients, in order to prevent adverse drug events that may affect quality of life.

Even though SC drug administration is usually well-tolerated, there are substances among the 17 investigated drugs that are associated with severe adverse drug reactions (e.g., haloperidol) that can be misinterpreted as symptoms (e.g., extrapyramidal movements). Potentially life-threatening adverse drug reactions (e.g., qt-time prolongation) may occur.⁶⁰ More severe local side effects were also described in individual patients where initiation of antibiotic treatment was required, albeit occurring

rarely.^{25,26,33,37,47,54,62,63,66} A potentially resulting prescribing cascade must be avoided as this is somewhat contradictory to the approach of maintaining quality of life.

The number of randomized double-blind placebo-controlled or cross-over clinical trials, which are considered to provide the highest level of evidence, was scarce. This study design is particularly difficult to perform in patients of hospice and palliative care due to the complexity and high frailty of this patient population.⁸⁰ Randomization into different treatment arms is impractical and blinding is often unethical. Comparison among drugs is nearly impossible due to the high inter-patient variability and required daily doses vary greatly between patients.⁷⁶ Drug therapy regimens are adapted to current requirements in symptom control and thus, can change frequently. This lack of high-level evidence results in a deficit of structured guidelines for evidence-based clinical decision-making. As a result, current recommendations on dosage and route of administration to guide drug choice and/or dose tailoring to individual patients are rarely supported by high-level evidence.⁸¹

Guidelines may also be based on well-documented clinical experience shared among institutions. In Switzerland, no database to facilitate the exchange of clinical experience pertaining to SC drug administration among institutions is available. As a result, most off-label prescriptions and SC drug administration remain low in evidence and are often limited to clinical experience at an institutional level. Available guidelines (e.g., *BIGORIO* Best Practice Guidelines) cover only a part of the broad spectrum of safety and effectiveness of drug administrations in palliative care.⁸²

The identified information gaps establish a basis for further research to support clinical decision-making. To provide evidence that subcutaneous drug administration, especially used off-label, is safe and effective in hospice and palliative care patients, studies providing pharmacokinetic data are required.

Limitations

The basis for selection of the drugs of interest was a previously performed survey study in Swiss hospices and hospice-like institutions. Hospice and palliative care physicians and nurses were asked to list all drugs that are subcutaneously administered in their institutions. Findings are therefore mainly of interest for institutions that use a similar list of SC drugs in hospice and palliative care patients. The clinical trial register was not searched for ongoing studies; therefore, the low number of intervention studies potentially underrepresents current progress in research on this topic.

Conclusion

To our knowledge, this is the first scoping review that provides an overview of clinical aspects on subcutaneous drug administration and off-label use in hospice and palliative care. Evidence on tolerability and effectiveness is limited, resulting in a lack of structured guidelines. Although both are common practices, in-depth knowledge is deficient, and the scoping review revealed a need to close existing information gaps, especially on pharmacokinetic properties of commonly used drugs.

Disclosures and Acknowledgments

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. We acknowledge the support received from Marc von Gernler, PhD, of the library at University of Bern when planning and performing the literature review. There are no conflicts to disclosure. All authors declare no competing interests nor personal financial interests.

Supplemental file: Search strategy PubMed

((("terminal patient"[Title/Abstract]) OR (((((((((Hospice*[Title/Abstract]) OR ("Palliative Care*[Title/Abstract])) OR ("Palliative Treatment*[Title/Abstract])) OR ("Palliative Therap*[Title/Abstract])) OR ("Palliative Nursing*[Title/Abstract])) OR ("Terminal Care*[Title/Abstract])) OR ("End of Life Care*[Title/Abstract])) OR ("End-Of-Life Care*[Title/Abstract])) OR ("Palliative Supportive Care*[Title/Abstract])) OR (("terminally ill"[Title/Abstract]) OR ("hospice patient"[Title/Abstract])))) OR ((terminally ill[MeSH Terms]) OR (((((((("Terminal Care"[Mesh:NoExp]) OR (hospices [MeSH Terms])) OR (hospice care[MeSH Terms])) OR (hospice and palliative care nursing[MeSH Terms])) OR (palliative medicine[MeSH Terms])) OR (palliative care[MeSH Terms])))) AND (((("Infusions, Subcutaneous"[Mesh:NoExp]) OR ("Injections, Subcutaneous"[Mesh:NoExp]) OR (((((((((((((((("subcutaneous administration"[Title/Abstract]) OR ("s.c. administration"[Title/Abstract])) OR ("sc administration"[Title/Abstract])) OR ("Subcutaneous Infusion"[Title/Abstract])) OR ("s.c. infusion"[Title/Abstract])) OR ("sc infusion"[Title/Abstract])) OR ("Continuous subcutaneous infusion"[Title/Abstract])) OR ("CSCI"[Title/Abstract])) OR ("Subcutaneous injection"[Title/Abstract])) OR ("s.c. injection"[Title/Abstract])) OR ("sc injection"[Title/Abstract])) OR ("Subcutaneous bolus injection"[Title/Abstract])) OR ("Subcutaneous application"[Title/Abstract])) OR ("Subcutaneous Access"[Title/

Abstract])) OR ("Subcutaneous drug"[Title/Abstract])) OR ("Subcutaneous medic*[Title/Abstract])) OR ("Subcutaneous route"[Title/Abstract])) OR ("Subcutaneous deliver*[Title/Abstract])) OR ("Subcutaneous dosage"[Title/Abstract])) OR ("Subcutaneous dose"[Title/Abstract])) OR ("Subcutaneous therap*[Title/Abstract])) OR ("Syringe pump*[Title/Abstract])) OR ("Syringe driver"[Title/Abstract])))).

References

1. Dickman A, Schneider J. The syringe driver: continuous subcutaneous infusions in palliative care. 4th ed. Oxford: Oxford University Press; 2016.
2. Thomas T, Barclay S. Continuous subcutaneous infusion in palliative care: a review of current practice. *Int J Palliat Nurs* 2015;21: 60, 62-4.
3. Drummond SH, Peterson GM, Galloway JG, et al. National survey of drug use in palliative care. *Palliat Med* 1996;10:119-124.
4. Herndon CM, Fike DS. Continuous subcutaneous infusion practices of United States hospices. *J Pain Symptom Manage* 2001;22:1027-1034.
5. O'Doherty CA, Hall EJ, Schofield L, et al. Drugs and syringe drivers: a survey of adult specialist palliative care practice in the United Kingdom and Eire. *Palliat Med* 2001;15:149-154.
6. Hanlon JT, Weinberger M, Samsa GP, et al. A randomized, controlled trial of a clinical pharmacist intervention to improve inappropriate prescribing in elderly outpatients with polypharmacy. *Am J Med* 1996;100:428-437.
7. Mekonnen AB, McLachlan AJ, Brien JA. Effectiveness of pharmacist-led medication reconciliation programmes on clinical outcomes at hospital transitions: a systematic review and meta-analysis. *BMJ Open* 2016;6:e010003.
8. Eychmüller S. [Essentials of palliative medicine: The 1x1 of palliative care]. 2nd ed. Bern: Hogrefe AG; 2019.
9. Rémi C, Redmann C. [Palliative pharmacy: The pharmacist in the palliative care team]. 1st ed. Stuttgart: Deutscher Apotheker Verlag; 2017.
10. European pharmacopeia (Ph.Eur.), C.o. Europe, Editor. 2020: Strasbourg.
11. Hagemann V, Bausewein C, Remi C. Drug use beyond the licence in palliative care: a systematic review and narrative synthesis. *Palliat Med* 2019;33:650-662.
12. Wilcock A, Howard P, Charlesworth S. Palliative care formulary seventh edition. London: Pharmaceutical Press; 2020.
13. SAMW, [Differentiation between standard therapy and experimental therapy in individual cases]. 2014 (adapted 2015).
14. [Federal court decision 134 IV 175]. 24.4.2008 Swiss Federal Court.
15. Cantonal Pharmacists Association, [Position paper: Recommendations on off label use of medicines in0007]. 2016, St Gallen.
16. SAMW Swiss Academy of Medical Sciences and FMH Swiss Medical Association, [Legal principles in everyday medical practice: a guide for practitioners]. 2020.

17. Aronson JK, Ferner RE. Unlicensed and off-label uses of medicines: definitions and clarification of terminology. *Br J Clin Pharmacol* 2017;83:2615–2625.
18. Turner S, Longworth A, Nunn AJ, et al. Unlicensed and off label drug use in paediatric wards: prospective study. *Bmj* 1998;316:343–345.
19. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
20. Wernli U, Dürr F, Kobleder A, Meyer-Masseti C. *[Safe and rational medication in the hospice setting - a new role for pharmacists?]*. palliative ch 2021: 48–52. Bern.
21. Bottomley DM, Hanks GW. Subcutaneous midazolam infusion in palliative care. *J Pain Symptom Manage* 1990;5:259–261.
22. Bottomley D, Hanks G. Controlling restlessness in advanced cancer patients. *Am J Nurs* 1992;92:72–74.
23. Amesbury B, Dunphy K. The use of subcutaneous midazolam in the home care setting. *Palliat Med* 1989;3.
24. Back I, Taubert M. Akathisia and an unusual symptomatic treatment: a case report. *Palliat Med* 2007;21:713–715.
25. Birch F, Boam E, Parsons S, et al. Subcutaneous furosemide in advanced heart failure: service improvement project. *BMJ Support Palliat Care* 2021.
26. Brown A, Westley K, Robson J, et al. Furosemide in end-stage heart failure: community subcutaneous infusions. *BMJ Support Palliat Care* 2020.
27. de Sousa E, Jepson BA. Midazolam in terminal care. *Lancet* 1988;1:67–68.
28. Hatton MQ, McMurray A, Harnett AN. Ulcerative skin reaction from subcutaneous infusion of isotonic methotrimeprazine and diamorphine. *Clin Oncol (R Coll Radiol)* 1995;7:268–269.
29. Kennett A, Hardy J, Shah S, et al. An open study of methotrimeprazine in the management of nausea and vomiting in patients with advanced cancer. *Support Care Cancer* 2005;13:715–721.
30. McNamara P, Minton M, Twycross R. Use of midazolam in palliative care. *Palliat Med* 1991;5:244–249.
31. Mulvenna PM, Regnard CF. Subcutaneous ondansetron. *Lancet* 1992;339:1059.
32. Murray-Brown FL, Stewart A. Remember Keppra: seizure control with subcutaneous levetiracetam infusion. *BMJ Support Palliat Care* 2016;6:12–13.
33. Sutherland A, Meldon C, Harrison T, et al. Subcutaneous levetiracetam for the management of seizures at the end of life: an audit and updated literature review. *J Palliat Med* 2021;24:976–981.
34. Sutherland AE, Curtin J, Bradley V, et al. Subcutaneous levetiracetam for the management of seizures at the end of life. *BMJ Support Palliat Care* 2018;8:129–135.
35. Wells GH, Mason LD, Foreman E, et al. Continuous subcutaneous levetiracetam in the management of seizures at the end of life: a case report. *Age Ageing* 2016;45:321–322.
36. Wilcock A, Twycross R. Midazolam for intractable hiccup. *J Pain Symptom Manage* 1996;12:59–61.
37. Zacharias H, Raw J, Nunn A, et al. Is there a role for subcutaneous furosemide in the community and hospice management of end-stage heart failure? *Palliat Med* 2011;25:658–663.
38. Goenaga MA, Millet M, Sánchez E, et al. Subcutaneous furosemide. *Ann Pharmacother* 2004;38:1751.
39. López-Saca JM, Vaquero J, Larumbe A, et al. Repeated use of subcutaneous levetiracetam in a palliative care patient. *J Pain Symptom Manage* 2013;45:e7–e8.
40. Negro S, Azuara ML, Sánchez Y, et al. Physical compatibility and in vivo evaluation of drug mixtures for subcutaneous infusion to cancer patients in palliative care. *Support Care Cancer* 2002;10:65–70.
41. Negro S, Martín A, Azuara L, et al. Compatibility and stability of ternary admixtures of tramadol, haloperidol, and hyoscine N-butyl bromide: retrospective clinical evaluation. *J Palliat Med* 2010;13:273–277.
42. Negro S, Reyes R, Azuara ML, et al. Morphine, haloperidol and hyoscine N-butyl bromide combined in s.c. infusion solutions: compatibility and stability. Evaluation in terminal oncology patients. *Int J Pharm* 2006;307:278–284.
43. Picazo Sánchez M, Castillo Lorenzo MN, Cuxart Pérez M, et al. Palliative use of furosemide in continuous subcutaneous infusion in a renal transplant patient with heart failure. *Nefrologia (Engl Ed)* 2018;38:438–439.
44. Porcel JM, Salud A, Porta J, et al. Antiemetic efficacy of subcutaneous 5-HT₃ receptor antagonists in terminal cancer patients. *J Pain Symptom Manage* 1998;15:265–266.
45. Desmidt T, Constans T. Subcutaneous infusion of esomeprazole in elderly patients in palliative care: a report of two cases. *J Am Geriatr Soc* 2009;57:1724–1725.
46. Michelon H, Souchu H, Chauvron-Defilippi B, et al. Subcutaneous pantoprazole in an elderly, palliative care patient. *BMJ Support Palliat Care* 2019.
47. Roubaud-Baudron C, Forestier E, Fraisse T, et al. Tolerance of subcutaneously administered antibiotics: a French national prospective study. *Age Ageing* 2017;46:151–155.
48. Bartz L, Klein C, Seifert A, et al. Subcutaneous administration of drugs in palliative care: results of a systematic observational study. *J Pain Symptom Manage* 2014;48:540–547.
49. Rémi C, Lorenzl S, Vyhnaek B, et al. Continuous subcutaneous use of levetiracetam: a retrospective review of tolerability and clinical effects. *J Pain Palliat Care Pharmacother* 2014;28:371–377.
50. Schlunk T, Friess D, Winterhalder D. Continuous subcutaneous application of peripherally and centrally acting analgesic agents. *Medizinische Welt* 1994;45:553–558.
51. Jensen JJ, Sjøgren P. Administration of label and off-label drugs by the subcutaneous route in palliative care: an observational cohort study. *BMJ Support Palliat Care* 2020.
52. Ventafridda V, Ripamonti C, Caraceni A, et al. The management of inoperable gastrointestinal obstruction in terminal cancer patients. *Tumori* 1990;76:389–393.
53. Furtado I, Gonçalves F, Gonçalves J, et al. Continuous subcutaneous levetiracetam in end-of-life care. *BMJ Case Rep* 2018;2018.
54. Bell RF. Low-dose subcutaneous ketamine infusion and morphine tolerance. *Pain* 1999;83:101–103.
55. Agar M, Webster R, Lacey J, et al. The use of subcutaneous omeprazole in the treatment of dyspepsia in palliative care patients. *J Pain Symptom Manage* 2004;28:529–531.

56. Bleasel MD, Peterson GM, Dunne PF. Plasma concentrations of midazolam during continuous subcutaneous administration in palliative care. *Palliat Med* 1994;8:231–236.
57. Burke AL, Diamond PL, Hulbert J, et al. Terminal restlessness—its management and the role of midazolam. *Med J Aust* 1991;155:485–487.
58. Gremaud G, Zulian GB. Indications and limitations of intravenous and subcutaneous midazolam in a palliative care center. *J Pain Symptom Manage* 1998;15:331–333.
59. Hardy J, Quinn S, Fazekas B, et al. Randomized, double-blind, placebo-controlled study to assess the efficacy and toxicity of subcutaneous ketamine in the management of cancer pain. *J Clin Oncol* 2012;30:3611–3617.
60. Hardy JR, O'Shea A, White C, et al. The efficacy of haloperidol in the management of nausea and vomiting in patients with cancer. *J Pain Symptom Manage* 2010;40:111–116.
61. Hunt R, Fazekas B, Thorne D, et al. A comparison of subcutaneous morphine and fentanyl in hospice cancer patients. *J Pain Symptom Manage* 1999;18:111–119.
62. Jackson K, Ashby M, Howell D, et al. The effectiveness and adverse effects profile of “burst” ketamine in refractory cancer pain: The VCOG PM 1-00 study. *J Palliat Care* 2010;26:176–183.
63. Jackson K, Ashby M, Martin P, et al. “Burst” ketamine for refractory cancer pain: an open-label audit of 39 patients. *J Pain Symptom Manage* 2001;22:834–842.
64. Miller RS, Peterson GM, Abbott F, et al. Plasma concentrations of fentanyl with subcutaneous infusion in palliative care patients. *Br J Clin Pharmacol* 1995;40:553–556.
65. Paix A, Coleman A, Lees J, et al. Subcutaneous fentanyl and sufentanil infusion substitution for morphine intolerance in cancer pain management. *Pain* 1995;63:263–269.
66. Philpot CR. Ondansetron by subcutaneous infusion. *Med J Aust* 1993;159:213.
67. Bruera E, Brenneis C, Michaud M, et al. Continuous Sc infusion of metoclopramide for treatment of narcotic bowel syndrome. *Cancer Treat Rep* 1987;71:1121–1122.
68. Bruera E, MacDonald N, Brenneis C, et al. Metoclopramide infusion with a disposable portable pump. *Ann Intern Med* 1986;104:896.
69. Stiefel F, Fainsinger R, Bruera E. Acute confusional states in patients with advanced cancer. *J Pain Symptom Manage* 1992;7:94–98.
70. Watanabe S, Pereira J, Hanson J, et al. Fentanyl by continuous subcutaneous infusion for the management of cancer pain: a retrospective study. *J Pain Symptom Manage* 1998;16:323–326.
71. Elsayem A, Bush SH, Munsell MF, et al. Subcutaneous olanzapine for hyperactive or mixed delirium in patients with advanced cancer: a preliminary study. *J Pain Symptom Manage* 2010;40:774–782.
72. Farless LB, Steil N, Williams BR, et al. Intermittent subcutaneous furosemide: parenteral diuretic rescue for hospice patients with congestive heart failure resistant to oral diuretic. *Am J Hosp Palliat Care* 2013;30:791–792.
73. Ramani S, Karnad AB. Long-term subcutaneous infusion of midazolam for refractory delirium in terminal breast cancer. *South Med J* 1996;89:1101–1103.
74. Eisenclas JH, Garrigue N, Junin M, et al. Low-dose levomepromazine in refractory emesis in advanced cancer patients: an open-label study. *Palliat Med* 2005;19:71–75.
75. Barbosa MG, Delfino RS, Sarin LM, et al. Repeated subcutaneous esketamine administration for depressive symptoms and pain relief in a terminally ill cancer patient: a case report. *Palliat Med* 2020;34:822–825.
76. Papa P, Oricchio F, Ginés M, et al. Pharmacokinetics of Subcutaneous Levetiracetam in Palliative Care Patients. *J Palliat Med* 2021;24:248–251.
77. Thorsen AB, Yung NS, Leung AC. Administration of drugs by infusion pumps in palliative medicine. *Ann Acad Med Singap* 1994;23:209–211.
78. Oshima E, Tei K, Kayazawa H, Urabe N. Continuous subcutaneous injection of ketamine for cancer pain. *Can J Anaesth* 1990;37:385–386.
79. WHO model list of essential medicines (22nd). World Health Organization; 2021.
80. Kamal AH, Peppercorn JM. The generalizability paradox within palliative care clinical trials. *Ann Palliat Med* 2013;2:101–104.
81. Masman AD, van Dijk M, Tibboel D, et al. Medication use during end-of-life care in a palliative care centre. *Int J Clin Pharm* 2015;37:767–775.
82. palliative.ch, *BIGORIO Best Practice Guidelines*, w. BIGORIO, Editor. 2003-2013.