Journal Pre-proofs

Letter to the Editor

Heterogeneity of Teaching Approaches to Determine Hand Position for Adult Chest Compressions among European Basic Life Support Instructors

Nino Fijačko, Ruth Masterson Creber, Sander Van Goor, Matej Strnad, Robert Greif

PII: DOI: Reference:	S0300-9572(22)00661-X https://doi.org/10.1016/j.resuscitation.2022.09.005 RESUS 9591
To appear in:	Resuscitation
Received Date: Accepted Date:	3 September 20226 September 2022



Volume 164, July 2021

RESUSCITATION

OFFICIAL IOURNAL OF THE EUROPEAN RESUSCITATION COUNCIL

ELSEVIER

186N 0200 0572

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Elsevier B.V. All rights reserved.

Heterogeneity of Teaching Approaches to Determine Hand Position for Adult Chest Compressions among European Basic Life Support Instructors

Nino Fijačko a,b,c

Ruth Masterson Creber ^d

Sander Van Goor ^e

Matej Strnad f,g,h

Robert Greif ^{b,i,j}

^a University of Maribor, Faculty of Health Sciences, Maribor, Slovenia

^b ERC Research Net, Niels, Belgium

^c Ljubljana Community Health Centre, Ljubljana, Slovenia

^d Department of Population Health Sciences, Division of Health Informatics, Weill Cornell Medicine, New York, NY, USA

^e Ambulance Service, Municipal Health Service Haaglanden, Den Haag, Netherlands

^f University of Maribor, Faculty of medicine, Maribor, Slovenia

^g Maribor University Medical Centre, Maribor, Slovenia

^h Community Healthcare Center dr. Adolf Drolc Maribor, Maribor, Slovenia

ⁱ Department of Anaesthesiology and Pain Medicine, Bern University Hospital, University of Bern,

Bern, Switzerland

^j School of Medicine, Sigmund Freud University Vienna, Vienna, Austria

Corresponding author: Nino Fijačko, Žitna ulica 15, 2000 Maribor, +386 2 30 04 764, nino.fijacko@um.si, University of Maribor, Faculty of Health Sciences, Maribor, Slovenia, Twitter: @NinoFijacko

Key words: basic life support; cardiopulmonary resuscitation, chest compressions, hand placement site, teaching

Word count: 490/500

To the Editor,

During cardiac arrest, high-quality chest compression is crucial for organ perfusion and is dependent on the correct hand position. The European Resuscitation Council simplified its basic life support (BLS) guidelines and now recommends that the correct hand position is in the center of the chest and on the lower half of the sternum¹, with no specific landmark specified on the victim's chest². When teaching BLS to the lay public, specific visual landmarks are relied on upon³. This letter presents different BLS lay teaching approaches for teaching the correct position of adult chest compressions to the lay public from BLS-instructors in two European countries. In May 2022, Slovenian and Dutch BLS-instructors participated in an observational study in which

they were asked to demonstrate their teaching approach for determining the chest compression landmark on a low fidelity manikin (Rescue Anne QCPR, Laerdal Medical, Stavanger, Norway). The manikin was lying on the ground with clothes on and arms down.

A total of 43 BLS-instructors, 22 from Slovenia (SLO) and 21 from the Netherlands (NL) were invited to participated in this observational study. Twenty-four (56%) were registered nurses (10 SLO, 14 NL), 19 (44%) physicians (12 SLO, 7 NL); 18 (42%) were women (5 SLO, 13 NL); 41±9.6 years; with 10±7.4 years of experience in BLS-teaching. All Slovenian BLS instructors used a combination of theoretical (e.g., figures/photos in PowerPoint presentation) and practical (e.g., manikin) teaching approaches for demonstrating hand position for adult chest compressions, whereas Dutch BLS instructors relied primarily on practical approaches. Before demonstrating their hand position, 19 (86%) Slovenian BLS-instructors exposed the manikin's chest, whereas only 8 (38%) Dutch BLS-instructors did (p>0.001). More than half (n=25; 58%) of BLS-instructors

Journal Pre-proofs

were kneeling on the right side of a manikin. Most Slovenian BLS-instructors first determined the chest size from the clavicles to the posterior costal arch aiming to locate the chest's center and the lower half of the sternum (n=18, 82%). Three different teaching approaches were demonstrated (Figure 1A-C). Less than half (n=6, 42%) of Dutch BLS-instructors did not use any method and just placed their hand in the middle of the manikin's chest, 4 (24%) used the "internipple" line. Another 3 (14%) used the knee to shoulder⁴ (Figure 1D), or arm under armpit approaches (Figure 1F) to determine the correct hand position for adult chest compressions. All BLS-instructors targeted the center of the chest and the lower half of the sternum; however, Slovenian BLS instructors demonstrated a combined approach of vertical and horizontal chest size measurement to determine the correct compression points.

This observational study highlights the heterogeneity of teaching approaches to determine correct hand position for adult chest compressions used by instructors from Slovenia and the Netherlands. Overall, few Slovenian and Dutch BLS-instructors rely on previous BLS-recommendations (e.g., the internipple line) as an anatomic landmark for hand position in adult chest compressions⁵. Future BLS-guidelines should provide further clarity on the most evidence-based location for chest compressions among adults⁶⁻⁸.

Conflict of interest:

Nino Fijačko is a member of the ERC BLS Science and Education Committee and mentee of ILCOR Task Force Education Implementation and Team. Sander van Goor is co-chair of the ERC BLS Science and Education Committee. Robert Greif is ERC Director of Guidelines and ILCOR, and ILCOR Task Force chair Education Implementation and Team. Matej Strnad and Ruth Masterson Creber declare that they have no conflict of interest.

3

References

- Olasveengen, T.M.; Semeraro, F.; Ristagno, G.; Castren, M.; Handley, A.; Kuzovlev, A.; Monsieurs, K.G.; Raffay, V.; Smyth, M.; Soar, J.; et al. European Resuscitation Council Guidelines 2021: Basic Life Support. Resuscitation 2021, 161, 98–114.
- Kang, J., Lee, K.L., Hwang, S.O., Kim, Y.W., Kim, T.H., Jung, W.J., Kim, O.H., Cha, Y.S., Kim, H., Lee, K.H. and Cha, K.C., 2014. Lay rescuer CPR: Exposing the Chest Facilitates more Accurate and Safer Chest Compression. Journal of the Korean Society of Emergency Medicine, pp.550-556.
- Secher, N., Grove, E.L., Adelborg, K. and Løfgren, B., 2011. Visual-aided directions are superior to verbal instruction only in obtaining hand position for cardiopulmonary resuscitation. The American journal of emergency medicine, 29(9), pp.1178-1181.
- Birkenes, T.S., Myklebust, H. and Kramer-Johansen, J., 2013. New pre-arrival instructions can avoid abdominal hand placement for chest compressions. Scandinavian journal of trauma, resuscitation and emergency medicine, 21(1), pp.1-6.
- Koster, R.W., Baubin, M.A., Bossaert, L.L., Caballero, A., Cassan, P., Castrén, M., Granja, C., Handley, A.J., Monsieurs, K.G., Perkins, G.D. and Raffay, V., 2010. European Resuscitation Council Guidelines for Resuscitation 2010 Section 2. Adult basic life support and use of automated external defibrillators. Resuscitation, 81(10), pp.1277-1292.
- Kim, H., Chon, S.B., Yoo, S.M., Choi, H. and Park, K.Y., 2020. Optimum chest compression point might be located rightwards to the maximum diameter of the right ventricle: A preliminary, retrospective observational study. Acta Anaesthesiologica Scandinavica, 64(7), pp.1002-1013.
- Olszynski, P.A., Bryce, R., Hussain, Q., Dunn, S., Blondeau, B., Atkinson, P. and Woods, R., 2021. A novel anatomic landmark to target the left ventricle during chest compressions in cardiac arrest. Cureus, 13(3).

 Usawasuraiin, P., Wittayachamnankul, B., Chenthanakij, B., Euathrongchit, J., Phinyo, P. and Tangsuwanaruk, T., 2022. Optimal Landmark for Chest Compressions during Cardiopulmonary Resuscitation Derived from a Chest Computed Tomography in Arms-Down Position. Journal of Cardiovascular Development and Disease, 9(4), p.100.

Fig 1: Figures above present five teaching approaches for determining the center of the chest and the lower half of the sternum for performing adult chest compression: A) The little finger-thumb, B) The hand-thumb, C) The X-cross, D) The knee to shoulder, and E) The arm under armpit techniques.



For better visualization of the teaching approaches for determining the center of the chest we

prepared the video:

https://www.youtube.com/watch?v=MNx60sWjZcg&feature=youtu.be&ab_channel=NinoFija%C

4%8Dko

Conflict of interest:

Nino Fijačko is a member of the ERC BLS Science and Education Committee and mentee of ILCOR Task Force Education Implementation and Team. Sander van Goor is co-chair of the ERC BLS Science and Education Committee. Robert Greif is ERC Director of Guidelines and ILCOR, and ILCOR Task Force chair Education Implementation and Team. Matej Strnad and Ruth Masterson Creber declare that they have no conflict of interest.