

Hakim Arsany (Orcid ID: 0000-0001-9431-1069)

## Preferential Perfusion to The Posterior Circulation in Hypoxia

**Running head:** preferential perfusion in hypoxia

**Authors:**

Arsany Hakim <sup>1</sup>, MD, Franca Wagner <sup>1</sup>, MD

1 University Institute of Diagnostic and Interventional Neuroradiology, Bern

University Hospital, Inselspital, University of Bern.

Freiburgstrasse 10, CH-3010, Bern, Switzerland

**Corresponding author:**

Arsany Hakim

Telephone number: 0041 31 63 26696

e-mail: [arsany.hakim@insel.ch](mailto:arsany.hakim@insel.ch)

Address: Freiburgstrasse 10, CH-3010, Bern, Switzerland

Fax: 0041 31 632 48 72

**Number of characters:**

Title: 55

Running head: 30


**Number of words:**

Text: 387

Figure legend: 105

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ana.26539](https://doi.org/10.1002/ana.26539)

This article is protected by copyright. All rights reserved.



*Figure 1 Nonenhanced computed tomography (NECT) (A and E), CBF (B and F), DWI (C and G) and ADC (D and H) at the level of the basal ganglia (upper row) and of the cerebellum (lower row). NECT, obtained directly after resuscitation, shows diffuse effacement of the sulci, but the gray-white matter differentiation, as well as the differentiation of the basal ganglia, can still be seen. CBF shows global perfusion reduction of the gray matter (cortex and basal ganglia), but the perfusion of the cerebellum is preserved. DWI and ADC obtained 2 days after resuscitation showing diffuse restricted diffusion in the cortex and basal ganglia.*

Neuroimaging plays an important role in evaluating patients after cardiac arrest. CT is frequently used due to its wide availability and is preferred over MRI for examining intubated patients. However, CT has limited sensitivity for the detection of early brain damage.

CT may show diffuse cerebral edema, loss of gray-white matter differentiation, decreased deep gray matter attenuation, and reversal signs, such as white cerebellum sign.<sup>1 2</sup> This sign appears as increased attenuation of the cerebellum relative to the edematous supratentorial brain. Its presence reflects severe injury and indicates a poor prognosis.<sup>2 3</sup>

Various theories have been proposed to explain the pathophysiology of this sign, including

increased capillary proliferation induced by hypoxia; petechial hemorrhage; transtentorial herniation that causes central preservation of perfusion; distension of the medullary veins as a result of increased intracranial pressure; or preferential flow to the posterior circulation.<sup>2 3</sup>

We encountered a case of a 21-year-old female who was resuscitated after strangulation in a suicide attempt. The patient underwent CT with CT perfusion an hour after resuscitation. Non-enhanced CT showed diffuse brain swelling with effacement of the sulci, but the differentiation between the gray and white matter as well as the basal ganglia was preserved. CT perfusion, however, showed diffuse CBF and CBV reduction in the cortex, and basal ganglia while the perfusion to the cerebellum was preserved (Figure 1).

After 24 hours in intensive care under a normothermia protocol, the patient had not regained consciousness and brainstem reflexes were absent. Serial EEG showed myoclonic status epilepticus. Two days after resuscitation DWI showed alterations typical of severe hypoxic-ischemic injury (Figure 1).<sup>4</sup> The patient died on the third day.

The pattern, in this case, is very similar to the poor-prognostic indicator “white cerebellum sign” in the non-enhanced CT.<sup>2</sup> This supports the theory of redistribution of blood which occurs during anoxia, with preferential supply to the posterior circulation.<sup>2 1</sup>

Except for edema, no other signs of hypoxic-ischemic encephalopathy were visible on the NECT, though the perfusion reversal was seen early on in this phase, suggesting that this sign most probably developed earlier than the classical signs of hypoxia on CT. To the best of our knowledge, this perfusion pattern has not been described before. Further studies are

needed to investigate the usefulness of this sign to predict patient outcome after resuscitation.

Potential Conflicts of Interest:

Nothing to report.

Author contributions:

All authors contributed to the conception and design of the study; the acquisition and analysis of data; and drafting the text and preparing the figure.

## References:

1. Han BK, Towbin RB, de Courten-Myers G, et al. Reversal sign on CT: effect of anoxic/ischemic cerebral injury in children. [Internet]. *AJNR Am J Neuroradiol* 1989;10(6):1191–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2512781>
2. Huang BY, Castillo M. Hypoxic-Ischemic brain injury: Imaging findings from birth to adulthood. *Radiographics* 2008;
3. Han BK, Towbin RB, de Courten-Myers G, et al. Reversal sign on CT: effect of anoxic/ischemic cerebral injury in children. [Internet]. *American Journal of Roentgenology* 1990;154(2):361–368. Available from: <http://www.ajronline.org/doi/10.2214/ajr.154.2.2105031>
4. Barth R, Zubler F, Weck A, et al. Topography of MR lesions correlates with standardized EEG pattern in early comatose survivors after cardiac arrest. *Resuscitation* 2020;149:217–224.