



Pushing the boundaries of minimally invasive surgery

Fabrizio Minervini^{1^}, Sven Oliver Schumann², Gregor J. Kocher^{3^}

¹Department of Thoracic Surgery, Lucerne Cantonal Hospital, Lucerne, Switzerland; ²Department of General Internal Medicine, Hospital Bülach, Bülach, Switzerland; ³Division of General Thoracic Surgery, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

Correspondence to: Fabrizio Minervini, MD, PhD. Department of Thoracic Surgery, Lucerne Cantonal Hospital, Spitalstrasse, 6000 Lucerne, Switzerland. Email: fabrizio.minervini@hotmail.com.

Comment on: Chen J, Ang KL, Wang C, *et al.* Minimally invasive carinal reconstruction using bronchial flap and omental flap reinforcement. *Ann Thorac Surg* 2021. [Epub ahead of print]. doi: 10.1016/j.athoracsur.2021.06.014.

Submitted Sep 27, 2021. Accepted for publication Oct 13, 2021.

doi: 10.21037/tlcr-21-795

View this article at: <https://dx.doi.org/10.21037/tlcr-21-795>

The bronchial circulation plays a crucial role in the vascularization of tracheal and bronchial wall.

Tracheal resection or bronchial dissection as part of lung transplantation or sleeve resection could potentially induce ischemia, leading to mucosal necrosis or dehiscence with consecutive air leak of the respective segment.

Circumferential airway resection with end-to-end anastomosis is the most commonly used procedure to treat tumors that invade the distal trachea or carina, even if some surgeons prefer to perform a non-circumferential airway resection with bronchial flap reconstructions aiming to reduce the anastomotic tension (1-3).

For decades, an open surgical approach was considered the gold standard, but lately there has been an extraordinary growth in thoracic surgery with the introduction and development of minimally invasive procedures (4-7).

We have read with great interest the case report published by Chen and colleagues. They described in their paper a non-circumferential resection and reconstruction using bronchial flap (performed with a thoracoscopic approach) and omental flap harvested via laparoscopy (8). There is to date no consensus about both the optimal flap for reconstruction during thoracic surgery procedures, and its harvesting technique.

The authors didn't mention if they performed the carinal resection/reconstruction using a multiport or uniport approach, but regardless of that the procedure, even for expert surgeons, is very demanding. Wright and colleagues did not include

patients with carinal resection in their paper because they believed that those patients "present a unique complication profile" (9). Further to the technical difficulties *per se*, induction therapy could make planes' dissection very challenging.

Several authors have published case series of VATS tracheal and carinal resection and reconstruction with outstanding results (10-13). With regards to the abdominal part, laparoscopic harvest of an omental flap does usually not require advanced surgical skills if dissection of gastroepiploic vessels is not necessary. In case of thin omentum, the division of gastroepiploic is mandatory and the procedure could be difficult for a thoracic surgeon without advanced laparoscopic skills (14).

The case report published by Chen along with other previous case series could help the thoracic surgery community in pushing the boundaries of minimally invasive surgery, performing a very challenging procedure even in expert hands with an open approach.

Nevertheless, it should be kept in mind that, according to the hippocratic "primum non nocere", the safety of the patient must hold the first place in every surgeon's attitude and therefore a multidisciplinary setting with a high expertise in the field of tracheal surgery performed via minimally invasive approach is mandatory.

Acknowledgments

Funding: None.

[^] ORCID: Fabrizio Minervini, 0000-0003-3595-7307; Gregor J. Kocher, 0000-0001-6531-2919.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Translational Lung Cancer Research*. The article did not undergo external peer review.

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://dx.doi.org/10.21037/tlcr-21-795>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Grillo HC. Carinal reconstruction. *Ann Thorac Surg* 1982;34:356-73.
2. Shin S, Park JS, Shim YM, et al. Carinal resection and reconstruction in thoracic malignancies. *J Surg Oncol* 2014;110:239-44.
3. Peng Q, Zhang L, Ren Y, et al. Reconstruction of Long Noncircumferential Tracheal or Carinal Resections With Bronchial Flaps. *Ann Thorac Surg* 2019;108:417-23.
4. Minervini F, Bertolaccini L, Patrini D, et al. Non-intubated awake uniportal VATS: how to start? Video-assist Thorac Surg 2018;3:27.
5. Nakanishi R, Yamashita T, Muranaka K, et al. Thoracoscopic carinal resection and reconstruction in a patient with mucoepidermoid carcinoma. *J Thorac Cardiovasc Surg* 2013;145:1134-5.
6. Jiao W, Zhu D, Cheng Z, et al. Thoracoscopic tracheal resection and reconstruction for adenoid cystic carcinoma. *Ann Thorac Surg* 2015;99:e15-7.
7. Li J, Wang W, Jiang L, et al. Video-Assisted Thoracic Surgery Resection and Reconstruction of Carina and Trachea for Malignant or Benign Disease in 12 Patients: Three Centers' Experience in China. *Ann Thorac Surg* 2016;102:295-303.
8. Chen J, Ang KL, Wang C, et al. Minimally invasive carinal reconstruction using bronchial flap and omental flap reinforcement. *Ann Thorac Surg* 2021. [Epub ahead of print]. doi: 10.1016/j.athoracsur.2021.06.014.
9. Wright CD, Grillo HC, Wain JC, et al. Anastomotic complications after tracheal resection: prognostic factors and management. *J Thorac Cardiovasc Surg* 2004;128:731-9.
10. Yamamoto K, Miyamoto Y, Ohsumi A, et al. Surgical results of carinal reconstruction: an alternative technique for tumors involving the tracheal carina. *Ann Thorac Surg* 2007;84:216-20.
11. Li J, Wang W, Jiang L, et al. Video-Assisted Thoracic Surgery Resection and Reconstruction of Carina and Trachea for Malignant or Benign Disease in 12 Patients: Three Centers' Experience in China. *Ann Thorac Surg* 2016;102:295-303.
12. Gonzalez-Rivas D, Soultanis KM, Garcia A, et al. Uniportal video-assisted thoracoscopic lung sparing tracheo-bronchial and carinal sleeve resections. *J Thorac Dis* 2020;12:6198-209.
13. Sekhniadze D, Gonzalez-Rivas D, Kononets P, et al. Uniportal video-assisted thoracoscopic carinal resections: technical aspects and outcomes. *Eur J Cardiothorac Surg* 2020;58:i58-64.
14. Puma F, Fedeli C, Ottavi P, et al. Laparoscopic omental flap for the treatment of major sternal wound infection after cardiac surgery. *J Thorac Cardiovasc Surg* 2003;126:1998-2002.

Cite this article as: Minervini F, Schumann SO, Kocher GJ. Pushing the boundaries of minimally invasive surgery. *Transl Lung Cancer Res* 2021;10(11):4308-4309. doi: 10.21037/tlcr-21-795