Van Sundert Kevin (Orcid ID: 0000-0001-6180-3075) Chang Scott (Orcid ID: 0000-0002-7624-439X) De Kauwe Martin Gerard (Orcid ID: 0000-0002-3399-9098) Langley Adam (Orcid ID: 0000-0001-5164-4760) Ma Zilong (Orcid ID: 0000-0002-0145-814X) Mariën Bertold (Orcid ID: 0000-0003-1136-2480) Reynaert Simon (Orcid ID: 0000-0003-4690-0955) Vanuytrecht Eline (Orcid ID: 0000-0003-4690-0955) Vanuytrecht Eline (Orcid ID: 0000-0003-0631-1232) Yue Kai (Orcid ID: 0000-0002-7709-8523) Vicca Sara (Orcid ID: 0000-0001-9812-5837)

When things get MESI: the Manipulation Experiments Synthesis Initiative – a coordinated effort to synthesize terrestrial global change experiments

Running title: terrestrial global change experiment database

Kevin Van Sundert^{1,2,3,4†*} (0000-0001-6180-3075) Sebastian Leuzinger^{5†} (0000-0001-9306-5281) Martin K.-F. Bader⁶ (0000-0002-3742-9762) Scott X. Chang⁷ (0000-0002-7624-439X) Martin G. De Kauwe⁸ (0000-0002-3399-9098) Jeffrey S. Dukes⁹ (0000-0001-9482-7743) J. Adam Langley¹⁰ (0000-0001-5164-4760) Zilong Ma¹¹ (0000-0002-4621-5766) Bertold Mariën¹ (0000-0003-1136-2480) Simon Reynaert¹ (0000-0003-4690-0955) Jingyi Ru¹² (0000-0001-8675-0723) Jian Song¹² (0000-0001-9957-6533) Benjamin Stocker^{13,14} (0000-0003-2697-9096) César Terrer² (0000-0002-5479-3486) Joshua Thoresen^{7,15} (0000-0002-8653-7110) Eline Vanuytrecht^{16,17} (0000-0002-1247-6183) Shiqiang Wan¹² (0000-0003-0631-1232) Kai Yue^{18,19} (0000-0002-7709-8523) Sara Vicca¹ (0000-0001-9812-5837)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.16585

¹Research Group PLECO (Plants and Ecosystems), Global Change Ecology Centre of Excellence, Biology Department, University of Antwerp, Wilrijk, Belgium

²Climate and Ecological Synthesis Lab, Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA, USA

³Department of Earth System Science, Doerr School of Sustainability, Stanford University, Stanford, CA, USA

⁴Ecological Synthesis Lab, School of Informatics, Computing and Cyber Systems, Northern Arizona University, Flagstaff, AZ, USA

⁵School of Science, Auckland University of Technology, Auckland, New Zealand

⁶Department of Forestry and Wood Technology, Linnaeus University, Växjö, Sweden

⁷Department of Renewable Resources, University of Alberta, Edmonton, Alberta, T6G 2E3, Canada ⁸School of Biological Sciences, University of Bristol, Bristol, BS8 1TQ, UK

⁹Department of Global Ecology, Carnegie Institution for Science, Stanford, CA, USA

¹⁰Department of Biology and Center for Biodiversity and Ecosystem Stewardship, Villanova University, Villanova, PA, USA

¹¹State Key Laboratory of Biocontrol, School of Ecology, Sun Yat-sen University, Guangzhou, China ¹²School of Life Sciences, Institute of Life Science and Green Development, Hebei University, Baoding, China

¹³Institute of Geography, University of Bern, Bern, Switzerland

¹⁴Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

¹⁵Wildland Consultants, Auckland, New Zealand

¹⁶Division of Soil & Water Management, Faculty of Bioscience Engineering, KU Leuven, Leuven, Belgium ¹⁷Climate Change Adaptation, European Environment Agency, Copenhagen, Denmark

¹⁸Key Laboratory for Humid Subtropical Eco-Geographical Processes of the Ministry of Education, School of Geographical Sciences, Fujian Normal University, Fuzhou 350007, Fujian, China;

¹⁹Fujian Sanming Forest Ecosystem National Observation and Research Station, Sanming 365002, Fujian, China

[†]These authors contributed equally to this work

*e-mail: kvansundert.ac@gmail.com

Responses of the terrestrial biosphere to rapidly changing environmental conditions are a major source of uncertainty in climate projections. In an effort to reduce this uncertainty, a wide range of global change experiments have been conducted that mimic future conditions in terrestrial ecosystems, manipulating CO₂, temperature, nutrient and water availability. Syntheses of results across experiments provide a more general sense of ecosystem responses to global change, and help to discern the influence of background conditions such as climate and vegetation type in determining global change responses. Several independent syntheses of published data have yielded distinct databases for specific objectives. Such parallel, uncoordinated initiatives carry the risk of producing redundant data collection efforts and have led to contrasting outcomes without clarifying the underlying reason for divergence. These problems could be avoided by creating a publicly available, updatable, curated database. Here, we report on a global effort to collect and curate 57,089 treatment responses across 3,644 manipulation experiments at 1,145 sites, simulating elevated CO₂, warming, nutrient addition and precipitation changes. In the resulting Manipulation Experiments Synthesis Initiative (MESI) database, effects of experimental global change drivers on carbon and nutrient cycles are included, as well as ancillary data such as background climate, vegetation type, treatment magnitude, duration, and, unique to our database, measured soil properties. Our analysis of the database indicates that most experiments are short-term (one or few growing seasons), conducted in the USA, Europe or China, and that the most abundantly reported variable is aboveground biomass. We provide the most comprehensive multifactor global change database to date, enabling the research community to tackle open research questions, vital policymaking. The MESI database. freelv accessible to alobal at doi.org/10.5281/zenodo.7153253, opens new avenues for model evaluation and synthesisbased understanding of how global change affects terrestrial biomes. We welcome contributions to the database on GitHub.

KEYWORDS

meta-analysis, climate change, CO₂, warming, drought, nitrogen, precipitation, manipulation experiment

1 | INTRODUCTION

One of our most important tools to make predictions about the response of terrestrial ecosystems to global change is ecosystem experimentation. Manipulative experiments are especially valuable for acquiring mechanistic insights into, and quantifying the influence of, individual drivers and their interactions in terrestrial ecosystems – which is typically not possible with ecosystem monitoring.

By combining data from many such experiments, meta-analyses can reveal broader-scale patterns. Several meta-analyses are based on collecting data from the primary literature, on the initiative of a particular research group or international collaboration, and with a particular set of research questions in mind. Starting data collection anew for each study avoids the possibility of a consistent database creation bias across studies, that is, researcher-specific selection of eligible publications, data interpretation, extraction methods etc. However, this approach of starting each meta-analysis from scratch is expensive and time-consuming. Moreover, it can lead to contrasting conclusions of different synthesis efforts without being clear about the exact reason behind the divergence (see e.g., Yue et al., 2017 versus Dieleman et al., 2012).

Meta-analyses of large numbers of experiments can also provide valuable input to validate, improve and parametrize ecosystem models applicable to the global scale. However, while data from individual smaller sets of global change experiments are indeed increasingly used to test hypotheses embodied by ecosystem models (Zaehle et al., 2014), a wider adoption of model-data syntheses remains hampered by the current lack of comprehensive integration, homogenization and accessibility of global change experimental data.

We propose that synthesis-related research questions, and the difficulties underlying model-data fusion, would be addressed by using a freely accessible homogenized database (or set of databases) that can be updated with recent experiments or ancillary data (e.g., treatment magnitude, soil properties, climate regime), as necessary. With such an approach, especially when different original constituent databases are combined, fewer experiments are overlooked. This approach will improve the power of analyses, particularly in poorly sampled regions, and for rare combinations of global change drivers. It also facilitates the identification of drivers behind

contrasting outcomes and allows researcher (or data extractor) bias to be quantified when analyses explicitly account for different constituent databases.

Here, we present the MESI database, which represents the most comprehensive database of terrestrial ecosystem manipulation experiments to date. This database results from a coordinated intercontinental effort to combine four independently created constituent databases of elevated atmospheric CO₂ (eCO₂), warming, nutrient addition and precipitation manipulation experiments. The combined, freely accessible and machine-readable database (doi.org/10.5281/zenodo.7153253 – Van Sundert et al., 2022) helps bridge the experimental and modeling communities, facilitating the exploitation of the many opportunities such data syntheses have to offer.

2 | DESCRIPTION AND COMPARISON OF THE FOUR CONSTITUENT DATABASES

2.1 | General overview

The Manipulation Experiments Synthesis Initiative (MESI) was constructed as a new database building on four global change databases, previously compiled at the University of Antwerp, Sichuan Agricultural University, Hebei University, and the University of Alberta. All four constituent databases include ecosystem-, species- and individual-level responses (means, standard deviations, number of replicates; aggregation level specified through a separate column – Table 1) of mostly C and nutrient cycling to single and combined experimental manipulations of CO₂, temperature, nutrient and water availability, with wide coverage around the globe and across climate zones (Figs. 1 and S1). Data from 1,145 sites were collected from a variety of ecosystem types (e.g., grassland, forest, cropland, shrubland, desert, wetland, tundra) and mainly from field experiments (with natural or planted communities), but, depending on database-specific inclusion and exclusion criteria specified below, also from experiments conducted in greenhouses and growth chambers. Variables include treatment type, treatment magnitude (ppm CO₂, °C of warming etc.), response name and value, moderators (mean annual temperature and precipitation, age, ecosystem type, experiment type) and other ancillary data such as location coordinates, sampling dates and years (with multiple records for repeated measures), experiment start and end dates, dominant species, fumigation or warming type, citation, etc. (Table 1). A large majority of data were collected from the peer-reviewed literature. A smaller share of data is from unpublished sources (953 out of 57,089 records) and was provided by principal investigators at each study site. Data from the literature were extracted from supplementary information and from

data files accompanying respective publications, and from figures, tables and text, with various programs used for figure extraction (specified below per database).

Data from the constituent databases were combined into a single table, in comma-separated values (csv) format, with the specification of the original database for each row. Six additional tables are included within MESI: (i) metadata (explanation of the columns), (ii) description of response variable abbreviations, (iii) methodology of response variable measurements including further comments relevant to data interpretation, (iv) full references, (v) a template for new data contributions, and (vi) additional in situ determined soil data extracted from the original publications (soil texture type and particle size distribution, C:N ratio, organic matter concentration, pH), relevant to water and nutrient availability (Tolk, 2003; Van Sundert et al., 2018), ecosystem function (Vicca et al., 2018) and potentially responses thereof to global change (Cable et al., 2008; Canarini et al., 2017). The database is organized in a "wide" format. That is, observations made under two treatment levels are provided in the same row to facilitate data analysis (e.g., the calculation of response ratios) in a majority of cases. For multifactorial experiments, the value provided for the 'control' represents an "absolute" control, where no manipulation is applied. Therefore, responses of the values representing the control treatment are repeated in multiple rows. The long format (with separate rows per treatment including control, but no comparison per row) may be more practical for some applications. R code to translate between wide and long format is provided with the database at doi.org/10.5281/zenodo.7153253 (Van Sundert et al., 2022).

2.2 | The databases within MESI

2.2.1 | University of Antwerp database

With 47,196 observations, the database from the University of Antwerp contains more than 80% of the records in the MESI database. Included are data from 608 sites, originating from 1,120 publications collected over the last decade until 2021, using the Web of Science and Google Scholar search engines. Carbon cycle and nutrient cycle responses to global change factors (eCO₂, warming, N, P and K addition, drought, irrigation and respective combinations) were collected. C cycle responses include both pools (above- and belowground and total C and biomass, soil C, microbial biomass C etc.) and fluxes (net primary production, soil respiration, net ecosystem exchange etc.). Common nutrient cycle-related responses in the database are soil organic and inorganic N, plant tissue nutrient concentrations and stoichiometric ratios. Extensive background information about the site and the experiments is provided as well, including

ecosystem and community type, treatment start date, end date and intensity, experimental facility, MAT and MAP, dominant species, unit of the response variable and experiment duration. This database also comes with a separate table containing records of soil properties, measured at 365 experimental studies, providing information on soil texture (USDA classes, and % sand, silt and clay), soil organic matter and carbon concentrations, soil C:N ratios and pH. Data presented in figures were extracted with the Engauge Digitizer software (Free Software Foundation, Inc., Boston, MA, USA). Studies using earlier versions of the database, alone or in combination with other data, include Dieleman et al. (2010; 2012), Leuzinger et al. (2011), Terrer et al. (2016; 2019; 2021) and Ogle et al. (2021).

2.2.2 | Sichuan Agricultural University database

Yue et al. (2017) compiled a database of plant community, soil and microbial C pools in response to single and combined experimental eCO₂, warming, N addition, P addition, drought and irrigation. Data from 518 sites were extracted from 612 peer-reviewed publications found through the Web of Science and Google Scholar search engines in October 2016, resulting in 3,478 observations in the MESI database. Ancillary data in the original database were treatment intensity, ecosystem type, experimental facility, MAT and MAP (extracted from published text or the WorldClim database - http://:www.worldclim.org). For the MESI initiative, additional data were collected on community type (planted vs natural), dominant species, start and end date within seasons, unit of the response variables and USDA soil texture classes. Data presented in figures were extracted with the Engauge Digitizer software (Free Software Foundation, Inc., Boston, MA, USA).

2.2.3 | Hebei University database

The Hebei University database presents C stock and flux responses to single and combined eCO2, warming, N and N+P addition, drought, and irrigation. Data originate from 445 publications on outdoor experiments at 350 sites, and were collected from the Web of Science up to 13 December 2016 for the initial version, plus websites of experimental networks and ecological laboratories (Song et al., 2019; 2020). Cropland and indoor studies were excluded from this database. When available in the publication, MAT and MAP, the experiment duration, elevation, ecosystem type and treatment intensity were extracted along with response values. Where MAT and MAP were not available, Song et al. (2019) performed a look-up in Climate Model Intercomparison Project phase 5 data (CMIP5 – https://esgf-node.llnl.gov/projects/cmip5/). Within

the MESI initiative, we added to the database information on sampling dates, community type (natural vs planted), dominant species, start and end dates of treatments within seasons, and USDA soil texture classes. SigmaScan 5.0 (Systat Software Inc., San Jose, CA, USA) was used to extract means and standard deviations from figures. The database occupies 4,895 rows in the MESI database, including both unique and duplicate records overlapping with the other constituent databases.

2.2.4 | University of Alberta database

Ma et al. (2020) created a database on aboveground, belowground, plant organ and total biomass, and C stocks and production in response to multifactor global change. Data were collected from 115 peer-reviewed studies at 101 sites by searching the Web of Science and Google Scholar up to 1 August 2018. The inclusion criteria dictated that only results with mean, standard deviation and number of replicates were retained, and at least two of the global change factors eCO₂, warming, N addition, drought, irrigation or species richness were combined per experiment. Originally collected ancillary data were experiment duration (years), treatment intensity, ecosystem type, and unit of the response variable. In our effort to bring together the different databases, we added information on sampling date, community type (natural vs planted), dominant species, background climate (MAT, MAP), USDA soil texture classes, and start and end date of treatments within seasons (e.g., dates of precipitation exclusion). SigmaScanPro 5.0 (Systat Software Inc., San Jose, CA, USA) was used to extract means and standard errors or deviations from publications and supplemental figures. The database contributes 1,520 records to our MESI database.

2.3 | Comparison of the databases

A common feature among the original four databases is the worldwide coverage of experimental C cycle responses to global change in a variety of terrestrial ecosystem types (grassland, forest etc.) (Figs. 1 and S1). The main contrasts are in the type of C cycle variables collected, the collection of nutrient cycle responses and the availability of ancillary soil data. While the Sichuan and Alberta databases focused on C pools only, both pools and fluxes were collected for the Antwerp and Hebei databases. Only the database from the University of Antwerp has nutrient cycle responses and information on various soil properties beyond soil texture. Some differences also occur in the type of global change factors included: only the Alberta database explicitly

includes species richness as a manipulated factor (although lines with specified but always equal richness for control and treatment do occur in the Antwerp and Hebei databases), and only the Antwerp database contains data from full-factorial experiments of N, P, and K addition.

Complementarity among the databases exists not only in terms of the variables collected, but also in coverage of data from different sites (Figs. 1, 2 and S1). Comparing all experimental sites of all databases, irrespective of the manipulation or response variables, limited overlap exists among the four databases: 73%, or 841 out of 1145 sites are unique to one database (Fig. 2a). Complementarity among the four constituent databases is also evident when looking at aboveground biomass - the most common variable across the databases: 75%, or 386 out of 518 sites are unique to one database (Fig. 2b). This complementarity in sites emphasizes the value of our data integration effort and the potential for more powerful analyses using the combined dataset, emerging here from our MESI project.

3 | DISCUSSION

3.1 | Applications and examples

Earlier meta-analyses based on individual databases adopted specific scopes. For example, several studies have examined responses to simultaneous manipulation of multiple factors (Leuzinger et al., 2011; Dieleman et al., 2012; Ma et al., 2020), the role of moderators such as treatment duration and magnitude (Leuzinger et al., 2011; Ma et al., 2020), types of mycorrhizal symbionts present in investigated plots (Ainsworth et al., 2002; Terrer et al., 2016; 2021), and the influence of soil N and P on above- and belowground C cycle to (primarily) CO₂ (de Graaff et al., 2006; van Groenigen et al., 2006; Terrer et al., 2019 - Table 2). Notably, contrasting results were sometimes reported among studies using different constituent databases. For example, Dieleman et al. (2012) suggested, based on an earlier version of the University of Antwerp database, that antagonistic interactions between global change factors (e.g., CO₂ and warming) would be common, while Yue et al. (2017) found mostly additive effects in the Sichuan database. Such contrasting results should be explored further to unravel the cause of the discrepancy, e.g., the choice of global change drivers and response variables considered, the number and spatial spread of data points, or data interpretation and extraction methods while using the same primary literature.

Our comparison of the four global change databases points to their complementarity with respect to experimental sites covered (Fig. 2). While the original databases can be used separately,

Accepted Articl

combining the databases can reduce the uncertainty of ecosystem responses in understudied regions, where even a few additional data points represent a substantial increase in the available information given the scarcity of multifactor global change studies in some regions. Also, combining data from highly sampled regions can prove useful in tackling more detailed unresolved questions. For example, a broader combined dataset could allow researchers to test the role of background gradients in determining global change responses or identify whether researcher bias could have led to differences among databases. In this regard, we recommend analyses on the combined database, provided that the different origins of the four databases are taken into account. That is, data duplicates can be transparently handled, e.g., in a sensitivity analysis, thanks to the information provided in the MESI database, i.e., a column identifying 6938 potential duplicates that were flagged based on identical experiment name and response variables, and by using information such as dates of experiment and sampling, units etc. (Table 1).

Some studies have compared ecosystem model outputs against data from one or a few global change experiments (Zaehle et al., 2014; Wu et al., 2018; Paschalis et al., 2020). However, a lack of data integration and homogenization has thus far been an obstacle to true multi-experiment synthesis in combination with modeling. That is, comparing treatment effects on multiple response variables across many experiments to test hypotheses brought forward by models, comparing the performance of alternative model structures, reduce uncertainties, and parametrize based on the synthesis results (Keenan et al., 2011; LeBauer et al., 2013). Questions arising from models can further inform relevant new variables to measure in situ (Medlyn et al., 2015), and, based on uncertainty quantifications, on what variables should be sampled more intensively and on what locations or in which types of ecosystems (Dietze et al., 2013). In order to enable such modeldata synthesis, clear, quantitative information is required on the database side on treatments and their magnitudes, ancillary and response variables, and distinction among individual experiments and studies. Ideally, data from many related response variables are collected and reported in this way, such that hypotheses can be tested on, for example, why a model performs (apparently) well for one variable (e.g., NPP), but not for another (e.g., N uptake – Zaehle et al., 2014). MESI opens doors for data-model synthesis by providing such carefully homogenized, accessible, and integrated data.

Many large-scale studies have considered the role of background climate (especially MAT and MAP) in influencing global change responses. These studies often found a significant role in climate; for example, site aridity interacting with eCO₂ (Lu et al., 2016; also see De Kauwe et al., 2021). However, if such background gradients covary with other background variables that are

not considered, it may lead to incomplete or erroneous conclusions. One frequently neglected factor is nutrient availability or soil properties in general (Vicca et al., 2018). Soil properties influence nutrient availability (Van Sundert et al., 2018; 2020b; Du et al., 2020), such that part of the variation related to climate may actually be attributed to a gradient in nutrient availability (Vicca et al., 2012).

In order to facilitate the incorporation of soil and nutrient information in future database analyses, we collected new in situ measured data on various soil properties as part of the MESI effort. Soil texture data were collected for all constituent databases. The soil C:N ratio is available both as a response variable and as background soil information for many of the experiments. In addition, multiple soil properties such as organic matter (SOM) concentration and pH were provided with the University of Antwerp database. While it is well established that such soil properties and nutrient availability play a key role in the structure and functioning of ecosystems (Vitousek et al., 1991; Cleveland et al., 2011; Vicca et al., 2012; Van Sundert et al., 2020b), these are often ignored in meta-analyses, except for some studies on the role of nutrient availability gradients in response to eCO₂ (Terrer et al., 2019). The soil data we collected should thus enable the disentangling of the role of soil properties and nutrients in determining ecosystem responses to not only eCO₂, but also warming, nutrient addition and precipitation manipulation. Analyses spanning gradients in nutrient availability can then be compared to those from experiments manipulating nutrients in combination with other global change factors.

The standardized structure of the MESI database facilitates the addition of new records and moderators as new or supplementary data become available. Coupling to databases of other initiatives is also possible based on the experiment identifiers. Biodiversity, for instance, is not included as a global change factor in our set of constituent databases (except for some data on species richness), while aspects of biodiversity, such as plant community and functional group composition, species richness and evenness, are important determinants of productivity, the overall functioning of ecosystems (Hooper et al., 2005), and responses to climate extremes (Kreyling et al., 2017; Van Sundert et al., 2021a) and to gradual global change (Komatsu et al., 2019). Databases such as the Community Responses to Resource Experiments (CORRE – https://corredata.weebly.com/publications.html) database could be used to further unravel the role of biodiversity in large-scale ecosystem responses to global change. Additional leaf-level measurements, as well as more ecophysiological and hydraulic variables, are further possibilities for database extension.

The Alberta, Antwerp, Hebei, and Sichuan databases contain mean treatment responses with standard deviations to manipulated global changes at 1,145 sites and 3,644 experiments extracted from the scientific literature. This includes data from individual initiatives of site principal investigators, as well as published data from coordinated global change networks (Fraser et al., 2013) such as the Nutrient Network (NutNet – Borer et al., 2017) and DroughtNet (Knapp et al., 2017). However, more plot-level data exist for these networks that are not publicly available. Compared to our approach, such data from standardized networks are easier to compare across sites within the network and, therefore, easier to interpret. In NutNet, for instance, standardized quantities of N, P, K and micronutrients are added annually at all sites of the network (Borer et al., 2017). In DroughtNet, severe 1-in-100-year chronic drought is imposed by passively intercepting a site-specific percentage of precipitation (Yahdjian & Sala, 2002; Knapp et al., 2017). While such standardization facilitates cross-site comparisons of responses to a common driver in a common framework, data from coordinated distributed experiments alone do not cover the full range of available data from diverse global change experiments (e.g., acute droughts in different seasons). Where overlap exists in research questions addressable with both network and literature databases such as ours, results of both types of databases can be compared. Research questions that cannot be answered with network databases can be tackled with a unified database such as the one presented here.

3.2 | Remaining gaps in data coverage

MESI represents the most complete database of global change experiments to date in terms of studies, factorial combinations, ancillary data and response variables covered. Plotting the data across spatial, climatic and temporal dimensions provides insights into existing gaps in experimental coverage, and can help in deciding on future experimental locations, designs and sampling strategies.

Spatial, climatic and biome coverage

As reported in earlier studies (Martin et al., 2012), a substantial share of experimental sites are concentrated in temperate grasslands (n = 306) and forests (n = 150) of North America (n = 158), Europe (n = 137) and East Asia (n = 128) (Figs. 1 and S1), with sparser representation of the remaining geographical space. Particularly understudied are tropical rainforests, especially in the (African) paleotropics (n = 4), which potentially function differently than neotropical forests (Hubau et al., 2020). When distinguishing among global change manipulation types (Figs. 3 and S2), the low number of studies in the tropics becomes even more apparent for warming, eCO₂, and, to a

lesser degree, precipitation manipulation experiments. A substantial share of experiments in the tropics has focused on nutrient limitation, with particular emphasis on the role of P versus N (n = 20). In this regard, we note that, given the importance of, and uncertainty around, the tropical carbon sink and its responses to eCO₂ and climate change (Harris et al., 2021; Crezee et al., 2022; Okello et al., 2022), more global change experiments are being set up or have recently started, such as AmazonFACE that will investigate eCO₂ effects on a mature Amazonian rainforest (Fleischer et al., 2019). Also, specifically for croplands, only a few global change experiments are found in tropical regions (Fig. S3), despite the particularly important socio-economic role of the primary sector here, and the vulnerability of food security to climate change (Lobell et al., 2008). Tropical ecosystems are thus vastly understudied, whether natural, semi-natural or agricultural.

Warming experiments have been prioritized in the colder regions, i.e., boreal and tundra biomes (n = 64), as opposed to eCO₂ (n = 14) and precipitation manipulation experiments (n = 31). This prioritization logically follows from the observed and projected faster-than-global warming at high latitudes and elevations (Wang et al., 2016), and the often carbon-rich soils in these ecosystems may be susceptible to loss of carbon in a fast-warming climate, providing a potential positive feedback loop to the climate system (Cao & Woodward, 1998). In contrast to temperature, precipitation is considered non-limiting to biomass production in the colder regions (Bergh et al., 1999 - but see Nilsson, 1997), and most of these regions are becoming wetter (Box et al., 2019). Consequentially, fewer irrigation and drought studies have been performed here. However, recent extreme events (e.g., the 2018 European drought that impacted parts of Scandinavia – Buras et al., 2020) and publications suggest that at least in the southern fractions of the boreal biome, severe seasonal droughts may become more common because of changing circulation patterns (Mann et al., 2017), indicating relevance for experimental and other studies with focus on water availability also in boreal ecosystems.

Coverage of experiment duration

215 out of 693 studies with specified treatment duration in MESI are one-year experiments, or longer-term experiments from which only first-year data were reported and collected (Fig. 4a). While valuable, such short-term reports on global change impacts are prone to unstable initial responses (e.g., only initial soil carbon loss under warming - Verbrigghe et al., 2022), and effects may exhibit a multi-year lag because of gradual shifts in plant community composition (Langley & Megonigal, 2010), plant-soil feedbacks (Van Sundert et al., 2021b) etc. With increasing experiment duration, the share of single-factor experiments - especially fertilization studies - increases in MESI. The few longer-term multifactorial experiments in the database exemplify the

relevance of concurrent manipulation of global change drivers over longer time scales. For instance, some eCO₂ x N experiments in forests (Norby et al., 2010) and grasslands (Reich & Hobbie, 2013) exhibited a weakening of the CO₂ fertilization effect on NPP over time under ambient but not under elevated soil fertility. We recommend the further establishment of longer-term, bi- or multifactorial experiments that identify concurrent vs lagged, and direct vs indirect effects of global change on terrestrial ecosystems.

Coverage of multifactorial treatment combinations

Albeit generally of shorter duration, multifactorial experiments are quite common in the MESI database (30% with two or more factors manipulated – Fig. 4b). In these 466 multifactorial studies, croplands are relatively overrepresented with a share of 33% (154 studies), as opposed to single-factor experiments where 907 out of 1062 studies occurred in (semi-)natural grasslands, forests or shrublands. The underrepresentation of non-cropland multifactor studies is illustrated by the $eCO_2 x$ drought experiments in the database: out of 37 studies manipulating at least the CO_2 level and reducing precipitation, only 13 were in grassland, shrubland or forest. Such $eCO_2 x$ drought experiments in grasslands and forests are relevant for constraining models: despite long-known effects of eCO_2 on stomatal closure, much uncertainty still remains on under what circumstances (e.g., duration, atmospheric water demand) eCO_2 mitigates drought stress (De Kauwe et al., 2021). Such gaps identified by modeling, on key moderators influencing organism and ecosystem function, should more often inform the design of (multifactorial) experiments (here e.g., bifactorial, longer-term and with regression-design treatment levels – see also Collins et al., 2022) as well as what variables to monitor (here e.g., vapor pressure deficit, leaf area).

Coverage of (coupled) response variables

MESI describes a total of 262 response variables, of which 111 are related to carbon and 140 to nutrient cycling. The most commonly reported responses are above- and belowground biomass (AGB and BGB), soil respiration (R_SOIL) and soil organic carbon concentration (SOC) (Figs. 4c and 5). These variables have regularly been measured together to test hypotheses on plant carbon allocation (AGB vs BGB – Verlinden et al., 2018) and the ecosystem carbon balance (AGB + BGB vs SOC, R_SOIL – Terrer et al., 2021). MESI facilitates syntheses of such carbon allocation and balance studies to further disentangle context dependence of these coupled responses, for example, on how background gradients such as in climate and nutrient availability modify the relationships. While many studies present both AGB and BGB, more common reporting on both

soil (SOC) and whole-plant (AGB + BGB) carbon or dry mass at the same experiment would be useful to this end.

In general, no striking differences among studies manipulating different factors appear in terms of their reporting of commonly collected carbon pool, carbon flux and nitrogen pool responses (Figs. 5 and S4). One notable exception is a relative overrepresentation of R SOIL and an underrepresentation of SOC data at precipitation manipulation experiments: 40% of studies that report R SOIL manipulated the availability of water, alone or in combination with other factors, whereas for SOC this was only 19%. The most probable explanation for the preference of reporting soil-related flux rather than pool data is in the design of drought and irrigation experiments: many manipulate water availability only for a number of weeks within growing seasons, in contrast to usually 'continuous' nutrient addition, eCO2, and warming experiments. Because of the shorter duration of these events, chances of (at least short-term and first-year) significant changes in SOC are lower here than for the other manipulation types. Fluxes, on the other hand, often respond strongly during and shortly after imposed shifts in water availability (Van Sundert et al., 2020a). We recommend that experimentalists clearly report on sampling dates (before, during, or after experiment, average over manipulation period, or growing season etc.). In specified columns of our MESI database, we provide sampling dates and show the start and end of manipulation periods within the growing season (Table 1), such that immediate, lagged and seasonally averaged responses can be distinguished in analyses. Furthermore, a thorough assessment of the effect of precipitation changes on SOC would require more SOC data, from a greater diversity of experiments. This would make it more straightforward to verify if precipitation change impacts on SOC (if any) found in experiments are less pronounced than under naturally occurring deviations, analogous to what Kröel-Dulay et al. (2022) recently found for AGB.

3.3 | MESI as a dynamic database – data management, use and citation

Within the MESI initiative (Fig. 6), we follow the FAIR (Findable, Accessible, Interoperable, Reusable) and TRUST (Transparency, Responsibility, User focus, Sustainability, Technology) principles for data stewardship and repositories (Wilkinson et al., 2016; Lin et al., 2020; Kim et al., 2022). We host the MESI database on GitHub (github.com/MESI-organization/mesi-db), from where versions are managed, tagged and released to Zenodo (Van Sundert et al., 2022 - doi.org/10.5281/zenodo.7153253), under open access license CC-BY-4, meaning that the database can be freely used and edited, provided that the present study and the database at Zenodo are properly cited. We invite the research community to suggest updates to MESI through

pull requests and the issue tracker on GitHub, or by simply emailing the lead authors of the database and present study. Contributions from the community may include additional data and experiments, the combination of MESI with other databases of similar form and scope, or the highlighting of issues, gapfilling, and suggestions for improvements. Substantial contributions to the current version (v1.0.2) are acknowledged with co-authorship on the next citable Zenodo data release. Researchers intending to use the database for their own meta-analyses are particularly encouraged to suggest improvements to the MESI database, as questions emerge during the process of preparing a study, and new data may be collected. For reuse, it is relatively straightforward to perform basic meta-analyses, but we strongly advise researchers to carefully consider the exact meaning of the data, including particularities of individual experiments, the meaning and interpretation of variables, etc. Therefore, we encourage involving MESI team members in future studies to advise data use, processing, and interpretation.

4 | CONCLUSIONS

Our manipulation experiments synthesis initiative (MESI) addresses a key gap in global change research by providing a platform to store all past and future global change manipulation experimental results. This facilitates the synthesis of global response patterns in an unprecedented way, allowing updatable, dynamic information extraction using standardized protocols. We invite research teams around the globe concerned with meta-analyses of global change experiments to add their data to the MESI database on GitHub (github.com/MESI-organization/mesi-db; doi.org/10.5281/zenodo.7153253 - Van Sundert et al., 2022). We also propose that funding agencies consider the importance of supporting initiatives such as ours over long periods (decades) to ensure the continued curation of such overarching databases.

Acknowledgements

This initiative was made possible with support from Auckland University of Technology and iLEAPS.

KVS, SR and SV acknowledge support from the Fund for Scientific Research (FWO), Flanders (Belgium). KVS was further funded by the Belgian American Educational Foundation (BAEF) and the Fulbright Commission in Belgium and Luxembourg. MGDK acknowledges support from the Australian Research Council Discovery Grants (DP190101823, DP190102025). JS is supported by the National Natural Science Foundation of China (32101346). SQW is supported by the National Natural Science Foundation of China (31830012) and Hebei Natural Science Foundation (C2022201042). BDS was funded by the Swiss National Science Foundation grant PCEFP2_181115.

We wish to thank the following contributors to data collection: Jonas Torfs, Dries Vrijens, Oberon Geunens, Adrita Ballal, Thomas D'heer, Amélie De La Rocha, Gert-Jan Goeminne, Jaime Escobar, Robin Halffman, Lena Kuperus, Pieter Luys and Quinten Versmissen.

Several of our institutions sit on indigenous peoples' lands. These include but are not limited to MIT, Stanford University and Northern Arizona University. MIT acknowledges Indigenous Peoples as the traditional stewards of the land, and the enduring relationship that exists between them and their traditional territories. The land on which MIT sits is the traditional unceded territory of the Wampanoag Nation. We acknowledge the painful history of genocide and forced occupation of their territory, and we honor and respect the many diverse indigenous people connected to this land on which we gather from time immemorial. Stanford sits on the ancestral land of the Muwekma Ohlone Tribe. This land was and continues to be of great importance to the Ohlone people. Consistent with Stanford's values of community and inclusion, the university has a responsibility to acknowledge, honor, and make visible its relationship to Native peoples. Northern Arizona University sits at the base of the San Francisco Peaks, on homelands sacred to Native

Americans throughout the region. We honor their past, present, and future generations, who have lived here for millennia and will forever call this place home.

REFERENCES

Ainsworth, E. A., Davey, P. A., Bernacchi, C. J., Dermody, O. C., Heaton, E. A., Moore, D. J., ... Long, S. P. (2002). A meta-analysis of elevated [CO2] effects on soybean (Glycine max) physiology, growth and yield. Global Change Biology, 8, 695-709.

Bergh, J., Linder, S., Lundmark, T., & Elfving, B. (1999). The effect of water and nutrient availability on the productivity of Norway spruce in northern and southern Sweden. Forest Ecology and Management, 119, 51–62.

Borer, E. T., Grace, J. B., Harpole, W. S., MacDougall, A. S., & Seabloom, E. W. (2017). A decade of insights into grassland ecosystem responses to global environmental change. Nature Ecology & Evolution, 1, 118.

Box, J. E., Colgan, W. T., Christensen, T. R., Schmidt, N. M., Lund, M., Parmentier, F.-J. W., Brown, R., Bhatt, U. S., Euskirchen, E. S., & Romanovsky, V. E. (2019). Key indicators of Arctic climate change: 1971–2017. Environmental Research Letters, 14, 045010.

Buras, A., Rammig, A., & Zang, C. S. (2020). Quantifying impacts of the 2018 drought on European ecosystems in comparison to 2003. Biogeosciences, 17, 1655–1672.

Cable J. M., Ogle. K., Williams D. G., Weltzin, J. F., & Huxman, T. E. (2008). Soil texture drives responses of soil respiration to precipitation pulses in the Sonoran Desert: implications for climate change. Ecosystems, 11, 961–979.

Canarini, A., Kiær, L. P., & Dijkstra, F. A. (2017). Soil carbon loss regulated by drought intensity and available substrate: a meta-analysis. Soil Biology & Biochemistry, 112, 90–99.

Cao, M., & Woodward, F. I. (1998). Dynamic responses of terrestrial ecosystem carbon cycling to global climate change. Nature, 393, 249–252.

Cleveland, C. C., Townsend, A. R., Taylor, P., Alvarez-Clare, S., Bustamante, M. M. C. et al. (2011). Relationships among net primary productivity, nutrients and climate in tropical rain forest a pan-tropical analysis. Ecology Letters, 14, 939–947.

Collins, S., Whittaker, H., & Thomas, M. K. (2022). The need for unrealistic experiments in global change biology. Current Opinion in Microbiology, 68, 102151.

Crezee, B., Dargie, G. C., Ewango, C. E. N., Mitchard, E. T. A., B., O. E., T., J. K., Bola, P., Ndjango, J.-B. N., Girkin, N. T., Bocko, Y. E., Ifo, S. A., Hubau, W., Seidensticker, D., Batumike,

R., Imani, G., Cuní-Sanchez, A., Kiahtipes, C. A., Lebamba, J., Wotzka, H.-P., ... Lewis, S. L. (2022). Mapping peat thickness and carbon stocks of the central Congo Basin using field data. Nature Geoscience, 15, 639–644.

de Graaff, M. A., van Groenigen, K. J., Six, J., Hungate, B., & van Kessel, C. (2006). Interactions between plant growth and soil nutrient cycling under elevated CO2: a meta-analysis. Global Change Biology, 12, 2077–2091.

De Kauwe, M. G., Medlyn, B. E., & Tissue, D. T. (2021). To what extent can rising CO2 ameliorate plant drought stress? New Phytologist, 231, 2118–2124.

Dieleman, W. I. J., Luyssaert, S., Rey, A., de Angelis, P., Barton, C. V. M., Broadmeadow, M. S. J., Broadmeadow, S. B., Chigwerewe, K. S., Crookshanks, M., Dufrêne, E., Jarvis, P. G., Kasurinen, A., Kellomäki, S., Le Dantec, V., Liberloo, M., Marek, M., Medlyn, B., Pokorný, R., Scarascia-Mugnozza, G., ... Janssens, I. A. (2010). Soil [N] modulates soil C cycling in CO2-fumigated tree stands: a meta-analysis. Plant, Cell & Environment, 33, 2001–2011.

Dieleman, W. I. J., Vicca, S., Dijkstra, F. A., Hagedorn, F., Hovenden, M. J., Larsen, K. S., Morgan, J. A., Volder, A., Beier, C., Dukes, J. S., King, J., Leuzinger, S., Linder, S., Luo, Y., Oren, R., De Angelis, P., Tingey, D., Hoosbeek, M. R., & Janssens, I. A. (2012). Simple additive effects are rare: a quantitative review of plant biomass and soil process responses to combined manipulations of CO2 and temperature. Global Change Biology, 18, 2681–2693.

Dietze, M. C., Lebauer, D. S., & Kooper, R. (2013). On improving the communication between models and data. Plant, Cell & Environment, 36, 1575–1585.

Fleischer, K., Rammig, A., De Kauwe, M. G., Walker, A. P., Domingues, T. F., Fuchslueger, L., Garcia, S., Goll, D. S., Grandis, A., Jiang, M., Haverd, V., Hofhansl, F., Holm, J. A., Kruijt, B., Leung, F., Medlyn, B. E., Mercado, L. M., Norby, R. J., Pak, B., ... Lapola, D. M. (2019). Amazon forest response to CO2 fertilization dependent on plant phosphorus acquisition. Nature Geoscience, 12: 736–741.

Fraser, L. H., Henry, H. A. L., Carlyle, C. N., White, S. R., Beierkuhnlein, C., Cahill, J. F., Casper, B. B., Cleland, E., Collins, S. L., Dukes, J. S., Knapp, A. K., Lind, E., Long, R., Luo, Y., Reich, P. B., Smith, M. D., Sternberg, M., & Turkington, R. (2013). Coordinated distributed experiments: an emerging tool for testing global hypotheses in ecology and environmental science. Frontiers in Ecology and the Environment, 11, 147–155.

Gao, C.-H., Yu, G., & Cai, P. (2021). ggVennDiagram: An intuitive, easy-to-use, and highly customizable R package to generate Venn diagram. Frontiers in Genetics, 12, 1598.

Harris, N. L., Gibbs, D. A., Baccini, A., Birdsey, R. A., de Bruin, S., Farina, M., Fatoyinbo, L., Hansen, M. C., Herold, M., Houghton, R. A., Potapov, P. V., Suarez, D. R., Roman-Cuesta, R. M., Saatchi, S. S., Slay, C. M., Turubanova, S. A., & Tyukavina, A. (2021). Global maps of twenty-first century forest carbon fluxes. Nature Climate Change, 11, 234–240.

Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P. et al. (2005). Effects of biodiversity on ecosystem functioning: a consensus of current knowledge. Ecological Monographs, 75, 3-35.

Hubau, W., Lewis, S. L., Phillips, O. L., Affum-Baffoe, K., Beeckman, H., Cuní-Sanchez, A., Daniels, A. K., Ewango, C. E. N., Fauset, S., Mukinzi, J. M., Sheil, D., Sonké, B., Sullivan, M. J.

P., Sunderland, T. C. H., Taedoumg, H., Thomas, S. C., White, L. J. T., Abernethy, K. A., Adu-Bredu, S., ... Zemagho, L. (2020). Asynchronous carbon sink saturation in African and Amazonian tropical forests. Nature, 579, 80–87.

Keenan, T. F., Carbone, M. S., Reichstein, M., & Richardson, A. D. (2011). The model-data fusion pitfall: assuming certainty in an uncertain world. Oecologia, 167, 587–597.

Kim, A. Y., Herrmann, V., Bareto, R., Calkins, B., Gonzalez-Akre, E., Johnson, D. J., Jordan, J. A., Magee, L., McGregor, I. R., Montero, N., Novak, K., Rogers, T., Shue, J., & Anderson-Teixeira, K. J. (2022). Implementing GitHub Actions continuous integration to reduce error rates in ecological data collection. Methods in Ecology and Evolution.

Knapp, A. K., Avolio, M. L., Beier, C., Carroll, C. J. W., Collins, S. L., Dukes, J. S., Fraser, L. H., Griffin-Nolan, R. J., Hoover, D. L., Jentsch, A., Loik, M. E., Phillips, R. P., Post, A. K., Sala, O. E., Slette, I. J., Yahdjian, L., & Smith, M. D. (2017). Pushing precipitation to the extremes in distributed experiments: recommendations for simulating wet and dry years. Global Change Biology, 23, 1774–1782.

Komatsu, K. J., Avolio, M. L., Lemoine, N. P., Isbell, F., Grman, E., Houseman, G. R., Koerner, S. E., Johnson, D. S., Wilcox, K. R., Alatalo, J. M., Anderson, J. P., Aerts, R., Baer, S. G., Baldwin, A. H., Bates, J., Beierkuhnlein, C., Belote, R. T., Blair, J., Bloor, J. M. G., ... Zhang, Y. (2019). Global change effects on plant communities are magnified by time and the number of global change factors imposed. Proceedings of the National Academy of Sciences of the United States of America, 116, 17867–17873.

Kreyling, J., Dengler, J., Walter, J., Velev, N., Ugurlu, E., Sopotlieva, D., Ransijn, J., Picon-Cochard, C., Nijs, I., Hernandez, P., Güler, B., von Gillhaussen, P., De Boeck, H. J., Bloor, J. M. G., Berwaers, S., Beierkuhnlein, C., Arfin, M. A., Apostolova, I., Altan, Y., ... Jentsch, A. (2017). Species richness effects on grassland recovery from drought depend on community productivity in a multisite experiment. Ecology Letters, 20, 1405–1413.

Kröel-Dulay, G., Mojzes, A., Szitár, K., Bahn, M., Batáry, P., Beier, C., Bilton, M., De Boeck, H. J., Dukes, J. S., Estiarte, M., Holub, P., Jentsch, A., Schmidt, I. K., Kreyling, J., Reinsch, S., Larsen, K. S., Sternberg, M., Tielbörger, K., Tietema, A., Vicca, S., & Peñuelas, J. (2022). Field experiments underestimate aboveground biomass response to drought. Nature ecology & evolution, 6, 540–545.

Langley, J. A., & Megonigal, P. J. (2010). Ecosystem response to elevated CO2 levels limited by nitrogen-induced plant species shift. Nature, 466, 96–99.

LeBauer, D. S., Wang, D., Richter, K. T., Davidson, C. C., & Dietze, M. (2013). Facilitating feedbacks between field measurements and ecosystem models. Ecological Monographs, 83, 133–154.

Leuzinger, S., Luo, Y., Beier, C., Dieleman, W., Vicca, S., & Körner, C. (2011). Do global change experiments overestimate impacts on terrestrial ecosystems? Trends in Ecology & Evolution, 26, 236–241.

Lin, D., Crabtree, J., Dillo, I., Downs, R. R., Edmunds, R., Giaretta, D., De Giusti, M., L'Hours, H., Hugo, W., Jenkyns, R., Khodiyar, V., Martone, M. E., Mokrane, M., Navale, V., Petters, J.,

Sierman, B., Sokolova, D. V., Stockhause, M., & Westbrook, J. (2020). The TRUST Principles for digital repositories. Scientific Data, 7, 1.

Lobell, D. B., Burke, M. B., Tebaldi, C., Mastrandrea, M. D., Falcon, W. P., & Naylor, R. L. (2008). Prioritizing climate change adaptation needs for food security in 2030. Science, 319, 607–610.

Lu, X., Wang, L., & McCabe, M. F. (2016). Elevated CO2 as a driver of global dryland greening. Scientific Reports, 6, 1.

Ma, Z., Chen, H. Y. H., Li, Y., & Chang, S. X. (2020). Interactive effects of global change factors on terrestrial net primary productivity are treatment length and intensity dependent. Journal of Ecology, 108, 2083–2094.

Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K., & Coumou, D. (2017). Influence of anthropogenic climate change on planetary wave resonance and extreme weather events. Scientific Reports, 7, 45242.

Martin, L. J., Blossey, B., & Ellis, E. (2012). Mapping where ecologists work: biases in the global distribution of terrestrial ecological observations. Frontiers in Ecology and the Environment, 10, 195–201.

Medlyn, B. E., Zaehle, S., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hanson, P. J., Hickler, T., Jain, A. K., Luo, Y., Parton, W., Colin Prentice, I., Thornton, P. E., Wang, S., Wang, Y.-P., Weng, E., Iversen, C. M., McCarthy, H. R., Warren, J. M., Oren, R., & Norby, R. J. (2015). Using ecosystem experiments to improve vegetation models. Nature Climate Change, 5, 528–534.

Nilsson, L.-O. (1997). Manipulation of conventional forest management practices to increase forest growth—results from the Skogaby project. Forest Ecology and Management, 91, 53–60.

Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E., & McMurtrie, R. E. (2010). CO2 enhancement of forest productivity constrained by limited nitrogen availability. Proceedings of the National Academy of Sciences of the United States of America, 107, 19368–19373.

Ogle, K., Liu, Y., Vicca, S., & Bahn, M. (2021). A hierarchical, multivariate meta-analysis approach to synthesising global change experiments. New Phytologist, 231, 2382–2394.

Okello, J., Bauters, M., Verbeeck, H., Kasenene, J., & Boeckx, P. (2022). Response of Afromontane soil organic carbon, nitrogen, and phosphorus to in situ experimental warming along an elevational gradient. Frontiers in Soil Science, 2.

Paschalis, A., Fatichi, S., Zscheischler, J., Ciais, P., Bahn, M., Boysen, L., Chang, J., De Kauwe, M., Estiarte, M., Goll, D., Hanson, P. J., Harper, A. B., Hou, E., Kigel, J., Knapp, A. K., Larsen, K. S., Li, W., Lienert, S., Luo, Y., ... Zhu, Q. (2020). Rainfall manipulation experiments as simulated by terrestrial biosphere models: Where do we stand? Global Change Biology, 26, 3336–3355.

Reich, P. B., & Hobbie, S. E. (2013). Decade-long soil nitrogen constraint on the CO2 fertilization of plant biomass. Nature Climate Change, 3, 278–282.

Song, J., Ru, J., Zheng, M., Wang, H., Fan, Y., Yue, X., Yu, K., Zhou, Z., Shao, P., Han, H., Lei, L., Zhang, Q., Li, X., Su, F., Zhang, K., & Wan, S. (2020). A global database of plant production and carbon exchange from global change manipulative experiments. Scientific Data, 7, 323.

Song, J., Wan, S., Piao, S., Knapp, A. K., Classen, A. T., Vicca, S., Ciais, P., Hovenden, M. J., Leuzinger, S., Beier, C., Kardol, P., Xia, J., Liu, Q., Ru, J., Zhou, Z., Luo, Y., Guo, D., Adam Langley, J., Zscheischler, J., ... Zheng, M. (2019). A meta-analysis of 1,119 manipulative experiments on terrestrial carbon-cycling responses to global change. Nature Ecology & Evolution, 3, 1309–1320.

Terrer, C., Jackson, R. B., Prentice, I. C., Keenan, T. F., Kaiser, C. et al. (2019). Nitrogen and phosphorus constrain the CO2 fertilization of global plant biomass. Nature Climate Change, 9, 684-+.

Terrer, C., Philips, R. P., Hungate, B. A., Rosende, J., Pett-Ridge, J. et al. (2021). A trade-off between plant and soil carbon storage under elevated CO2. Nature, 591, 599–603.

Terrer, C., Vicca, S., Hungate, B. A., Phillips, R. P., & Prentice, I. C. (2016). Mycorrhizal association as a primary control of the CO₂ fertilization effect. Science, 353, 72–74.

Tolk, J. A. (2003). Plant available soil water. In: Stewart, B. A., Howell, T. A., eds. Encyclopedia of water science. New York, USA: Marcel-Dekker.

Valentin, S., & Levin, S. (2022). Plotbiomes: plot Whittaker biomes with ggplot2. R package version 0.0.0.9001.

van Groenigen, K. J., Six, J., Hungate, B. A., de Graaff, M. A., van Breemen, N., & van Kessel, C. (2006). Element interactions limit soil carbon storage. Proceedings of the National Academy of Sciences of the United States of America, 103, 6571–6574.

Van Sundert, K., Arfin Khan, M. A. S., Bharath, S., Buckley, Y. M., Caldeira, M. C. et al. (2021a). Fertilized graminoids intensify negative drought effects on grassland productivity. Global Change Biology, 27, 2441–2457.

Van Sundert, K., Brune, V., Bahn, M., Deutschmann, M., Hasibeder, R., Nijs, I., & Vicca, S. (2020a). Post-drought rewetting triggers substantial K release and shifts in leaf stoichiometry in managed and abandoned mountain grasslands. Plant and Soil, 448, 353–368.

Van Sundert, K., Horemans, J. A., Stendahl, J., & Vicca, S. (2018). The influence of soil properties and nutrients on conifer forest growth in Sweden, and the first steps in developing a nutrient availability metric. Biogeosciences, 15, 3475–3496.

Van Sundert, K., Leuzinger, S., Bader, M. K.-F., Chang, S. X., De Kauwe, M. G., Dukes, J. S., Langley, J. A., Ma, Z., Mariën, B., Reynaert, S., Ru, J., Song, J., Stocker, B., Terrer, C., Thoresen, J., Vanuytrecht, E., Wan, S., Yue, K., & Vicca, S. (2022). MESI: a database of terrestrial global change experiments. doi.org/10.5281/zenodo.7153253

Van Sundert, K., Linder, S., Marshall, J. D., Nordin, A., & Vicca, S. (2021b). Increased tree growth following long-term optimised fertiliser application indirectly alters soil properties in a boreal forest. European Journal of Forest Research, 140, 241–254.

Van Sundert, K., Radujković, D., Cools, N., De Vos, B., Etzold, S., Fernández-Martínez, M., Janssens, I. A., Merilä, P., Peñuelas, J., Sardans, J., Stendahl, J., Terrer, C., & Vicca, S. (2020b). Towards comparable assessment of the soil nutrient status across scales-Review and development of nutrient metrics. Global Change Biology, 26, 392–409.

Verbrigghe, N., Leblans, N. I. W., Sigurdsson, B. D., Vicca, S., Fang, C., Fuchslueger, L., Soong, J. L., Weedon, J. T., Poeplau, C., Ariza-Carricondo, C., Bahn, M., Guenet, B., Gundersen, P., Gunnarsdóttir, G. E., Kätterer, T., Liu, Z., Maljanen, M., Marañón-Jiménez, S., Meeran, K., ... Janssens, I. A. (2022). Soil carbon loss in warmed subarctic grasslands is rapid and restricted to topsoil. Biogeosciences, 19, 3381–3393.

Verlinden, M. S., Ven, A., Verbruggen, E., Janssens, I. A., Wallander, H., & Vicca, S. (2018). Favorable effect of mycorrhizae on biomass production efficiency exceeds their carbon cost in a fertilization experiment. Ecology, 99, 2525–2534.

Vicca, S., Luyssaert, S., Peñuelas, J., Campioli, M., Chapin, F. S., 3rd, Ciais, P., Heinemeyer, A., Högberg, P., Kutsch, W. L., Law, B. E., Malhi, Y., Papale, D., Piao, S. L., Reichstein, M., Schulze, E. D., & Janssens, I. A. (2012). Fertile forests produce biomass more efficiently. Ecology Letters, 15, 520–526.

Vicca, S., Stocker, B. D., Reed, S., Wieder, W. R., Bahn, M., Fay, P. A., Janssens, I. A., Lambers, H., Peñuelas, J., Piao, S., Rebel, K. T., Sardans, J., Sigurdsson, B. D., Van Sundert, K., Wang, Y.-P., Zaehle, S., & Ciais, P. (2018). Using research networks to create the comprehensive datasets needed to assess nutrient availability as a key determinant of terrestrial carbon cycling. Environmental Research Letters, 13, 125006.

Vitousek, P. M. & Howarth, R. W. (1991). Nitrogen limitation on land and in the sea: how can it occur? Biogeochemistry, 13, 87–115.

Wang, Q., Fan, X., & Wang, M. (2016). Evidence of high-elevation amplification versus Arctic amplification. Scientific Reports, 6, 19219.

Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., ... Mons, B. (2016). The FAIR Guiding Principles for scientific data management and stewardship. Scientific Data, 3, 160018.

Whittaker, R. H. (1970). Communities and Ecosystems. New York, USA: Macmillan.

Wu, D., Ciais, P., Viovy, N., Knapp, A. K., Wilcox, K., Bahn, M., Smith, M. D., Vicca, S., Fatichi, S., Zscheischler, J., He, Y., Li, X., Ito, A., Arneth, A., Harper, A., Ukkola, A., Paschalis, A., Poulter, B., Peng, C., ... Piao, S. (2018). Asymmetric responses of primary productivity to altered precipitation simulated by ecosystem models across three long-term grassland sites. Biogeosciences, 15, 3421–3437.

Yahdjian, L., & Sala, O. E. (2002). A rainout shelter design for intercepting different amounts of rainfall. Oecologia, 133, 95–101.

Yue, K., Fornara, D. A., Yang, W., Peng, Y., Peng, C., Liu, Z., & Wu, F. (2017). Influence of multiple global change drivers on terrestrial carbon storage: additive effects are common. Ecology Letters, 20, 663–672.

Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., Luo, Y., Wang, Y.-P., El-Masri, B., Thornton, P., Jain, A., Wang, S., Warlind, D., Weng, E., Parton, W., Iversen, C. M., Gallet-Budynek, A., McCarthy, H., Finzi, A., ... Norby, R. J. (2014). Evaluation of

11 terrestrial carbon-nitrogen cycle models against observations from two temperate Free-Air CO2 Enrichment studies. New Phytologist, 202, 803–822.

TABLE 1 Content description of the combined MESI database.

able category Field name Description

	db id duplicate_id citation	Database name (antwerp, sichuan, hebei, alberta) Unique identifier per data point As id, but common identifier for data points potentially shared among databases based on identical experiment name 'exp' and response variable name 'response' Reference from which data were extracted
Site characterization	site study exp lat lon elevation mat map ecosystem_type vegetation_type experiment_type community_type dominant_species growth_form age disturbance_type	Site name (e.g., euroface) Individual study with one or more manipulation types (e.g., euroface_populusalba) Individual experiment with manipulation (e.g., euroface_populusalba_cf) Latitude (°, negative for south, positive for north) Longitude (°, negative for west, positive for east) m a.s.l. M ean annual temperature (°C) M ean annual precipitation (mm) e.g., grassland, cropland, forest, desert e.g., meadow steppe, serpentine grassland, humid tropical forest e.g., field, greenhouse, outdoor chamber Natural, planted Latin genus + species name(s) Herbaceous, woody Ecosystem age in years (years since planting or disturbance) M owing, tilling, defoliation, grazing, fire (e.g., 10000 = mowing only)
Treatment characterization	treatment npk w_t1	Global change factor(s) manipulated (e.g., $c = CO_2$, $w = warming$, $cw = CO_2 + warming$) N, P and or K addition (e.g., $100 = N$ only, $010 = P$ only, $011 = P$ and K) Warming method: soil, air, open top chamber, infrared (e.g., $1000 = soil warming only)$
Treatment details	$\begin{array}{c} c_{-}c \\ c_{-}t \\ d_{-}t \\ d_{-}t \\ n_{-}c \\ n_{-}c \\ n_{-}c \\ p_{-}t \\ k_{-}c \\ k_{-}t \\ i_{-}c \\ k_{-}t \\ i_{-}c \\ k_{-}t \\ i_{-}t \\ i_{-}t \\ s_{-}c \\ s_{-}t \\ s_{-}c \\ s_{-}t \\ s_{-}c \\ s_{-}t \\ w_{-}t \\ s_{-}c \\ s_{-}t \\ s_{-}c \\ s_{-}t \\ w_{-}t \\ s_{-}c \\ s_{-}t \\ s_{-}$	CO ₂ concentration of control treatment (ppm) CO ₂ concentration of CO ₂ addition treatment (ppm) Targeted precipitation reduction during experimental drought (e.g., $0.6 = 60\%$ reduction) Precipitation exclusion during drought treatment or per year for continuous treatment (mm) N addition in control treatment (g N/m ² y) P addition in treatment (g N/m ² y) P addition in treatment (g P/m ² y) K addition in control treatment (g K/m ² y) K addition in treatment (g K/m ² y) Water addition in control treatment of irrigation experiment (mm/d) Water addition during irrigation treatment of irrigation experiment (mm/d) Targeted treatment irrigation relative and in addition to control (e.g., $0.6 = +60\%$ addition) Species richness of control treatment Air warming (°C) Year when experimental treatment began Number of years the experiment was running Days between start and end of treatment (e.g., drought period) CO ₂ addition method (FACE, greenhouse, OTC, SACC, tunnels) Start of treatment (d/m/yyyy or continuous)

TABLE 1 (Cont	tinued; colors	represent	different	database	tables))
---------------	----------------	-----------	-----------	----------	---------	---

category Field name Description

M easurement info	response sampling_year sampling_depth aggregation_level x_c x_t x_units sd_c sd_t se_c se_t rep_c rep_t sampling_date	Response variable (e.g., anpp, bnpp, leaf_n, mineral_soil_cn) Year of sampling from start of experiment (1 = in start year, 2 = first year after start) Depth of soil sampling (cm) Representative of plot-level (community) or for species within a community (species) Response value of control treatment Response value of global change treatment Unit of response value Standard deviation of control response value Standard deviation of treatment response value Standard error of control response value Standard error of treatment response value Number of control replicates Number of treatment replicates Date of sampling in d/m/yyyy
	bibliography	Table with full references
5	soil	Table with soil properties as ancillary data
	metadata	Table with an explanation of columns
	response variable abbreviations	Table with response variable abbreviations of the main table explained
	methods_comments	Table with methodological information on how response variables were measured per data point, and further comments for data interpretation
1	template	Empty table for data contributions, with main table headers

MESI constituent database	Example research topics
Antwerp, Sichuan, Hebei, Alberta: analyses on single and combined databases	 Single and multifactor global change effects on C cycling^{1,2,3,4} M oderating role of background climate gradients, ecosystem type, treatment duration and magnitude^{1,5} M odel-data synthesis
Antwerp	 Role of soil and nutrients in global change responses^{6,7,8} Global change effects on leaf, plant and soil stoichiometry Comparison of global change responses among spatial scales (e.g., leaf vs ecosystem⁵)
Alberta	- Influence of species richness alone on C cycling, and in combination with global change factors ¹
Expansions (in combination with other databases)	 Role of biodiversity, incl. species composition, in determining responses to global change⁹ Comparison against data from global change experimental networks Comparison of terrestrial vs aquatic vs marine global change responses
 ¹ Ma et al., 2020 ² Dieleman et al., 2012 ³ Song et al., 2019 ⁴ Yue et al., 2017 ⁵ Leuzinger et al., 2011 ⁶ Terrer et al., 2016 ⁷ Terrer et al., 2019 ⁸ Terrer et al., 2021 	

⁹ CORRE database: https://corredata.weebly.com/publications.html

FIGURE 1 Spatial (a) and climatic (b) distribution of global change manipulation experimental sites in databases of the Universities of Antwerp (n = 608 sites), Sichuan (n = 518 sites), Hebei (n = 350sites) and Alberta (n = 101 sites). The Whittaker biome plot (Whittaker, 1970) was created using the R package plotbiomes (Valentin & Levin, 2022). MAT = mean annual temperature, MAP = mean annual precipitation.

FIGURE 2 Site overlap among the global change constituent databases. Panel a was based on all variables and treatments in the databases, panel b was based on all treatments but aboveground biomass only. Diagrams were made using the ggVenndiagram R package (Gao et al., 2021).

FIGURE 3 Spatial and climatic distribution of 1145 global change manipulation experimental sites per manipulation type. Whittaker biome plots (Whittaker, 1970) were created using R package plotbiomes (Valentin & Levin, 2022). MAT = mean annual temperature, MAP = mean annual precipitation.

FIGURE 4 Distribution of experiment duration (n = 693), factors investigated (n = 1528), and some commonly measured responses (n = 1128) across studies in MESI. For display purposes, precipitation manipulation in panel b refers to both irrigation and drought experiments. $c = eCO_2$, d = drought, f = fertilization, w = warming, i = irrigation, AGB = aboveground biomass, BGB = belowground biomass, R_SOIL = soil respiration, SOC = soil organic carbon concentration.

FIGURE 5 Availability of some commonly measured C cycle pool, flux and N pool responses across 1536 unique studies in MESI, stratified by factors that were manipulated. $c = eCO_2$, d = drought, f =fertilization, w = warming, i = irrigation, AGB = aboveground biomass, BGB = belowground biomass, SOC = soil organic C concentration, MBC = microbial biomass C, ANPP = aboveground net primary production, BNPP = belowground net primary production, R_SOIL = soil respiration, NEP = net ecosystem production, SOIL_TN = soil total N concentration, SOIL_iN = soil inorganic N (N-NH4⁺ and N-NO3⁻), AGB_N = aboveground plant N concentration, BGB_N = belowground plant N concentration, LEAF_N = mass-based leaf N concentration, MBN = microbial biomass N.

FIGURE 6 Protocol for MESI data maintenance, use and citation, including a non-exhaustive list of suggestions for updates and data uses.







Boreal forest 📕 Temperate rain forest

Tropical seasonal forest/savanna 📒 Temperate grassland/desert







