

Pesonen Maria (Orcid ID: 0000-0003-0356-7064)
Caroppo Francesca (Orcid ID: 0000-0003-3583-0816)
Larese Filon Francesca (Orcid ID: 0000-0002-7717-0417)
Geier Johannes (Orcid ID: 0000-0002-5047-8948)
Giménez-Arnau Ana M (Orcid ID: 0000-0001-5434-7753)
Napolitano Maddalena (Orcid ID: 0000-0003-3309-8190)
Simon Dagmar (Orcid ID: 0000-0001-8965-9407)
Schuttelaar Marie Louise Anna (Orcid ID: 0000-0002-0766-4382)
Spiewak Radoslaw (Orcid ID: 0000-0001-5968-0555)
Stingeni Luca (Orcid ID: 0000-0001-7919-8141)
Wilkinson Mark (Orcid ID: 0000-0001-7253-3461)
Uter Wolfgang (Orcid ID: 0000-0002-4498-3710)
Bauer Andrea (Orcid ID: 0000-0002-4411-3088)

Occupational contact allergy: The European perspective – analysis of patch test data from ESSCA between 2011-2020

Corresponding author

Univ.-Prof. Dr. med. Andrea Bauer, MPH
Department of Dermatology, University Hospital Carl Gustav Carus
Technical University Dresden
Fetscherstr. 74
01307 Dresden
Germany
e-mail: Andrea.Bauer@ukdd.de

Conflict of Interests

W.U. has accepted travel reimbursement and research funds from the cosmetic industry association IFRA. T.R. has participated in patch test studies from SmartPractice and has given lectures for ALK Abello, Leo Pharma, Novartis, Sanofi Genzyme, UCB Pharma and Ypsomed. All other authors have no pertinent conflict of interests to declare.

Andrea Bauer (ORCID-ID: 0000-0002-4411-3088), Department of Dermatology, University Hospital Carl Gustav Carus, Technical University, Dresden, Germany

Maria Pesonen (ORCID-ID: 0000-0003-0356-7064), Occupational Health Unit, Finnish Institute of Occupational Health, Helsinki, Finland

Richard Brans (ORCID-ID: 0000-0002-1245-024X), Department of Dermatology, Environmental Medicine and Health Theory, University of Osnabrück, Osnabrück, Germany

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1111/cod.14280](https://doi.org/10.1111/cod.14280)

and Institute for Interdisciplinary Dermatologic Prevention and Rehabilitation (iDerm) at the University of Osnabrück, Osnabrück, Germany

Francesca Caroppo (ORCID-ID: 0000-0003-3583-0816), Unit of Clinic Dermatology, Department of Medicine DIMED, University of Padova, Italy

Heinrich Dickel (Orcid-ID:0000-0001-6837-4402), Department of Dermatology, Venereology and Allergology, St. Josef Hospital, University Medical Center, Ruhr University Bochum, Bochum, Germany

Aleksandra Dugonik (orcid.org/0000-0001-8433-7716) Department of Dermatology, University Medical Centre, Maribor, Slovenia

Francesca Larese Filon (ORCID-ID: 0000-0002-7717-0417), Department of Public Health, Occupational Medicine, University of Trieste, Trieste, Italy

Johannes Geier (Orcid-ID:0000-0002-5047-8948) Information Network of Departments of Dermatology (IVDK), Institute at the University Medical Center Göttingen, Göttingen, Germany

Ana M. Gimenez-Arnau (Orcid-ID:0000-0001-5434-7753) Department of Dermatology, Hospital del Mar, Barcelona, Spain

Maddalena Napolitano (ORCID-ID: 0000-0003-3309-8190), Department of Medicine and Health Sciences Vincenzo Tiberio, University of Molise, Campobasso, Italy.

Cataldo Patruno (ORCID-ID: 0000-0001-8661-2822) Department of Health Sciences, University Magna Graecia of Catanzaro, Catanzaro, Italy

Thomas Rustemeyer (Orcid-ID: 0000-0001-7580-0684), Department of Dermatology and Allergology, Amsterdam University Medical Centers, Amsterdam, The Netherlands

Dagmar Simon (Orcid-ID:0000-0001-8965-9407), Department of Dermatology, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

Marie L.A. Schuttelaar (Orcid-ID:0000-0002-0766-4382), Department of Dermatology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

Radoslaw Spiewak (Orcid-ID:0000-0001-5968-0555), Department of Experimental Dermatology and Cosmetology, Jagiellonian University Medical College, Krakow, Poland.

Luca Stingeni (ORCID-ID: 0000-0001-7919-8141), Dermatology Section, Department of Medicine and Surgery, University of Perugia, Italy

Marko Vok (ORCID: 0000-0002-8912-1233), Dermatovenerološka ambulanta, Izola, Slovenia

Elke Weisshaar (Orcid-ID: 0000-0002-7016-0224), Occupational Dermatology, Department of Dermatology, University Hospital Heidelberg, Heidelberg, Germany

Mark Wilkinson (ORCID-ID: 0000-0001-7253-3461), Leeds Teaching Hospitals NHS Trust, Leeds, UK

Skaidra Valiukeviciene (ORCID-ID: 0000-0003-1219-5928), Department of Skin and Venereal Diseases, Lithuanian University of Health Sciences, Kaunas, Lithuania, Kaunas, Lithuania

Wolfgang Uter (ORCID-ID: 0000-0002-4498-3710), Department of Medical Informatics, Biometry and Epidemiology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany

Collaborators:

Birger Kränke, Department of Dermatology, Medical University of Graz, Graz, Austria, Graz, Austria (birger.kraenke@medunigraz.at); Kathrin Scherer Hofreiter, Department of Dermatology, Allergy Unit, University Hospital Basel, Basel, Switzerland, Basel, Switzerland (Kathrin.scherer@usb.ch); Alexander Navarini, NA, Basel, Switzerland (alexander.navarini@usb.ch); Jürgen Grabbe, Department of Dermatology, Kantonsspital Aarau, Aarau, Switzerland, Aarau, Switzerland (juergen.grabbe@ksa.ch); Philip Spring, Dermatologie et vénéréologie FMH, Center Médical d'Epalinges, Epalinges, Switzerland, Lausanne/Epalinges, Switzerland (springphil@hotmail.com); Ulrike Beiteke, Department of Dermatology, Dortmund, Germany, Dortmund, Germany (ulrike.beiteke@klinikumdo.de); Timo Buhl, Department of Dermatology, University Medicine, Göttingen, Germany, Göttingen, Germany (timo.buhl@med.uni-goettingen.de); Cecilia Dietrich, Department of Dermatology, University of Schleswig-Holstein, Campus Kiel, Kiel, Germany, Kiel, Germany (cdietrich@dermatology.uni-kiel.de); Sibylle Schliemann, Department of Dermatology and Allergology, University Hospital Jena, Jena, Germany, Jena, Germany (schliemann@derma-jena.de); Detlef Becker, Department of Dermatology, University of Mainz, Mainz, Germany, Mainz, Germany (detlef.becker@unimedizin-mainz.de); Swen Malte John, Institute for Interdisciplinary Dermatologic Prevention and Rehabilitation (iDerm), University of Osnabrück, Lower Saxony Institute for Occupational Dermatology (NIB), Osnabrück, Germany, Osnabrück, Germany (sjohn@uni-osnabrueck.de); Nicola Wagner, Department of Dermatology, University of Erlangen-Nürnberg, Erlangen, Germany, Erlangen, Germany (nicola.wagner@uk-erlangen.de); Thomas Werfel, Medizinische Hochschule Hannover, Department of Immunodermatology and experimental Allergy, Hannover, Germany, Hannover, Germany (Werfel.Thomas@mh-hannover.de); Javier Sánchez-Pérez, Department of Dermatology, Hospital Universitario la Princesa, Madrid, Spain, Madrid/Princesa, Spain (jsanchezperez@aedv.es); Juan Fco. Silvestre, Department of Dermatology, Hospital General Universitario de Alicante, Alicante, Spain, Alicante, Spain (silvestre_jfr@gva.es); José C. Amario-Hita, Department of Dermatology, University Hospital of Puerto Real, Cádiz, Spain

, Cádiz, Spain (jcarmarioh@gmail.com); Pedro Mercader, Dermatology Department, Hospital General Universitario Morales Meseguer, Murcia, Spain, Murcia, Spain (pedromercader@hotmail.com); Inmaculada Ruiz, Department of Dermatology, Complejo Asistencial de León, C/Altos de Nava s/n, León, Spain, León, Spain (iruizgon@saludcastillayleon.es); Juan García-Gavín, Clínica Pérez & Gavín de Rmalogos, Vigo, Spain (juangavin@gmail.com); Maria Teresa Corradin, Dermatology Unit, Santa Maria degli Angeli Hospital, Pordenone, Italy, Pordenone, Italy (teresa.corradin@aas5.sanita.fvg.it); Rosella Gallo, Clinica Dermatologica, IRCCS - AOU San Martino - IST and Department of Health Sciences, University of Genoa, 16132 Genoa, Italy, Genova, Italy (rs.gallo@unige.it); Magdalena Czarnecka-Operacz, Dermatology Department, University of Medical Sciences, Poznan, Poland, Poznan, Poland (czarnecka.operacz@gmail.com); Mojca Simoncic Godnic, Department of dermatology, Novo mesto, (please edit), Slovenia, Novo Mesto, Slovenia (mojcasg78@gmail.com); Susan M. Cooper, Churchill Hospital, Oxford, U.K., Oxford, United Kingdom (sm_nc@btopenworld.com); Cathy M. Green, Department of Dermatology, Ninewells Hospital and Medical School, Dundee, U.K., Dundee, United Kingdom (cathy.green@nhs.net); Sharizan Ghaffar, Department of Dermatology, Ninewells Hospital and Medical School, Dundee, U.K., Dundee, United Kingdom (NA); Philippa Cousen, James Cook University Hospital, Middlesbrough, U.K., Middlesbrough, United Kingdom (Philippa.Cousen@stees.nhs.uk); Graham A. Johnston, Department of Dermatology, Leicester Royal Infirmary, UK, Leicester, United Kingdom (graham.johnston@uhl-tr.nhs.uk); Mahbub M.U. Chowdhury, The Welsh Institute of Dermatology, University Hospital of Wales, Cardiff, U.K., Cardiff, United Kingdom (Mmu.Chowdhury@CardiffandVale.wales.nhs.uk); Natalie Stone, Aneurin Bevan Health Board, Newport, Gwent, U.K., Newport, United Kingdom (Natalie.Stone@wales.nhs.uk); Ian R. White, St. John's Institute of Dermatology, Guy's Hospital, London, U.K., London, United Kingdom (ian.white@kcl.ac.uk);

Funding

Partial funding by EADV Grant PPRC-2018-8. Open access publication was funded by DEAL agreement.

Summary

Background. Occupational skin diseases have led the occupational disease statistics in Europe for many years. Especially occupational allergic contact dermatitis is associated with a poor prognosis and low healing rates leading to an enormous burden for the affected individual and for society.

Objectives. To present the sensitization frequencies to the most relevant allergens of the European baseline series in patients with occupational contact dermatitis (OCD) and to compare sensitization profiles of different occupations.

Methods. The data of 16022 patients considered having OCD after patch testing within the

ESSCA network between January 2011 and December 2020 were evaluated. Patients (n=46652) in whom an occupational causation was refuted served as comparison group.

Results. The highest percentages of OCD were found among patients working in agriculture, fishery and related workers, metal industry, chemical industry, followed by building and construction industry, health care, food and service industry. Sensitizations to rubber chemicals (thiurams, carbamates, benzothiazoles) and epoxy resins were associated with at least a doubled risk of OCD. After a decline from 2014 onwards, the risks to acquire an occupation-related sensitization to methyl(chloro)isothiazolinone (MCI/MI) and especially to methylisothiazolinone (MI) seem to increase again. Sensitization rates to formaldehyde were stable, and to methyldibromo glutaronitrile (MDBGN) slightly decreasing over time.

Conclusions. Among allergens in the European Baseline Series, occupational relevance is most frequently attributed to rubber accelerators, epoxy resins and preservatives.

Key words

Baseline series, occupational contact allergy, clinical epidemiology, surveillance, patch testing, RRID:SCR_001905

1. Introduction

Up to 35% of all occupational diseases in Europe affect the skin. Worldwide incidence rates vary due to different definitions of occupational skin diseases (OSD), data collection methods, and reporting systems, and occupational exposures.¹⁻³ Economic costs of occupational skin diseases in the European Union (EU) amount up to approximately 5 billion €/year for treatment, pension and compensation payments and loss of productivity.⁴ Moreover, occupational contact dermatitis OCD might be associated with significant reduction in quality of life, unemployment and social decline.⁵⁻⁹ Occupationally relevant sensitizations associated with an at least doubled risk for OCD were shown to be caused by thiurams, epoxy resins, and preservatives, such as

Accepted Article

methylchloroisothiazolinone/methylisothiazolinone (MCI/MI), methyldibromoglutaronitrile (MDBGN) and formaldehyde.^{10,11} Poor prognosis and low healing rates were associated with sensitization to occupationally relevant contact allergens and the number of positive patch test reactions.¹²

A prerequisite for targeted prevention in skin risk occupations is the knowledge of sensitizing agents at the workplaces, changes in sensitization rates over time and the discovery of emerging allergens. This can be obtained by continuous surveillance of contact allergy by contact allergy surveillance systems like the European Surveillance System on Contact Allergies (ESSCA)¹³⁻¹⁵ or the data base of the North American Contact Dermatitis Group (NACDG) and other national registries or data bases.¹⁶⁻¹⁹

The aim of this study was to update the knowledge on the most relevant baseline series allergens and their sensitization frequencies in patients with OCD in Europe and to compare the sensitization profiles of different occupations.

2. Methods

The objective of the ESSCA is the clinical surveillance of contact allergy.¹³⁻¹⁵ To this end, contributing departments (**Online supplemental Table S1**) submit either all patch test results, or just patch test results of the European baseline series (or national or local adaptations thereof), obtained following ESCD standards,²⁰ to the data center in Erlangen. This is accompanied by important demographic and clinical information, ranging from “MOAHLFA index”²¹ characteristics to a wider range of information according to the ESSCA “minimal dataset” definition.^{13,22} Data are quality checked, providing an “internal report” for each contributing department for scrutiny and approval before pooling of the respective data.¹⁴ Data from contributing departments are delivered in an anonymous format or partly, following national network standards and software usage (WinAlldat software),²² in a pseudonymized format, where the pseudonym cannot be related to actual personal data except in the contributing department itself. Only in those departments using WinAlldat software, pseudonymized data enabled the elimination of duplicate patients (presently, in 1438 patients from Austria, Germany, Poland, Spain, Switzerland, and The Netherlands). In

such cases, a random selection of one of the multiple consultations of a patient was made. For the present analysis, further constraints were employed:

- only patients aged between 16 and 68 (inclusive) were included.
- a valid occupational code, in this case ISCO-88 still used by most departments as well as additions by ESSCA allowing to document activities beyond formal employments, such as homemaker, unemployed, etc. needed to be documented (coding "+": denotes job titles on the 3-digit level of groups in ISCO-88, for which a finer categorization is generally available, but had not been used in individual cases or when job titles were aggregated for analysis; all other job titles reported are on the 4-digit level of groups);
- patients with occupation related contact dermatitis tagged with "occ" in MOAHLFA index after patch testing by the examining dermatologist/clinician were considered as cases of OCD. Patients in whom occupational causation was explicitly negated were tagged with "non-occ" and these comprised the non-OCD group.
- patients were tested with the baseline series as used locally.
- The association of OCD with any sensitization to an EBS allergen was investigated.

Data management and analysis was performed with the R software package (<https://www.r-project.org>; RRID:SCR_001905), version 4.0.3. For the calculation of 95% confidence intervals (CIs) to zero proportions an approximation to an exact CI was used.²³ The in- or decrease of risk to be sensitized to a particular allergen associated with OCD, compared to those patients in whom occupational causation of contact dermatitis was explicitly denied, was estimated with log-binomial regression analysis, adjusting for age (dichotomized in < 40 vs. older) and sex, deriving a prevalence ratio (PR) accompanied by a 95% confidence interval (95% CI).

3. Results

Between January 2011 and December 2020, 89777 patients fulfilled above-mentioned inclusion criteria. Among these, 16022 were diagnosed with OCD, while in 46652 occupational causation was explicitly negated. In 12754 the occupational causation was unclear, and in the remainder (n=14349) information was altogether missing; these latter

two groups were excluded from the comparisons. Contributing countries and respective patient numbers included are summarized in **Table 1**.

3.1 Patients

The proportion of men was significantly higher in patients with OCD (44.5%) than in the non-OCD group (29.2%; **Table 2**). The proportion of patients over 40 years of age was significantly lower in the OCD group compared to the non-OCD group (52.9% vs. 62.6%). Patients with current or previous atopic dermatitis were slightly overrepresented in OCD patients (32.4% vs. 28.7%). Hand was the predominant eczema location in OCD patients (73.1% vs. 21.2%).

3.2 Risk of OCD in different skin risk occupations

In **Online supplemental table S2**, the proportion of patients with OCD among all patients patch tested in the strata of occupational groups is shown. The highest risk for OCD with a share of >65% in occupational groups comprising more than 50 patients was seen in metal workers, agricultural, fishery and related laborers, as well as in the rubber and chemical industry.

3.3 Characteristics of patients working in occupations with a high occurrence of OCD

For patients working in occupations with a high occurrence of OCD, the distribution of ACD, ICD, and both diagnoses combined are shown in **Table 3**. Overall, the share of patients with ACD was 33.5%, of ICD 14.4% (both combined in 1.9%), and of current atopic eczema 11.8%, in 1.5% combined with ICD. ACD prevailed in building finishers and related trade workers (65.3%), nursing and midwifery professionals (56.0%), dentists (50%) and hairdressers, barbers, beauticians, and related workers (48.8%). ICD was most common in toolmakers and related workers (41.4%), plant and machine operators and assemblers (38.0%), machine-tool setters and setters-operators (37.6%) followed by nursing and midwifery associate professionals (37.5%). A combined etiology (ACD and ICD) was seen most frequently in food processing and related trades workers (8.1%), hairdressers, barbers, beauticians, and related workers (7.0%), plant and machine operators and assemblers (6.3%) as well as metal wheel-grinders, polishers, and tool sharpeners (6.1%).

3.4 EBS allergens and association to OCD

The association of OCD with sensitizations to allergens of the European baseline series quantified with the prevalence ratio (PR) with accompanying 95% confidence interval is shown in **Table 4**. Occupationally relevant allergens with a doubled risk for occupational sensitization were rubber chemicals like thiuram mix, (PR 4.48, 95% CI 4.03-4.98), carba mix (PR 2.8, 95% CI 2.44-3.22), mercaptobenzothiazole and mercapto mix (PR 2.55, 95% CI 2.08-3.12/2.35, 95% CI 1.8-3.0) and epoxy resin (diglycidyl ether of bisphenol A) (PR 3.58, 95% CI 3.14-4.09). The frequency of carba mix positivity (6.7%) was higher than that of thiuram mix sensitization (5.4%). (**Table 4**).

A comparison with the previous data analysis from Pesonen et al¹¹ is shown in **Online supplemental table 3 A-C**. The most relevant data are reported here. Concerning thiuram-mix and mercaptobenzothiazoles, no relevant changes in overall sensitization rates and prevalence ratios (PR for positivity in OCD vs non-OCD cases) compared to the analysis period from 2002 to 2010 were seen (Table S3 A). For Carba mix, no data from the former analysis period are available. N-isopropyl-N'-phenyl-p-phenylenediamine (IPPD) sensitization showed a downward trend in occupational relevance, as PR fell from 2.62 to 1.85.

No significant changes in epoxy resin sensitization rates were seen. However, there was a slight increase from 3.1. to 3.5%.

Sensitization rates to MCI/MI and MI showed a considerable increase as compared to the analysis period from 2002-2010, as MCI/MI increased from 4.0% to 8.6% in OCD patients and from 2.5% to 5.7% in non OCD patients, and MI increased from 1.9% to 9.0% in OCD patients and from 0.5% to 6.4% in non OCD patients. The likelihood of epidemiological relevance remained nearly the same for MCI/MI and clearly decreased for MI (PR 2.36 vs PR 1.49). Sensitization rates in OCD and non-OCD patients peaked 2014 and fell until 2018. Since then a slight upward trend with a peak in 2019 which is more pronounced in non-OCD patients, driven by MI sensitizations was seen. (**Figure 1a and b**). (**Table S3 C (biocides)**).

Occupations with the highest risk of contact allergy to selected allergens are shown in **Online supplemental table S4 A-F**.

High rates of thiuram mix sensitization were seen in the construction and food industry. Highest sensitization rates were seen in builders (17.9%, 95% CI 8.9-30.4), followed by concrete placers, concrete finishers, and related workers (17.1%, 95% CI 6.6-33.6) as well as in butchers, fishmongers, and food preparers (14.1%, 95% CI 7.3-23.8) (**Table S4A**).

High rates of carba mix positivity were recorded in medical professions, construction, and food industry. Highest positivity rates were seen in health professionals (except nursing) (26.0%, 95% CI 16.5-37.6), building finishers and related trades workers (19.4%, 95% CI 10.4-31.4), and cooks (13.1%, 95% CI 8.4-19.2) (**Table S4B**).

Formaldehyde sensitization was observed in teaching professionals and in the metal industry. Highest sensitization rates were seen in teaching professionals (6.9%, 95% CI 2.6-14.4), machine-tool setters and setters-operators (6.7%, 95% CI 3.5-11.5) as well as turners (6.2%, 95% CI 3.1-10.8) (**Table S4C**).

MCI/MI sensitizations were found predominately in market gardeners and crop growers (31.2%, 95% CI 16.1-50), building finishers and related trades workers (21.7%, 95% CI 12.7-33.3) as well as in painters, building structure cleaners and related trades (17.6%, 95% CI 6.8-34.5) (**Table S4D**).

MI sensitizations were most prominent in building finishers and related trades workers (27.8%, 95% CI 17.9-39.6), painters and related workers (20.1%, 95% CI 13.8-27.8) as well as in chemical-processing-plant operators (20.0%, 95% CI 9.6-34.6) (**Table S4E**).

Sensitizations to epoxy resin were most prominent in craft occupations. Highest sensitization rates were seen in building construction laborers (34.1%, 95% CI 20.5-49.9), Floor layers and tile setters (27.2%, 95% CI 17.9-38.2) as well as in plumbers and pipe fitters (26.2%, 95% CI 18-35.8) (**Table S4F**).

Sensitization rates against MDBGN were less prominent in occupational (3.6%, 95% CI 3.3-3.9) and non-occupational settings (2.8%, 95% CI 2.7-3.0).

4. Discussion

Delayed type sensitizations to work related allergens of the European baseline series in patients with OCD patch tested within the ESSCA network from 2011-2020 are dominated by sensitizations to rubber chemical accelerators as well as by epoxy resin. The occupational importance of sensitizations to preservatives has declined, compared to the previous analysis period from 2002-2010.¹¹ Concerning rubber chemical accelerators as well as epoxy resin these results are in line with recent data from the NACDG. However, occupational sensitization to preservatives, especially to MI, is still of great importance in the US and in Canada, where sensitization rates still rise²⁴

4.1. Rubber chemicals

Accelerators added to rubber vulcanization processes represent the most common cause of contact sensitizations to rubber chemicals.^{10,25-27}

Occupational rubber chemical sensitizations are mainly acquired by the use of occlusive rubber gloves.²⁸⁻³¹ Other allergen sources include rubber boots, rubber bands, rubber insulation material, car tires, machine belts, and rubber tool handles.³²⁻³⁵

Carbamates are the most common accelerators in medical rubber gloves nowadays, while thiurams are hardly used in their production anymore.³⁰ The persistent high rate of sensitization to the thiuram mix in patients with OCD is probably linked to occupational use of carbamate-containing gloves as thiurams and carbamates are a redox pair and thiurams are considered a better marker for a thiuram/carbamate sensitization.³¹ In recent years, an overall increase in carba mix positivity,³² especially in the health care sector was shown.^{10,27,28,36} These results are supported by our data and the recent analyses by the NACDG.²⁴ Medical professionals with OCD showed positivity rates to carba mix from 7.2% in nursing and midwifery associates up to 26% in health professionals (except nursing). However, these results must be interpreted with caution because carba mix contains not only carbamates but 1,3-DPG, a problematic test substance frequently causing false-positive irritant reactions, which may explain the high positivity rate.^{32,37-39} Therefore, the clinical relevance of carba mix sensitizations should be proven by testing the components of the

carba mix , especially 1,3-DPG and by testing the thiuram mix or its components as well as patient's own material. A false positive reaction to carba mix has major consequences for the patient's occupational opportunities, because the selection of protective gloves is severely compromised when carbamates must be avoided..⁴⁰ However, one has to keep in mind that true allergic reactions to 1,3-DPG can occur, which was recently shown in health care workers wearing rubber gloves .^{41,42}

In contrast, sensitization rates to mercaptobenzothiazoles derivatives remained low in occupations with high OCD risk. Compared to the data of Pesonen et al 2015 even a slight downward trend was apparent.¹¹ However, the risk of acquiring occupational sensitizations to mercapto-compounds is still significantly increased. Highest rates were found in service and food industry (data not shown).

4.2. Epoxy resin

Epoxy resin, e.g., in epoxy paints, adhesives and protective coatings, is the emerging contact allergen of the last two decades due to the widespread use in craft trades and its high sensitization potential.^{11,24,33,43-45}

Overall sensitization rates to epoxy resin (diglycidyl ether of bisphenol A) in Europe rose from 3.1 during 2001-2010 to 3.5% during 2011-2020. The NACDG reported sensitization rates to epoxy resin (bisphenol A epoxy resin) as high as 5.6% in OSD patients.²⁴

Differences in sensitization rates might be explained by real differences in exposure but also by the respective definition of OSD as well as by different patient selection criteria.

Contamination of the skin occurs by direct skin contact, contaminated surfaces, inappropriate personal protective equipment, and by handling of uncured or not fully cured epoxy materials.⁴⁶ Sensitization rates to epoxy resin in exposed occupations further increased in the last decade. Most frequently sensitized were building and construction workers (34.1%) as well as floor layers and tile setters (27.2%). This is not surprising, since high sensitization rates in construction coating as well as by using adhesives, e.g. in tile setters are well known.⁴⁷ A dramatic increase compared to the previous analysis period from 2002-2010 in sensitization rates from 7.7% ¹¹ to 26.2% was seen in plumbers and pipe fitters. High allergen exposure especially in epoxy injection casting or pipe lining techniques

is likely responsible for the increasing sensitization rates in this occupational group.⁴⁸ Further occupations with a dramatic increase in epoxy resin (diglycidyl ether of bisphenol A) sensitization rates in the present and the previous analysis of ESCCA data are plastic-products machine operators (from 14.5% to 24.3%) and painters (from 16.5% to 22.6%). Practical guidelines for prevention of epoxy allergies are highly needed.⁴⁹

Other components of epoxy resin systems namely hardeners, and reactive diluents, are potent allergens, too.⁵⁰⁻⁵² To detect these sensitizations, patch testing of an epoxy resin series besides the baseline series is highly recommended, because up to 20% of patients sensitized to a reactive diluent and up to 60% of the patients sensitized to an epoxy resin hardener show no positive patch test to epoxy resin.⁵¹

4.3 Preservatives

MCI/MI and MI

As a result of the guidance of Cosmetics Europe, the association of the cosmetic industry, as well as European legislation on the use of MI in cosmetic products the sensitization to MI and MCI/MI decreased in Europe from 2014 onwards^{53,54}, while in the US and Canada, where legal regulations for consumer products are yet lacking, the upward trend is still ongoing.²⁴

However, our data indicate that this downward trend ended in 2018. Since then, sensitization rates for MCI/MI and MI seem to increase again. Separate analyses for MCI/MI and MI showed that the peak in 2019 and the slight upward trend in non-occupational settings from 2018 onwards is again mainly driven by non-occupational MI sensitizations. The latter is most probably due to ongoing exposures to MCI/MI and MI in wall paints, household products such as wash machine powders and liquids, dish washing fluids, floor cleaning agents, shoeshine, etc.⁵⁵⁻⁵⁸

Besides private exposure, there is still pronounced occupational relevance for MCI/MI and MI. In the last 10 years, the spectrum of occupations at risk for MCI/MI and MI sensitizations changed and sensitization rates rose dramatically in distinct occupations. Pesonen et al 2015 reported floor layers and tile setters, chemical processing plant

operators and precision worker in metal and related materials to have the highest risk to acquire occupational MCI/MI sensitization.¹¹ In the current analysis, we found highest sensitization rates for MCI/MI in market gardeners and crop growers, in which almost one third of the patch test population was sensitized, followed by building finishers and related trades as well as painters, building structure cleaners and related workers. For MI sensitization, building finishers and painters showed highest sensitization rates of 27.8% and 20.1%, respectively. In the recent analysis of the NACDG occupational relevance was assumed in 52.5% of positive reactions to MI.²⁴

MCI/MI and MI exposure in painters and cleaners occurs via preserved water based wall paints and cleaning products, respectively.^{57,59,60} Market gardeners and crop growers might be exposed to MCI/MI and MI treated plants, fruits and vegetable used to reduce food borne illnesses as well as microbial spoilage.^{61,62} Occupational exposure of building finishers and related trades (roofer, floor layer and tile setters, plasterers, insulation workers, glaziers, plumbers, and tile fitters, building and related electricians), might be related to occupational exposure to products like paints, adhesives, wall and joint mortars and cleansers which may contain isothiazolinones as preservatives. Non- and incorrect labelling of isothiazolinones still lead to considerable problems to identify the culprit allergen.^{57,58}

Formaldehyde

Formaldehyde and formaldehyde releasers are still frequently used as preservatives in household detergents, cosmetic and pharmaceutical products, and various industrial products.^{63,64} Sensitization rates to formaldehyde (1% and 2% aq.) were lower in Europe (2.7% (95% CI 2.5–3%) compared to the US and Canada 3.9%).²⁴ Continuous exposure might contribute to the constant level of sensitization rates to formaldehyde over the last 2 decades.¹¹ Current sensitization rates were highest in teaching professionals (6.9%), followed by metal workers (machine tool setters, turners: 5.6- 6.7%) and lower in health care (nurses and midwives: 3.6%). When it comes to teaching professionals at vocational schools, sensitization might have been acquired by using formaldehyde releaser-containing liquid soaps or other substances (e.g. metal working fluids) at the workplace. However, all the patients (n=6) concerned were women between 37 and 50 years of age and they might

as well have acquired formaldehyde sensitization by personal use of formaldehyde releaser containing cosmetics or household products.⁶⁵

High formaldehyde sensitization rates in metal workers are plausible due to regular contact to metal working fluids commonly preserved with a variety of different formaldehyde-releasers.^{66,67}

MDBGN

Sensitization rates to MDBGN slightly dropped in occupational and non-occupational settings by approximately 1%. Banned in cosmetics by the EU from 2007,⁶⁸ MDBGN may still be used in all product-type 6 (PT6) categories (preservatives for products during storage). PT6 sub-categories⁶⁹ are technical fluids, including washing and cleaning fluids, paints and coatings, metal working fluids, and glues and adhesives. Labelling is not mandatory if concentration limits are not exceeding 0.1%. Other possible exposures are medical devices and medicinal products, where MDBGN is permitted.⁶⁹

4.4. Labelling

Due to the lack of full labelling of many products including technical and cleaning fluids, glues, adhesives and medical devices determining relevance of a sensitization as well as consecutive allergen avoidance is difficult. Legal regulations for full labelling are highly needed.^{15,69,70}

5. Limitations and strength

A limitation of our study is, that the risk of sensitization is calculated from all positive patch tests, and not only from the relevant ones. The correct assessment of exposure at work is definitive to really confirm the occupational relevance.⁷¹ However, in our data set we cannot ensure that the information on relevance is entirely reliable (e.g. based on workplace exposure analysis in every case), therefore we opted for analysis of the entire data set.

Our data provide a broad epidemiological basis of the sensitization in Europe. However, aggregation of occupations within ISCO classification may have the effect, that important allergens in specific occupations could be overlooked. For these cases, occupation-specific analyses should be intended. Moreover, to comprehensively diagnose occupational allergic contact dermatitis in individual cases additional testing of occupation-related patch test series as well as patients' own products is commonly inevitable.

6. Conclusion

From the EBS allergens, rubber chemicals – especially thiurams – and epoxy resin (DGEBA) were associated with a pronounced risk of OCD. High rates of sensitization continued to be detectable against preservatives, especially MCI/MI and MI. Since the previous analysis of OCD patients in ESSCA in 2002-2010, the MCI/MI and MI sensitization rates have increased in both OCD and non-OCD patients. Time will show whether the observed rise in their sensitization rate in 2019 which was largely driven by MI sensitizations represents a trend. Due to the necessity to test the single constituents anyway, testing of carba mix in the EBS should be reconsidered.

Tables

Table 1: Contributing countries.

Country	N(patients)	N(occ.)	%(occ.)
Austria	1711	349	20.4
Switzerland	9234	1383	15
Germany	16064	6210	38.7
Spain	6090	1022	16.8
Finland	1170	721	61.6
Italy	10507	873	8.3
Lithuania	2885	624	21.6
The Netherlands	9674	1805	18.7
Poland	5437	1154	21.2
Slovenia	8119	137	1.7
United Kingdom	18886	1744	9.2
Total	89777	16022	17.8

Legend: N: number, %: percent, occ: patients with occupational contact dermatitis

Table 2: Demographic and clinical characteristics according to the MOAHLFA index⁴.

Factor	N(occ.)	%(occ.)	N(non-occ.)	%(non-occ.)	P-value
Male	7126	44.48	13609	29.2	< .0001
Occupational	16022	100	0	0	
Atopic Eczema	5145	32.37	13370	28.7	< .0001
Site: Hand	11517	73.11	9383	21.2	< .0001
Site: Leg	242	1.54	2359	5.3	< .0001
Site: Face	1113	7.06	10082	22.8	< .0001
Age 40+	8470	52.86	29224	62.6	< .0001

Legend: N: number, %: percent, occ: patients with occupational contact dermatitis, non-occ: patients without occupational contact dermatitis

Table 3: Characteristics of patients with OCD in occupational subgroups with a high level of occurrence of OCD (defined as > 50 OCD cases and a relative frequency of at least 50% OCD) in detail, i.e., on the four-digit ISCO level.

Occupation	N(occ.)	Mean age (years)	% males	% with ACD	% with ICD	% ACD+ICD
------------	---------	------------------	---------	------------	------------	-----------

Hairdressers, barbers, beauticians, and related workers	1329	33.1	7.1	48.8	24.0	7.0
Nursing and midwifery associate Professionals	1137	39.9	10.1	20.4	37.5	4.8
Machinery mechanics and fitters	679	39.2	95.1	24.8	36.9	3.3
Cooks	437	35.6	54.9	28.7	31.1	5.4
Nursing and midwifery Professionals	420	42.4	9.8	56.0	19.2	5.6
Machine-tool setters and setter-operators	382	45.9	92.1	27.9	37.6	4.6
Physiotherapists and related associate professionals	232	37.8	19.0	21.8	34.8	5.2
Painters, Varnishers and related workers	232	41.2	87.1	38.3	23.8	2.8
Bakers, pastry-cooks and confectionery makers	222	34.2	52.3	35.3	26.4	2.5
Gardeners, horticulturists and nursery growers, NEC*	178	40.1	34.3	41.8	23.1	4.0
Metal- and mineral-products machine operators	170	44.0	88.8	29.6	33.8	4.1
Bricklayers and stonemasons	143	41.6	97.9	40.7	22.9	3.0
Chemical and physical science technicians	132	37.6	29.5	33.1	32.3	1.5
Electrical and electronic equipment mechanics and fitters	124	36.5	90.3	25.8	32.3	1.6
Dental assistants	119	33.3	0.8	34.2	29.1	3.4
Plumbers and pipe fitters	108	40.4	99.1	36.4	23.4	1.3
Metal-, rubber- and plastic-products assemblers	103	45.1	68.0	23.3	25.3	5.8
Other machine operators and assemblers	100	45.1	84.0	24.5	32.7	5.1
Wood treaters, cabinetmakers and related trades workers	94	39.8	86.2	29.7	21.2	4.3
Toolmakers and related workers	88	42.6	93.2	19.6	41.4	1.1
Dentists	87	39.6	23.0	50.0	18.3	3.7
Floor layers and tile setters	87	42.6	93.1	35.8	19.4	4.5

Welders and flame cutters	87	45.8	90.8	26.3	23.6	5.3
Metal wheel-grinders, polishers, and tool sharpeners	86	42.4	87.2	26.8	34.1	6.1
Carpenters and Joiners	84	44.1	90.5	40.3	29.9	0.0
Butchers, fishmongers, and related food preparers	84	40.1	51.2	20.8	34.2	2.4
Plant and machine operators and assemblers	82	44.7	92.7	11.4	38.0	6.3
Metal, machinery, and related trades workers	80	39.9	86.2	32.9	26.0	5.5
Precision workers in metal and related materials	73	41.2	38.4	34.3	35.7	2.7
Plastic-products machine operators	72	42.4	75.0	44.8	25.4	1.5
Building finishers and related trades workers	70	43.7	94.3	65.3	13.1	4.3
Precision-instrument makers and repairers	70	39.8	81.4	30.0	31.4	2.9
Modern health associate Professionals (except nursing)	61	44.0	23.0	47.1	30.1	5.7
Building frame and related trades workers	60	43.4	91.7	48.0	32.0	0.0
Food processing and related trades workers, NEC	60	40.3	40.0	43.3	21.6	8.1
Chemical-processing-plant operators	58	42.6	89.7	47.3	10.5	5.3
Mechanical engineering technicians	55	41.1	98.2	29.6	31.4	1.9
Baked-goods, cereal, and chocolate-products machine operators	54	39.0	3.7	18.5	27.7	1.9

Legend: N: number, %: percent, occ: patients with occupational contact dermatitis, ICD: irritant contact dermatitis, ACD: allergic contact dermatitis, NEC: Not elsewhere classified

Table 4: Risk of occupational contact dermatitis (OCD) associated with contact sensitization to allergens of the European baseline series, quantified with the prevalence ratio (PR) with accompanying 95% confidence interval (CI).

Allergen (in petrolatum except where otherwise specified)	OCD-tested	OCD-positive	non-OCD-tested	non-OCD-positive	PR (95% CI)
Thiuram mix 1%	15550	5.4% (5–5.7%)	46139	1.2% (1.1–1.4%)	4.48 (4.03–4.98)
Epoxy resin (DGEBA) 1%	14922	3.5% (3.2–3.8%)	44635	0.9% (0.8–1%)	3.58 (3.14–4.09)
Carba mix 10%	4762	6.7% (6–7.4%)	22818	2.2% (2–2.4%)	2.8 (2.44–3.22)
Mercaptobenzothiazole 0.5%	15629	1.2% (1–1.3%)	46135	0.4% (0.4–0.5%)	2.55 (2.08–3.12)
Mercapto mix 1% and 2%	10997	1.2% (1–1.4%)	28600	0.5% (0.4–0.6%)	2.35 (1.84–3)
IPPD 0.1%	13525	1.2% (1–1.4%)	37859	0.6% (0.5–0.7%)	1.85 (1.51–2.27)
Budesonide (0.1 and 0.01%)	7611	1% (0.8–1.3%)	26728	0.6% (0.5–0.7%)	1.76 (1.34–2.32)
Formaldehyde 1% and 2%	15466	2.7% (2.5–3%)	45055	1.7% (1.6–1.9%)	1.66 (1.47–1.87)
Textile dye mix 6.6%	1530	5.9% (4.8–7.2%)	5311	3.8% (3.3–4.4%)	1.61 (1.26–2.05)
<i>p</i> -Phenylenediamine 1.0%	8773	5.1% (4.6–5.5%)	35284	3.4% (3.3–3.6%)	1.6 (1.44–1.79)
MCI/MI 0.02% and 0.01% aq.	15615	8.6% (8.2–9%)	46153	5.7% (5.5–6%)	1.57 (1.47–1.68)
Colophonium 20%	15616	4.2% (3.8–4.5%)	46132	2.8% (2.7–3%)	1.55 (1.41–1.71)
Pot. Dichromate 0.5%	15364	5.8% (5.5–6.2%)	45103	3.8% (3.6–4%)	1.53 (1.42–1.66)

Quaternium 15 1.0%	7548	1.2% (0.9–1.4%)	26052	0.8% (0.7–0.9%)	1.51 (1.18–1.94)
MI 0.2% and 0.05% aq.	11021	9% (8.5–9.6%)	36105	6.4% (6.2–6.7%)	1.49 (1.38–1.6)
ptBFR 1%	10600	1% (0.8–1.2%)	36486	0.8% (0.7–0.8%)	1.37 (1.09–1.72)
MDBGN 0.5%, 0.2% and 0.3%	14202	3.6% (3.3–3.9%)	42503	2.8% (2.7–3%)	1.36 (1.22–1.5)
Cobalt chloride 1%	15485	7.8% (7.4–8.2%)	45967	6.2% (6–6.4%)	1.32 (1.24–1.41)
Paraben mix 16%	14145	0.8% (0.6–0.9%)	44490	0.6% (0.5–0.6%)	1.31 (1.04–1.65)
SL mix 0.1%	7862	0.9% (0.7–1.1%)	27627	0.8% (0.7–0.9%)	1.23 (0.93–1.62)
Neomycin sulfate 20%	8143	1.1% (0.9–1.4%)	31199	1% (0.9–1.1%)	1.22 (0.97–1.54)
HICC 2.5%	14189	1.8% (1.6–2%)	42831	1.6% (1.5–1.8%)	1.19 (1.03–1.38)
Nickel sulfate 5%	15241	19.4% (18.8–20%)	45898	21.3% (20.9–21.7%)	1.09 (1.05–1.13)
Fragrance mix II 14%	15013	4% (3.7–4.3%)	43944	4.1% (3.9–4.3%)	1.06 (0.97–1.16)
Fragrance mix I 8%	15122	7% (6.5–7.4%)	45531	7.3% (7–7.5%)	1.05 (0.98–1.13)
<i>Myroxylon pereirae</i> 25%	15155	5.5% (5.1–5.9%)	45349	5.8% (5.6–6%)	1.01 (0.93–1.09)
Lanolin (wool alcohols) 30%	15466	1.9% (1.7–2.2%)	44956	2% (1.9–2.1%)	0.98 (0.86–1.12)

Tixocortol pivalate (0.1% and 1%)	8651	0.7% (0.5–0.9%)	33078	0.8% (0.7–0.9%)	0.9 (0.68–1.19)
Primin 0.01%	5096	0.2% (0.1–0.4%)	17728	0.3% (0.3–0.4%)	0.6 (0.31–1.17)
Benzocaine 5%	3842	0.5% (0.3–0.8%)	11274	0.9% (0.7–1.1%)	0.59 (0.36–0.97)

OCD: Occupational Contact Dermatitis; non-OCD: OCD negated; PR: Prevalence Ratio; 95%CI: 95% Confidence Interval

Aq: aqua; DGEBA: diglycidyl ether of bisphenol A; MCI: methylchloroisothiazolinone; MI: methylisothiazolinone; MDBGN: Methylidibromo glutaronitrile; ptBFR: *p-tert*-butylphenol-formaldehyde resin; IPPD: N-isopropyl-N'-phenyl-*p*-phenylenediamine; pot.: potassium; HICC: Hydroxylisohexyl 3-cyclohexene carboxaldehyde; SL: sesquiterpenlactone mix: alantolactone, dehydrocostus lactone and costunolide; thiuram mix: tetramethylthiuram monosulfide, tetraethylthiuramdisulfide, tetramethylthiuramdisulfide, dipentamethylene thiuramdisulfide; carba mix: 1.3-diphenylguanidine, zinc dibutyldithiocarbamate, zinc diethyldithiocarbamate; mercapto mix (1% pet.): N-cyclohexylbenzothiazylsulfenamide, dibenzothiazylsulfide, morpholinylmercaptobenzothiazole

Figure legends

Figure 1 a: Time trend of MCI/MI contact sensitization in patients with occupational and non-occupational contact dermatitis, 2011-2020

Figure 1b: Time trend of and MI contact sensitization in patients with occupational and non-occupational contact dermatitis, 2011-2020

Figures

Figure 1 a: Time trend of MCI/MI contact sensitization in patients with occupational and non-occupational contact dermatitis, 2011-2020





Figure 1b: Time trend of and MI contact sensitization in patients with occupational and non-occupational contact dermatitis, 2011-2020



References

1. Uter W. Surveillance in occupational contact dermatitis. In: John SM, Johansen JD, Rustemeyer T, Elsner P, Maibach HI, eds. *Kanerva's Occupational Dermatology*. Springer; 2020.
2. Keegel T, Moyle M, Dharmage S, Frowen K, Nixon R. The epidemiology of occupational contact dermatitis (1990-2007): a systematic review. *Int J Dermatol*. Jun 2009;48(6):571-8. doi:10.1111/j.1365-4632.2009.04004.x
3. Stocks SJ, McNamee R, van der Molen HF, et al. Trends in incidence of occupational asthma, contact dermatitis, noise-induced hearing loss, carpal tunnel syndrome and upper limb musculoskeletal disorders in European countries from 2000 to 2012. *Occup Environ Med*. Apr 2015;72(4):294-303. doi:10.1136/oemed-2014-102534
4. Augustin M. Pharmacoeconomics of Occupational Diseases. In: Rustemeyer T, Elsner P, John S-M, Maibach HI, eds. *Kanerva's Occupational Dermatology*. Springer Berlin Heidelberg; 2012:19-26.
5. Dickel H, Bruckner T, Bernhard-Klimt C, Koch T, Scheidt R, Diepgen TL. Surveillance scheme for occupational skin disease in the Saarland, FRG. First report from BKH-S. *Contact Dermatitis*. Apr 2002;46(4):197-206. doi:10.1034/j.1600-0536.2002.460403.x
6. Dickel H, Kuss O, Blesius CR, Schmidt A, Diepgen TL. Report from the register of occupational skin diseases in northern Bavaria (BKH-N). *Contact Dermatitis*. Apr 2001;44(4):258-9. doi:10.1034/j.1600-0536.2001.440409-14.x
7. Blanciforti LA. Economic burden of dermatitis in US workers [corrected]. *J Occup Environ Med*. Nov 2010;52(11):1045-54. doi:10.1097/JOM.0b013e3181f475b2
8. McCall BP, Horwitz IB, Feldman SR, Balkrishnan R. Incidence rates, costs, severity, and work-related factors of occupational dermatitis: a workers' compensation analysis of Oregon, 1990-1997. *Arch Dermatol*. Jun 2005;141(6):713-8. doi:10.1001/archderm.141.6.713
9. Ofenloch RF, Diepgen TL, Popielnicki A, et al. Severity and functional disability of patients with occupational contact dermatitis: validation of the German version of the Occupational Contact Dermatitis Disease Severity Index. *Contact Dermatitis*. Feb 2015;72(2):84-9. doi:10.1111/cod.12302
10. Warburton KL, Bauer A, Chowdhury MM, et al. ESSCA results with the baseline series, 2009-2012: rubber allergens. *Contact Dermatitis*. Nov 2015;73(5):305-12. doi:10.1111/cod.12454
11. Pesonen M, Jolanki R, Larese Filon F, et al. Patch test results of the European baseline series among patients with occupational contact dermatitis across Europe - analyses of the European Surveillance System on Contact Allergy network, 2002-2010. *Contact Dermatitis*. Mar 2015;72(3):154-63. doi:10.1111/cod.12333
12. Macan J, Rimac D, Kežić S, Varnai VM. Occupational and non-occupational allergic contact dermatitis: a follow-up study. *Dermatology*. 2013;227(4):321-9. doi:10.1159/000354763

13. Uter W, Schnuch A, Giménez-Arnau A, Orton D, Statham B. Databases and Networks. The Benefit for Research and Quality Assurance in Patch Testing. In: JD J, V M, J-P L, PJ F, eds. *Contact Dermatitis*. 6 ed. 2020:1-16.
14. Uter W, Schnuch A, Wilkinson M, Dugonik A, Dugonik B, Ganslandt T. Registries in Clinical Epidemiology: the European Surveillance System on Contact Allergies (ESSCA). *Methods Inf Med*. 2016;55(2):193-9. doi:10.3414/me15-01-0099
15. Uter W, Werfel T, Lepoittevin JP, White IR. Contact Allergy-Emerging Allergens and Public Health Impact. *Int J Environ Res Public Health*. Apr 1 2020;17(7):2404. doi:10.3390/ijerph17072404
16. Holness DL. Occupational Dermatoses. *Curr Allergy Asthma Rep*. Jul 27 2019;19(9):42. doi:10.1007/s11882-019-0870-6
17. Bensefa-Colas L, Telle-Lamberton M, Paris C, et al. Occupational allergic contact dermatitis and major allergens in France: temporal trends for the period 2001-2010. *Br J Dermatol*. Dec 2014;171(6):1375-85. doi:10.1111/bjd.13133
18. Aalto-Korte K, Koskela K, Pesonen M. Allergic contact dermatitis and other occupational skin diseases in health care workers in the Finnish Register of Occupational Diseases in 2005-2016. *Contact Dermatitis*. Apr 2021;84(4):217-223. doi:10.1111/cod.13753
19. Carøe TK, Ebbelhøj N, Agner T. A survey of exposures related to recognized occupational contact dermatitis in Denmark in 2010. *Contact Dermatitis*. Jan 2014;70(1):56-62. doi:10.1111/cod.12134
20. Johansen JD, Aalto-Korte K, Agner T, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing - recommendations on best practice. *Contact Dermatitis*. Oct 2015;73(4):195-221. doi:10.1111/cod.12432
21. Schnuch A, Geier J, Uter W, et al. National rates and regional differences in sensitization to allergens of the standard series. Population-adjusted frequencies of sensitization (PAFS) in 40,000 patients from a multicenter study (IVDK). *Contact Dermatitis*. Nov 1997;37(5):200-9. doi:10.1111/j.1600-0536.1997.tb02435.x
22. Uter W, Arnold R, Wilkinson J, et al. A multilingual European patch test software concept: WinAlldat/ESSCA. *Contact Dermatitis*. Nov 2003;49(5):270-1. doi:10.1111/j.0105-1873.2003.0225n.x
23. Gefeller O, Pfahlberg AB, Uter W. What can be learnt from nothing? - A statistical perspective. *Contact Dermatitis*. Dec 2013;69(6):350-4. doi:10.1111/cod.12112
24. DeKoven JG, DeKoven BM, Warshaw EM, et al. Occupational contact dermatitis: Retrospective analysis of North American Contact Dermatitis Group Data, 2001 to 2016. *J Am Acad Dermatol*. Apr 2022;86(4):782-790. doi:10.1016/j.jaad.2021.03.042
25. Bergendorff O, Hansson C. Contact dermatitis to a rubber allergen with both dithiocarbamate and benzothiazole structure. *Contact Dermatitis*. May 2007;56(5):278-80. doi:10.1111/j.1600-0536.2007.01076.x
26. Bergendorff O, Persson C, Lütcke A, Hansson C. Chemical changes in rubber allergens during vulcanization. *Contact Dermatitis*. Sep 2007;57(3):152-7. doi:10.1111/j.1600-0536.2007.01194.x
27. Warburton KL, Urwin R, Carder M, Turner S, Agius R, Wilkinson SM. UK rates of occupational skin disease attributed to rubber accelerators, 1996-2012. *Contact Dermatitis*. May 2015;72(5):305-11. doi:10.1111/cod.12356

28. Cao LY, Taylor JS, Sood A, Murray D, Siegel PD. Allergic contact dermatitis to synthetic rubber gloves: changing trends in patch test reactions to accelerators. *Arch Dermatol*. Sep 2010;146(9):1001-7. doi:10.1001/archdermatol.2010.219
29. Liskowsky J, Geier J, Bauer A. Contact allergy in the cleaning industry: analysis of contact allergy surveillance data of the Information Network of Departments of Dermatology. *Contact Dermatitis*. Sep 2011;65(3):159-66. doi:10.1111/j.1600-0536.2011.01937.x
30. Geier J, Lessmann H, Mahler V, Pohrt U, Uter W, Schnuch A. Occupational contact allergy caused by rubber gloves--nothing has changed. *Contact Dermatitis*. Sep 2012;67(3):149-56. doi:10.1111/j.1600-0536.2012.02139.x
31. Hansson C, Pontén A, Svedman C, Bergendorff O. Reaction profile in patch testing with allergens formed during vulcanization of rubber. *Contact Dermatitis*. May 2014;70(5):300-8. doi:10.1111/cod.12168
32. Uter W, Warburton K, Weisshaar E, et al. Patch test results with rubber series in the European Surveillance System on Contact Allergies (ESSCA), 2013/14. *Contact Dermatitis*. Dec 2016;75(6):345-352. doi:10.1111/cod.12651
33. Bauer A, Geier J, Mahler V, Uter W. [Contact allergies in the German workforce : Data of the IVDK network from 2003-2013]. *Hautarzt*. Sep 2015;66(9):652-64. Kontaktallergien bei Erwerbstätigen in Deutschland : Ergebnisse des IVDK-Netzwerkes 2003-2013. doi:10.1007/s00105-015-3660-7
34. Bauer A, Geier J, Elsner P. Type IV allergy in the food processing industry: sensitization profiles in bakers, cooks and butchers. *Contact Dermatitis*. Apr 2002;46(4):228-35. doi:10.1034/j.1600-0536.2002.460408.x
35. Bauer A, Schubert S, Geier J, Mahler V. [Type IV contact allergies in the food processing industry: an update]. *Hautarzt*. Jun 2018;69(6):443-448. Typ-IV-Kontaktallergien in Nahrungsmittelberufen: Eine aktuelle Übersicht. doi:10.1007/s00105-018-4174-x
36. Warburton KL, Uter W, Geier J, et al. Patch testing with rubber series in Europe: a critical review and recommendation. *Contact Dermatitis*. Apr 2017;76(4):195-203. doi:10.1111/cod.12736
37. Aalto-Korte K, Pesonen M. Patterns of simultaneous patch test reactions to thiurams and dithiocarbamates in 164 patients. *Contact Dermatitis*. Dec 2016;75(6):353-357. doi:10.1111/cod.12687
38. Geier J, Lessmann H, Uter W, Schnuch A. Occupational rubber glove allergy: results of the Information Network of Departments of Dermatology (IVDK), 1995-2001. *Contact Dermatitis*. Jan 2003;48(1):39-44. doi:10.1034/j.1600-0536.2003.480107.x
39. Warshaw EM, Gupta R, Silverberg JI, et al. Positive Patch Test Reactions to Carba Mix and Thiuram Mix: The North American Contact Dermatitis Group Experience (1994-2016). *Dermatitis*. May-Jun 01 2021;32(3):173-184. doi:10.1097/der.0000000000000648
40. Hansen A, Brans R, Sonsmann F. Allergic contact dermatitis to rubber accelerators in protective gloves: Problems, challenges, and solutions for occupational skin protection. *Allergol Select*. 2021;5:335-344. doi:10.5414/alx02265e
41. Dejonckheere G, Herman A, Baeck M. Allergic contact dermatitis caused by synthetic rubber gloves in healthcare workers: Sensitization to 1,3-diphenylguanidine is common. *Contact Dermatitis*. Sep 2019;81(3):167-173. doi:10.1111/cod.13269

42. Hamnerius N, Pontén A, Bruze M, Svedman C, Bergendorff O. Focus Session 57 (FS-57): Decline of diphenylguanidin contact allergy among HCW after changed pattern of medical glove procurement. *Contact Dermatitis*. 2022;86(S1):7-38. doi:<https://doi.org/10.1111/cod.14095>
43. Geier J, Krautheim A, Uter W, Lessmann H, Schnuch A. Occupational contact allergy in the building trade in Germany: influence of preventive measures and changing exposure. *Int Arch Occup Environ Health*. Apr 2011;84(4):403-11. doi:10.1007/s00420-010-0581-8
44. Prodi A, Rui F, Fortina AB, Corradin MT, Filon FL. Occupational sensitization to epoxy resins in Northeastern Italy (1996-2010). *Int J Occup Environ Health*. 2015;21(1):82-7. doi:10.1179/2049396714y.0000000095
45. Radillo L, Riosa F, Mauro M, Fortina AB, Corradin MT, Larese Filon F. Contact Dermatitis in Construction Workers in Northeastern Italian Patch Test Database Between 1996 and 2016. *Dermatitis*. Nov-Dec 01 2021;32(6):381-387. doi:10.1097/der.0000000000000552
46. Suuronen K, Bäck B, Aalto-Korte K, et al. Skin exposure to epoxy chemicals in construction coating, assessed by observation, interviews, and measurements. *Contact Dermatitis*. Jan 2019;80(1):18-25. doi:10.1111/cod.13122
47. Aalto-Korte K, Pesonen M, Suuronen K. Occupational allergic contact dermatitis caused by epoxy chemicals: occupations, sensitizing products, and diagnosis. *Contact Dermatitis*. Dec 2015;73(6):336-42. doi:10.1111/cod.12445
48. Anveden Berglind I, Lind ML, Lidén C. Epoxy pipe relining - an emerging contact allergy risk for workers. *Contact Dermatitis*. Aug 2012;67(2):59-65. doi:10.1111/j.1600-0536.2011.02028.x
49. Aalto-Korte K, Bäck B, Henriks-Eckerman M, et al. EPOXY FOLDER - Chemical safety in construction coating work. Finnish Institute of Occupational Health. Accessed 5th May 2022, 2022. <https://www.ttl.fi/en/themes/tyoturvallisuus/altistuminen-tyoympariston-haittatekijöille/kemiallisten-tekijöiden-hallinta-tyopaikalla/epoxy-folder-chemical-safety-construction-coating-work>
50. Geier J, Lessmann H, Hillen U, Skudlik C, Jappe U. Sensitization to reactive diluents and hardeners in epoxy resin systems. IVDK data 2002-2011. Part II: concomitant reactions. *Contact Dermatitis*. Feb 2016;74(2):94-101. doi:10.1111/cod.12490
51. Geier J, Lessmann H, Hillen U, Skudlik C, Jappe U. Sensitization to reactive diluents and hardeners in epoxy resin systems. IVDK data 2002-2011. Part I: reaction frequencies. *Contact Dermatitis*. Feb 2016;74(2):83-93. doi:10.1111/cod.12491
52. Suomela S, Pesonen M, Ylinen K, Aalto-Korte K, Suuronen K. Characterization of patients with occupational allergy to two new epoxy hardener compounds. *Contact Dermatitis*. Jul 2022;87(1):81-88. doi:10.1111/cod.14109
53. Recommendation on MIT. Accessed 29th June 2022, 2022. https://cosmeticseurope.eu/files/3614/7634/5470/Recommendation_on_MIT.pdf
54. (SCCS) SCoCS. OPINION ON Methylisothiazolinone (P94) Submission II (Sensitisation only). Accessed 29th June 2022, http://ec.europa.eu/health/scientific_committees/consumer_safety/docs/sccs_o_145.pdf
55. Marrero-Alemán G, Borrego L, Antuña AG, Macías Montes A, Pérez Luzardo O. Isothiazolinones in cleaning products: Analysis with liquid chromatography tandem mass spectrometry of samples from sensitized patients and market. *Contact Dermatitis*. Feb 2020;82(2):94-100. doi:10.1111/cod.13430

56. Garcia-Hidalgo E, Schneider D, von Goetz N, Delmaar C, Siegrist M, Hungerbühler K. Aggregate consumer exposure to isothiazolinones via household care and personal care products: Probabilistic modelling and benzisothiazolinone risk assessment. *Environ Int.* Sep 2018;118:245-256. doi:10.1016/j.envint.2018.05.047
57. Schnuch A, Schubert S, Lessmann H, Geier J. The methylisothiazolinone epidemic goes along with changing patients' characteristics - After cosmetics, industrial applications are the focus. *Contact Dermatitis.* Feb 2020;82(2):87-93. doi:10.1111/cod.13414
58. Aerts O, Meert H, Goossens A, Janssens S, Lambert J, Apers S. Methylisothiazolinone in selected consumer products in Belgium: Adding fuel to the fire? *Contact Dermatitis.* Sep 2015;73(3):142-9. doi:10.1111/cod.12449
59. Urwin R, Craig S, Latheef F, Wilkinson M. Methylisothiazolinone: the epidemic is declining - but not gone. *Contact Dermatitis.* May 2017;76(5):301-302. doi:10.1111/cod.12750
60. Schubert S, Bauer A, Hillen U, Werfel T, Geier J, Brans R. Occupational contact dermatitis in painters and varnishers: Data from the Information Network of Departments of Dermatology (IVDK), 2000 to 2019. *Contact Dermatitis.* Nov 2021;85(5):494-502. doi:10.1111/cod.13935
61. Pablos C, Romero A, de Diego A, et al. Novel antimicrobial agents as alternative to chlorine with potential applications in the fruit and vegetable processing industry. *Int J Food Microbiol.* Nov 20 2018;285:92-97. doi:10.1016/j.ijfoodmicro.2018.07.029
62. García S, Heredia N. Microbiological Safety of Fruit and Vegetables in the Field, During Harvest, and Packaging: A Global Issue. 2017:27-48.
63. Goossens A, Aerts O. Contact allergy to and allergic contact dermatitis from formaldehyde and formaldehyde releasers: A clinical review and update. *Contact Dermatitis.* Jul 2022;87(1):20-27. doi:10.1111/cod.14089
64. Aalto-Korte K, Pesonen M. Patterns of positive patch test reactions to formaldehyde and formaldehyde releasers at the Finnish Institute of Occupational Health from 2007 to 2020. *Contact Dermatitis.* Oct 2021;85(4):429-434. doi:10.1111/cod.13876
65. Hauksson I, Pontén A, Isaksson M, Persson L, Hamada H, Bruze M. Formaldehyde in skin care products and cosmetics in dermatitis patients. *Contact Dermatitis.* 2012;66(Suppl 2):47.
66. Schubert S, Brans R, Reich A, et al. Assessment of occupational exposure and spectrum of contact sensitization in metalworkers with occupational dermatitis: results of a cohort study within the OCCUDERM project. *J Eur Acad Dermatol Venereol.* Jul 2020;34(7):1536-1544. doi:10.1111/jdv.16130
67. Schubert S, Geier J, Skudlik C, et al. Relevance of contact sensitizations in occupational dermatitis patients with special focus on patch testing of workplace materials. *Contact Dermatitis.* Dec 2020;83(6):475-486. doi:10.1111/cod.13688
68. Commission Directive 2007/17/EC of 22 March 2007 amending Council Directive 76/768/EEC, concerning cosmetic products, for the purposes of adapting Annexes III and VI thereto to technical progress (Text with EEA relevance). In: Commission E, editor. 2007.
69. Lidén C, White IR. Comment on MDBGN/DBDCB, the European baseline series, and EU legislation. *Contact Dermatitis.* Nov 2021;85(5):607-610. doi:10.1111/cod.13937
70. Herman A, Uter W, Rustemeyer T, et al. Position statement: The need for EU legislation to require disclosure and labelling of the composition of medical devices. *J Eur Acad Dermatol Venereol.* Jul 2021;35(7):1444-1448. doi:10.1111/jdv.17238

71. Alfonso JH, Bauer A, Bensefa-Colas L, et al. Minimum standards on prevention, diagnosis and treatment of occupational and work-related skin diseases in Europe - position paper of the COST Action StanDerm (TD 1206). *J Eur Acad Dermatol Venereol*. Jun 2017;31 Suppl 4:31-43. doi:10.1111/jdv.14319

Occupational contact allergy: The European perspective – analysis of patch test data from ESSCA between 01/2011-12/2020

Tested population

- 16022 patients diagnosed with OCD versus
- 46652 patients in whom occupational causation was refuted after patch testing

High risk occupations

1. Agriculture, fishery and related workers
2. Metal industry
3. Chemical industry
4. Building and construction industry
5. Health care
6. Food and service industry

Occupational relevance of EBS Allergens

was most frequently attributed to:

- Rubber accelerators
- Epoxy resins
- Preservatives




Figure 1 a: Time trend of MCI/MI contact sensitization in patients with occupational and non-occupational contact dermatitis, 2011-2020

Among allergens in the European Baseline Series, patch tested within the ESSCA network from 2011-2020 in patients with occupational contact dermatitis, occupational relevance is most frequently attributed to rubber accelerators, epoxy resins and preservatives.