


Journal Pre-proof

A New Computed Tomography Classification for Acute Mesenteric Ischaemia: More Than a “Gut Feeling”?

Jos C. van den Berg



PII: S1078-5884(23)00212-5

DOI: <https://doi.org/10.1016/j.ejvs.2023.03.005>

Reference: YEJVS 8707

To appear in: *European Journal of Vascular & Endovascular Surgery*

Received Date: 27 February 2023

Accepted Date: 7 March 2023

Please cite this article as: van den Berg JC, A New Computed Tomography Classification for Acute Mesenteric Ischaemia: More Than a “Gut Feeling”?, *European Journal of Vascular & Endovascular Surgery* (2023), doi: <https://doi.org/10.1016/j.ejvs.2023.03.005>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

1 <Short title>Invited Commentary

2 **INVITED COMMENTARY**

3

4 **A New Computed Tomography Classification for Acute Mesenteric Ischaemia: More Than a**
5 **“Gut Feeling”?**

6

7 **Jos C. van den Berg** ^{a,b,*}

8

9 ^a Centro Vascolare Ticino, Ospedale Regionale di Lugano, sede Civico, Bern, Switzerland

10 ^b Universitätsinstitut für Diagnostische, Interventionelle und Pädiatrische Radiologie
11 Inselspital, Universitätsspital Bern, Bern, Switzerland

12

13 * Corresponding author. Centro Vascolare Ticino, Ospedale Regionale di Lugano, sede Civico, Via
14 Tesserete 46, 6903 Lugano, Bern, Switzerland.

15 jos.vandenberg@eoc.ch (Jos C. van den Berg).

16

17 Knowledge of the anatomy remains a cornerstone in medicine; accordingly, a proper understanding
18 of anatomical variations is essential in the treatment of (acute) mesenteric ischaemia (AMI). In
19 this issue of *EJVES*, Tual *et al.* propose an anatomical classification based on computed
20 tomography angiography (CTA) that may help to describe occlusions of the superior mesenteric
21 artery (SMA) in patients with AMI better.¹ Over the last few decades, CTA has replaced the “gold
22 standard” of intra-arterial angiography in the evaluation of the mesenteric circulation in patients
23 with chronic and acute mesenteric ischemia. With current scan technology, that is able to provide
24 isotropic imaging, three dimensional images in multiple planes can be obtained, which are useful
25 in the depiction of the complex mesenteric vasculature.² The mesenteric vascularisation is
26 characterised by a significant variation in anatomy and the presence of various collateral pathways.

27 For example the “standard” coeliac trunk branching pattern occurs in only 70% of cases, the right
28 colic artery that can arise from the SMA is absent in up to 80% of individuals, and the number of
29 jejunal arteries may vary from four to six.²

30 When developing a new anatomical classification, given the significant number of
31 anatomical variants, it is important to include as many patients as possible. The authors have
32 chosen to limit the study to patients presenting with mesenteric ischaemia, thus eliminating the
33 possibility of establishing a large pool of (variant) anatomies. Although it is important to have a
34 cohort of patients with mesenteric ischaemia to allow for a correlation with the clinical outcome,
35 the evaluation of anatomical segments can also be done on patients with non-related disease (e.g.,
36 abdominal aneurysms). It is also remarkable that the diagnosis of AMI was not confirmed
37 clinically, but on radiological grounds only. Furthermore, it is not clear whether all CTAs were
38 evaluated interactively on a three dimensional workstation (using all possible reconstruction
39 planes). This may be an issue as a significant number of examinations was obtained from external
40 institutions, with a heterogeneous protocol, as indicated by the authors, and it is known that three
41 dimensional evaluation of a CTA increases the diagnostic accuracy.²

42 Looking specifically at the SMA variations of the colic patterns, another group has
43 proposed a classification identifying four patterns for the SMA anatomy.³ This (purely anatomical)
44 study was based on a cadaveric ($n = 50$) and a radiological sample ($n = 560$; CTA performed for
45 various pathologies, not specifically related to mesenteric ischaemia), and showed a huge
46 difference in incidence of the various patterns in both samples. The sample size as used by Tual *et*
47 *al.* ($n = 95$)¹ may therefore not be sufficiently large to allow for application in the general
48 population. This issue could have been overcome by developing the classification on a larger
49 cohort of patients that underwent CTA for other indications than mesenteric ischaemia and would
50 also have overcome the problem of eliminating patients because of an insufficient quality (lack of
51 arterial phase enhancement) of the CTA examination.

52 As the authors rightfully state, in other vascular territories (e.g., cerebral) the benefit of an
53 anatomical description/classification of collateral pathways is well known. These descriptions
54 typically include all arteries that contribute to the supply of the “end organ”, which is not done in
55 this study having excluded the contribution of the coeliac trunk and inferior mesenteric artery.

56 As mentioned, this anatomical classification should be validated in a larger cohort (“real
57 world” scenario) with multiple observers, preferably in “all comers” (not necessarily related to
58 mesenteric ischaemia). The classification should then also be applied to CTAs in patients with
59 mesenteric ischaemia where an arterial phase is not available (as is oftentimes the case), in order
60 to make this a valuable addition to the work up of patients with AMI. Of even more importance is
61 to correlate the findings with clinical outcomes and establish whether this classification can be
62 used in a decision tree to define optimal management. This should be the goal of a very “granular”
63 classification as described.

64 The gold standard in this study was a single, specialist observer, with a verification of the
65 interobserver agreement with only one, junior, observer. This incurs a risk of the apprentice
66 emulating the same methodology, which in the absence of a correlation with angiography (still
67 considered the gold standard)⁴ may not be a proper methodology.

68 Although the incidence of various patterns is reported differently in several studies, the
69 mere fact of the huge anatomical variation is important to keep in mind: depending on the distal
70 ramification patterns and connections of SMA branches with coeliac trunk, inferior mesenteric
71 arteries, or internal iliac arteries, clinical presentation and therapy may differ.

72 A classification needs to be simple, reproducible, and allow for a rapid learning curve, and
73 this seems to be the case for the classification described given the outcomes of the second reading
74 session. In the current classification a correlation with surgical and/or endovascular treatment and
75 subsequent outcome is lacking. This will be the main question that needs to be answered as an
76 occlusion at the same level and of the same extent in a patient with an “acute on chronic” occlusion
77 may not have the same clinical course as a patient with an embolic occlusion given the potential
78 lack of collaterals in the latter as it is not clear what to do with multiple segment and multiple level
79 occlusions. Given the small sample size, and the significant variability of anatomy, it will probably
80 be difficult to determine the true value of this classification in daily practice. The proof of the
81 pudding will therefore be in the eating.

82

83 **REFERENCES**

- 84 1 Tual A, Garzelli L, Nuzzo A, Corcos O, Castier Y, Ben Abdallah I, Ronot M. Strengthening the
85 description of superior mesenteric artery occlusions in acute mesenteric ischaemia: proposition for
86 an anatomical classification. *Eur J Vasc Endovasc Surg* 2023 [This Issue].
- 87 2 Horton KM, Fishman EK. CT angiography of the mesenteric circulation. *Radiol Clin North Am*
88 2010;**48**:331–45.
- 89 3 Gamo E, Jimenez C, Pallares E, Simón C, Valderrama F, Sañudo JR, Arrazola J. The superior
90 mesenteric artery and the variations of the colic patterns. A new anatomical and radiological
91 classification of the colic arteries. *Surg Radiol Anat.* 2016;**38**:519–27.
- 92 4 Ofer A, Abadi S, Nitecki S, Karram T, Kogan I, Leiderman M, et al. Multidetector CT
93 angiography in the evaluation of acute mesenteric ischemia. *Eur Radiol* 2009;**19**:24–30.

94