Processing of semantic incongruency at the onset of sleep: An auditory N400 evoked potential study



Sarah Diezig, Melissa Spaar, Simone Denzer, Peter Achermann, Fred W. Mast, Thomas Koenig

PII:	S0167-8760(23)00052-1
DOI:	https://doi.org/10.1016/j.ijpsycho.2023.03.006
Reference:	INTPSY 12194
To appear in:	International Journal of Psychophysiology
Received date:	3 January 2023
Revised date:	22 February 2023
Accepted date:	15 March 2023

Please cite this article as: S. Diezig, M. Spaar, S. Denzer, et al., Processing of semantic incongruency at the onset of sleep: An auditory N400 evoked potential study, *International Journal of Psychophysiology* (2023), https://doi.org/10.1016/j.ijpsycho.2023.03.006

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier B.V.

Processing of Semantic Incongruency at the Onset of Sleep: An Auditory N400 Evoked Potential Study

Sarah Diezig^{a,b,*}, Melissa Spaar^a, Simone Denzer^{b,c}, Peter Achermann^d, Fred W. Mast^c, Thomas Koenig^a

^a Translational Research Center, University Hospital of Psychiatry and Psychotherapy, University of Bern, Switzerland

^b Graduate School for Health Sciences, University of Bern, Switzerland

^c Department of Psychology, University of Bern, Switzerland

^d Institute of Pharmacology and Toxicology, University of ⁷ rich, Switzerland

* Corresponding author. Translational Research Center, University h. poital of Psychiatry and Psychotherapy, University of Bern, Switzerland. E-mail address: sarah.diezig@1pd.u iibe.ch

1

Abstract

The ability to organize self-generated thought into coherent, meaningful semantic representations is a central aspect of human cognition and undergoes regular alterations throughout the day. To investigate whether changes in semantic processing might explain the loss of coherence, logic, and voluntary control over thinking typically accompanying the transition to sleep, we recorded N400 evoked potentials from 44 healthy subjects. Auditory word pairs with varying semantic distance were presented while they were allowed to fall asleep. Using semantic distance and wakefulness level as regressors, we found that semantic distance reliably elicited an N400, and lower wakefulness levels were associated with increased frontal negativity within a similar time range. Additionally, and contrary to our initial hypothesis, the results showed an interaction of semantic distance and wakefulness that is best interpreted as an increased N400 effect with decreasing wakefulness. While these results do not rule out a possible role of semantic processes in the generation of the use along the transition to sleep, we discuss the possibility of additional brain methods and thought control during the transition to sleep, we discuss the possibility of additional brain methods and thought constrain the inner stream of consciousness during wakefulness.

Keywords: Transition to sleep, Event-related potential Sepantic processing, N400, Formal thought disorder, Self-generated thought

2

1. Introduction

The transition to sleep, that is, the period between relaxed wakefulness and the onset of sleep (first occurrence of sleep stage 2) is a dynamic process, with a series of physiological, behavioral, and psychological changes accompanying this state (Goupil & Bekinschtein, 2012; Ogilvie, 2001; Yang et al., 2010). This was already recognized early on by Kleitman, who stated: "Whereas it is easy to distinguish between the conditions of alertness or being wide awake, and definite sleep, the passage from one to the other involves a succession of intermediate states, part wakefulness and part sleep in varying proportions - what is designated in Italian as dormiveglia, sleep-waking" (Kleitman, 1987, p. 71). Among the most prominent indicators of this state are various neurophysiological, cardiovascular, and respiratory changes, the appearance of slow eye movements and sleep-specific components of event-related potentials (ERPs) like the N350, N550, or P900 component, a decrease in muscle strength, and a loss of behavioral responsiveness (Ogilvi 2001).

Besides the effects of decreased wakefulness on different physiologi al at 1 behavioral parameters, changes can also be observed on a level of subjective experience, making the trandition to sleep a unique state of mind (Vaitl et al., 2005). Changes in subjective experiences paralleling the physiological and behavioral alterations include the gradual loss of awareness of external stimuli, thought processes as well as sensory/perceptual experiences (e.g., hypnagogic imagery) (Siclari et al., 2013; Yang coal., 2010). In this context, especially the loss of coherence of thought, logic and voluntary control over thinking are most determinant for the correct identification of having fallen asleep (Yang et al., 200), and at the same time, they suggest that changes in mental activity across the sleep onset process are related to changes in underlying neurophysiological activity (Fox & Girn, 2017). In support of this notion, the frequency of hypnagogic imagery has been shown to correlate with specific spectral EEG parameters (e.g., Fourkes & Vogel, 1965; Hayashi et al., 1999). To date, however, the processes mediating diminished thought for the data the transition to sleep have not been investigated systematically.

As altered thought processes are considered a core feature of pathological conditions such as schizophrenia (Roche et al., 2015; Strik et al., 2017, one might assume that the loosening of associations characterizing the transition to sleep share some com ion features with the emergence of formal thought disorder in schizophrenia. Deficits in semantic processes is have been proposed as a possible cause of bizarre associations and disorganized speech in formal thought disorder, as shown by studies investigating language-related evoked potentials (see e.g., Kumar & Debruille, 2004). Thus, the loosening of associations at the transition to sleep might imply that the brain mechanisms associated with semantic processing are modified during this state. However, unlike formal thought disorder in psychosis, the transition to sleep is a regular, reversible state, which therefore provides a unique opportunity to investigate the neurophysiological correlates of spontaneous variations in the evaluation of the coherence of thought.

In wakefulness, a classical index for the activity of brain mechanisms involved in semantic processing is the N400 of evoked potentials. The N400 effect is defined as a negative potential difference that is obtained by comparing ERPs elicited by stimuli that are semantically incompatible with the preceding context against ERPs elicited by stimuli that are compatible with such a context. The N400 effect is typically observed within a time range between around 200 to 600 ms, most prominent at centro-parietal electrode sites (Kutas & Hillyard, 1980).

Together with the altered logic and coherence of thinking during the transition to sleep, we, therefore hypothesized that the N400 effect systematically varies within the range of full wakefulness to the onset of sleep.

Other studies investigating ERPs during the transition to sleep and sleep in general indicate that the neurophysiological response to specific stimuli persists with the progression towards sleep but also shows increasing modifications (for review see Atienza et al., 2001; Ibáñez et al., 2009). For example, the amplitudes of the N100 and P300 components have been shown to be decreased in slow-wave sleep compared with wakefulness or rapid eye movement (REM) sleep, suggesting that the information received by the cortex is less processed during non-REM sleep than during REM sleep (e.g., Bastuji et al., 1995; de Lugt et al., 1996). Similar to this, the mismatch negativity (MMN) could be elicited during all sleep stages but showed a diminished amplitude during sleep compared with wakefulness (Atienza et al., 2000). In contrast, the amplitude of other components like the N2 and P2, and later components, that have been a sociated with the occurrence of K-complexes, showed consistent increases with sleep progression (e.g., de Lv_{cs} , t a., 1996; Harsh et al., 1994).

Of particular relevance for the present study, previous investigations showed that the N400 could be elicited during non-REM and REM sleep, indicating an at lease partially preserved detection of semantic incongruency during these states (e.g., Brualla et al., 1998; Ib/nez et al., 2006; Perrin et al., 2002). However, none of these studies directly compared the size of the N400 effect between different levels of wakefulness or sleep stages. The current study aimed to fill this gap and locused on the transition to sleep, where variability in the coherence of thinking is particularly prominent.

Regarding the directionality and causing noclanisms underlying the N400 effect, several theories have been put forward, the two most influential of which are the theory of semantic integration and the theory of facilitated lexical access. In the semantic integration view, the N400 reflects the integration of a target word within the longstanding semantic network an trecently activated sentential context (Lau et al., 2008), which is based on previous inputs, world knot ledge, and thus evolving expectations (Kutas & Federmeier, 2011). According to this view, the N400 is the integration of a target word within the increased N400 peak the remantically incongruent stimuli is then explained by the effort needed to properly integrate an implaus ble or mismatching stimulus into an existing sentential context (Lau et al., 2008). Kutas and Federmeier (2011, highlighted the strength of this view as its capacity to account for the multi- and cross-modal nature of the N400. This is based on the assumption that all modes of access to a particular semantic information (auditory, visual, pictorial, lexical) converge to similar conceptual representations (Kutas & Federmeier, 2011). Moreover, the N400 has been argued to occur too late to only represent facilitated lexical access (Hauk et al., 2006).

In contrast, the theory of facilitated lexical access defines the process represented by the N400 as automatic and prelexical activation of brain regions associated with the target word in long-term memory (Brown & Hagoort, 1993). The increased N400 after semantically incongruent stimuli is then explained by the additional effort necessary to access the meaning of a word that has not been primed. Therefore, all conditions that ease the access to the representation of the target word, including prior congruent prime words, result in a reduced amplitude of the N400. This is argued to be based on the pre-activation of certain parts of the semantic network by the prime, facilitating the retrieval of the target word. In line with this notion, Lau et al. (2008) provided support for this view by demonstrating connections between brain regions associated with lexical access and

evoked N400 potentials (Lau et al., 2008). While this data rules out a purely integrative, i.e., post-lexical role of the N400, integration can still play an important role in facilitating lexical access. More evidence that the N400 cannot fully be explained by post-lexical processes was presented by Hoeks et al. (2004). They found that sentence endings, that were semantically related to the previous word, but made the sentence meaningless, elicited no N400 but a later occurring component called the P600, and concluded that the N400 corresponds with the lexical retrieval of a word, while the P600 represents its integration into a broader semantic context (Brouwer et al., 2012; Hoeks et al., 2004).

Despite conceptual differences, the aforementioned theories predict that the N400 effect will be reduced when subjects process word pairs with varying semantic associations while falling asleep. More specifically, based on the semantic integration theory, one needs to make assumptions about the effect of sleep onset on semantic integration. The most likely prediction is that integration is less demanding, as conceptual representations are less constrained during this state. As a consequence, find end of the integration of incongruent stimuli is expected to be smaller, thus leading to a reduced N400. This hypothesis is supported by the fact the N400 is proportionally reduced to the degree of loosened issociations in formal thought disorder or the recovery status of psychosis (Jackson et al., 2014; Kostova et al., 2005; Kumar & Debruille, 2004; Wang et al., 2011). On the other side, when N400 changes during sleep on et are predicted in terms of the theory of facilitated access, one has to make assumptions about the effect or sleep onset, it is likely that pre-activation by prime words is systematically diminished with declassing wakefulness. Therefore, the assumed priming effect is reduced, yielding an increased N400 for congruent is get words and consequently a reduced N400 effect.

By directly studying the N400 potential a ring the transition to sleep, we tried to answer whether changes in the processes associated with the N4() n, also be useful to explain phenomena such as the loss of coherence, logic, and voluntary control over thinking that are typically characterizing this state. While subjects were allowed to fall asleep during the refording, we presented them with auditory word pairs with varying degrees of semantic associations. Because of the effect of different word characteristics on the neurophysiological processing of auditory stimuli (O'Rourke & Holcomb, 2002), the N400 evoked potentials were analyzed in reference to the recognition point, that is, the time elapsed after the onset of an auditory presented stimulus that is sufficient for correct identification of the semantic content of the stimulus. On a confirmatory basis, we expected to find a consistent effect of semantic distance that is characterized by a parietal negativity associated with increasing semantic distance, and the appearance of sleep-specific ERP components such as a frontocentral negativity around 300-500 ms (Atienza et al., 2001). Moreover, we expected the N400 effect to be systematically modified with decreasing wakefulness.

2. Methods

2.1. Subjects

Forty-four healthy, native German-speaking subjects were recruited from undergraduate University students and personal acquaintances. Nine of them had to be excluded from the data analysis because of the absence of a detectable alpha rhythm with eyes closed, which was required for reliable classification of wakefulness levels (see section 2.6.2). One additional subject was excluded because of technical problems with the EEG, leaving a final sample of 34 subjects (5 males, mean age 23.50 ± 3.34 years, range 19-33 years).

Subjects were screened for exclusion of neurological, psychiatric and sleep disorders before the experiment using a semi-structured interview. Further, subjects presenting with severe uncorrected hearing problems, psychoactive or hypnotic substance use, shift work, or incompatible wake-sleep rhythm due to other reasons were excluded from the study. There were no complaints of excessive daytime sleepiness as assessed by the Epworth Sleepiness Scale (ESS; Johns, 1991; scores ≤ 10), insomnia assessed by the Regensburg Insomnia Scale (RIS; Crönlein et al., 2013; scores ≤ 12), or sleep disturbances determined by the Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1988 scores ≤ 5). Participants were instructed to follow a regular sleep schedule and to refrain from the use of alcohol the day before the experiment. We decided against any sleep restriction on the night before the experiment to avoid effects of sleep deprivation on behavior, particularly falling asleep too quickly. To minimize potential factors that prevent subjects from being able to fall asleep, they were additionally instructed to avoid daytime napping and to refrain from caffeine and nicotine approximately four hours before the experiment. The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the University of Bern (approval no. 2019-04-0 003, Written informed consent was obtained and undergraduate students received course credits for their partic bation.

2.2. Procedure

Participants were invited for the screening of exclusion crite. A several days prior to the experiment. They further filled out a set of questionnaires to assess sleeping r abits and potential sleeping problems respectively (see section 2.1). All questionnaires were presented via the browser-based survey platform Qualtrics XM (Qualtrics, 2019).

During the experiment, participants were lying in a dark, sound-attenuated room on a comfortable reclining chair. While being presented with the auditory somantic association task, they were informed that they were allowed to fall asleep. The only task was to keer the eyes closed and follow the natural course of one's mind. As the recording was part of a larger study an interval of approximately 1.5 h of resting-state EEG preceded the actual recording (Diezig et al., 2022). The presentation of the task lasted for about 35 min. The timing of the stimulus presentation was chosen to maximize the time for participants to adapt to the setting and to be able to relax irrespective of the aud tory stimulation. Moreover, the experiment took place in the early evening to increase the probability that perturbations were able to fall asleep.

2.3. Stimulus Material

The stimuli consisted of a pre-existing list of 180 pairs of semantically associated German nouns with a length of 4-5 letters, which were rated regarding their semantic distance and matched for visual, orthographic and phonological similarity (Dimigen et al., 2012). A female native speaker of German unknown to the participants spoke the words with neutral intonation and constant voice intensity. The duration of the spoken stimuli ranged from 405-918 ms (mean 628 ms). The recording and processing of the stimuli were performed using Audacity 2.2.2 (http://www.audacityteam.org/).

As subjects were allowed to sleep during the stimulus presentation, we implemented the measurements of the N400 potentials in a passive listening condition. Each trial started with the presentation of a prime word, followed by an inter-stimulus interval of 1000 ms and the presentation of a target word with varying degrees of semantic distance. The interval between two trials was set to 1500 ms. After this interval, the next trial started

without any further intervention of the subject. The order of the word pairs was pseudo-randomized to prevent randomly generated semantic associations between the trials. Each word was presented twice (one time as a prime and one time as a target), resulting in a total of 360 trials. The stimulus pairs were presented via loudspeakers at a constant volume of 50 dB at the loudspeaker level using PsychoPy2 Version 1.90.2 (Peirce et al., 2019). The loudspeakers were placed on the right and left side behind the subject, at a distance of 65 cm from the subject's head. The onset of the prime and target words was marked in the EEG.

2.4. EEG Recordings

Multichannel EEG was recorded throughout the experiment using a 64-channel actiCAP snap electrode system (Brain Products GmbH, Gilching, Germany) placed according to the extended 10-20 system (Jasper, 1958) and referenced against FCz. Active electrodes were chosen because of their improved signal quality and lowered preparation time compared to passive electrode systems. The signals were sampled at 500 Hz and stored for offline analysis using BrainVision Recorder software and BrainAmp DC amplifiers (BrainProducts GmbH, Gilching, Germany). Impedances were improved until $\leq 10 \text{ k}\Omega$. A term to resting state EEG with alternating opening and closing of the eyes was conducted before the start of the experiment to be able to identify eye movements for artifact rejection. No additional EOG or EMG channets were recorded.

2.5. Estimation of word Recognition points

To correct the effect of different word characteristics in the timing of the processing of auditory stimuli (O'Rourke & Holcomb, 2002), recognition points were determined for all the stimuli in an additional experiment using a gated word recognition task (Grosjean, 1, 80). To this end, each word was presented in fragments, starting with the first 40% of the individual word duration and continuing with the addition of 50 ms for each subsequent presentation. The stimuli were presented to the participants through headphones using PsychoPy2 Version 1.90.2 (Peirce et al., 2019). Premembers is the meaning of the presented word. The trial terminated after three consecutive rounds with correct identification of the word, or when the word ended before the participant reached the required number of context responses. Recognition points for each word were defined as the mean duration of gates across a.' bat ticir ants required for the first correct identification of the word, followed by two additional correct responses. B sed on this definition, the stimulus onset of each target word was replaced by its individual recognition point offline in the EEG. On average, the recognition point across all stimuli was 363.59 ms (SD = 79 ms).

2.6. EEG Data Processing

2.6.1. Preprocessing

EEG data preprocessing was performed using BrainVision Analyzer 2.2 (BrainProducts GmbH, Gilching, Germany) and MATLAB R2020b (Mathworks Inc. Natick, MA, USA). An independent component analysis (ICA) was applied to the resting state EEG and components typical for EOG and ECG signals were removed from the data. Remaining segments presenting physiological or technical artifacts were marked manually. Channels containing excessive artifacts were interpolated by use of spherical splines interpolation and the EEG was recomputed to average reference. In a further step, wakefulness levels were estimated (see section 2.6.2).

Finally, as the N400 is a relatively slow cognitive component, the signals were filtered between 0.3-7 Hz to improve the signal-to-noise ratio and to suppress unwanted variance in the alpha band.

2.6.2. Classification of Wakefulness levels (VIGALL)

Wakefulness levels were estimated offline by use of the Vigilance Algorithm Leipzig (VIGALL 2.1 plugin for BrainVision Analyzer, Hegerl et al., 2014). The algorithm assigns one out of seven wakefulness levels (0 = 'alert', A1/A2/A3 = 'relaxed', B1/B23 = 'drowsy', C = K-complexes or sleep spindles) at consecutive 1-s intervals based on the power and cortical distribution of the spectral EEG and the occurrence of sleep graphoelements (i.e. K-complexes and sleep spindles). The required processing steps include a manual scoring of Kcomplexes and sleep spindles and a reduction of the data to a predefined nontage. We generally followed the recommended procedure (Hegerl et al., 2016) except for some minor devicions. The horizontal eye movements (HEOG) were reconstructed based on a bipolar derivation of the prefronta, che anels F7-F8 and the threshold for detection of slow eye movements (SEM) was set to 100 μ V. Additionally we removed all segments indicating open eyes during the trials by visual inspection. The main reason μ r this was to improve the accuracy of the classification algorithm. The data was reduced to 25 channel. Bar dpass filtered between 0.5-70 Hz with an additional notch filter at 50 Hz and down-sampled to 200 Hz prior to the classification. The obtained VIGALL markers were then imported into the data with all recorded channels (see section 2.6.1).

2.7. Statistical Analysis

The present study aimed at testing whether the auditory N400 effect varied, in a trial-by-trial way, as a function of momentary wakefulness levels. For this purpose, the EEG data were segmented into trials from 100 ms before to 1000 ms after the target stimulu: ac cording to the recognition point markers. Trials containing EEG periods marked as artifacts were excluted. Each trial was then assigned to the wakefulness level of the nearest VIGALL marker.

Because of large inter-individual differences in the distribution of wakefulness levels across trials, a classical comparison of a reraged voked potentials between different wakefulness levels was not available in reasonably large subsamples or our data. Instead, we used a regression approach where the different wakefulness levels (as quantified by the VIGALL algorithm) and the semantic distance between prime and target words served as rank-scaled linear predictors of ERP variance across trials. As we were mainly interested in the interaction of wakefulness level and semantic distance, a third predictor was constructed by multiplying the previous two predictors, yielding a set of three predictor values for each trial per subject (wakefulness, semantic distance, and semantic distance x wakefulness).

To obtain the scalp field data that can be explained by the variance of these three predictors, so-called covariance maps were computed (Koenig et al., 2008), which can be considered as channel- and time-wise beta values obtained by regressing the trial-by-trial ERP data against the predictors. Since the predictors were barely correlated (all r < 0.02), we applied single instead of multiple regression (Krzywinski & Altman, 2015). As a result, for each predictor and subject, a channel-by-time matrix of beta values was obtained that was labeled individual covariance maps of semantic distance, wakefulness level, and their interaction.

To establish whether and when in the analysis window the obtained individual covariance map series generalize across the sample of subjects, we applied a Topographic Consistency Test (TCT). This procedure assesses the probability of the presence of a constant scalp field across individual observations by comparing the global field power of the averaged scalp field across subjects to the global field power of the averaged scalp field after randomly shuffling the electrode order in each data set (Koenig & Melie-García, 2010). The TCT was computed with 5'000 randomization runs and a p-threshold of 0.05. To control for effects of multiple testing, additional testing based on the duration of continuous periods of significance observed in the data was performed (Koenig & Melie-García, 2010). Moreover, to test whether the interaction effect of semantic distance x wakefulness level matches the effect of semantic distance in terms of spatial similarity, spatial correlation between the covariance maps was assessed. Topographic analysis of variance (TANOVA) of the covariance maps obtained for semantic distance alone against the semantic distance x wakefulness level were computed to confirm the similarity of the maps on a level of significance testing. Finally, un ERP data were split according to high/low semantic distance and high/low wakefulness in the 33rd percentile to show ERP waveforms of the different conditions. The result of this additional analysis is presented in the supplementary material. All data were analyzed using the software RAGU (Koenig et al., 2011) and MALLAB R2020b (Mathworks Inc. Natick, MA, USA).

3. Results

3.1. Evolution of Wakefulness

In general, the data showed an overall vr.iab ity of wakefulness levels and a noticeable decrease on



average across session time (see Fig. 1).

Fig. 1 Individual and mean course of wakefulness across trials. Grey values indicate the number of subjects assigned to a particular wakefulness level as quantified by the VIGALL algorithm (0 = `alert', A1/A2/A3 = `relaxed', B1/B23 = `drowsy', C = K-complexes or sleep spindles) for each trial. The red line shows wakefulness averaged across subjects over time.

3.2. ERP Analysis

The first aim of the ERP analysis was to establish whether and when in the analysis window the three predictors (wakefulness, semantic distance, and semantic distance x wakefulness) yielded covariance maps that generalize across the sample of subjects. The TCT showed significant consistent activation from 194 to 432 ms for the effect of semantic distance (see Fig. 2a), which sustained the duration threshold test. Additionally, time windows with consistent topographies that did not comply with the duration criterion (min duration required for significant effect = 116 ms) were found from 34 to 96 ms and from 564 to 676 ms. For the wakefulness effect depicted in Fig. 2b, topographies were significantly consistent between 262 and 556 ms. Further consistencies were found from 32 to 86 ms and from 150 to 210 ms. However, these two shorter effects were not significant when correcting for multiple testing (min duration required for significant effect = 140 ms). The effect of both predictors (semantic distance x wakefulness) combined is illustrated in Fig. 2c. The TCT showed a period of consistency from 362 to 420 ms, which nearly reached significance in the duration threshold test (p = 0.077).

On the confirmatory side of the analysis, the observed effect of semi-rite distance from 194 to 432 ms showed a left-lateralized centro-parietal negativity, confirming the existe, ∞ of a classical N400 effect in the present data. Because we analyzed the data with respect to the word β cognition points and not the word onset, the latency of the N400 effect was shifted forward. Moreover, the effect of wakefulness observed from 262 to 556 ms showed an increased frontal negativity with decreasing wakefulness, which is well in line with the pre-existing literature on sleep-specific ERP components.

Finally, when testing whether the interaction effect of semantic distance x wakefulness level matches the effect of semantic distance in terms of spatial similarity, we found that the topographic pattern of the interaction effect showed a considerable spatial similarity (r = 0.70) and temporal overlap with the effect described by semantic distance, indicating an increase of the N400 effect with decreasing wakefulness. The TANOVA of the covariance maps obtained for semantic distance alone against the semantic distance x wakefulness level covariance in the identified time window receased no significant difference between the two covariance maps (p = 0.171). Thus, while the analysis of the interaction of semantic distance and wakefulness per se did not yield fully sufficient evidence for a system. Fic effect, taking also into account the topographic and temporal similarity of the effect with the main effect of semantic distance made us conclude that wakefulness level has a systematic effect on the N400. However, Vis funding needs further validation and should be interpreted accordingly.



Fig. 2 Group averaged topographic map series (contour lines displayed in step. of ¹ μ V; red positive and blue negative electric potential) and p-values (color bar at right; p-values uncorrected) of t te top ographic consistency test (TCT) at every time point (depicted by color scales) for each of the three conditions *seman. c distance* (a), *wakefulness* (b) and *wakefulness x semantic distance* (c). Covariance maps are displayed every 50 ms betv een v and 600 ms after the target stimulus recognition point.

4. Disc 15.10.1

The present study examined the effect of a creating wakefulness on the expression of the N400 component. Given that coherence, logic and vol. tary control over thinking are frequently lost during the transition to sleep, and considering the most prominent current explanatory approaches of the N400 component, we expected the N400 effect (increasing n:entroparietal electrodes with increasing incongruency) to be systematically reduced with decreasing wakefulness. By using a passive auditory semantic association task and correcting the or set of the evoked potentials of each trial for its word recognition point, we were able to elicit an N400 potent. 1 characterized by a centro-parietal negativity associated with increasing semantic distance between 194 and 422 ms after the target recognition point while subjects were falling asleep. Moreover, there was a significant main effect of wakefulness, confirming the existence of additional sleepspecific ERP components in the data (Atienza et al., 2001). The association of decreased wakefulness with a late frontal negativity has repeatedly been linked to the presence of K-complexes (Colrain, 2005). Since some but not all of our subjects reached VIGALL state C (which by definition contains K-complexes) during the recording, this seems quite likely but not conclusive. On the contrary, the hypothesis of a decrease in the N400 effect when falling asleep could not be confirmed. Instead, we found an increase in N400 with decreasing wakefulness, which overlapped with the main effect of semantic distance in time and spatial configuration. Importantly, this effect cannot be explained by the mere repetition of word pairs. As word repetition is known to attenuate the N400 (Hohlfeld, Ullsperger, & Sommer, 2008), and repetitions accumulate over time, one would expect a decreased N400 over time. However, the observation that the N400 increased with lower wakefulness, and that wakefulness was lower towards the end of the experiment (Fig. 1) implies that the N400 rather increased over time, contradicting an explanation of the effect by word repetition. Thus, unlike other conditions where a reduction of the N400 effect has been reported to be associated with impaired thought processes like formal thought disorder or delusions (Kiang & Gerritsen, 2019; Kostova et al., 2005), we could not extend these

findings to the condition of falling asleep - a state with similar phenomenological characteristics (Goupil & Bekinschtein, 2012). Therefore, these findings offer novel and interesting insights into the neuro- and psychophysiology of falling asleep.

As the N400 potential is thought to represent an index of semantic processing, this raises the question of the possible mechanisms that contribute to the observation of an enhanced N400 effect during a state, which rather implicates deficient thought processes. At the neural level, the processing of semantic information (McKiernan et al., 2006; Seghier & Price, 2012; Wirth et al., 2011; Zhang et al., 2022) but also the generation of spontaneous, task-unrelated thoughts (Binder et al., 1999, 2009; Smallwood et al., 2012) have been shown to critically depend on the activity of several large-scale networks, including the so-called default mode network (DMN). Regarding the N400 component, the involvement of these networks has been found to depend on whether retrieval is usefully constrained or miscued by the surrounding context (Jacob et al., 2019; Lanzoni et al., 2020). Whereas a 'semantic control network' (including the inferior '... 'ta, and posterior middle temporal gyrus) is recruited for the processing of incongruent stimuli, DMN active. Las been specifically related to the processing of congruent stimuli. This suggests a facilitatory role c^c the DMN for priming (Jackson, 2021; Lanzoni et al., 2020; Lau et al., 2008). Moreover, DMN ac. vity inversely correlated with unusual thoughts/thought disorder in schizophrenia, which led to the assumption that the DMN may support the combination of concepts into meaningful representations (Harc et al., 2019; Jacob et al., 2019). At the same time, spontaneous, stimulus-independent thoughts, like the on " that might characterize mental activity during the transition to sleep, have been supposed to be ess ... al., semantic because they depend on the activation and manipulation of acquired knowledge about the world (Binder et al., 1999, 2009).

In line with this notion, activation and functional connectivity in the DMN (Horovitz et al., 2008; Picchioni et al., 2008) but not the posterior cingulate control (PCC) (Kinreich et al., 2014) were found to increase during light sleep (but see Larson-Prior et al., 2014) tor loss of DMN connectivity). Together with the fact that DMN activity is involved in semantic proming, this might explain why the N400 effect, contrary to our initial expectation, did not decrease but rather increased when subjects were falling asleep. According to the lexical retrieval account of the N400, pro-accuvation or priming by congruent prime words might not be systematically diminished during the transition to sleep but rather increased as a result of predominant DMN activity, yielding a decreased N400 for congruent target words and therefore an increased N400 effect. However, whereas DMN activity might also drive the occurrence of spontaneous thoughts during the transition to sleep in general, it does not provide an explanation for the loss of control and coherence over thinking characterizing this state.

On the other side, DMN activity has also been proposed to be involved in the post-lexical integration of semantic stimuli. Based on the theory of semantic integration, the smaller N400 effect often reported in schizophrenia is usually thought to result from a smaller N400 amplitude for incongruent targets, facilitating semantic integration of inappropriate content (Besche-Richard et al., 2014; Mohammad & de Lisi, 2013). In contrast, the increase of the N400 effect observed in our study might therefore suggest that the effort made for the integration of incongruent stimuli rather becomes larger during the associatively loosened state of falling asleep, thus leading to an increased N400 effect. However, since DMN activity has been shown to be stronger when multiple congruent primes were presented in a semantic priming task compared with a single prime (Lanzoni et al., 2020; Vatansever et al., 2015), the increased N400 effect might also be the result of facilitated integration of congruent targets, while processing of incongruent targets remains unaffected. As we used the

semantic distance between prime and target words as a linear predictor for ERP variance between trials and it is not possible to control for an equal distribution of congruent and incongruent trials across wakefulness levels, we are not able to test which of the above options is accurate. The supplementary analysis of ERP waveforms split by high/low semantic distance and high/low wakefulness (see supplementary figures S3) revealed a more negative potential for incongruent targets as well as a more positive potential for congruent targets.

In schizophrenia, both, deficient early automatic semantic activation and late post-lexical processes have been put forward to explain deficits in semantic processing. Whereas it has been argued that these views are rather complementary than in conflict (Wang et al., 2011), they may also be partly dependent on differences in experimental task properties such as stimulus onset asynchrony (SOA) or the type of paradigm used to measure the N400 response to semantic processing (Kiang & Gerritsen, 2019; Kreher et al., 2009; Mohammad & de Lisi, 2013). In turn, these factors either might favor the predominance of early or late processes (Kuperberg et al., 2007). Longer SOAs for instance, that is, the time passed between the or set or the prime and the onset of the target stimulus, have been associated with a more negative N400, whereas priming paradigms using a short SOA produced reduced N400 (Mohammad & de Lisi, 2013). Similarly the explicit tasks, patients showed reduced direct and indirect semantic priming in comparison with healthy controls whereas in implicit tasks, they showed normal or, in positively thought-disordered patients, increated cirect and indirect N400 priming effects compared with healthy controls (Kreher et al., 2009).

As we used a task with a long SOA, one might angue that processes involving semantic integration dominated in the present study. However, the conclusion that the integration of incongruent stimuli either recruits more resources when falling asleep or alter. Atively, facilitated integration of congruent targets leads to an increased N400 effect both contradict the pecific phenomenological characteristics observed during this state. Moreover, activity of the DMN has been suggested to be critical for organizing thoughts into coherent, meaningful utterances (Hare et al., 20 9). However, this does not apply to the transition to sleep, as DMN activity is increased, but logic and where, ce of thinking decrease during that time. Therefore, it is likely that DMN activity cannot account for the effect alone but additional brain mechanisms that possibly mediate the generation of the N400 effect ... the eenhanced or disinhibited during the transition to sleep, providing some further explanatory value. Fo. example, the semantic control network has been shown to be activated when retrieval is miscued by the su rounding context (Lanzoni et al., 2020), and is thought to be involved in the selection of relevant aspects of knowledge while irrelevant information is suppressed (Jackson, 2021), but its interaction with wakefulness has not been systematically investigated so far. Moreover, the presence of a specific EEG microstate (reflecting whole-brain network dynamics) with sources in frontoparietal brain regions was associated with subjective experiences such as diminished thought control during the transition to sleep (Diezig et al., 2022) but it remains unclear whether and how this network is involved in the processing of semantic information.

5. Conclusion

The present study investigated brain mechanisms underlying the loss of logic, coherence and voluntary control over thinking observed during the transition to sleep. Based on the hypothesis that semantic processing might be important in this context, we expected the N400 effect of the auditory event-related potentials to be reduced when falling asleep. However, the results showed an enhanced N400 effect with decreasing

wakefulness. Default mode network activity is involved in semantic processing, it increases when falling asleep, and it might partly explain these results but cannot account for the phenomenological experiences characterizing this state. While our findings do not exclude that semantic processing has a part in deficient thought processes, they stress the involvement of additional brain mechanisms that usually constrain the inner stream of consciousness.

Acknowledgments

The authors kindly thank Anja Baumberger for helping with the recording of the stimulus material.

Author contributions

Sarah Diezig: methodology, conceptualization, software, stimulus material, investigation, data curation, software, formal analysis, visualization, and writing – original draft. Meli sa Spaar: writing N400 literature – original draft, formal analysis. Simone Denzer: conceptualization, stimulus material, writing – review and editing. Peter Achermann: supervision, writing – review and editing. Fre l W Mast: funding acquisition, conceptualization, resources, writing – review and editing. Thomas Ko, vig: conceptualization, methodology, formal analysis, writing – review and editing, resources and supervision.

Declarations

Funding

This work was supported by the Interfaculty Research 'or peration (IRC) Grant "Decoding Sleep: From Neurons to Health and Mind" funded by the University on Pern. The funding source was not involved in the study design, collection, analysis or interpretation of .ne data, the writing or publication of the report.

Del'arations of interest

None.

Ethics approval

This study was conducted in acco. Jance with the principles of the Declaration of Helsinki. This study was approved by the Ethics Committee of the University of Bern (approval no. 2019-04-00003).

Consent to participate

Informed constant was obtained from all individual participants included in the study.

Data availability

The datasets generated in this study are available from the corresponding author upon reasonable request.

References

- Atienza, M., Cantero, J. L., & Escera, C. (2001). Auditory information processing during human sleep as revealed by event-related brain potentials. *Clinical Neurophysiology*, 112(11), 2031–2045. https://doi.org/10.1016/S1388-2457(01)00650-2
- Atienza, M., Cantero, J. L., & Gómez, C. M. (2000). Decay time of the auditory sensory memory trace during wakefulness and REM sleep. *Psychophysiology*, 37(4), 485–493. https://doi.org/10.1111/1469-8986.3740485
- Bastuji, H., García-Larrea, L., Franc, C., & Mauguière, F. (1995). Brain processing of stimulus deviance during slow-wave and paradoxical sleep: a study of human auditory evoked responses using the oddball

paradigm. Journal of Clinical Neurophysiology : Official Publication of the American Electroencephalographic Society, 12(2), 155–167. https://doi.org/10.1097/00004691-199503000-00006

- Besche-Richard, C., Iakimova, G., Hardy-Baylé, M. C., & Passerieux, C. (2014). Behavioral and brain measures (N400) of semantic priming in patients with schizophrenia: Test-retest effect in a longitudinal study. *Psychiatry and Clinical Neurosciences*, 68(5), 365–373. https://doi.org/10.1111/pcn.12137
- Binder, J. R., Desai, R. H., Graves, W. W., & Conant, L. L. (2009). Where Is the Semantic System? A Critical Review and Meta-Analysis of 120 Functional Neuroimaging Studies. *Cerebral Cortex*, 19(12), 2767– 2796. https://doi.org/10.1093/CERCOR/BHP055
- Binder, J. R., Frost, J. A., Hammeke, T. A., Bellgowan, P. S. F., Rao, S. M., & Cox, R. W. (1999). Conceptual Processing during the Conscious Resting State: A Functional MRI Study. *Journal of Cognitive Neuroscience*, 11(1), 80–93. https://doi.org/10.1162/089892999563265
- Brouwer, H., Fitz, H., & Hoeks, J. (2012). Getting real about Semantic Illusions: Rethinking the functional role of the P600 in language comprehension. *Brain Research*, *1446*, 127–112. https://doi.org/10.1016/J.BRAINRES.2012.01.055
- Brown, C., & Hagoort, P. (1993). The Processing Nature of the N400: Evicine from Masked Priming. *Journal of Cognitive Neuroscience*, 5(1), 34–44. https://doi.org/10.1162/ OCi V.1993.5.1.34
- Brualla, J., Romero, M. F., Serrano, M., & Valdizán, J. R. (1998). A. ⁴itory event-related potentials to semantic priming during sleep. *Electroencephalography and Clinic & Ne worphysiology Evoked Potentials*, 108(3), 283–290. https://doi.org/10.1016/S0168-5597(97)00102-0
- Buysse, D. J., Reynolds III, C. F., Monk, T. H., Berman, S. K., & Kupfer, D. J. (1988). The Pittsburgh Sleep Quality Index: A New Instrument for Psychiatric Praerice and Research. *Psychiatry Research*, 28, 193–213.
- Colrain, I. M. (2005). The K-Complex: A 7-Decal Yistory. *Sleep*, 28(2), 255–273. https://doi.org/10.1093/SLEEP/28.2.255
- Crönlein, T., Langguth, B., Popp, R., Lukesci, H., Pieh, C., Hajak, G., & Geisler, P. (2013). Regensburg Insomnia Scale (RIS): a new short rat.n⁻, scale for the assessment of psychological symptoms and sleep in insomnia; Study design: development and validation of a new short self-rating scale in a sample of 218 patients suffering from insomnic and 94 healthy controls. *Health and Quality of Life Outcomes*, *11*(65). http://www.hqlo.com/content/11/⁻¹/65
- de Lugt, D. R., Loewy, D. H., & Composell, K. B. (1996). The effect of sleep onset on event related potentials with rapid rates of stimulus presentation. *Electroencephalography and Clinical Neurophysiology*, 98(6), 484–492. https://doi.org/10.101/0013-4694(96)94726-4
- Diezig, S., Denzer, S., Acherm Inn, P., Mast, F. W., & Koenig, T. (2022). EEG Microstate Dynamics Associated with Dream-Like Experiences During the Transition to Sleep. *Brain Topography*. https://doi.org/10.1007/s10548-022-00923-y
- Dimigen, O., Kliegl, R., & Sommer, W. (2012). Trans-saccadic parafoveal preview benefits in fluent reading: A study with fixation-related brain potentials. *NeuroImage*, 62(1), 381–393. https://doi.org/10.1016/j.neuroimage.2012.04.006
- Foulkes, D., & Vogel, G. (1965). Mental activity at sleep onset. *Journal of Abnormal Psychology*, 70(4), 231–243.
- Fox, K. C. R., & Girn, M. (2017). Neural correlates of self-generated imagery and cognition throughout the sleep cycle. In *The Oxford Handbook of Spontaneous Thought* (pp. 371–384).
- Goupil, L., & Bekinschtein, T. A. (2012). Cognitive processing during the transition to sleep. Archives Italiennes de Biologie, 150(2–3), 140–154. https://doi.org/10.4449/aib.v150i2.1247
- Grosjean, F. (1980). Spoken word recognition processes and the gating paradigm. *Perception & Psychophysics*, 28(4), 267–283.

- Hare, S. M., Ford, J. M., Mathalon, D. H., Damaraju, E., Bustillo, J., Belger, A., Lee, H. J., Mueller, B. A., Lim, K. O., Brown, G. G., Preda, A., van Erp, T. G. M., Potkin, S. G., Calhoun, V. D., & Turner, J. A. (2019). Salience–Default Mode Functional Network Connectivity Linked to Positive and Negative Symptoms of Schizophrenia. *Schizophrenia Bulletin*, 45(4), 892–901. https://doi.org/10.1093/SCHBUL/SBY112
- Harsh, J., Voss, U., Hull, J., Schrepfer, S., & Badia, P. (1994). ERP and behavioral changes during the wake/sleep transition. *Psychophysiology*, *31*(3), 244–252. https://doi.org/10.1111/J.1469-8986.1994.TB02213.X
- Hauk, O., Davis, M. H., Ford, M., Pulvermüller, F., & Marslen-Wilson, W. D. (2006). The time course of visual word recognition as revealed by linear regression analysis of ERP data. *NeuroImage*, 30(4), 1383–1400. https://doi.org/10.1016/J.NEUROIMAGE.2005.11.048
- Hayashi, M., Katoh, K., & Hori, T. (1999). Hypnagogic imagery and EEG activity. *Perceptual and Motor Skills*, 88, 676–678.
- Hegerl, U., Hensch, T., Böttger, D., & Sander, C. (2014). VIGALL 2.0: Ana. ring different functional brain states and their regulation during resting states.
- Hegerl, U., Sander, C., Ulke, C., Böttger, D., Hensch, T., Huang, J., Mauch, N., & Olbrich, S. (2016). Vigilance Algorithm Leipzig (VIGALL) Version 2.1 Manual.
- Hoeks, J. C. J., Stowe, L. A., & Doedens, G. (2004). Seeing words in context: the interaction of lexical and sentence level information during reading. *Cognitive Brai*. *Re. 2arch*, *19*(1), 59–73. https://doi.org/10.1016/J.COGBRAINRES.2003.10.022
- Hohlfeld, A., Ullsperger, P., & Sommer, W. (2008). How *Loc* 5 the incrementality of auditory word perception interplay with episodic and semantic memory? *Journ. Lo. Neurolinguistics*, 21(4), 279–293. https://doi.org/10.1016/J.JNEUROLING.2007 06. 02
- Horovitz, S. G., Fukunaga, M., de Zwart, J. A., va. Celderen, P., Fulton, S. C., Balkin, T. J., & Duyn, J. H. (2008). Low frequency BOLD fluctuations durn.³ resting wakefulness and light sleep: A simultaneous EEG-fMRI study. *Human Brain Mapping*, ⁹⁹(6), 671–682. https://doi.org/10.1002/HBM.20428
- Ibáñez, A., López, V., & Cornejo, C. (2006, 'J.R 's and contextual semantic discrimination: Degrees of congruence in wakefulness and sleep. *Prain and Language*, 98(3), 264–275. https://doi.org/10.1016/j.bandl.?00505
- Ibáñez, A., San Martín, R., Hurtado, Γ, & López, V. (2009). ERPs studies of cognitive processing during sleep. International Journal of Psychology, 44(4), 290–304. https://doi.org/10.1080/00207590802194234
- Jackson, F., Foti, D., Kotev, F., Pelman, G., Mathalon, D. H., & Proudfit, G. H. (2014). An incongruent reality: The N400 in relation to psychosis and recovery. *Schizophrenia Research*, *160*(1–3), 208–215. https://doi.org/10.1016/J SCHRES.2014.09.039
- Jackson, R. L. (2021). The neural correlates of semantic control revisited. *NeuroImage*, 224, 117444. https://doi.org/10.1016/J.NEUROIMAGE.2020.117444
- Jacob, M. S., Ford, J. M., Roach, B. J., Calhoun, V. D., & Mathalon, D. H. (2019). Aberrant activity in conceptual networks underlies N400 deficits and unusual thoughts in schizophrenia. *NeuroImage: Clinical*, 24, 101960. https://doi.org/10.1016/J.NICL.2019.101960
- Jasper, H. (1958). Report of the committee on methods of clinical examination in electroencephalography. *Electroencephalogr Clin Neurophysiol*, *10*(2), 370–375. https://doi.org/10.1016/0013-4694(58)90053-1
- Johns, M. W. (1991). A New Method for Measuring Daytime Sleepiness: The Epworth Sleepiness Scale. *Sleep*, 14(6), 540–545. https://academic.oup.com/sleep/article/14/6/540/2742871
- Kiang, M., & Gerritsen, C. J. (2019). The N400 event-related brain potential response: A window on deficits in predicting meaning in schizophrenia. *International Journal of Psychophysiology*, 145, 65–69. https://doi.org/10.1016/J.IJPSYCHO.2019.04.005

- Kinreich, S., Podlipsky, I., Jamshy, S., Intrator, N., & Hendler, T. (2014). Neural dynamics necessary and sufficient for transition into pre-sleep induced by EEG NeuroFeedback. *NeuroImage*, 97, 19–28. https://doi.org/10.1016/J.NEUROIMAGE.2014.04.044
- Kleitman, N. (1987). Sleep and Wakefulness. Chicago, IL: University of Chicago Press.
- Koenig, T., Kottlow, M., Stein, M., & Melie-García, L. (2011). Ragu: A free tool for the analysis of EEG and MEG event-related scalp field data using global randomization statistics. *Computational Intelligence and Neuroscience*, 2011. https://doi.org/10.1155/2011/938925
- Koenig, T., & Melie-García, L. (2010). A method to determine the presence of averaged event-related fields using randomization tests. *Brain Topography*, 23(3), 233–242. https://doi.org/10.1007/s10548-010-0142-1
- Koenig, T., Melie-García, L., Stein, M., Strik, W., & Lehmann, C. (2008). Establishing correlations of scalp field maps with other experimental variables using covariance analysis and resampling methods. *Clinical Neurophysiology*, 119(6), 1262–1270. https://doi.org/10.1016/j.clinph.2007.12.023
- Kostova, M., Passerieux, C., Laurent, J.-P., & Hardy-Baylé, M.-C. (2005). N+30 anomalies in schizophrenia are correlated with the severity of formal thought disorder. *Schizophren'a Kesearch*, 78(2–3), 285–291. https://doi.org/10.1016/j.schres.2005.05.015
- Kreher, D. A., Goff, D., & Kuperberg, G. R. (2009). Why all the corfution' Experimental task explains discrepant semantic priming effects in schizophrenia under "au omatic" conditions: Evidence from Event-Related Potentials. *Schizophrenia Research*, 111(1–3), 17(–18). https://doi.org/10.1016/J.SCHRES.2009.03.013
- Krzywinski, M., & Altman, N. (2015). Multiple linear regression. *Nature Methods*, *12*(12), 1103–1104. https://doi.org/10.1038/NMETH.3665
- Kumar, N., & Debruille, ; J Bruno. (2004). Semar LCs and 13400: insights for schizophrenia. J Psychiatry Neurosci, 29(2).
- Kuperberg, G. R., Deckersbach, T., Holt, D. J., Goff, D., & West, W. C. (2007). Increased Temporal and Prefrontal Activity in Response to Ser. in 'ic Associations in Schizophrenia. Archives of General Psychiatry, 64(2), 138–151. https://dci.org/10.1001/ARCHPSYC.64.2.138
- Kutas, M., & Federmeier, K. D. (2011). Thirty years and counting: Finding meaning in the N400 component of the event related brain potential (ERt). Annual Review of Psychology, 62, 621. https://doi.org/10.1146/ANNU.PEV.PSYCH.093008.131123
- Kutas, M., & Hillyard, S. A. (1000). Reading Senseless Sentences: Brain Potentials Reflect Semantic Incongruity. *Science* 207(44-7), 203–205. https://doi.org/10.1126/SCIENCE.7350657
- Lanzoni, L., Ravasio, D., Thon pson, H., Vatansever, D., Margulies, D., Smallwood, J., & Jefferies, E. (2020). The role of default mode network in semantic cue integration. *NeuroImage*, 219, 117019. https://doi.org/10.1016/J.NEUROIMAGE.2020.117019
- Larson-Prior, L. J., Power, J. D., Vincent, J. L., Nolan, T. S., Coalson, R. S., Zempel, J., Snyder, A. Z., Schlaggar, B. L., Raichle, M. E., & Petersen, S. E. (2011). Modulation of the brain's functional network architecture in the transition from wake to sleep. *Progress in Brain Research*, 193, 277–294. https://doi.org/10.1016/B978-0-444-53839-0.00018-1
- Lau, E. F., Phillips, C., & Poeppel, D. (2008). A cortical network for semantics: (de)constructing the N400. *Nature Reviews Neuroscience 2008 9:12*, 9(12), 920–933. https://doi.org/10.1038/nrn2532
- McKiernan, K. A., D'Angelo, B. R., Kaufman, J. N., & Binder, J. R. (2006). Interrupting the "stream of consciousness": An fMRI investigation. *NeuroImage*, 29(4), 1185–1191. https://doi.org/10.1016/J.NEUROIMAGE.2005.09.030
- Mohammad, O. M., & de Lisi, L. E. (2013). N400 in schizophrenia patients. *Current Opinion in Psychiatry*, 26(2), 196–207. https://doi.org/10.1097/YCO.0B013E32835D9E56

- Ogilvie, R. D. (2001). The process of falling asleep. *Sleep Medicine Reviews*, 5(3), 247–270. https://doi.org/10.1053/smrv.2001.0145
- O'Rourke, T. B., & Holcomb, P. J. (2002). Electrophysiological evidence for the efficiency of spoken word processing. *Biological Psychology*, *60*, 121–150.
- Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, 51(1), 195–203. https://doi.org/10.3758/S13428-018-01193-Y
- Perrin, F., Bastuji, H., & Garcia-Larrea, L. (2002). Detection of verbal discordances during sleep. *NeuroReport*, 13(10), 1345–1349. https://doi.org/10.1097/00001756-200207190-00026
- Picchioni, D., Fukunaga, M., Carr, W. S., Braun, A. R., Balkin, T. J., Duyn, J. H., & Horovitz, S. G. (2008). fMRI differences between early and late stage-1 sleep. *Neuroscience Letters*, 441(1), 81–85. https://doi.org/10.1016/J.NEULET.2008.06.010
- Roche, E., Creed, L., Macmahon, D., Brennan, D., & Clarke, M. (2015). The Lyidemiology and Associated Phenomenology of Formal Thought Disorder: A Systematic Review Sci. zophrenia Bulletin, 41(4), 951– 962. https://doi.org/10.1093/schbul/sbu129
- Seghier, M. L., & Price, C. J. (2012). Functional heterogeneity within the dufault network during semantic processing and speech production. *Frontiers in Psychology*, 3(1) UG), 281. https://doi.org/10.3389/FPSYG.2012.00281/BIBTEX
- Siclari, F., LaRocque, J. J., Postle, B. R., & Tononi, G. (2013). Assessing sleep consciousness within subjects using a serial awakening paradigm. *Frontiers in Psy. ncicgy*, 4. https://doi.org/10.3389/fpsyg.2013.00542
- Smallwood, J., Brown, K., Baird, B., & Schooler, J. W. '2012). Cooperation between the default mode network and the frontal-parietal network in the production of an internal train of thought. *Brain Research*, *1428*, 60–70. https://doi.org/10.1016/J.BRAINRE_2.11.03.072
- Strik, W., Stegmayer, K., Walther, S., & Dierk, T. (2017). Systems Neuroscience of Psychosis: Mapping Schizophrenia Symptoms onto Brain System. *Neuropsychobiology*, 75(3), 100–116. https://doi.org/10.1159/000485221
- Vaitl, D., Gruzelier, J., Jamieson, G. A., Jehmann, D., Ott, U., Sammer, G., Strehl, U., Birbaumer, N., Kotchoubey, B., Kübler, A., M. tner, W. H. R., Pütz, P., Strauch, I., Wackermann, J., & Weiss, T. (2005). Psychobiology of altered states of consciousness. In *Psychological Bulletin* (Vol. 131, Issue 1, pp. 98– 127). https://doi.org/10.105./0033-2909.131.1.98
- Vatansever, D., Menon, Γ K. Mar ktelow, A. E., Sahakian, B. J., & Stamatakis, E. A. (2015). Behavioral/Cognitive L fault Mode Dynamics for Global Functional Integration. https://doi.org/10.1523/J .EUROSCI.2135-15.2015
- Wang, K., Cheung, E. F. C., Gong, Q. yong, & Chan, R. C. K. (2011). Semantic Processing Disturbance in Patients with Schizophrenia: A Meta-Analysis of the N400 Component. *PLOS ONE*, 6(10), e25435. https://doi.org/10.1371/JOURNAL.PONE.0025435
- Wirth, M., Jann, K., Dierks, T., Federspiel, A., Wiest, R., & Horn, H. (2011). Semantic memory involvement in the default mode network: A functional neuroimaging study using independent component analysis. *NeuroImage*, 54(4), 3057–3066. https://doi.org/10.1016/J.NEUROIMAGE.2010.10.039
- Yang, C., Han, H., Yang, M., Su, W., & Lane, T. (2010). What subjective experiences determine the perception of falling asleep during sleep onset period ? *Consciousness and Cognition*, 19(4), 1084–1092. https://doi.org/10.1016/j.concog.2009.12.017
- Zhang, G., Hung, J., & Lin, N. (2022). Coexistence of the social semantic effect and non-semantic effect in the default mode network. *Brain Structure and Function*, 1, 1–19. https://doi.org/10.1007/S00429-022-02476-Z/FIGURES/3

Highlights

- Logic and control over thoughts are frequently lost at the onset of sleep
- Auditory N400 evoked potentials were investigated during the transition to sleep
- The N400 effect increased with decreasing wakefulness
- The results contradict N400 findings in diminished thought control in psychosis