

Current Eye Research



ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/icey20

Analysis of Biomechanical Response After Corneal Crosslinking with Different Fluence Levels in Porcine Corneas

Isaak Fischinger, Sophia A. Reifeltshammer, Theo G. Seiler, Malavika H. Nambiar, Maria A. Komninou, Philippe Büchler, Jascha Wendelstein, Achim Langenbucher & Matthias Bolz

To cite this article: Isaak Fischinger, Sophia A. Reifeltshammer, Theo G. Seiler, Malavika H. Nambiar, Maria A. Komninou, Philippe Büchler, Jascha Wendelstein, Achim Langenbucher & Matthias Bolz (2023): Analysis of Biomechanical Response After Corneal Crosslinking with Different Fluence Levels in Porcine Corneas, Current Eye Research, DOI: 10.1080/02713683.2023.2205612

To link to this article: <u>https://doi.org/10.1080/02713683.2023.2205612</u>

9

© 2023 The Author(s). Published with license by Taylor & Francis Group, LLC

ď	-	-0		
			Ľ	

Published online: 05 May 2023.



Submit your article to this journal 🕝



View related articles 🗹



View Crossmark data 🗹

Taylor & Francis Taylor & Francis Group

OPEN ACCESS Check for updates

Analysis of Biomechanical Response After Corneal Crosslinking with Different Fluence Levels in Porcine Corneas

Isaak Fischinger^{a,b,c}, Sophia A. Reifeltshammer^{b,c}, Theo G. Seiler^{d,e,f}, Malavika H. Nambiar^{g,h}, Maria A. Komninou^{g,h}, Philippe Büchler^g, Jascha Wendelstein^{b,c} , Achim Langenbucherⁱ, and Matthias Bolz^{b,c}

^aAugentagesklinik Spreebogen Berlin, Berlin, Germany; ^bKlinik für Augenheilkunde, Kepler Universitätsklinikum, Linz, Austria; ^cMedizinische Fakultät, Johannes Kepler Universität Linz, Linz, Austria; ^dKlinik für Augenheilkunde, Universitätsklinkum Düsseldorf, Düsseldorf, Germany; ^eInstitut für Refraktive und Ophthalmo-Chirurgie (IROC), Zurich, Switzerland; ^fDepartment of Ophthalmology, Inselspital Bern, University of Bern, Bern, Switzerland; ^gARTORG Center for Biomedical Engineering Research, University of Bern, Bern, Switzerland; ^hGraduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland; ⁱDepartment of Experimental Ophthalmology, Saarland University, Homburg, Germany

ABSTRACT

Purpose: To evaluate corneal stiffening of porcine corneas induced by corneal crosslinking (CXL) with constant irradiance as a function of total fluence.

Methods: Ninety corneas from freshly enucleated porcine eyes were divided into five groups of 18 eyes. Groups 1–4 underwent epi-off CXL using a dextran-based riboflavin solution and an irradiance of 18 mW/cm², group 5 served as the control group. Groups 1 to 4 were treated with a total fluence of 20, 15, 10.8, and 5.4 J/cm², respectively. Thereafter, biomechanical measurements were performed on 5 mm wide and 6 mm long strips using an uniaxial material tester. Pachymetry measurements were performed on each cornea.

Results: At 10% strain, the stress was 76, 56, 52, and 31% higher in groups 1–4, respectively compared to the control group. The Young's modulus was 2.85 MPa for group 1, 2.53 MPa for group 2, 2.46 MPa for group 3, 2.12 MPa for group 4, and 1.62 MPa for the control group. The difference between groups 1 to 4 and the control group 5 were statistically significant (p = <0.001; p = <0.001; p = <0.001; p = 0.021). In addition, group 1 showed significantly more stiffening than group 4 (p = <0.001), no other significant differences were found. Pachymetry measurements revealed no statistically significant differences among the five groups.

Conclusion: Additional mechanical stiffening can be achieved by increasing the fluence of the CXL. There was no threshold detected up to 20 J/cm². A higher fluence could compensate the weaker effect of accelerated or epi-on CXL procedures.

ARTICLE HISTORY Received 1 March 2022

Accepted 14 April 2023

KEYWORDS

Corneal crosslinking; keratoconus; fluence; corneal biomechanics; accelerated

Introduction

To date corneal crosslinking (CXL) is the only treatment that halts the progression of keratoconus by increasing the stiffness of the weakened corneal stroma.¹ Therefore, riboflavin is applied to the cornea and subsequently irradiated by UV-A light to induce new bonds within the corneal extracellular matrix.^{2,3} During this process, reactive oxygen species (ROS) are generated, which are crucial for the process, but also harmful to vulnerable keratocytes, corneal endothelial cells, and more posterior structures. Therefore, safety limits, such as minimal corneal thickness and total applied UV-energy were postulated when CXL was introduced. These limits have been valid for many years.4,5 Recent studies, however, showed that the riboflavin concentration at the endothelium level is substantially lower than anticipated in the initial safety calculations.⁶ Accordingly, the traditional fluence limit of 5.4 J/cm² has been violated in newer

protocols.^{7–9} Hence, thinner corneas have been treated by adjusting the fluence levels to the minimal thickness, newer epi-on treatments were evaluated,^{10,11} or customized energy levels inducing a stiffening gradient were used to improve corneal regularization.⁹ The clinical results of CXL using higher fluence are promising, and to date, no endothelial decompensation has been reported for these protocols.

These studies have shown that higher CXL fluence is clinically safe, but the effect of higher fluence on corneal mechanical stiffness has been only poorly studied so far. To our knowledge, there are only two studies that measure CXL stiffening of CXL for various fluence levels up to 10.8 J/cm².^{12,13}

To determine the biomechanical response of CXL, uniaxial stress-strain measurements on porcine corneas are used by many groups in the past,^{14,15} therefore, allowing good comparability of results.

CONTACT Isaak Fischinger 😒 isaak.fischinger@hotmail.com 🗈 Medical Faculty, Johannes Kepler University Linz, Altenberger Strasse 69, Linz, 4040, Austria © 2023 The Author(s). Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

The aim of this study is to analyze the biomechanical stiffening after accelerated epi-off CXL, with a fluence of 5.4 up to 20 J/cm^2 in porcine corneas using uniaxial extensiometry.

Methods

Corneal preparation and CXL procedure

Ninety freshly enucleated porcine eyes were collected from the local slaughterhouse. All eyes were examined for integrity and pachymetry (SP-100; Tomey Corporation) was performed to exclude damaged eyes. The eyes were randomly divided into five groups. All eyes were de-epithelialized with a hockey knife. A solution containing 0.1% riboflavin (riboflavin-5monophosphate in 0.9% NaCl) dissolved in 16% dextran was applied on the corneal surface every 2 min for 30 min. Then, the eyes of group 1-4 were continuously irradiated with UVA-light ($\lambda = 365$ nm) (Mosaic, Avedro, Inc., Waltham, MA, USA) at an irradiance of 18 mW/cm², applying a total fluence of 20, 15, 10.8, and 5.4 J/cm² in group 1, 2, 3, and 4, respectively. Group five was not irradiated and served as a control group. After the treatment, corneal discs were excised and stored in a 16% dextran bath for 2h to ensure uniform hydration conditions. Pachymetry was then measured again and 5 mm wide strips were excised along the vertical axis.

Stress-strain measurement

Corneal biomechanics was quantified using uniaxial tensile tests. The samples were mounted with two clamps on the testing device equipped with a 1 N load cell (UStrech Cellscale, Ontario, Canada). A preload of 5×10^3 Pa prestress was applied to each specimen for 2 min to ensure uniform tension of the tissue at the beginning of each measurement. The strain was applied at a rate of 0.035 mm/s, similar to previous biomechanical measurements.¹⁶⁻¹⁸ The force on the sample was measured during the extension until a strain of 10%. Both force and displacement were recorded every 10 ms during the measurement. The distance between the two clamps was 6 mm at the beginning of the measurement, similar to previous experiments.^{17,18} The samples were immersed in a bath containing 16% dextran solution to maintain hydration during the measurement. The stress and strain were calculated at each measurement point, while the Young's modulus was calculated at 6, 8, and 10% strain.

Numerical evaluation

Data was exported to Excel MS Software (Microsoft Corporation) where descriptive analysis was performed. The

Table 1. Stress (kPa) at the strain of 6, 8, and 10%.

cross-section area of the samples required to determine the stress was calculated by multiplying the sample thickness measured by pachymetry by its width of 5 mm. The Young's modulus (*E*) describing the relationship between stress (σ) and strain (ε) was calculated as:

$$E = \sigma/\varepsilon$$

Then, the mean of the Young's modulus at 10% strain were compared using the unifactorial ANOVA using SPSS (version 26.0 IBM Corporation). *Post-hoc* Holm-Bonferroni correction for multiple comparisons was performed. Pachymetry of the five groups was analyzed using Kruskal-Wallis. A *p*-value of <.05 was considered statistically significant.

Results

Eighty-six (96%) of 90 eyes could be considered for the statistical evaluation. Four eyes of had to be excluded due to a failure of the software of the mechanical testing device during the measurement. Accordingly, 16 corneas from group 3, 17 from groups 2 and 4, and 18 from groups 1 and 5 were included in the data analysis.

The average stress of groups 1–5 at 6, 8, and 10% strain is shown in Table 1. The stress at 10% strain was increased by 76% in group 1, 56% in group 2, 52% in group 3, and by 31% in group 4 compared to control group 5. The linear regression analysis at a strain of 10% showed a high correlation between fluence and stress ($R^2 = 0.955$; p = 0.004).

The Young's modulus at 10% strain was calculated for the five groups and is shown in Figure 1. The Young's modulus was compared using *post-hoc* Bonferroni adjusted Analysis of Variance (ANOVA), resulting p values are shown in Table 2. Groups 1–4 showed a statistically significant stiffening compared with group 5 (p = <0.001; p = <0.001; p = <0.001; p = 0.021). In addition, group 1 showed significantly more stiffening than group 4 (p = <0.001), no other significant differences were found.

The average pachymetry before stress-strain measurement was 731 ± 57 , 725 ± 59 , 713 ± 52 , 711 ± 62 , and $741 \pm 62 \,\mu\text{m}$ in groups 1–5, respectively. There was no statistically significant difference between the five groups (p = 0.538).

Discussion

The most important finding of this study is the significant dose-response relationship of the stiffening effect after epioff CXL with increasing applied fluence from 5.4 to 20 J/cm^2 .

The first studies reporting the biomechanical response after CXL using the Dresden protocol with a total fluence of 5.4 J/cm^2 showed an increased stress by a factor of 1.8 at a

	Group 1 (20 J/cm ²)	Group 2 (15 J/cm ²)	Group 3 (10.8 J/cm ²)	Group 4 (5.4 J/cm ²)	Group 5 (control)
6%	116.9±33.6	93.9 ± 20.6	97.8 ± 23.9	81.0 ± 23.5	59.9 ± 34.0
8%	193.9 ± 44.4	166.8 ± 25.9	166.7 ± 36.0	141.0 ± 32.2	103.6 ± 46.8
10%	284.5 ± 51.5	252.8 ± 32.1	246.0 ± 46.8	212.1 ± 39.1	162.0 ± 58.9



Young's Modulus at Strain 10%

Figure 1. Young's modulus at a strain of 10%. Group 1 = corneal crosslinking (CXL) with a fluence of 20 J/cm², group 2 = CXL with a fluence of 15 J/cm²; group 3 = CXL with a fluence of 10.8 J/cm²; group 4 = CXL with a fluence of 5.4 J/cm²; group 5 = control group untreated corneas.

Table 2. p-Values resulting from post-hoc Bonferroni adjusted analysis of variance (ANOVA) between stress (kPa) at the strain of 10% after CXL with different fluences.

	Group 1 (20 J/cm ²)	Group 2 (15 J/cm ²)	Group 3 (10.8 J/cm ²)	Group 4 (5.4 J/cm ²)	Group 5 (control)
Group 1 (20 J/cm ²)	Х	0.475	0.210	<0.001	<0.001
Group 2 (15 J/cm ²)	0.475	х	1.000	0.129	< 0.001
Group 3 (10.8 J/cm ²)	0.210	1.000	х	0.304	< 0.001
Group 4 (5.4 J/cm ²)	<0.001	0.129	0.304	х	0.021
Group 5 (Control)	<0.001	<0.001	<0.001	0.021	х

strain of 10%.³ Previous studies by our group using similar experimental settings and the same mechanical tester showed a similar increased stress of up to 80% at a strain of 10% using an accelerated protocol (9 mW/cm², 10 min).^{16,17} The current results of the accelerated treatment show a much smaller increase of stress of only 31% compared to the control group. However, in this study, the standard fluence of 5.4 J/cm² was induced by an irradiance of $18 \,\mathrm{mW/cm^2}$ for $5 \,\mathrm{min}$, whereas the irradiance of the Dresden protocol is only 3 mW/cm². Similar findings of reduced stiffening in accelerated protocols were reported by our group¹⁷ and other authors.¹⁵ Hammer et al. found an even smaller and non-significant (p = 0.065) stiffening of about 12% in their experiments using 18 mW/cm² with a standard fluence of 5.4 J/cm² too.¹⁵ This effect can be explained by the hypothesis, that high irradiances are associated with a rapid depletion of oxygen in the anterior layers of the cornea, which limits oxygen availability in the

posterior layers, and consequently leads to a more superficial crosslinking effect.^{19,20} This is consistent with the more superficial demarcation lines after CXL with high irradiance.²¹

Although higher fluences than 5.4 up to 15 J/cm² have been used clinically for several years,^{10,22} to our knowledge, there are only two recently published studies^{12,13} reporting biomechanical measurements using fluences up to 10.8 J/cm² for CXL. Biomechanical investigations for higher fluences do not exist to date but are relevant, as they are used clinically.

We found that doubling the fluence increased the stiffening effect from 31 to 53%, which is comparable to the 22 and 49% stiffening reported by Abrishamchi et al. using a similar setting to measure the corneal biomechanics. However, the CXL protocol used differed from our study as they used an irradiance of 9 mW/cm^2 for 10 min to reach the fluence of 5.4 J/cm^2 and an irradiation of 18 mW/cm^2 for 9 min and 15 s, corresponding to a slightly smaller total fluence of 10 J/cm² than in our study. The authors also did not specify which riboflavin solution was used in their experiments. The use of a HPMC based solution could explain the somewhat weaker stiffening effect, as dextranbased solutions which were used in our study are known to produce stronger stiffening.¹⁶ Boschetti et al. reported more stiffening for higher fluences using inflation tests to measure the biomechanics, what makes a comparison of the results difficult.¹³

A consequence of the data presented here is that stiffening can be increased by additional treatment time and fluences above 10.8 J/cm², a significant difference was found between 5.4 and 20 J/cm². A longer treatment allows more oxygen to diffuse in the cornea and therefore enables more CXL. Based on the present results, we cannot determine which factor has the greatest influence, the additional treatment time or energy. Clinical data of customized CXL report deeper demarcation lines in areas with higher applied fluence and treatment duration^{9,23} suggesting that some rediffusion of oxygen in deeper layers, after an initial saturation of the anterior stroma, is responsible for this deepening of the demarcation line. This theory is supported by the slight increase in oxygen after 10 min during continuous UV-irradiation in a depth of 200 microns, found by Kamaev et al. in 2012.²⁴ However, only future oxygen measurements in combination with biomechanical tests can confirm this assumption and determine the upper threshold for induced crosslinks. An increased anaerobic CXL²⁴ might also serve as a potential explanation for the deeper demarcation line and the increased stiffening.

The classic Dresden protocol with a total fluence of 5.4 J/cm^2 applies an uniform energy to the treated area of the cornea, stopping the progression of ectatic corneas with good outcomes and patients safety.^{25,26} Already a few years after the introduction of CXL, the procedure has been successfully accelerated by increasing the irradiance and varying the riboflavin solutions,^{27–30} but the stiffening effect, as well as the regularization of the cornea, were not noticeably improved for a long time.^{31,32} But since the introduction of customized CXL, favorable results have been published regarding the regularization and flattening of the cornea are treated with a higher fluence to achieve a stronger flattening in these areas. The assumption, that higher fluences cause more stiffening is confirmed by our results.

Currently, the individual amount of the induced crosslinking that is required to stop the progression of an ectatic cornea is not known preoperatively, since the individual biomechanical state of the patient's cornea cannot be assessed accurately clinically. Therefore, it might be inappropriate to apply more energy in every CXL treatment, as the success rates of current protocols are very high. In addition, clinicians have to be aware that in rare cases, there is ongoing flattening even years after CXL,³³ so not all patients will benefit from a very strong CXL.

Though the former fluence threshold of 5.4 J/cm^2 is no longer valid and higher radiant exposures can be used,⁶ fluences higher than 10.8 J/cm^2 should be used reluctant, since

little data about safety exists.²² In thinner corneas higher fluences are not recommended at all, but a reduction should be considered.¹¹ Epi-on CXL, where the epithelium absorbs additional UV light and probably causes additional safety for the endothelium might be a field of application for higher fluences.

One limitation of this study is the use of porcine corneas. Because they are much more readily available and have similar dimensions to human corneas, porcine specimens are commonly used in experimental CXL studies, although previous studies have found a smaller CXL effect in porcine than in human corneas.³ Nevertheless, it is believed that the results are transferable to human eyes. In our experiments, we used riboflavin dissolved in 16% dextran, as opposed to the 20% dextran used by clinicians and other ex vivo studies. The 16% solution was chosen to minimize corneal thinning which is reduced by using a lower concentration of dextran. The uniaxial stress-strain measurements could be identified as an additional disadvantage. Cutting the strips compromises the integrity of the cornea. In addition, the strips are pulled flat in the extensometer, so that the superficial fibers are stressed less than the deeper layers. However, many other studies have used a similar experimental design, so it is a standard protocol that allows the comparison of results with the existing literature.

In conclusion, the stiffening effect of the accelerated protocol based on $18 \,\mathrm{mW/cm^2}$ with a standard fluence of $5.4 \,\mathrm{J/cm^2}$ is lower than earlier reported for the standard CXL that relies on $3 \,\mathrm{mW/cm^2}$. The biomechanical response after accelerated CXL can be increased by extending the treatment time causing a higher fluence. According to our results, there seems to be no threshold up to $20 \,\mathrm{J/cm^2}$. A higher fluence could compensate the weaker effect of accelerated or epi-on CXL procedures.

Disclosure statement

Dr. Seiler is a scientific advisor for Avedro/Glaukos and receives material support for research projects. The other authors report no conflicts of interest.

ORCID

Jascha Wendelstein (D) http://orcid.org/0000-0003-4145-2559

Data availability statement

The data that support the findings of this study are available from the corresponding author, [IF], upon reasonable request.

References

- Seiler T, Spoerl E, Huhle M, Kamouna A. Conservative therapy of keratoconus by enhancement of collagen cross-links. Invest Ophthalmol Vis Sci. 1996;37:1017.
- Santhiago MR, Randleman JB. The biology of corneal cross-linking derived from ultraviolet light and riboflavin. Exp Eye Res. 2021;202:108355. doi:10.1016/j.exer.2020.108355.
- 3. Wollensak G, Spoerl E, Seiler T. Stress-strain measurements of human and porcine corneas after riboflavin-ultraviolet-A-

induced cross-linking. J Cataract Refract Surg. 2003;29(9):1780-1785. doi:10.1016/s0886-3350(03)00407-3.

- 4. Wollensak G, Spörl E, Reber F, Pillunat L, Funk R. Corneal endothelial cytotoxicity of riboflavin/UVA treatment *in vitro*. Ophthalmic Res. 2003;35(6):324–328. doi:10.1159/000074071.
- Wollensak G, Spoerl E, Wilsch M, Seiler T. Endothelial cell damage after riboflavin-ultraviolet-A treatment in the rabbit. J Cataract Refract Surg. 2003;29(9):1786–1790. doi:10.1016/s0886-3350(03)00343-2.
- Seiler TG, Batista A, Frueh BE, Koenig K. Riboflavin concentrations at the endothelium during corneal cross-linking in humans. Invest Ophthalmol Vis Sci. 2019;60(6):2140–2145. doi:10.1167/ iovs.19-26686.
- Lang PZ, Hafezi NL, Khandelwal SS, Torres-Netto EA, Hafezi F, Randleman JB. Comparative functional outcomes after corneal crosslinking using standard, accelerated, and accelerated with higher total fluence protocols. Cornea. 2019;38(4):433–441. doi: 10.1097/ICO.000000000001878.
- Shetty R, Pahuja N, Roshan T, Deshmukh R, Francis M, Ghosh A, Sinha Roy A. Customized corneal cross-linking using different UVA beam profiles. J Refract Surg. 2017;33(10):676–682. doi:10. 3928/1081597X-20170621-07.
- Seiler TG, Fischinger I, Koller T, Zapp D, Frueh BE, Seiler T. Customized corneal cross-linking: one-year results. Am J Ophthalmol. 2016;166:14–21. doi:10.1016/j.ajo.2016.02.029.
- Mazzotta C, Sgheri A, Bagaglia SA, Rechichi M, Di Maggio A. Customized corneal crosslinking for treatment of progressive keratoconus: clinical and OCT outcomes using a transepithelial approach with supplemental oxygen. J Cataract Refract Surg. 2020;46(12):1582–1587. doi:10.1097/j.jcrs.000000000000347.
- Hafezi F, Kling S, Gilardoni F, Hafezi N, Hillen M, Abrishamchi R, Gomes JAP, Mazzotta C, Randleman JB, Torres-Netto EA. Individualized corneal cross-linking with riboflavin and UV-A in ultrathin corneas: the Sub400 protocol. Am J Ophthalmol. 2021; 224:133–142. doi:10.1016/j.ajo.2020.12.011.
- Abrishamchi R, Abdshahzadeh H, Hillen M, Hafezi N, Torres-Netto EA, Aslanides IM, Chen S, Randleman JB, Hafezi F. Highfluence accelerated epithelium-off corneal cross-linking protocol provides dresden protocol-like corneal strengthening. Transl Vis Sci Technol. 2021;10(5):10. doi:10.1167/tvst.10.5.10.
- Boschetti F, Conti D, Soriano EM, Mazzotta C, Pandolfi A. Experimental *in-vitro* investigation on epi-off-crosslinking on porcine corneas. PLOS One. 2021;16(4):e0249949. doi:10.1371/ journal.pone.0249949.
- Spoerl E, Huhle M, Seiler T. Induction of cross-links in corneal tissue. Exp Eye Res. 1998;66(1):97–103. doi:10.1006/exer.1997. 0410.
- Hammer A, Richoz O, Arba Mosquera S, Tabibian D, Hoogewoud F, Hafezi F. Corneal biomechanical properties at different corneal cross-linking (CXL) irradiances. Invest Ophthalmol Vis Sci. 2014;55(5):2881–2884. doi:10.1167/iovs.13-13748.
- Fischinger I, Seiler TG, Wendelstein J, Tetz K, Fuchs B, Bolz M. Biomechanical response after corneal crosslinking with riboflavin dissolved in dextran solution versus hydroxy propyl methyl cellulose. J Refract Surg. 2021;37(9):631–635. doi:10.3928/ 1081597X-20210610-04.
- Seiler TG, Fischinger I, Senfft T, Schmidinger G, Seiler T. Intrastromal application of riboflavin for corneal crosslinking. Invest Ophthalmol Vis Sci. 2014;55(7):4261–4265. doi:10.1167/ iovs.14-14021.
- Lammer J, Laggner M, Pircher N, Fischinger I, Hofmann C, Schmidinger G. Endothelial safety and efficacy of *ex-vivo* collagen cross linking (CXL) of human corneal transplants. Am J Ophthalmol. 2020;214:127–133. doi:10.1016/j.ajo.2020.02.024.

- Seiler TG, Komninou MA, Nambiar MH, Schuerch K, Frueh BE, Buchler P. Oxygen kinetics during corneal cross-linking with and without supplementary oxygen. Am J Ophthalmol. 2021;223: 368–376. doi:10.1016/j.ajo.2020.11.001.
- Richoz O, Hammer A, Tabibian D, Gatzioufas Z, Hafezi F. The biomechanical effect of corneal collagen cross-linking (CXL) with riboflavin and UV-A is oxygen dependent. Transl Vis Sci Technol. 2013;2(7):6. doi:10.1167/tvst.2.7.6.
- 21. Brittingham S, Tappeiner C, Frueh BE. Corneal cross-linking in keratoconus using the standard and rapid treatment protocol: differences in demarcation line and 12-month outcomes. Invest Ophthalmol Vis Sci. 2014;55(12):8371–8376. doi:10.1167/iovs.14-15444.
- 22. Sachdev GS, Ramamurthy S, B S, Dandapani R. Comparative analysis of safety and efficacy of topography-guided customized cross-linking and standard cross-linking in the treatment of progressive keratoconus. Cornea. 2021;40(2):188–193. doi:10.1097/ ICO.000000000002492.
- Cassagne M, Pierne K, Galiacy SD, Asfaux-Marfaing MP, Fournie P, Malecaze F. Customized topography-guided corneal collagen cross-linking for keratoconus. J Refract Surg. 2017; 33(5):290–297. doi:10.3928/1081597X-20170201-02.
- Kamaev P, Friedman MD, Sherr E, Muller D. Photochemical kinetics of corneal cross-linking with riboflavin. Invest Ophthalmol Vis Sci. 2012;53(4):2360–2367. doi:10.1167/iovs.11-9385.
- Koller T, Mrochen M, Seiler T. Complication and failure rates after corneal crosslinking. J Cataract Refract Surg. 2009;35(8): 1358–1362. doi:10.1016/j.jcrs.2009.03.035.
- Wollensak G, Spoerl E, Seiler T. Riboflavin/ultraviolet-A-induced collagen crosslinking for the treatment of keratoconus. Am J Ophthalmol. 2003;135(5):620–627. doi:10.1016/s0002-9394(02)02220-1.
- Wen D, Li Q, Song B, Tu R, Wang Q, O'Brart DPS, McAlinden C, Huang J. Comparison of standard versus accelerated corneal collagen cross-linking for keratoconus: a meta-analysis. Invest Ophthalmol Vis Sci. 2018;59(10):3920–3931. doi:10.1167/iovs.18-24656.
- Males JJ, Viswanathan D. Comparative study of long-term outcomes of accelerated and conventional collagen crosslinking for progressive keratoconus. Eye. 2018;32(1):32–38. doi:10.1038/eye. 2017.296.
- Thorsrud A, Hagem AM, Sandvik GF, Drolsum L. Superior outcome of corneal collagen cross-linking using riboflavin with methylcellulose than riboflavin with dextran as the main supplement. Acta Ophthalmol. 2019;97(4):415–421. doi:10.1111/aos. 13928.
- Lenk J, Herber R, Oswald C, Spoerl E, Pillunat LE, Raiskup F. Risk factors for progression of keratoconus and failure rate after corneal cross-linking. J Refract Surg. 2021;37(12):816–823. doi: 10.3928/1081597X-20210830-01.
- Rapuano PB, Mathews PM, Florakis GJ, Trokel SL, Suh LH. Corneal collagen crosslinking in patients treated with dextran versus isotonic hydroxypropyl methylcellulose (HPMC) riboflavin solution: a retrospective analysis. Eye Vis. 2018;5:23. doi:10. 1186/s40662-018-0116-z.
- 32. Shajari M, Kolb CM, Agha B, Steinwender G, Muller M, Herrmann E, Schmack I, Mayer WJ, Kohnen T. Comparison of standard and accelerated corneal cross-linking for the treatment of keratoconus: a meta-analysis. Acta Ophthalmol. 2019;97(1): e22–e35. doi:10.1111/aos.13814.
- Noor IH, Seiler TG, Noor K, Seiler T. Continued long-term flattening after corneal cross-linking for keratoconus. J Refract Surg. 2018;34(8):567–570. doi:10.3928/1081597X-20180607-01.