

OPEN

# The burden of cardiovascular risk in individuals with spinal cord injury (SCI) and its association with rehabilitation outcomes – Results from the Swiss SCI Cohort

Peter Francis Raguindin, MD, PhD 1,2,3,4; Gabi Mueller, PhD 1,5; Jivko Stoyanov, PhD 1, 2; Inge Eriks-Hoogland, MD, PhD 5; Xavier Jordan, MD 6; Stevan Stojic, MSc 1; Margret Hund-Georgiadis, MD, PhD 7; Taulant Muka, MD, PhD 2; Gerold Stucki, MD 1,8; Marija Glisic, MD, PhD 1,2; on behalf of the SwiSCI Study Group

1. Swiss Paraplegic Research, Nottwil, Switzerland
2. Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland
3. Graduate School for Health Sciences, University of Bern, Bern, Switzerland
4. Faculty of Health Sciences and Medicine, University of Lucerne, Lucerne, Switzerland
5. Swiss Paraplegic Centre, Nottwil, Switzerland
6. Clinique romande de readaptation, Sion, Switzerland
7. REHAB Basel, Basel, Switzerland
8. Center for Rehabilitation in Global Health Systems, Faculty of Health Sciences and Medicine, University of Lucerne, Lucerne, Switzerland

## Corresponding Author

Peter Francis Raguindin, MD, PhD

Faculty of Health Sciences and Medicine

University of Lucerne

Frohburgstrasse 3, 6002 Lucerne, Switzerland

Email: peter.raguindin@unilu.ch

Phone: +41 41 229 59 69

### **Funding**

Peter Francis Raguindin has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 801076, through the SSPH+ Global PhD Fellowship Programme in Public Health Sciences (GlobalP3HS) of the Swiss School of Public Health.

### **Author Disclosures**

Authors declare no conflicts of interest. The partial study results were submitted at the 61<sup>st</sup> Annual Scientific Meeting of the International Spinal Cord Society (ISCOS), 15-18 September 2022, Vancouver, Canada.

### **Acknowledgment**

We thank the SwiSCI Steering Committee with its members Fabienne Reynard (Clinique Romande de Réadaptation, Sion); Michael Baumberger, Luca Jelmoni (Swiss Paraplegic Center, Nottwil); Armin Curt, Martin Schubert (Balgrist University Hospital, Zürich); Laurent Prince (Swiss Paraplegic Association, Nottwil); Daniel Joggi (Representative of persons with SCI); Mirjana Bosnjakovic (Parahelp, Nottwil); Mirjam Brach (Swiss Paraplegic Research, Nottwil); Carla Sabariego (SwiSCI Coordination Group at Swiss Paraplegic Research, Nottwil). Also, we would like to extend our gratitude to Christine Thyrian, Wolfgang Segerer, and Beat Gurtner of the SwiSCI Study Center for their assistance in data management and cleaning.

### **Author contributions**

PFR and MG conceptualized the study. PFR analyzed the data. GM, JS, IEH, SS, MHG, JX, TM, GS interpreted the data and drafted the manuscript. MG and TM supervised the study conduct. All authors approved the final version of the manuscript.

### **Data availability statements**

Dataset utilized in the analysis are available from the SwiSCI Study Center upon reasonable request (contact@swisci.ch)

Peter Francis Raguindin is in training.

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## **Abstract**

### Objectives

To determine the cardiovascular risk burden rehabilitation discharge, and explore the association between recovery during rehabilitation and CVD-risk profile

### Methods

We included adults without CVD history admitted for rehabilitation. We evaluated rehabilitation outcomes on admission and discharge. CVD-risk was assessed by Framingham risk score (FRS), high-density lipoprotein (HDL), and fasting glucose level.

### Results

We analyzed data from 706 participants (69.55% men) with median age of 53.5 years. The median time-since-injury was 14 days and the admission length was 5.2 months. Majority had paraplegia (53.26%), and motor incomplete injury (53.68%). One-third of the cohort had high cardiovascular risk profile pre-discharge. At discharge, poorer anthropometric measures were associated with higher FRS and lower HDL levels. Individuals with higher forced vital capacity (>2.72 L) and peak expiratory flow (>3.4L/min) had 0.16 mmol/L and 0.14 mmol/L higher HDL compared to those with lower respiratory function, respectively. Individuals with higher mobility score (>12.5) and functional independence score (>74) had 0.21 mmol/L and 0.18 mmol/L higher HDL compared to those with lower scores.

### Conclusion

There is high cardiometabolic syndrome burden and CVD-risk upon rehabilitation discharge. Higher respiratory function, mobility, and overall independence were associated with better

CVD profile, although with study design limitations and short follow-up. Future studies should explore whether rehabilitation outcomes could be used to prioritize screening.

**MeSH/Keywords**

cardiovascular risk, paraplegia, tetraplegia, cardiovascular disease, lung volumes, spirometry, obesity, functioning

ACCEPTED

### What is Known

Individuals with spinal cord injury are at high-risk for cardiovascular mortality and morbidity. However, cardiovascular screening is rarely done before discharge.

### What is New

One-third of the cohort had a high cardiovascular risk profile. Lower lung function, mobility, and overall functioning have higher cardiovascular disease risk and should be targeted for routine screening.

ACCEPTED

## Introduction

Spinal cord injury (SCI) results in dramatic changes in an individual's functioning that changes across the injury trajectory <sup>1</sup>. The injury causes skeletal muscle paralysis and spasticity that limit their independence over activities of daily living and their physical activity. The injury also results in lower cardiorespiratory function secondary to impaired neurologic control (autonomics) or respiratory muscle paralysis <sup>2</sup>. Furthermore, changes in metabolic milieu were reported within weeks post-injury as the body shift from a catabolic state in response to acute injury to an anabolic state in subacute injury which promotes fat accumulation <sup>3</sup>. Therefore, subacute injury is a critical phase characterized by dynamic physiologic adaptation. A recent study on the subacute phase observed no deterioration of cardiovascular health during inpatient rehabilitation <sup>4</sup>. Thus, the inpatient rehabilitation program could have a beneficial effect on cardiometabolic health.

Rehabilitation programs are tailored to shorten hospital stays, decrease re-admissions and reduce the risks of secondary complications<sup>5-7</sup> by improving physical capacity (e.g., respiratory function, and muscle strength) and functional independence <sup>8</sup>, which have been crucial for daily physical activities and social integration <sup>9,10</sup>. Among other activities, these programs focus on muscle strengthening, preventing spasticity, and improving cardiorespiratory fitness through exercises <sup>8</sup>. Few studies have associated rehabilitation outcomes with cardiovascular health in this group. A study has previously associated poorer lung volumes with higher odds of type 2 diabetes and hypertension <sup>11</sup>. Another study observed an association between lower extremity spasticity and fasting glucose and adiposity <sup>12</sup>. Understanding these changes during inpatient rehabilitation could be important in cardiovascular disease (CVD) prevention since deterioration in lipid or glucose profile could lead to long-term complications and morbidity <sup>13</sup>. In a previous study, at discharge from

initial rehabilitation almost one-third of patients can be classified as high risk for CVD<sup>4</sup>. Despite being considered for a high CVD risk group because of risk factors clustering as early as within weeks post-injury<sup>14,15</sup>, CVD screening has not been routinely included prior to rehabilitation discharge<sup>16,17</sup>. Previous study has shown 20% of individuals with SCI are being screened for lipid profile and fasting glucose upon discharge<sup>18</sup> even though 30-60% of them are at high cardiovascular risk<sup>4</sup>. The main determinants of poorer CVD risk profile in subacute injury phase remain largely unknown. We hypothesize that better functional recovery during initial rehabilitation is associated with more favorable CVD risk profile at discharge and may be used to triage CVD screening in inpatient setting. Thus, for this study, we aimed to: (i) determine the CVD risk and cardiometabolic syndrome burden during initial inpatient rehabilitation, (ii) explore longitudinal changes in rehabilitation outcomes, and (ii) associate these rehabilitation outcomes with cardiovascular risk profile prior to discharge from initial rehabilitation risk.

## **Methods**

This is a cohort study on rehabilitation outcomes over time (longitudinal analysis) and the association of these measures to cardiovascular risk profile at discharge during inpatient rehabilitation. Specific details on study procedures and data analysis can be found in the **Online Appendix (Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>).**

### *Study setting and population*

The Swiss Spinal Cord Injury cohort (SwiSCI) is a multicenter study across all four rehabilitation centers in Switzerland, namely, Swiss Paraplegic Center (Nottwil), Clinique Romande de Readaptation (Sion), Balgrist Spine Center (Zurich), and Basel Rehabilitation Clinic (Basel). The additional details of the study cohort can be found in previous



publications<sup>19,20</sup>. In summary, this study enrolled individuals with a new diagnosis of SCI from each participating center. We collected demographic and clinical profiles, with detailed injury characteristics and additional information related to overall wellbeing and functioning, using the prescribed data collection form of the International Spinal Cord Society<sup>21,22</sup>. Data were collected upon admission and before discharge.

We enrolled adults ( $\geq 18$  years) admitted for inpatient rehabilitation in one of the participating centers from May 2013 (cohort inception date) to September 2021. We excluded individuals with acute inflammatory and infectious causes of the injury and terminal cancers or those in palliative care. We also excluded those with a history of cardiovascular disease to estimate the participant's risk for the first cardiovascular event. Participants were followed from admission until discharge to rehabilitation centers.

#### *Definition of rehabilitation outcomes*

Rehabilitation measure was divided into five components: anthropometric measures, respiratory function, spasticity, muscle strength (including frailty), and functional independence. All body anthropometrics were measured in the supine position. Waist circumference (WC) was measured after bowel care, at the end of normal expiration, approximately between the lower margin of the last palpable rib and the top of the iliac crest. It was measured using a pliable tape measure with a precision of 0.5 cm. Weight was measured using an electric wheelchair scale. The total weight of the subject with the wheelchair was subtracted from the wheelchair's weight to determine the subject's weight expressed in kilograms (kg). Both waist circumference and weight were measured once per assessment. Body mass index (BMI) was computed using the standard formula [weight in kilograms/(height in meters)<sup>2</sup>].

The respiratory function was assessed through functional vital capacity (FVC), forced expiratory volume in 1 second (FEV1), and peak expiratory flow (PEF). Lung volumes (FVC and FEV1) are measured in liters (L), and the flow rate (PEF) was measured in L/min. These measure the extent of respiratory muscle innervation and function of the participants due to the injury. The best of the 3 efforts the participant made for respiratory function test was recorded and used in analysis. Spasticity was assessed on the biceps and gastrocnemius of the upper and lower extremities from both sides. A rehabilitation specialist measured spasticity through modified Ashworth scale (0 as no spasticity to 5 maximum score). Handgrip strength was measured on both sides using a hand dynamometer using REACT protocol.

Functional independence was measured using the Spinal Cord Independence Measure (SCIM III)<sup>23</sup>. The score has a minimum of 0 and a maximum 100 score. It is a composite of three main domains: self-care, respiration and sphincter management, and mobility. The mobility subcomponent has a maximum score of 40 and has six components, namely, indoor mobility (0-8), moderate distance 10-100 meters mobility (0-8), outdoor mobility >100 meters (0-8), mobility in stairs (0-3), wheelchair to car transfer (0-2), and wheelchair to ground (0-1) We used the functional independence (total score) and mobility subcomponent score for the analysis.

#### *Definition of cardiovascular risk and cardiometabolic syndrome*

Cardiovascular risk profile was assessed using the cardiovascular risk score, dyslipidemia, impaired fasting glucose, hypertension, overweight (high-risk obesity), and cardiometabolic syndrome. The Framingham risk score (FRS) predicted the risk of having the first cardiovascular event within ten years, and was based on a cohort of 5,209 adults in Framingham, Massachusetts, US.<sup>24</sup> For this study, we computed the FRS of each patient at

discharge for participants using the following variables: (a) age, (b) sex, (c) systolic blood pressure, (d) total cholesterol (units to be converted to mg/dL), (e) high-density lipoprotein (units to be converted to mg/dL), (f) diabetes, and (g) current smoking. The risk of the first cardiovascular event within 10 years is considered low when it is <10% and medium to high risk with those having  $\geq 10\%$ .<sup>24</sup> FRS was chosen in the main analysis to be comparable with the previous attempt in validating the score with hard outcomes.<sup>25</sup> We also iterated cardiovascular risk score using SCORE2 (adjusted for Swiss population-low risk region)<sup>26</sup> and WHO CVD risk score (adjusted for Swiss population and the 2017 Global Burden of Disease).<sup>27</sup> The details of SCORE2 and WHO CVD risk score can be found in separate publications and have not yet been validated in the SCI population.

Venous blood samples were obtained from each participant after an overnight fast. The blood was sent to respective hospital laboratories, which measured high-density lipoprotein and glucose. Dyslipidemia was defined as those with HDL at < 1.0 mmol/L in males or < 1.3 mmol/L in females, or with total cholesterol  $\geq 5.5$  mmol/L.<sup>28</sup> Impaired fasting glucose (high risk for diabetes) was defined as those with  $\geq 7.0$  mmol/L from an international consensus.<sup>29</sup>

Hypertension was defined as having resting systolic BP  $\geq 130$  mmHg or diastolic BP  $\geq 90$  mmHg based from a clinical guideline.<sup>30</sup> Overweight (high risk for obesity) was defined as those having BMI  $> 22$  kg/m<sup>2</sup> or waist circumference  $\geq 86.5$  cm according to SCI-specific cut-off<sup>31,32</sup>, and BMI  $> 25$  kg/m<sup>2</sup> or waist circumference  $> 102$  cm (males)/  $> 88$  cm (females) according to the cut-off for general population.<sup>33</sup> Finally, cardiometabolic syndrome is classified by having any three of the following (based on 2005 AHA/NHLBI guideline): (a)

overweight (SCI-specific cut-off, (b) elevated triglycerides, (c) low HDL, (d) impaired fasting glucose, and (e) hypertension.

#### *Definition of covariates*

The level of injury was classified as tetraplegia (at level C2-C7) and paraplegia (level T1-S5), and the completeness of injury into complete motor injury (AIS A and B) and incomplete (AIS C and D) based on the International Standards for Neurological Classification of Spinal Cord Injury (ISNCSCI)<sup>34</sup>. SCI validated questionnaires were used to obtain some covariates. The dietary pattern is obtained using the Swiss Health Survey questionnaire<sup>35</sup>, and subsequently, a nutritional score was computed using the recommendations of the German Society for Nutrition<sup>36</sup>. Physical activity is measured through Physical Activity Scale for Individuals with Physical Disabilities<sup>37</sup>.

#### *Data analysis*

We summarized the rehabilitation measures using median and interquartile range (IQR) as prescribed by ISCOS Standards of Data Analysis and Reporting<sup>22</sup>. We determined the changes in rehabilitation outcomes from admission to discharge using the Wilcoxon signed-rank test for paired samples. We summarized the rehabilitation outcomes by injury level and injury completeness. We iterated longitudinal analysis to determine if injury characteristics affect the rehabilitation outcome changes. We also performed linear mixed model using random intercept and individuals as clusters, adjusted for age, sex, and injury level.

We selected cardiovascular risk score (FRS), HDL, and fasting glucose as main dependent variables based on previous literature on cardiometabolic risk in SCI. We used multivariable linear regression adjusted for age, sex, injury level, and body mass index (a surrogate for

obesity). We drafted directed acyclic graphs (DAGs) to determine the minimum covariate adjustments in assessing the effect of independent variables ([www.daggity.net](http://www.daggity.net)) (**Figure S1, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). We used the rehabilitation outcomes both as continuous and dichotomous variables. The use of independent variables (rehabilitation outcomes) as a continuous variable was done to determine the strength of association to the dependent variables (cardiovascular risk). Furthermore, we dichotomized the independent variables (rehabilitation outcomes) to determine the unit change in cardiovascular risk (dependent variables) for those with high versus lower rehabilitation measure (independent variables). We used SCI-adjusted cutoffs for waist circumference and body mass index. For other variables of rehabilitation, we divided the group into those above or below the medians. Furthermore, we log-transformed FRS prior to fitting in the regression model. No transformations were done for HDL and glucose prior to regression. We tested the normality of the residuals after regression fitting.

#### *Sensitivity analyses*

Detailed results of the sensitivity analyses can be found in the **Online Appendix (Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>)**. To determine selection bias, we compared the baseline characteristics of those with complete data with those with missing dependent variables using the chi-squared test or Wilcoxon test. We performed a post hoc test (Dunn's test with Bonferroni adjustment) to confirm the differences between uneven group comparisons. We also fitted our main models with interaction term of sex and rehabilitation outcome (independent variable) to test for effect modification (sex). For statistically significant interactions, we iterated a subgroup estimate for each sex. We also restricted the analyses to those with paraplegia, motor complete injury, and traumatic SCI to create a more homogenous population based on injury characteristics. We performed

subgroup analyses according to age (adults  $\leq 50$  years and elderly  $>50$  years). We performed Bonferroni corrections to adjust for multiple comparisons.

We also explored the association of longitudinal changes in rehabilitation outcomes and cardiovascular risk profile through a repeated measures approach. We used a multilevel mixed model using random intercept of each individual trajectory by residual maximum likelihood estimation. The longitudinal model was also adjusted for age, sex, injury level, and body mass index.

All analyses were performed using two-tailed tests, with p values  $< 0.05$  considered statistically significant. All analyses were performed using Stata 16.1 (StataCorp, Texas, US).

#### *Ethical considerations*

Written informed consent was obtained from all participants before data collection. The study is compliant with the Swiss Human Research Act (810.30 Federal Act of September 30, 2011, on Research involving Human Beings) and Federal Regulations on Data Protection (235.1 Federal Act of June 19, 1992, on Data Protection). The study has been cleared by ethics committees of participating centers, namely, Ethics Committee northwest/central Switzerland (EKNZ): PB\_2016-00183, Ethics Committee Vaud (CEVD): 032/13 (CEVS), Ethics Committee Zurich (KEKZH): 2013-0249]. To protect the confidentiality of the participants, the investigators and data analysts only accessed de-identified datasets. The dataset was kept under the secured servers of the Swiss Paraplegic Research. This study conforms to STROBE guidelines for reporting (see **Supplementary Checklist, Supplemental Digital Content 2**, <http://links.lww.com/PHM/C55>).

## RESULTS

A total of 706 individuals with SCI were included in the analysis (**Figure 1**). The median age is 53.5 years (Interquartile range, IQR 39-65), with the majority (59.02%) of the participants belonging to 46-60 years (32.72%) and 61-75 years (26.63%) age groups (**Table 1**). There were more males in the enrolled population compared to females (69.55% vs. 30.45%). The majority of individuals had paraplegia (53.26%), incomplete motor injury (53.68%), and traumatic injuries (60.20%). The list of non-traumatic causes can be found in the Appendix (**Table S1, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). The median length of rehabilitation was 5.2 months (3.1-7.4).

### *Cardiovascular risk profile and cardiometabolic syndrome at discharge*

Dyslipidemia was observed in 46.7% and impaired fasting glucose was seen in 26.1% of the cohort (**Figure 2**). Overweight (high-risk for obesity) was seen in 72.6% using SCI-specific BMI cut-off and 49.9% using the cut-off for the general population. Moreover, 37.2% were considered high cardiovascular risk (>10% on FRS) and 35.4% have cardiometabolic syndrome.

### *Longitudinal changes in rehabilitation outcomes*

Over the period of rehabilitation, we saw an overall improvement in respiratory function (**Table 3**) which was reflected in increasing FVC (from 2.65 L (1.97-3.3) to 2.72 L (IQR 2.12-3.33),  $p < 0.001$ ) and PEF (from 3.2 L/min (IQR 2.48-3.96) to 3.4 L/min (2.6-4.1).  $p < 0.001$ ). Similar improvement were seen in individuals with tetraplegia and paraplegia (**Table S2a, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**) and those with complete and incomplete injury (**Table S2b, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Changes in lung function and spasticity does not seem to

vary according to injury characteristics (**Table S2c, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Total SCIM score increased from 34 (IQR 21-58) to 74 (IQR 56-91) ( $p < 0.001$ ), and mobility SCIM increased from 4 (IQR 0-7) to 12.5 (IQR 7-26) ( $p < 0.001$ ). Individuals with incomplete injury has incremental change in functioning compared to those with complete injury (**Table S2c, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**).

There was an increase in spasticity in the lower extremities. At baseline, moderate to severe spasticity was observed in 9.28% of the left extremity and 9.58% of the right extremity, which increased to 19.94% and 18.63%, respectively (**Table 3**). We observed an increase in handgrip strength from 34.2 kg (IQR 25.2-45.9) to 34.6 (IQR 26-27.3) ( $p < 0.001$ ). We observed no statistically significant changes in body anthropometrics over the rehabilitation period.

#### *Rehabilitation outcomes and cardiovascular risk*

At discharge, almost one-third of study participants were considered to have a poor cardiovascular risk profile. Specifically, 37.2% (162/435) had high FRS ( $\geq 10\%$ ), 40.9% (200/488) had low HDL ( $< 1.0$  mmol/L in males or  $< 1.3$  mmol/L in females), and 26.1% (125/479) had high fasting glucose levels ( $> 7.0$  mmol/L) (**Table S3, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Cardiovascular risk scores were also computed based on SCORE2 and WHO risk score found in the Appendix (**Table S4, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Waist circumference ( $> 86.5$  cm), and higher BMI ( $> 22$  kg/m<sup>2</sup>) were associated with higher FRS compared to lower anthropometric measures (waist  $\beta$  0.28 95% CI 0.16, 0.40; BMI  $\beta$  0.32 95% CI 0.19, 0.46) (**Table 4**). Waist circumference was also associated with lower HDL-C by 0.10



mmoL/L and 0.09 mmoL/L, respectively. Crude and age-adjusted models were consistent with fully adjusted models (**Table S5a, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). We also fitted an association models using SCORE2 and WHO CVD Risk score with comparable findings (**Table S5b, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**)

Higher PEF (>3.4 L/min) and FEV1 (>0.8 L) were associated with 0.16 mmoL/L and 0.14 mmoL/L higher HDL-C compared to those with lower respiratory function (**Table 4**). These findings were consistent with age-, sex-, and medication-adjusted models (**Table S6, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Likewise, age-, sex- and medication-adjusted models of FVC showed higher HDL-C, but the fully adjusted model did not reach statistical significance.

Lastly, a high functional independence score (>74) and high mobility score (>12.5) are associated with 0.21 mmoL/L and 0.19 mmol/L higher HDL-C compared to those with lower scores. These associations are consistent across crude and incrementally adjusted models (**Table S6, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**).

We did not find any statistically significant association between rehabilitation outcomes and fasting glucose levels in fully adjusted models (**Table 3**) except for spasticity. Those with moderate-severe grade spasticity (+1 to 4/4) of the left lower extremity had 0.25 mmol/L higher fasting glucose compared to low-grade spasticity (0 to 1/4). Across crude and various adjusted models, no statistically significant association was seen (**Table S7, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**).

We explored the association between longitudinal changes in rehabilitation outcomes and cardiovascular risk profile by fitting a multilevel mixed model approach (**Table S8, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). An increase in BMI and waist circumference during rehabilitation was associated with increase in FRS and decrease in HDL (high CV risk). An increase in respiratory function (FVC and PEF) and functioning are associated with increase in HDL (low CV risk).

#### *Sensitivity analyses*

We iterated our models using independent variables treated as continuous data. The association between body anthropometrics, respiratory function, and functional independence scores with FRS and HDL-C were consistent with main analysis (**Tables S9 and S10, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Higher waist circumference as a continuous variable was associated with higher fasting glucose ( $\beta$  0.01 95% CI 0.002, 0.01), and were not observed in dichotomization (**Table S11, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**).

We restricted our analysis to specific groups according to injury characteristics (motor complete, traumatic injury, and individuals with paraplegia) (**Table S12, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). The anthropometric measures were consistently associated with higher FRS in most of these groups but not with HDL-C. The association of functional independence was also consistent with higher HDL-C across different subgroups. We found that higher spasticity is associated with higher fasting glucose for individuals with paraplegia and traumatic injury

We iterated our fully corrected models using sex as an interaction term (**Table S13, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). Sex-rehabilitation outcome interactions were statistically significant in fasting glucose-BMI, HDL-FVC, HDL-maximum grip, and HDL-spasticity. On fitting interaction terms in our fully adjusted models, statistically significant associations with respiratory function, spasticity, muscle strength, and functioning with HDL. We also found statistically significant association with anthropometric measures and FRS. Also, we performed subgroup analyses to determine the influence of age in our estimates (**Table S14, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). The association of anthropometrics with FRS is consistent with adults ( $\leq 50$  years). Also, the association between respiratory function and HDL was also consistent with adults. The association between independent functioning and HDL was consistent across different subgroups.

Missing dependent (**Table S15, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**) and independent (**Table S16 a-c, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**) variables were tabulated accordingly. We compared the clinical profile of individuals with missing dependent variables (**Table S16 a-c, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**). We observed a higher proportion of females with missing dependent variables. Those with missing FRS (or lacking one component from the composite score) were older (**Table S16a, Supplemental Digital Content 1, <http://links.lww.com/PHM/C54>**).

## DISCUSSION

We found that at least a third of the cohort are considered at high cardiovascular risk and majority are classified high-risk for obesity and cardiometabolic syndrome. Over the period

of rehabilitation, we found that individuals with SCI improved in their respiratory function, muscle strength and functioning. Furthermore, we showed an association between rehabilitation outcomes and cardiovascular risk profile. Future studies are to explore whether anthropometric measures, respiratory function, and functional independence may be used to prioritize screening in clinical settings when cardiovascular risk factors are not routinely measured. The summary of our main findings can be found in **Figure 2** (Graphical Abstract).

During on average 5.2 months in rehabilitation setting, our study participants improved respiratory function, muscle strength, and functional independence. This is in line with previous studies that reported improvements in gross motor and fine motor skills, a higher level of independence in activities of daily living, and higher motivation and outlook in life<sup>9,10,38</sup>. We also observed an increase in upper and lower extremities spasticity that is also physiologically expected as a long-term consequence of paralysis and immobility<sup>39</sup>. We did not see significant changes in any of the body anthropometrics measures across time. Dynamic changes in fat, lean mass, bone, and water distribution in an individual with SCI in a subacute phase could have contributed to this null finding<sup>40,41</sup>. Lean muscle mass declines below the level of injury because of the loss of neural control and disuse. Bone mass also decreases due to the lack of weight-bearing and physical stress that is crucial for mineral deposition and bone development. The acute stress results in higher energy requirements and sequestration of fat storage to aid in energy production<sup>3,40</sup>. However, upon physiologic stability, the body shifts from catabolic to anabolic processes, which promote fat deposition<sup>40</sup>. This is also similar to water accumulation as a response to stress and eventual diuresis after injury stabilization. All body anthropometrics (waist circumference and BMI) then may depend on which body components change during the injury spectrum.

Further, increase in anthropometric measures (BMI and waist circumference) were associated with poorer CV profile (increased FRS and decreased HDL) from our longitudinal models. Similarly, at discharge higher mobility and overall functional independence scores were associated with a better lipid profile, which is consistent over several sensitivity tests and upon fitting to a longitudinal model. This strong association is consistent with previous studies and systematic reviews on physical activity and cardiometabolic health <sup>42</sup>. Furthermore, longitudinal studies on chronic and subacute injuries have reiterated these findings <sup>43</sup>. Exercise or a high level of physical activity had been associated with higher HDL because of improved cholesterol efflux capacity that promotes the transport of LDLs back to the liver for excretion <sup>44</sup>. This is the purported mechanism on how anaerobic exercise confers its beneficial effects on atherosclerosis. As such, there is a current trend of standardizing the measurement and prescription of exercises or physical activity in individuals with SCI <sup>45</sup>.

Better respiratory function was associated with a better cardiovascular risk profile. Higher peak expiratory flow and expiratory volume is used as a measure of cardiorespiratory fitness, which directly influences cardiometabolic health <sup>46,47</sup>. Another hypothesis to support is the loss of elastic recoil and reduced chest wall compliance, which were both associated with reduced vascular elasticity related to hypertension <sup>47,48</sup>. Finally, reduced oxygen uptake and relative hypoxia are associated with systemic inflammation, with a concomitant increase in inflammatory cytokines (C-reactive protein and interleukin-6) that have a crucial role in lipid profile and atherogenesis <sup>46</sup>. This association was previously observed in 253 chronic SCI individuals (4 years from injury), such that lower FEV1 and FVC were associated with hypertension and diabetes in a univariate model <sup>11</sup>, but did not show any significant association with HDL. The previous study had a younger study population and thus, had a

lower prevalence of dyslipidemia compared to our group. Thus, the previous study may have been underpowered to detect any significant findings <sup>11</sup>.

A previous study found an association between severe spasticity with reduced adiposity and lower fasting glucose <sup>12</sup>. It was hypothesized that the higher muscle mass recruitment and contraction from severe spasticity were protective against fat cell deposition resulting in a lower total body fat percentage. Our fully-adjusted model showed the higher spasticity group had 0.25 mmol/L higher fasting glucose compared to the lower spasticity (**Table 3**). Our focus was more on subacute injury without prior history of cardiovascular disease, which could have led to the divergence in our findings and the past study. The past study was focused on chronic injury with a smaller sample size, and did not adjust for important confounders, although, the past study has broader range of individuals with mild spasticity to severe spasticity <sup>12</sup>. Our hypothesis, instead, is that increased physical capacity and activity as shown by higher functional independence results in a better glucose profile with an opposite relationship with spasticity.

Finally, the FRS was higher and HDL-C was lower in individuals whose waist circumference and BMI were higher than this SCI-specific cutoff as compared to individuals with lower BMI and waist circumference values. Current anthropometric measures seem to underestimate the fat composition in SCI population <sup>49,50</sup>. Studies have been done to validate the use of waist circumference and body mass index and the association of cardiovascular risk including cut-off range for high-risk <sup>31,32,51</sup>. Body mass index was validated using fat composition measured through bioelectrical impedance and C reactive protein to approximate the high-risk groups <sup>32</sup>. Waist circumference cutoff was established using fat composition measured through MRI and inflammatory biomarkers <sup>31,51</sup>. These studies were mostly done in

individuals with chronic SCI, and none, yet, has explored this SCI-specific cut-off in subacute SCI. Although our study did not aim to validate these cut-offs in subacute SCI, our findings seem to be consistent with the current literature.

Few studies, to date, have ventured on the role of sex or gender with cardiovascular risk factors in individuals with SCI<sup>52</sup>. By fitting sex-rehabilitation outcome interaction term in our models, we identified sex as an effect modifier in our main finding. This means that the association between cardiovascular risk and rehabilitation outcome differs according to sex. Males have poorer cardiovascular profile than females<sup>53</sup>. For rehabilitation outcome, a previous longitudinal assessment showed males having better improvement in functionality compared to females<sup>54</sup>. As such, our findings were expected. Our main analysis was driven mostly by the high proportion of males. The small number of females in our study is a main limitation to fully understand effect modification by sex in some of our dependent variables.

#### *Clinical implications and outlook*

Clinical guidelines recommended cardiometabolic screening of individuals with SCI at one to three-year intervals, yet, they do not specify the optimal timing for initial screening.<sup>55</sup> Our data showed that there is huge proportion of individuals with SCI being at high cardiovascular risk prior to discharge (e.g., high proportion of study population was with metabolic syndrome). And as such, early cardiometabolic screening, at least for those at-risk should be considered prior to rehabilitation discharge.

Discharge planning from inpatient rehabilitation is mostly based on physiologic stability, functional independence, and readiness to reintegrate into society. Although CVDs have been recognized as an increasingly important cause of mortality<sup>56,57</sup>, cardiovascular screening may

not be a medical priority before discharge. A retrospective assessment of hospital records participating in SwiSCI showed that only one-fifth of the study population was screened for lipid profile (51), even though more than one-third may be at increased CVD risk and 39.5% met criteria for metabolic syndrome <sup>4</sup>. Our study revealed that higher rehabilitation outcomes relate to better cardiovascular health. To put into context, the use of statins result to 3-5% increase in HDL (2 mg/dL or 0.05 mmol/L) <sup>58</sup>. In our study, a higher functional independence score has 0.2 mmol/L higher than those with lower score. The functional independence score we used has self-care, respiration, and sphincter function (bladder and bowel), and mobility subcomponents. A high score on self-care, respiration, bladder, and bowel function is extremely important to achieve before discharge as this usually translates to less need for specialized care provided by a rehabilitation facility. Individuals with lower mobility scores might require less specialized care that could prompt early discharge, and they would likely benefit from cardiovascular screening prior to discharge. Furthermore, individuals with lower respiratory ventilation should also be prioritized for screening. As of time of writing, only one study has explored this association <sup>11</sup>, and would further be useful if validated in other centers or countries with different rehabilitation programs. This information would be essential in healthcare settings with earlier discharge and shorter lengths of stay <sup>38,59,60</sup>.

In addition, future studies should explore sex differences on CV risk and CV diseases. More studies should report sex-disaggregated or female-specific results on the association of rehabilitation outcomes and CV risk. SCI epidemiology, as of now, still occurs more commonly in males. This is rapidly changing as there are more and more non-traumatic and elderly sustaining injuries, of which sex disparity is less evident. As such, the epidemiology of comorbidities in women with SCI are increasingly becoming more important.



Finally, although current risk prediction tools are able to predict cardiovascular events within five years, the true burden of the disease is grossly underestimated by the FRS<sup>25</sup>. Individuals with SCI have lower blood pressure and lower lipid profiles which computationally result to lower risk scores, yet remain at high-risk in reality<sup>61</sup>. Physical activity and baseline inflammation status are factors not considered and should be used to increase predictive ability. Thus, future studies should validate, adjust, or develop a new cardiovascular risk-scoring tool based on hard outcomes (cardiovascular events or cardiovascular deaths) in this group using long-term follow-up in this cohort. Novel cardiovascular biomarkers should also be explored to shed light on the pathophysiology and provide acuity in predictive tools.

#### *Strengths and Weaknesses*

We presented results from prospectively collected data from a multicenter study that has less selection bias compared to most studies that which are based on registries and administrative data from hospitals. The use of administrative data from hospitals are problematic as physicians decide when and what clinical outcomes to measure (confounding by indication). Using the SwiSCI cohort, we routinely collected and used SCI-specific dependent and independent variables on all participants. As one of the largest SCI cohorts in the literature, we provided a more precise estimate, adjusted for important confounding variables, performed subgroup analyses (by age and sex), and iterated restrictive analysis (according to injury characteristics). Our population is more homogenous as we removed all individuals with incident and prevalent CVDs, and because of the inpatient setting, which standardizes diet and physical activity to a healthy level. Our cohort has a huge proportion of elderly compared to other cohorts. This makes our cohort more ideal in determining cardiometabolic changes as we do not expect any abnormal findings in a cohort with young participants. Lastly, our final model was iterated, not only by different subgroups according to age, sex,

and injury characteristics, but also by longitudinal multilevel mixed model analysis with consistent findings. As such, our findings are robust as the beta estimates maintain their statistical significance across different scenarios.

Although our findings found important associations that may have implications for future clinical guidelines, these were based on cross-sectional study that are prone to reverse causation. This would mean that those with higher cardiovascular risk factors (independent variables) might be those with rehabilitation measure (dependent variables), and what we observed is a reversed cause-and-effect. Although we analyze longitudinally as part of sensitivity, the short latency period between dependent and independent variables could still make reverse causality possible. In addition, we used study sample-medians of rehabilitation outcomes since there are no SCI-specific cut-offs in the literature and since sample size would mostly likely be underpowered for comparisons of unequal groups. Except for SCI-specific BMI and waist circumference, a cutoff on other rehabilitation outcomes that predicts a cardiovascular outcome would be more useful. However, this is impossible with our prospectively collected data as we still lack hard outcomes (e.g., incident CVD events) in our cohort. Another important limitation is the enrollment bias that was previously established in another publication<sup>19</sup>. Underreporting of women, the elderly, and those with lower functional independence should be considered in interpreting our findings. Lesser women in our analysis group could make our findings less representative. It is also likely that the true association is underestimated, considering the underrepresentation of the elderly and more severe SCI. Our analysis was also limited to common laboratory measures available in the clinics. It would have been ideal to have high-sensitivity C reactive protein and other cardiovascular inflammatory makers that aids in characterizing the risk. The use of HDL-C for cardiovascular risk profiling could also be imprecise considering previous study on non-

linear relationship<sup>62</sup> and non-causal associations<sup>63</sup>. However, low HDL profile is more frequently seen in SCI compared to the general population<sup>64</sup>. This was hypothesized as the reason for high cardiovascular risk in SCI<sup>64,65</sup>. We also lack detailed interventions that a participant received during inpatient admission. Although we collected the diet and physical activity data, we lacked information on the detailed exercise regimen, occupational training, and psychological interventions. Our information on the rehabilitation regimen does not have the adequate granularity to point out a specific intervention that provided cardiovascular benefit.

Finally, we used cardiometabolic risk profile, FRS, WHO CV score, and SCORE 2 to approximate cardiovascular risk. We have observed discrepancies when comparing the change in proportion of individuals classified as high risk over rehabilitation stay when using the three scores. This can be driven by lack of validation of such scores in SCI population, inherent differences in risk estimation, and disparities in components. Cardiovascular health promotion focuses on preventing hard cardiovascular event such as myocardial infarction, coronary diseases or cardiovascular deaths, none of which were observed during our study period. As such, care should be taken in interpreting our findings not to over extrapolate in disease prevention.

## **Conclusion**

We observed a high burden of cardiometabolic syndrome and high CVD risk upon initial rehabilitation stay discharge. We also observed a significant improvement in rehabilitation outcomes (i.e., respiratory function, mobility, and SCIM score) during rehabilitation. Higher respiratory function, mobility, and functional independence score were associated with a better cardiovascular profile at discharge and sex was identified as effect modifier. Thus,

aside from established cardiovascular risk factors, rehabilitation outcomes may be useful to improve current CVD prediction models in the SCI population. Yet, this remains to be verified in future studies, considering our limitations in study design and short follow-up period.

ACCEPTED

## References

1. World Health Organization and International Spinal Cord Society. *International Perspectives on Spinal Cord Injury*. Geneva: WHO Press; 2013.
2. Zimmer MB, Nantwi K, Goshgarian HG. Effect of spinal cord injury on the respiratory system: basic research and current clinical treatment options. *J Spinal Cord Med*. 2007;30(4):319-330.
3. Felleiter P, Krebs J, Haeberli Y, Schmid W, Tesini S, Perret C. Post-traumatic changes in energy expenditure and body composition in patients with acute spinal cord injury. *Journal of rehabilitation medicine*. Jul 7 2017;49(7):579-584.
4. Raguindin PF, Stoyanov J, Eriks-Hoogland I, et al. Cardiometabolic risk profiling during spinal cord injury rehabilitation: A longitudinal analysis from Swiss Spinal Cord Injury cohort (SwiSCI). *PM R*. Jun 1 2022;n/a(n/a).
5. Denis AR, Feldman D, Thompson C, Mac-Thiong JM. Prediction of functional recovery six months following traumatic spinal cord injury during acute care hospitalization. *J Spinal Cord Med*. May 2018;41(3):309-317.
6. Cifu DX, Seel RT, Kreutzer JS, McKinley WO. A multicenter investigation of age-related differences in lengths of stay, hospitalization charges, and outcomes for a matched tetraplegia sample. *Arch Phys Med Rehabil*. Jul 1999;80(7):733-740.
7. Wilson JR, Davis AM, Kulkarni AV, et al. Defining age-related differences in outcome after traumatic spinal cord injury: analysis of a combined, multicenter dataset. *Spine J*. Jul 1 2014;14(7):1192-1198.
8. Burns AS, Marino RJ, Kalsi-Ryan S, et al. Type and Timing of Rehabilitation Following Acute and Subacute Spinal Cord Injury: A Systematic Review. *Global Spine J*. Sep 2017;7(3 Suppl):175S-194S.

9. Tooth L, McKenna K, Geraghty T. Rehabilitation outcomes in traumatic spinal cord injury in Australia: functional status, length of stay and discharge setting. *Spinal Cord*. Apr 2003;41(4):220-230.
10. Whalley Hammell K. Experience of rehabilitation following spinal cord injury: a meta-synthesis of qualitative findings. *Spinal Cord*. Apr 2007;45(4):260-274.
11. Koseoglu BF, Safer VB, Oken O, Akselim S. Cardiovascular disease risk in people with spinal cord injury: is there a possible association between reduced lung function and increased risk of diabetes and hypertension? *Spinal Cord*. Jan 2017;55(1):87-93.
12. Jung IY, Kim HR, Chun SM, Leigh JH, Shin HI. Severe spasticity in lower extremities is associated with reduced adiposity and lower fasting plasma glucose level in persons with spinal cord injury. *Spinal Cord*. Apr 2017;55(4):378-382.
13. Dallmeijer AJ, van der Woude LH, van Kamp GJ, Hollander AP. Changes in lipid, lipoprotein and apolipoprotein profiles in persons with spinal cord injuries during the first 2 years post-injury. *Spinal Cord*. Feb 1999;37(2):96-102.
14. Libin A, Tinsley EA, Nash MS, et al. Cardiometabolic risk clustering in spinal cord injury: results of exploratory factor analysis. *Top Spinal Cord Inj Rehabil*. Summer 2013;19(3):183-194.
15. Cragg JJ, Noonan VK, Krassioukov A, Borisoff J. Cardiovascular disease and spinal cord injury: results from a national population health survey. *Neurology*. Aug 20 2013;81(8):723-728.
16. Sabharwal S. Addressing cardiometabolic risk in adults with spinal cord injury: acting now despite knowledge gaps. *Spinal Cord Series and Cases*. 2019;5(96).

17. Yarar-Fisher C, Heyn P, Zanca JM, Charlifue S, Hsieh J, Brienza DM. Early Identification of Cardiovascular Diseases in People With Spinal Cord Injury: Key Information for Primary Care Providers. *Arch Phys Med Rehabil.* Jun 2017;98(6):1277-1279.
18. Boehl G. *Spinal cord injury: Associated variations in diagnostic and prognostic blood biomarkers.* Lucerne: Department of Health Sciences and Medicine, University of Lucerne.
19. Fekete C, Gurtner B, Kunz S, et al. Inception cohort of the Swiss Spinal Cord Injury Cohort Study (SwiSCI): Design, participant characteristics, response rates and non-response. *Journal of rehabilitation medicine.* Feb 26 2021;53(2):jrm00159.
20. Post MW, Brinkhof MW, von Elm E, et al. Design of the Swiss Spinal Cord Injury Cohort Study. *Am J Phys Med Rehabil.* Nov 2011;90(11 Suppl 2):S5-16.
21. Biering-Sørensen F, DeVivo MJ, Charlifue S, et al. International Spinal Cord Injury Core Data Set (version 2.0)-including standardization of reporting. *Spinal Cord.* Aug 2017;55(8):759-764.
22. DeVivo MJ, Biering-Sorensen F, New P, Chen Y, International Spinal Cord Injury Data S. Standardization of data analysis and reporting of results from the International Spinal Cord Injury Core Data Set. *Spinal Cord.* May 2011;49(5):596-599.
23. Catz A, Itzkovich M, Tesio L, et al. A multicenter international study on the Spinal Cord Independence Measure, version III: Rasch psychometric validation. *Spinal Cord.* Apr 2007;45(4):275-291.
24. D'Agostino RB, Sr., Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation.* Feb 12 2008;117(6):743-753.

25. Barton TJ, Low DA, Bakker EA, et al. Traditional Cardiovascular Risk Factors Strongly Underestimate the 5-Year Occurrence of Cardiovascular Morbidity and Mortality in Spinal Cord Injured Individuals. *Arch Phys Med Rehabil.* Jan 2021;102(1):27-34.
26. SCORE2 risk prediction algorithms: new models to estimate 10-year risk of cardiovascular disease in Europe. *Eur Heart J.* Jul 1 2021;42(25):2439-2454.
27. World Health Organization cardiovascular disease risk charts: revised models to estimate risk in 21 global regions. *The Lancet. Global health.* Oct 2019;7(10):e1332-e1345.
28. Stone NJ, Robinson JG, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* Jul 1 2014;63(25 Pt B):2889-2934.
29. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes care.* Jan 2003;26 Suppl 1:S5-20.
30. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *Jama.* Feb 5 2014;311(5):507-520.
31. Gill S, Sumrell RM, Sima A, Cifu DX, Gorgey AS. Waist circumference cutoff identifying risks of obesity, metabolic syndrome, and cardiovascular disease in men with spinal cord injury. *PLoS One.* 2020;15(7):e0236752.
32. Laughton GE, Buchholz AC, Martin Ginis KA, Goy RE, Group SSR. Lowering body mass index cutoffs better identifies obese persons with spinal cord injury. *Spinal Cord.* Oct 2009;47(10):757-762.



33. Klein S, Allison DB, Heymsfield SB, et al. Waist circumference and cardiometabolic risk: a consensus statement from Shaping America's Health: Association for Weight Management and Obesity Prevention; NAASO, The Obesity Society; the American Society for Nutrition; and the American Diabetes Association. *Am J Clin Nutr.* May 2007;85(5):1197-1202.
34. Marino RJ, Barros T, Biering-Sorensen F, et al. International standards for neurological classification of spinal cord injury. *J Spinal Cord Med.* Spring 2003;26 Suppl 1:S50-56.
35. Federal Statistics Office. *Swiss Health Survey 2007.* Neuchatel-CH: Federal Department of Home Affairs; 2008.
36. Deutsche Gesellschaft für Ernährung e. V. Vollwertig essen und trinken nach den 10 Regeln der DGE. 2017; [www.dge.de/10regeln](http://www.dge.de/10regeln), May 9 2022.
37. de Groot S, van der Woude LH, Niezen A, Smit CA, Post MW. Evaluation of the physical activity scale for individuals with physical disabilities in people with spinal cord injury. *Spinal Cord.* Jul 2010;48(7):542-547.
38. Kao YH, Chen Y, Deutsch A, Wen H, Tseng TS. Rehabilitation length of stay and functional improvement among patients with traumatic spinal cord injury. *Spinal Cord.* Mar 2022;60(3):237-244.
39. Adams MM, Hicks AL. Spasticity after spinal cord injury. *Spinal Cord.* Oct 2005;43(10):577-586.
40. Singh R, Rohilla RK, Saini G, Kaur K. Longitudinal study of body composition in spinal cord injury patients. *Indian journal of orthopaedics.* Mar 2014;48(2):168-177.
41. Silveira SL, Ledoux TA, Robinson-Whelen S, Stough R, Nosek MA. Methods for classifying obesity in spinal cord injury: a review. *Spinal Cord.* Sep 2017;55(9):812-817.

42. Itodo OA, Flueck JL, Raguindin PF, et al. Physical activity and cardiometabolic risk factors in individuals with spinal cord injury: a systematic review and meta-analysis. *Eur J Epidemiol.* Apr 7 2022.
43. de Groot S, Dallmeijer AJ, Post MW, Angenot EL, van den Berg-Emons RJ, van der Woude LH. Prospective analysis of lipid profiles in persons with a spinal cord injury during and 1 year after inpatient rehabilitation. *Arch Phys Med Rehabil.* Mar 2008;89(3):531-537.
44. Ruiz-Ramie JJ, Barber JL, Sarzynski MA. Effects of exercise on HDL functionality. *Curr Opin Lipidol.* Feb 2019;30(1):16-23.
45. Martin Ginis KA, van der Scheer JW, Latimer-Cheung AE, et al. Evidence-based scientific exercise guidelines for adults with spinal cord injury: an update and a new guideline. *Spinal Cord.* Apr 2018;56(4):308-321.
46. Wannamethee SG, Shaper AG, Rumley A, et al. Lung function and risk of type 2 diabetes and fatal and nonfatal major coronary heart disease events: possible associations with inflammation. *Diabetes care.* Sep 2010;33(9):1990-1996.
47. Wu Y, Vollmer WM, Buist AS, et al. Relationship between lung function and blood pressure in Chinese men and women of Beijing and Guangzhou. *International Journal of Epidemiology.* 1998;27(1):49-56.
48. Sparrow D, Weiss ST, Vokonas PS, Cupples LA, Ekerdt DJ, Colton T. Forced vital capacity and the risk of hypertension: the normative aging study *American Journal of Epidemiology.* 1988;127(4):734-741.
49. Raguindin PF, Bertolo A, Zeh RM, et al. Body Composition According to Spinal Cord Injury Level: A Systematic Review and Meta-Analysis. *Journal of Clinical Medicine.* 2021;10(17).

50. Nash MS, Gater DR, Jr. Cardiometabolic Disease and Dysfunction Following Spinal Cord Injury: Origins and Guideline-Based Countermeasures. *Phys Med Rehabil Clin N Am*. Aug 2020;31(3):415-436.
51. Sumrell RM, Nightingale TE, McCauley LS, Gorgey AS. Anthropometric cutoffs and associations with visceral adiposity and metabolic biomarkers after spinal cord injury. *PLoS One*. 2018;13(8):e0203049.
52. Raguindin PF, Muka T, Glisic M. Sex and gender gap in spinal cord injury research: Focus on cardiometabolic diseases. A mini review. *Maturitas*. May 2021;147:14-18.
53. Leinwand LA. Sex is a potent modifier of the cardiovascular system. *Journal of Clinical Investigation*. 2003;112(3):302-307.
54. Sipski ML, Jackson AB, Gomez-Marin O, Estores I, Stein A. Effects of gender on neurologic and functional recovery after spinal cord injury. *Arch Phys Med Rehabil*. Nov 2004;85(11):1826-1836.
55. Nash MS, Groah SL, Gater DR, Jr., et al. Identification and Management of Cardiometabolic Risk after Spinal Cord Injury: Clinical Practice Guideline for Health Care Providers. *Top Spinal Cord Inj Rehabil*. Fall 2018;24(4):379-423.
56. Brinkhof MW, Al-Khodairy A, Eriks-Hoogland I, et al. Health conditions in people with spinal cord injury: Contemporary evidence from a population-based community survey in Switzerland. *Journal of rehabilitation medicine*. Feb 2016;48(2):197-209.
57. Levi R, Hultling C, Seiger A. The Stockholm Spinal Cord Injury Study. 3. Health-related issues of the Swedish annual level-of-living survey in SCI subjects and controls. *Paraplegia*. Dec 1995;33(12):726-730.
58. McTaggart F, Jones P. Effects of statins on high-density lipoproteins: a potential contribution to cardiovascular benefit. *Cardiovasc Drugs Ther*. Aug 2008;22(4):321-338.

59. Günes S. Demographic and clinical profile and functional outcomes of patients with spinal cord injury after rehabilitation. *Türkiye Fiziksel Tıp ve Rehabilitasyon Dergisi*. 2017;63(1):66-71.
60. Shin JC, Kim DH, Yu SJ, Yang HE, Yoon SY. Epidemiologic change of patients with spinal cord injury. *Ann Rehabil Med*. Feb 2013;37(1):50-56.
61. Raguindin PF, Frankl G, Itodo OA, et al. The neurological level of spinal cord injury and cardiovascular risk factors: a systematic review and meta-analysis. *Spinal Cord*. Aug 20 2021.
62. Zhong GC, Huang SQ, Peng Y, et al. HDL-C is associated with mortality from all causes, cardiovascular disease and cancer in a J-shaped dose-response fashion: a pooled analysis of 37 prospective cohort studies. *European journal of preventive cardiology*. Jul 2020;27(11):1187-1203.
63. Voight BF, Peloso GM, Orho-Melander M, et al. Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *The Lancet*. 2012;380(9841):572-580.
64. Gilbert O, Croffoot JR, Taylor AJ, Nash M, Schomer K, Groah S. Serum lipid concentrations among persons with spinal cord injury - a systematic review and meta-analysis of the literature. *Atherosclerosis*. Feb 2014;232(2):305-312.
65. Wilt TJ, Carlson K, Goldish G, et al. Carbohydrate and lipid disorders and relevant considerations in persons with spinal cord injury. *Evid Rep Technol Assess*. 2008;163(163):1-95.

## Figure Legends

**Figure 1.** Flowchart of study participant enrollment

**Figure 2.** Cardiometabolic risk profile of individuals with SCI on discharge

**Figure 3.** Graphical abstract

ACCEPTED




Figure 1. Flowchart of study participants






Figure 2. Cardiometabolic risk profile of individuals with SCI at discharge

Figure 3





**Table 1. Baseline demographics and injury characteristics of study participants**

Parameters	Eligible population (n=706)
<b>Baseline demographics</b>	
Age, years, median (IQR)	53.5 (39-65)
Age groups, n (%)	
• 18-30	105 (14.87)
• 30-45	136 (19.26)
• 46-60	231 (32.72)
• 61-75	188 (26.63)
• 76+	46 (6.52)
Sex, n (%)	
• Male	491 (69.55)
• Female	215 (30.45)
<b>Injury characteristics (T4)</b>	
Injury level, n (%) <sup>1</sup>	
• Tetraplegia (C2-C8)	208 (29.46)
• Paraplegia (T1-S5)	376 (53.26)
• Unknown	122 (17.28)
Injury completeness, n (%) <sup>2</sup>	
• Motor complete	201 (28.47)
• Motor incomplete	379 (53.68)
• Unknown	126 (17.85)
Etiology of injury, n (%)	

<ul style="list-style-type: none"> <li>• Traumatic</li> </ul>	425 (60.20)
<ul style="list-style-type: none"> <li>• Non-traumatic</li> </ul>	281 (39.80)
AIS/ISNCSCI Impairment Scale, n (%) <sup>3</sup>	
<ul style="list-style-type: none"> <li>• A</li> </ul>	134 (22.95)
<ul style="list-style-type: none"> <li>• B</li> </ul>	67 (11.47)
<ul style="list-style-type: none"> <li>• C</li> </ul>	79 (13.53)
<ul style="list-style-type: none"> <li>• D</li> </ul>	299 (51.20)
<ul style="list-style-type: none"> <li>• E</li> </ul>	1 (0.17)
<ul style="list-style-type: none"> <li>• Unable to determine</li> </ul>	3 (0.51)
<ul style="list-style-type: none"> <li>• Unknown</li> </ul>	1 (0.17)
Time since injury, days, median (IQR)	14 (9-25)

<sup>1</sup>Injury level according to ISCOS standards of reporting

<sup>2</sup>Complete injury (ASIA A and B) and incomplete injury (ASIA C, D and E).

<sup>3</sup>Based on ISNCSCI, International Standards for Neurological Classification of Spinal Cord Injury and SCIM III, Spinal Cord Independence Measure version III

**Table 2. Lifestyle factors, health conditions and cardiovascular risk profile at baseline**

Parameters	Eligible population (n=706)
<b>Lifestyle factors (at T2)</b>	
Special diet, n (%)	
<ul style="list-style-type: none"> <li>• No special diet</li> <li>• On medical diet<sup>1</sup></li> </ul>	<p>479 (89.70)</p> <p>55 (10.30)</p>
Diet, n (%)	
<ul style="list-style-type: none"> <li>• Follow guideline on meat intake<sup>2</sup></li> <li>• Follows guidelines on fruit and vegetables intake<sup>3</sup></li> </ul>	<p>392 (73.82)</p> <p>475 (88.95)</p>
Nutritional score, median (IQR) <sup>4</sup>	4 (3-4)
Alcohol intake, n (%)	
<ul style="list-style-type: none"> <li>• Never</li> <li>• Drinker</li> </ul>	<p>116 (16.43)</p> <p>590 (83.57)</p>
Smoker, n (%)	
<ul style="list-style-type: none"> <li>• Never</li> <li>• Smoker</li> </ul>	<p>229 (43.45)</p> <p>298 (56.55)</p>
Physical activity (PASIPD), n (%) <sup>5</sup>	
<ul style="list-style-type: none"> <li>• Follows physical activity guidelines</li> <li>• Does not follow</li> </ul>	<p>305 (96.52)</p> <p>11 (3.48)</p>
<b>Health conditions</b>	
Medication intake, n (%)	
<ul style="list-style-type: none"> <li>• Steroid</li> <li>• Opioid use</li> </ul>	<p>24 (3.40)</p> <p>106 (15.01)</p>

● Anti-lipid medications/statins	62 (8.78)
Length of rehabilitation, month, median (IQR)	5.2 (3.1-7.4)
<b>Cardiovascular risk profile (at T4)</b>	
Framingham risk score, %, median (IQR) <sup>6</sup>	7.16 (2.5-13.6)
Fasting glucose, mmol/L, median (IQR)	5.0 (4.6-5.6)
Total cholesterol, mmol/L, median (IQR)	4.7 (3.98-5.4)
Triglycerides, mmol/L, median (IQR)	1.37 (1.0-1.89)
High density lipoprotein, mmol/L, median (IQR)	1.18 (0.96-1.4)
Low density lipoprotein, mmol/L, median (IQR)	2.8 (2.2-3.45)
HDL-C/Total cholesterol ratio, median (IQR)	0.25 (0.20-0.31)

<sup>1</sup>Has special indication for diet

<sup>2</sup>Recommended meat intake (<4 days/week) according to German Society for Nutrition (Deutsche Gesellschaft für Ernährung, 2013)

<sup>3</sup>Recommended fruits and vegetable intake (>3 portions/day) according to German Society for Nutrition (Deutsche Gesellschaft für Ernährung, 2013)

<sup>4</sup>Composite score computed using the dietary recommendation by the German Society for Nutrition (Deutsche Gesellschaft für Ernährung, 2013). This comprises the score for liquid intake, fruit and vegetables, and meat intake with an overall score of 6 as best and 0 as poor diet and nutrition.

<sup>5</sup>Physical Activity Scale for Individuals with Physical Disabilities (PASIPD) based on modification (Washburn RA, Zhu W, McAuley E, Frogley M, Figoni SF. The Physical Activity Scale for Individuals with Physical Disabilities: development and evaluation. Arch Phys Med Rehabil 2002;83:193-200). Physical activity was coded as active and sedentary based on the recommendation (30 mins of moderate or vigorous activity for 3x per day, or 90 mins cumulative for cardiometabolic health).

<sup>6</sup>10-year risk for first cardiovascular event based from D'Agostino RB, Sr., Vasan RS, Pencina MJ, Wolf PA, Cobain M, Massaro JM, et al. General cardiovascular risk profile for use in primary care: the Framingham Heart Study. Circulation. 2008;117(6):743-53.

**Table 3. Changes in the rehabilitation outcomes during rehabilitation period**

	Admission <sup>1</sup>	Discharge <sup>1</sup>	P value <sup>2</sup>	LMM ( Δ time )	P value <sup>2</sup>
<b>Anthropometric measure</b>					
• Body mass index kg/m <sup>2</sup>	24.7 (21.6-28.0)	24.7 (21.9-27.9)	0.564	0.01 (-0.18, 0.19)	0.95
• Waist circumference, cm	90.1 (81-100.3)	91.2 (82-100)	0.824	0.46 (-0.72, 1.65)	0.44
<b>Lung function</b>					
• Forced vital capacity (FVC), L	3.22 (2.48-3.96)	3.4 (2.6-4.1)	<0.001	-0.28 (-0.35,-0.22)	<0.01
• Peak expiratory flow (PEF), L/min	363 (271-456)	406 (308-499)	<0.001	-46.83 (-56.87,- 36.79)	<0.01
• Forced expiratory volume at 1 second (FEV1), L	2.65 (1.97-3.3)	2.72 (2.12-3.33)	<0.001	-0.20 (-0.25,-0.15)	<0.01
<b>Spasticity<sup>4</sup></b>					
• Left gastrocnemius	1 (1-2)	1 (1-2)	0.002	-	-
• Left gastrocnemius moderate-	34 (9.28)	73 (19.94)			

severe spasticity (+1-4/4), n, (%)					
• Right gastrocnemius	1 (1-2)	1 (1-2)	0.004	-	-
• Right gastrocnemius mild spasticity (+1-4/4), n, (%)	35 (9.58)	68 (18.63)			
<b>Muscle strength<sup>5</sup></b>					
• Right hand, kg,	34.1 (24-45.4)	34.4 (25.8-46.6)	<0.001	4.00 (-0.26, 8.26)	0.07
• Left hand, kg,	32.2 (22.6-42.7)	32.7 (23.3-43)	<0.001	2.99 (-0.60, 6.57)	0.10
• Best grip, kg,	34.8 (25.5-46.3)	36 (27.2-47.8)	<0.001		
<b>Functional Independence scale (SCIM III)<sup>6</sup></b>					
• Mobility total	4 (0-7)	12.5 (7-26)	<0.001	-8.58 (-9.20,-7.97)	<0.01
• Total SCIM	34 (21-58)	74 (56-91)	<0.001	-30.46 (-32.13,- 28.79)	<0.01

<sup>1</sup>Expressed in medians and IQR

<sup>2</sup>Wilcoxon signed rank test for paired samples

<sup>3</sup>Linear mixed model using individual patient as cluster and time as predictor variables.  $\beta$  (95% CI) is the change at discharge from baseline. Model is adjusted for age, sex, and injury level.

<sup>4</sup>Spasticity is based on Modified Ashworth scale comprising 6 grades of spasticity assessed by a rehabilitation specialist (0 increase in tone, and 5 rigidity in flexion or extension). Linear mixed models not provided as the scores did not satisfy regression assumptions.

<sup>5</sup>Hand grip strength is measured by hand dynamometer from 0-100 kg

<sup>6</sup>Spinal Cord Independence Measure (SCIM III) measures the functional independence score in different domains, namely, self-care, respiration and sphincter management, and mobility. Mobility subtotal comprise indoor mobility, mobility in moderate distances (10-100 m), outdoor mobility (>100 m), mobility using stairs and transfer from wheelchair to car.

ACCEPTED

**Table 4. Association of cardiovascular risk factors and rehabilitation parameters<sup>1</sup>**

Parameter (cut off)	Framingham risk score <sup>2</sup>	p value	HDL	p value	Glucose	p value
Anthropometric measure						
• Waist (86.5 cm) <sup>3</sup>	<b>0.28* [0.16, 0.40]</b>	<b>&lt;0.001</b>	<b>-0.10* [-0.20, -0.001]</b>	<b>0.047</b>	0.14 [-0.09, 0.37]	0.238
• Body mass index (22 kg/m <sup>2</sup> ) <sup>4</sup>	<b>0.32* [0.19, 0.46]</b>	<b>&lt;0.001</b>	-0.09 [-0.20, 0.02]	0.118	-0.004 [-0.24, 0.23]	0.973
Respiratory function						
• Forced vital capacity (2.72 L)	0.08 [-0.08, 0.24]	0.341	0.12 [-0.002, 0.25]	0.054	-0.03 [-0.27, 0.22]	0.842
• Peak expiratory flow (3.4 L/min)	0.02 [-0.14, 0.18]	0.806	<b>0.16* [0.03, 0.29]</b>	<b>0.013</b>	0.02 [-0.24, 0.27]	0.902
• Forced expiratory volume at 1 min (0.8 L)	-0.03 [-0.19, 0.13]	0.731	<b>0.14*[0.02, 0.26]</b>	<b>0.021</b>	0.07 [-0.19, 0.32]	0.605
Spasticity <sup>5</sup>						
• Left lower extremity spasticity (1/4)	-0.05 [-0.19, 0.09]	0.481	-0.10 [-0.20, 0.01]	0.075	<b>0.26* [0.03, 0.47]</b>	<b>0.023</b>



• Right lower extremity spasticity (1/4)	-0.02 [-0.15, 0.12]	0.784	-0.08 [-0.18, 0.02]	0.114	0.19 [-0.03, 0.40]	0.088
Muscle strength and frailty <sup>6</sup>						
• Left handgrip (33.3 kg)	0.02 [-0.19, 0.24]	0.834	0.13 [-0.04, 0.31]	0.128	-0.09 [-0.41, 0.23]	0.579
• Right handgrip (32.0 kg)	0.10 [-0.10, 0.31]	0.321	<b>0.16* [0.001, 0.32]</b>	<b>0.049</b>	0.01 [-0.30, 0.33]	0.926
• Best grip (34.63 kg)	0.13 [-0.10, 0.35]	0.270	-0.02 [-0.19, 0.16]	0.844	-0.05 [-0.39, 0.28]	0.758
Functioning <sup>7</sup>						
• Total SCIM (12.5)	0.09 [-0.05, 0.21]	0.383	<b>0.21*[0.11, 0.30]</b>	<b>&lt;0.001</b>	-0.008 [-0.21, 0.12]	0.937
• Mobility SCIM (74)	0.06 [-0.07, 0.18]	0.210	<b>0.19* [0.09, 0.29]</b>	<b>&lt;0.001</b>	0.003 [-0.20, 0.20]	0.980

<sup>1</sup>Adjusted for age, sex, injury level, medication use (statins and steroids) and body mass index (except for body anthropometrics as independent variables). Independent variables (rehabilitation outcomes) as dichotomous variables and dependent variables as continuous variables. \*p values < 0.05 in **BOLD**

<sup>2</sup>Log transformed values, values for blood pressure not shown as Framingham risk scores includes this dependent variable in the calculation.

<sup>3</sup>Waist adjusted for SCI cut off at 86.5 kg (Gill S, Sumrell RM, Sima A, Cifu DX, Gorgey AS (2020) Waist circumference cutoff identifying risks of obesity, metabolic syndrome, and cardiovascular disease in men with spinal cord injury. PLoS ONE 15(7): e0236752.

<https://doi.org/10.1371/journal.pone.0236752>)

<sup>4</sup>BMI adjusted for SCI cut off at 22 kg/m<sup>2</sup> (Laughton, G., Buchholz, A., Martin Ginis, K. et al. Lowering body mass index cutoffs better identifies obese persons with spinal cord injury. Spinal Cord 47, 757–762 (2009). <https://doi.org/10.1038/sc.2009.33>)

<sup>5</sup>Spasticity is based on Modified Ashworth scale comprising 6 grades of spasticity assessed by a rehabilitation specialist (0 increase in tone, and 5 rigidity in flexion or extension)

<sup>6</sup>Hand grip strength is measured by hand dynamometer from 0-100 kg

<sup>7</sup>Spinal Cord Independence Measure (SCIM III) measures the functional independence score in different domains, namely, self-care, respiration and sphincter management, and mobility. Mobility subtotal comprise indoor mobility, mobility in moderate distances (10-100 m), outdoor mobility (>100 m), mobility using stairs and transfer from wheelchair to car.

ACCEPTED