



# Article Contribution of the α5 nAChR Subunit and α5SNP to Nicotine-Induced Proliferation and Migration of Human Cancer Cells

Irida Papapostolou <sup>(D)</sup>, Daniela Ross-Kaschitza <sup>(D)</sup>, Florian Bochen <sup>(D)</sup>, Christine Peinelt <sup>(D)</sup> and Maria Constanza Maldifassi \*

> Institute of Biochemistry and Molecular Medicine, University of Bern, 3012 Bern, Switzerland; irida.papapostolou@unibe.ch (I.P.); daniela.ross@unibe.ch (D.R.-K.); florian.bochen@unibe.ch (F.B.); christine.peinelt@unibe.ch (C.P.)

\* Correspondence: maria.maldifassi@unibe.ch

Abstract: Nicotine in tobacco is known to induce tumor-promoting effects and cause chemotherapy resistance through the activation of nicotinic acetylcholine receptors (nAChRs). Many studies have associated the  $\alpha$ 5 nicotinic receptor subunit ( $\alpha$ 5), and a specific polymorphism in this subunit, with (i) nicotine administration, (ii) nicotine dependence, and (iii) lung cancer. The  $\alpha$ 5 gene CHRNA5 mRNA is upregulated in several types of cancer, including lung, prostate, colorectal, and stomach cancer, and cancer severity is correlated with smoking. In this study, we investigate the contribution of  $\alpha$ 5 in the nicotine-induced cancer hallmark functions proliferation and migration, in breast, colon, and prostate cancer cells. Nine human cell lines from different origins were used to determine nAChR subunit expression levels. Then, selected breast (MCF7), colon (SW480), and prostate (DU145) cancer cell lines were used to investigate the nicotine-induced effects mediated by  $\alpha$ 5. Using pharmacological and siRNA-based experiments, we show that  $\alpha 5$  is essential for nicotine-induced proliferation and migration. Additionally, upon downregulation of  $\alpha 5$ , nicotine-promoted expression of EMT markers and immune regulatory proteins was impaired. Moreover, the  $\alpha$ 5 polymorphism D398N ( $\alpha$ 5SNP) caused a basal increase in proliferation and migration in the DU145 cell line, and the effect was mediated through G-protein signaling. Taken together, our results indicate that nicotine-induced cancer cell proliferation and migration are mediated via  $\alpha$ 5, adding to the characterization of  $\alpha$ 5 as a putative therapeutical target.

**Keywords:** cancer; proliferation; migration; nicotinic acetylcholine receptors; α5 nicotinic receptor subunit; nicotine; ion channel

# 1. Introduction

Nicotinic acetylcholine receptors (nAChRs) are pentameric ligand-gated channels that, once activated, are permeable to sodium (Na<sup>+</sup>), potassium (K<sup>+</sup>), and calcium (Ca<sup>2+</sup>) ions [1–3]. In mammals, 16 different subunits have been identified [4]. Similarly to other pentameric receptors, such as  $\gamma$ -Aminobutyric acid type A (GABA-A) receptors [5,6], nAChRs can assemble as either homomeric or heteromeric channels, whereby the subunit composition determines the channel's kinetic and pharmacological characteristics [7–9]. Although nAChRs are known to be widely expressed through the peripheral nervous system (PNS) and central nervous system (CNS) [10], they are also found throughout the entire body, where they mediate diverse functions [11,12]. It is now recognized that apart from their very well-described ionotropic properties, these receptors also have metabotropic characteristics and activate several intracellular signaling pathways, not only in neurons but also in various other cell types [13–15]. Endogenous agonists, such as acetylcholine, and exogenous agonists, such as nicotine, activate these receptors by binding to very well-characterized sites found between subunits [16,17].

Citation: Papapostolou, I.; Ross-Kaschitza, D.; Bochen, F.; Peinelt, C.; Maldifassi, M.C. Contribution of the  $\alpha$ 5 nAChR Subunit and  $\alpha$ 5SNP to Nicotine-Induced Proliferation and Migration of Human Cancer Cells. *Cells* 2023, *12*, 2000. https:// doi.org/10.3390/cells12152000

Academic Editors: Michael X. Zhu and Yong Zhou

Received: 28 June 2023 Revised: 1 August 2023 Accepted: 2 August 2023 Published: 4 August 2023



**Copyright:** © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/). In the context of lung cancer, nicotine exposure is known to increase cell proliferation, cell invasion, migration, Ca<sup>2+</sup> influx, epithelial to mesenchymal transition (EMT), and to initiate specific signaling cascades because of nicotine's interaction with nAChRs [18–23]. Nicotine can also modify the expression of immune-regulatory proteins such as CD47 and PDL1 [24–29]. Several researchers have associated the nAChR subunits  $\alpha 5$ ,  $\alpha 3$ , and  $\beta 4$  with lung cancer [30–32]. Additionally, in lung-cancer-derived cell lines, the activation of  $\alpha 5$  induces proliferation, migration, invasion, and EMT transition events [33–35]. Interestingly,  $\alpha 5$  fails to contribute to agonist binding but acts as an accessory subunit [36–38], where  $\alpha 5$  can influence receptor traffic, sensitivity, efficacy, and Ca<sup>2+</sup> permeability, such as in the  $\alpha 4\beta 2\alpha 5$  and  $\alpha 3\beta 4\alpha 5$  receptors [39]. To date, however, how  $\alpha 5$  modulates the signaling properties of nAChRs in cancer remains an unresolved question [40]. Notably, the expression of receptors containing  $\alpha 5$  in the brain is important for dopamine release and attention [41,42] and is involved in nicotine's administration and withdrawal effects [43]. Consequently,  $\alpha 5$ -containing nAChRs are putative drug targets because of the association between lung cancer risk and nicotine's addictive effects.

Further studies have shown that a specific polymorphism in this subunit, which changes an aspartate, D, in position 398 to an asparagine, N, (referred to as  $\alpha$ 5SNP), is strongly linked with nicotine dependence, quantity of smoking, and lung cancer [44–47]. Several reports claim that patients who carry the  $\alpha$ 5SNP are more prone to chronic pulmonary disease (COPD) independent of their smoking status [48,49]. Humanized mouse models expressing  $\alpha$ 5SNP confirm these data and associate this polymorphism with altered airway epithelial remodeling, as well as with an increase in basal cellular proliferation [32]. In epithelial lung cells,  $\alpha$ 5SNP was found to cause an increase in proliferation dependent on adenylyl cyclase (AC) activation [32]. Furthermore, this  $\alpha$ 5SNP is located at the intracellular loop localized between the M3–M4 transmembrane domains, an area known to be relevant for G<sub>o</sub>-protein interactions [50] but with a largely unknown structure [38]. Likewise, it was proposed that  $\alpha$ 5 modulates the recovery from receptor desensitization, with its expression resulting in a higher rate of receptor signaling [40].

Apart from lung cancer, current data also associate cigarette smoking with other types of cancer, such as prostate cancer (PCa) [51,52], colorectal cancer (CRC) [53], and breast cancer [54,55]. Several reports show the expression of diverse nAChR subunits in these cancer types and indicate that the activation of nAChRs stimulates cellular migration and EMT [56,57]. To date, the relevance of  $\alpha$ 5 in these nicotine-induced effects is completely unknown. In the present study, we provide information on the function of  $\alpha 5$  in human cancer cell lines of different origins and suggest a general role of the  $\alpha$ 5 nicotinic receptor subunit in cancer cell migration and proliferation. Using datasets from The Cancer Genome Atlas (TCGA) and qPCR, we found that among nAChR subunits,  $\alpha$ 5 was prominently expressed throughout different types of cancer. Upon pharmacological blocking of nAChRs or siRNA-based silencing of  $\alpha$ 5, the nicotine-induced proliferation, migration, and EMT transition in breast, prostate, and colon cancer cells were reduced. In addition, the expression of  $\alpha$ 5SNP can enhance basal cell proliferation and migration in DU145 PCa cells, independent of nicotine exposure. In summary, our findings present valuable insights into human  $\alpha$ 5 signaling in nicotine-induced migration and proliferation, underlining  $\alpha$ 5's role as a future therapeutical target.

### 2. Materials and Methods

# 2.1. Cell Culture

The following human cancer cell lines were used: MCF7 (breast cancer; MEM; Gibco, Waltham, MA, USA); A549 (lung cancer; DMEM; Gibco, Waltham, MA, USA); PC9 (lung cancer; RPMI; Gibco, Waltham, MA, USA); SW480 (colorectal cancer; RMPI; Gibco, Waltham, MA, USA); SW620 (colorectal cancer; L15; Gibco, Waltham, MA, USA); CACO2 (colorectal cancer; DMEM; Gibco, Waltham, MA, USA); DU145 (prostate cancer; DMEM; Gibco, Waltham, MA, USA); PC3 (prostate cancer; RMPI; Gibco, Waltham, MA, USA); and LNCaP (prostate cancer; RMPI; Gibco, Waltham, MA, USA). All cell lines were obtained from

ATCC (Rockville, MD, USA). The corresponding media for each cell line was supplemented with 10% FBS (Gibco, Waltham, MA, USA), and additionally supplemented with 2 mM L-glutamine (Gibco, Waltham, MA, USA), 1% NEAA (Gibco, Waltham, MA, USA) in the case of DU145. Cells were grown in a humidified atmosphere containing 5%  $CO_2$  at 37 °C.

### 2.2. Drug Treatment

Nicotine and three nAChR antagonists (the non-specific inhibitor mecamylamine MEC, an  $\alpha$ 7 receptor inhibitor methyllycaconitine MLA, and a  $\beta$ 2 subunit containing receptor inhibitor dihydro- $\beta$ -erythroidine DHBE), together with the adenylyl cyclase inhibitor SQ22536 and the G<sub> $\alpha$ i</sub> signaling blocker pertussis toxin PTX, were evaluated in cellular proliferation and migration assays. Additionally, their ability to induce changes in the expression of certain proteins was assayed via qPCR. All antagonists and agonists were obtained from Tocris (Bristol, UK), and were dissolved in H<sub>2</sub>O according to the manufacturer's instructions. When antagonists' effects were evaluated, these effects were co-applied with nicotine or alone for the control experiments.

# 2.3. Proliferation Assay

Cell proliferation was evaluated as previously published by our group [58,59] using xCELLigence<sup>®</sup> E-Plates (ACEA Biosciences, San Diego, CA, USA). This system monitors changes in electrical impedance and records them as an increasing number of cells adhering to the surface of the plates. Depending on the cell type,  $2 \times 10^4$  or  $4 \times 10^4$  cells were plated in their corresponding complete growth medium with or without an agonist/antagonist, and proliferation was measured for 48–72 h in 15 min intervals. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (Version 9.3.1) software.

## 2.4. Quantitative Real-Time PCR (qPCR)

The expression of diverse genes was evaluated following previously published procedures from our group [58]. In brief, RNA from the diverse cell lines and treatments was obtained using a QIAshredder kit (Qiagen, Hilden, Germany) followed by an RNeasy Mini kit (Qiagen, Hilden, Germany). Afterward, reverse transcription of 2  $\mu$ g RNA was performed (High-Capacity cDNA Reverse Transcription Kit, Thermo Fisher Scientific, MA, USA), and gene expression was evaluated using a TaqMan Gene Expression Assay. All determinations were performed in triplicate. Results were analyzed with the  $\Delta$ Ct method, and expression levels were normalized to the housekeeping gene TATA-binding protein. When comparing relative changes in protein expression in treated cells, the expression was compared to non-treated cells. Table 1 shows the ThermoFisher (Waltham, MA, USA) primers/probes used to quantify expression levels:

Protein	Gene Name	Probe ID	
α3 subunit *	CHRNA3	Hs01088199_m1	
α4 subunit *	CHRNA4	Hs00181247_m1	
α5 subunit *	CHRNA5	Hs00181248_m1	
α7 subunit *	CHRNA7	Hs01063372_m1	
dupα7 subunit *	CHRFAM7A	Hs04189909_m1	
α9 subunit *	CHRNA9	Hs00395558_m1	
β2 subunit *	CHRNB2	Hs01114010_g1	
β4 subunit *	CHRNB4	Hs00609523_m1	
CD47	CD47	Hs00179953_m1	
PDL-1	CD274	Hs00204257_m1	

Table 1. Proteins and corresponding primers/probes used in the present study.

Table 1. Cont.

Protein	Gene Name	Probe ID	
Vimentin	VIM	Hs00958111_m1	
E-cadherin	CDH1	Hs01023895_m1	
TATA-binding protein	ТВР	Hs00427621_m1	

\* Corresponds to nAChR subunits. All probes are commercially available from ThermoFisher.

#### 2.5. Transwell Migration Assay

Transwell migration of cells was evaluated as previously published by our group [58] using the MCF7, DU145, and SW480 cell lines. Here,  $4 \times 10^4$  cells were seeded in 0% serum media in the top chamber of xCELLigence<sup>®</sup> CIM-Plates (ACEA Biosciences, San Diego, CA, USA). The respective culture media with 10% serum was added to the lowest chambers and served as a chemoattractant. For MCF7 cells, 1% serum was used in the top chamber. When compounds were being tested, these solutions were added to the top chamber. Afterward, migration was measured in 15 min intervals for 25 h. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with GraphPad Prism (Version 9.3.1) software.

### 2.6. Small Interfering RNA Transfection (siRNA)

We performed the siRNA transfections with a mix of 4 different human  $\alpha$ 5 siRNAs using a 1:1:1:1 ratio in a final concentration of 0.12 nmol siRNA. All siRNAs were obtained from Qiagen. The  $\alpha$ 5 siRNAs used were as follows: Hs\_*CHRNA5\_7*, target sequence CTGGTATCCGTATGTCACTTA; Hs\_*CHRNA5\_6*, target sequence CTGGACTC-CACCGGCAAACTA; Hs\_*CHRNA5\_5*, target sequence ATGGATCACAGGTTGATATAA; and Hs\_*CHRNA5\_4*, target sequence CTGAGTAACAGCTAATCTTTA. Control cells were transfected with non-silencing siRNA (ns-RNA) (Qiagen, Hilden, Germany), which presented no homology with any known mammalian gene. We performed the siRNA transfections using Interferin siRNA Transfection Reagent (Polyplus, Illkirch-Graffenstaden, France). Cells were incubated overnight with a transfection mix consisting of a combination of the corresponding serum-free media, the siRNA of interest or ns-RNA, and the Interferin solution. Afterward, the cells were exposed to treatments as specified in the drug treatment section, and the expression of diverse proteins was analyzed via qPCR. Treated cells were also used for proliferation and migration assays. Knockdown verification was performed 24 h or 48 h after siRNA transfection, and  $\alpha$ 5 expression was analyzed with qPCR.

#### 2.7. Constructs and Nucleofection

The  $\alpha$ 5.pcDNA3.1/V5 and  $\alpha$ 5D398N.pcDNA3.1/V5 epitope-tagged constructs used for transfection were a generous gift from Dr. Larry S. Barak and have been described elsewhere [40]. DU145 cells were transfected with either plasmid using an SE Nucleofection kit (Lonza, Basel, Switzerland). Briefly,  $1.5 \times 10^6$  cells suspended in nucleofection SE solution were transfected with the appropriate constructs using the DU145-specific program of the nucleofector. Next, cells were seeded and incubated for an additional 24 h before starting proliferation and migration experiments. Transfection verification was performed 24 h after nucleofection, and  $\alpha$ 5 expression was analyzed via qPCR and IF.

### 2.8. Immune Fluorescence (IF)

To positively ascertain the transfection of  $\alpha$ 5.pcDNA3.1/V5 and  $\alpha$ 5D398N.pcDNA3.1/V5 epitope-tagged constructs (a generous gift from Dr. Larry S. Barak) [40] into DU145 cells, specific immunofluorescence staining was performed on fixed cells. In brief, 24 h after nucleofection, the cells were fixed using 4% PFA in PBS (ThermoFisher, Waltham, MA, USA) for 15 min, and subsequently incubated with anti-V5 antibody (1:500, Cell Signaling, Danvers, MA, USA) overnight at 4 °C. Afterwards, cells were incubated with AlexaFluor 488 (1:1000, Invitrogen, Waltham, MA, USA) for 1 h. Samples were visualized using a

Nikon Eclipse Ti2-E Widefield microscope with a Photometrics BSI camera and SpectraX laser as light source, with either a  $40 \times$  or  $100 \times$  objective lens.

#### 3. Results

# 3.1. Expression of nAChR Subunits in Diverse Cancer Cell Lines: Focusing on the $\alpha 5$ Subunit

First, to assess the expression of diverse nAChR subunits in various cancers, we analyzed data from the Human Protein Atlas [60], which is based on the Cancer Genome Atlas TCGA project [61]. Figure 1A,B show that throughout different types of human cancers,  $\alpha$ 5 is a prominently expressed subunit.



**Figure 1.** Nicotinic acetylcholine receptor subunit (nAChRs) in different cancer types. Gene *CHRNA1*:  $\alpha$ 1 subunit; gene *CHRNA3*:  $\alpha$ 3 subunit; gene *CHRNA4*:  $\alpha$ 4 subunit; gene *CHRNA5*:  $\alpha$ 5 subunit; gene *CHRNA6*:  $\alpha$ 6 subunit; gene *CHRNA7*:  $\alpha$ 7 subunit; gene *CHRNA7*: dup $\alpha$ 7 subunit (partially duplicated  $\alpha$ 7 subunit isoform); gene *CHRNA9*:  $\alpha$ 9 subunit; gene *CHRNA10*:  $\alpha$ 10 subunit; gene *CHRNB2*:  $\beta$ 2 subunit; gene *CHRNB4*:  $\beta$ 4 subunit. (**A**) Heat map view of several nicotinic acetylcholine receptor (nAChR) subunit genes in human cancers (figure adapted from the Human Protein Atlas, www.proteinatlas.org; accessed on 1 May 2023). The heat map represents the median values per gene for each cancer type. The *CHRNA5* subunit is the most commonly expressed nAChR subunit in all cancer types. (**B**) *CHRNA5* expression levels in different human cancers. Each sample is shown as a circle on the graph, and the median is indicated with a red line. Sequencing data were reported as FPKM (fragments per kilo-base of transcript per million reads mapped). Data were generated by the Cancer Genome Atlas (Pathophysiology of *CHRNA5*, https://www.proteinatlas.org/ENSG0000016 9684-CHRNA5/pathology; accessed on 1 May 2023).

Next, we investigated the expression of  $\alpha$ 5 and other nAChR subunits that are known to mediate nicotine's pro-oncogenic effects in different human cancer cell lines. Despite the fact that the maximum expression of the  $\alpha$ 5 was in urothelial and testicular cancer, our study focused on the cancer types with the highest incidence and mortality [62], namely prostate, colon, and breast cancer. Correspondingly, we used the breast cancer cell line MCF7; the lung cancer cell lines A549 and PC9; the colorectal cancer cell lines SW480, SW620, and CACO2; and the prostate cancer cell lines DU145, PC3, and LNCAP. The gene expression profiles were studied via qPCR analysis, and the results are presented in Figure 2 and Table 2. The results show that nAChR mRNAs were differentially expressed throughout the different cell lines. In a comparison between the cell lines, the  $\alpha$ 3 and  $\alpha$ 4 subunits showed the strongest expression in SW480 cells; in a comparison between the subunits  $\alpha$ 5 was the most prominently expressed subunit in all tested cell lines. The  $\alpha$ 7 and dup $\alpha$ 7 transcripts were differentially expressed throughout all cell lines. Interestingly,  $\alpha$ 9 was barely detectable across all cells. Lastly,  $\beta$ 2 and  $\beta$ 4 were also expressed but differed between cell lines.



**Figure 2.** Cell-specific expression patterns of transcripts for nAChR subunits. Human cell lines examined: MCF7 (**A**), breast cancer), A549 (**B**), lung cancer), PC9 (**C**), lung cancer), SW480 (**D**), colorectal cancer), SW620 (**E**), colorectal cancer), CACO2 (**F**), colorectal cancer), DU145 (**G**), prostate cancer), PC3 (**H**), prostate cancer), and LNCAP (**I**), prostate cancer). The gene expression profiles were investigated via qPCR analysis. Bar graphs show the relative expression  $\pm$  SEM of each subunit normalized to endogenous TATA-binding protein expression. Measurements were made using 3–5 independent experiments performed in triplicate.

nAChR Subunit	MCF7	A549	PC9	SW480	SW620	CACO2	DU145	РС3	LNCAP
α3	$0.024 \pm 0.002$	$0.057 \pm 0.014$	n.d.	$\begin{array}{c} 0.261 \\ \pm 0.054 \end{array}$	$\begin{array}{c} 0.007 \\ \pm 0.002 \end{array}$	n.d.	$\begin{array}{c} 0.002 \\ \pm 0.0 \end{array}$	n.d.	$\begin{array}{c} 0.013 \\ \pm 0.0 \end{array}$
α4	n.d.	$0.005 \\ \pm 0.001$	n.d.	0.035 ±0.012	n.d.	0.004 ±0.002	n.d.	n.d.	n.d.
α5	1.431 ±0.001	3.396 ±0.743	$1.020 \pm 0.044$	$0.461 \pm 0.114$	3.739 ±0.284	0.698 ±0.301	$1.444 \pm 0.087$	0.321 ±0.075	0.469 ±0.052
α7	0.002 ±0.0	0.141 ±0.033	n.d.	0.010 ±0.004	0.007 ±0.002	$0.011 \pm 0.005$	0.013 ±0.002	$0.001 \\ \pm 0.0$	0.098 ±0.021
dup α7	n.d.	1.211 ±0.766	$0.009 \\ \pm 0.005$	0.002 ±0.001	0.058 ±0.021	$0.012 \pm 0.004$	$0.271 \pm 0.053$	0.077 ±0.025	$0.010 \\ \pm 0.002$
α9	n.d.	0.001 ±0.0	$0.001 \\ \pm 0.0$	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
β2	0.003 ±0.0	0.102 ±0.017	0.002 ±0.0	$0.055 \pm 0.016$	0.263 ±0.097	0.003 ±0.001	$0.001 \pm 0.0$	0.001 ±0.0	0.001 ±0.0
β4	0.031 ±0.007	0.107 ±0.006	0.009 ±0.003	$0.050 \\ \pm 0.019$	$0.013 \pm 0.005$	0.009 ±0.004	$\begin{array}{c} 0.02\\ 2\pm 0.0\end{array}$	0.001 ±0.0	n.d.

**Table 2.** Relative expression of nAChR subunits in diverse human cancer cell lines (refers to Figure 2, see above).

Results of the qPCR analysis of nAChR gene expression in the human breast cancer cell line MCF7; the lung cancer cell lines A549 and PC9; the colorectal cancer cell lines SW480, SW620, and CACO2; and the prostate cancer cell lines DU145, PC3, and LNCAP. The data relate to the relative expression  $\pm$  SEM (normalized to endogenous TATA-binding protein expression) calculated in 3–5 independent experiments performed with samples measured in triplicate; n.d. not determined.

Thus, the nAChR mRNA expression profiles demonstrate that  $\alpha 5$  is the most significantly expressed subunit in lung, breast, CRC, and PCa cancer cells, albeit at different levels. Previous reports showed that  $\alpha 5$  is upregulated in human lung cancer tissue [19], hepatocellular carcinoma tissue [63], gastric cancer tissue [64], and prostate cancer tissue [53]. Therefore, overexpression of the  $\alpha 5$  subunit may contribute to the general pathological role of nicotine in cancer cells.

# 3.2. Nicotine Increases Proliferation and Migration in Various Human Cancer Cell Lines through nAChRs

We next studied whether nicotine alters proliferation in several types of human cancer cell lines through the activation of nAChRs. We tested nicotine-induced proliferation in the breast cancer cell line MCF7, the CRC line SW480, and the PCa cell line DU145. For this analysis, cells were incubated with increasing concentrations of nicotine, from  $0.1 \ \mu$ M to  $10 \ \mu$ M, for 72 h. These concentrations were previously used to study nicotine as an inducer of proliferation in cancer cells [20,65]. To determine the nicotine-induced proliferation, we employed a label-free impedance-based xCELLigence system that was also used by our group in a previous study [59]. In this system, an increasing number of cells on a plate raises the electrical impedance, which can be displayed as an increase in the cell index parameter. Figure 3 shows that in the presence of 1  $\mu$ M nicotine, proliferation was significantly increased. In the MCF7 cell line (Figure 3A,B), this effect was already significant after 12 h, whereas the effect was significant in DU145 (Figure 3G,H) and SW480 (Figure 3D,E) cells after 24 h of incubation. Nicotine thus consistently increases proliferation but to a varying extent at different time points between cell lines tested. Our results are in line with previous observations in the lung cancer cell line A549 and the breast cancer line MCF7, in which the authors observed that nicotine augmented the proliferation of both cell types with the maximum effect observed at a concentration of 1  $\mu$ M nicotine [20].



**Figure 3.** Nicotine increases proliferation through nAChRs in MCF7, SW480, and DU145 human cancer cell lines. Proliferation was evaluated using a label-free impedance-based assay (xCELLigence). (**A**,**D**,**G**) The corresponding cells were treated with 0.1  $\mu$ M, 1  $\mu$ M, or 10  $\mu$ M nicotine. The cell proliferation index is plotted versus time. (**B**,**E**,**H**) Bar diagram of data (mean  $\pm$  SEM) at the indicated time points from the experiments in A-D-G, correspondingly. (**C**,**F**,**I**) The corresponding cells were treated with 1  $\mu$ M nicotine in the presence or absence of the inhibitors MEC, MLA, or DHBE. A bar diagram of the proliferation cell index (mean  $\pm$  SEM) at the indicated time points is shown. Measurements were made using 4–5 independent experiments performed in triplicate. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \* *p* ≤ 0.05, \*\* *p* ≤ 0.001.

To challenge the hypothesis that the increase in proliferation with 1  $\mu$ M nicotine is caused by nAChR activation, proliferation was tested with nicotine in the presence of the non-specific nAChR antagonist mecamylamine (MEC). Since earlier work showed the relevance of  $\alpha$ 7 in nicotine's pro-oncogenic effects [66], and because  $\alpha$ 7 mRNA was detected in the cell lines used (Figure 2), we also tested the role of  $\alpha$ 7 in nicotine-induced proliferation using a known  $\alpha$ 7 inhibitor (methyllycaconitine, MLA). Some cell lines also expressed detectable amounts of the  $\beta$ 2 transcript (SW480). Thus, the  $\beta$ 2 inhibitor dihydro-betaerythroidine (DHBE) was also employed. As outlined above, we used an impedance-based label-free xCELLigence assay and induced proliferation using 1 µM nicotine with or without 10 µM MEC, 10 nM MLA, or 10 µM DHBE. Previously, these concentrations were shown to inhibit the corresponding nAChR subtypes in different neuronal cell types [67]. As presented in Figure 3C,F,I, nicotine induced an increase in proliferation, reflected by an increase in the cell index. In the presence of MEC, nicotine failed to increase proliferation in all tested cell lines, demonstrating that nicotine's effect was mediated through activation of nAChRs. On the other hand, the presence of MLA or DHBE failed to inhibit nicotine-induced proliferation, showing that  $\alpha$ 7 and  $\beta$ 2 are not involved in nicotine-induced signaling. When applied without nicotine, the inhibitors did not modify the basal proliferation of SW480 or DU145 cells, demonstrating that they do not interfere with normal cellular growth mechanisms (Supplementary Figure S1).

Additionally, we evaluated whether nicotine, through nAChRs, affected migration in the DU145 cell line. Migration was investigated using an impedance-based assay with FBS as a chemoattractant [68]. As shown in Figure 4, when cells were incubated with 1  $\mu$ M or 10  $\mu$ M nicotine, we observed augmented cell migration at 24 h. This effect was significant for 1  $\mu$ M nicotine. As described above, we used the non-specific nAChR antagonist MEC to determine the contribution of nAChRs to this effect. Additionally, since it was shown that  $\alpha$ 7 is important for nicotine-induced migration in the A549 cell line [20], and because the qPCR analyses indicated that  $\alpha$ 7 is also expressed in DU145 cells, we also used the  $\alpha$ 7 receptor inhibitor MLA. The  $\beta$ 2 inhibitor DHBE was evaluated as well. Figure 4 shows that in the presence of MEC, migration was not affected by nicotine, reflecting the clear contribution of nAChRs to nicotine-induced migration. MLA failed to affect migration, and DHBE also impaired nicotine's effects.



**Figure 4.** Nicotine increased migration through nAChRs in the human prostate cancer cell line DU145. Analysis was performed using a label-free impedance-based migration assay (xCELLigence). (A) DU145 cells were treated with 0.1  $\mu$ M, 1  $\mu$ M, or 10  $\mu$ M nicotine in the presence or absence of the inhibitors MEC, MLA, or DHBE. The cell migration index is plotted versus time. (B) Bar diagram of the data (mean  $\pm$  SEM) at 12 and 24 h from the experiments in A. Measurements were made using 4–5 independent experiments performed in triplicate. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \*  $p \le 0.05$ , \*\*  $p \le 0.001$ .

When migration was studied in the SW480 cells (Supplementary Figure S2A), a more constrained migration was observed. Here, exposure to nicotine augmented the migration index, but because of the limited migration, no further studies were performed on these cells. In the MCF7 cell line, nicotine failed to induce migration at both tested concentrations (Supplementary Figure S2B).

# 3.3. siRNA Based Silencing of the $\alpha$ 5 Subunit Reduces Nicotine-Induced Proliferation and Migration in Several Cancer Cell Lines

As our qPCR studies indicated high expression levels of  $\alpha$ 5 in all cell lines, we sought to determine whether nicotine could modify proliferation and migration through the activation of  $\alpha$ 5-containing nAChRs in the analyzed cancer cell lines. As no specific agonist exists for  $\alpha$ 5-containing receptors [69], we studied the relevance of  $\alpha$ 5 by silencing the subunit using specific siRNAs. MCF7, SW480, and DU145 cells were transfected either with a mixture of four siRNAs specifically directed towards the  $\alpha$ 5 subunit ( $\alpha$ 5-siRNA) or with a non-silencing control RNA (ns-RNA). Proliferation and migration were also evaluated. The qPCR analysis in Figure 5 shows that siRNA-based knockdown of the  $\alpha$ 5 gene *CHRNA5* caused a ~90% decrease in transcripts in MCF7 and SW480 cells and a ~70% decrease in the DU145 cell line when evaluated at 24 h (Figure 5A) and 48 h (Figure 5B) post-transfection. As a control, we also determined the expression levels of other nAChR subunits (Supplementary Figure S3), and no considerable changes in expression levels were detected.



**Figure 5.** Knockdown of  $\alpha$ 5 suppressed the proliferative and migratory effects of nicotine on human cancer cell lines. (**A**,**B**) The  $\alpha$ 5 gene expression demonstrating siRNA knockdown (KD) of the subunit after 24 h (**A**,**B**) 48 h in MCF7, SW480, and DU145 cells. Data are shown as a percentage compared to expression of the subunit in the corresponding cell lines transfected with non-silencing ns-RNA (set to 100% expression, red dashed line). (**C**–**E**) Bar diagram of the cell proliferation index of the indicated cell line when silencing  $\alpha$ 5 gene expression.  $\alpha$ 5-siRNA- or ns-RNA-transfected cells were treated with 1  $\mu$ M nicotine. Data represent the mean  $\pm$  SEM in the indicated time points. (**F**) Migration cell index of DU145 cells as a result of silencing  $\alpha$ 5 gene expression.  $\alpha$ 5-siRNA- or ns-RNA-transfected cells were treated with 1  $\mu$ M nicotine. The bar diagram shows the mean  $\pm$  SEM 12 and 24 h after the treatment. Measurements were made using 3 independent experiments performed in triplicate. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \*  $p \leq 0.05$ , \*\*  $p \leq 0.001$ .

Next, we analyzed the contribution of  $\alpha$ 5 in the nicotine-induced proliferation in MCF7, SW480, and DU145 cells. Cells were transfected with  $\alpha$ 5 siRNA or ns-RNA, and proliferation was detected as described above. Figure 5C–E show that 1  $\mu$ M nicotine increased proliferation in ns-RNA-transfected cells, which is comparable to the nicotine-induced increase in proliferation in non-transfected cells. In cells transfected with  $\alpha$ 5 siRNA, nicotine failed to stimulate proliferation. When nicotine was absent, no statistically significant difference between ns-RNA- and  $\alpha$ 5-siRNA-transfected cells could be observed. This result indicates that in these cells, the presence of this subunit is not relevant for cellular proliferation in the absence of a nAChR agonist. Additionally, nicotine-induced proliferation is mediated via nAChRs, with the presence of  $\alpha$ 5 being indispensable.

In the previous sections, we showed that nAChRs mediate nicotine-induced migration. Subsequently, we tested if this effect is specific to  $\alpha$ 5-containing receptors. DU145 cells were transfected with  $\alpha$ 5-siRNA or ns-RNA, and after 24 h, migration was determined as described above. As shown in Figure 5F, 1  $\mu$ M nicotine promoted migration in ns-RNA-transfected cells at 24 h, an effect comparable to that observed in non-transfected cells (see above). When  $\alpha$ 5 was downregulated, nicotine failed to promote migration at 24 h. In line with our results, previous work by other researchers in the A549 lung cancer cell line could demonstrate that  $\alpha$ 5-containing receptors also seemed to exert a controlling effect upon nicotine-induced migration [34]. In summary, in MCF7, SW480, and DU145 cells, nicotine promoted proliferation and migration through the  $\alpha$ 5-subunit-containing nAChRs.

# 3.4. Downregulation of the $\alpha$ 5 nAChR Subunit Inhibits PDL1, CD47, and EMT Marker Expression Elicited by Nicotine

Previous reports showed that nicotine is able to induce the epithelial-mesenchymal transition (EMT) [20,70] and that nicotine stimulates the expression of the immunoregulatory proteins CD47 and PDL1 in lung bronchial and epithelial cells [26,29]. Consequently, we next explored whether knock down of the  $\alpha$ 5 subunit affects the nicotine-elicited expression of these immunoregulatory proteins and EMT markers. Here, we used 10  $\mu$ M of nicotine to ensure transcript expression of the analyzed RNAs, as it was previously shown that nicotine acts in a dose-dependent manner when analyzing the expression of these proteins [20]. For this test, cells were incubated with 10 µM nicotine for 48 h, and mRNA expression changes in epithelial (E-cadherin) and mesenchymal (vimentin) EMT markers as well as CD47 and PDL1 were analyzed via qPCR. We also used  $10 \,\mu M$  MEC to determine if nicotine-induced changes in mRNA expression levels are mediated via nAChRs. As a positive control of nicotine's effects, we used the lung cancer cell line A549. Figure 6A shows that in A549 cells, nicotine induced the mRNA expression of PDL-1, CD47, and vimentin and the downregulation of E-cadherin, clearly showing the EMT transition. In the MCF7 cell line (Figure 6B), nicotine induced EMT, as previously described [20], but, interestingly, it also upregulated CD47. In DU145 cells (Figure 6C), nicotine increased the mRNA expression of PDL1 but did not do so in SW480 cells (Figure 6D). Nicotine increased the mRNA expression of not only vimentin transcripts but also E-cadherin in SW480 and DU145 cells. When cells were stimulated with nicotine in the presence of MEC, the transcript levels of PDL1, CD47, and both EMT markers returned to their non-stimulated levels (Figure 6), showing that the effect was specific to nAChRs.

Next, we studied how the silencing of  $\alpha$ 5 affected the nicotine-induced mRNA expression of PDL1, CD47, and both EMT markers in these cell lines. Cells were transfected with  $\alpha$ 5-siRNA or ns-RNA, and after 24 h, cells were incubated, or not, with 10  $\mu$ M nicotine for 48 h before analyses. Figure 7 shows that upon downregulation of  $\alpha$ 5, nicotine did not stimulate the mRNA expression of PDL1, CD47, and EMT markers, indicating that  $\alpha$ 5 mediates nicotine-induced signaling in the immune regulatory and EMT pathways.

# 3.5. The $\alpha$ 5SNP Mutation D398N Increases Basal Proliferation and Migration in the DU145 Cell Line

The results above clearly demonstrate that  $\alpha$ 5-containing nicotinic receptors are essential for nicotine-induced effects. As previous reports showed that the D398N muta-

tion impairs  $\alpha 5'$ s function [40,71], we next investigated whether this mutation affects the nicotine-induced effects in proliferation and migration. In this set of experiments, DU145 cells were either transfected with the wild-type  $\alpha 5$  construct ( $\alpha 5$ .pcDNA3.1/V5,  $\alpha 5$ -WT) or with its mutated version  $\alpha 5D398N$  ( $\alpha 5D398N$ .pcDNA3.1/V5,  $\alpha 5$ -MUT). Both constructs had a V5 tag, and experiments were performed after 24 h. To verify transfection, qPCR and IF determinations were carried out. This cell line was chosen as it had a clear increase in proliferation and migration because of nAChRs activation by nicotine, as opposed to MCF7 and SW480 cells. As shown in Figure 8A, the transfection of  $\alpha 5$ -WT or  $\alpha 5$ -MUT increased the expression levels of this  $\alpha 5$  subunit in DU145 cells. Figure 8B shows transfection levels via anti-V5 antibodies. We also determined the expression of other nAChR subunits (Supplementary Figure S4) as a control. Here, no considerable changes in the expression of the other nAChR subunits were detected.



**Figure 6.** Effects of nicotine on the expression of EMT markers and immunosuppressive proteins in A549 (**A**), MCF7 (**B**), SW480 (**C**), and DU145 (**D**) cancer cell lines. Each figure reflects the expression of CD47, PDL1, VIM (vimentin), and ECAD (E-cadherin) at the mRNA level. The gene expression profile was investigated using qPCR analysis. Bar graphs show the relative expression  $\pm$  SEM of each subunit normalized to the corresponding expression in non-treated cells (NT; set to a value of 1; red dashed line. Meaningful changes of  $1 \pm 0.2$  is represented by a blue dashed line). Measurements were made using 3–5 independent experiments performed in triplicate. Cells were incubated with nicotine (10  $\mu$ M) for 48 h. If nicotine had an effect, cells were also incubated in the presence of the inhibitor MEC (10  $\mu$ M). Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \*  $p \le 0.05$ , \*\*  $p \le 0.001$ .



**Figure 7.** Knockdown of  $\alpha$ 5 renders cells nonresponsive to nicotine-induced upregulation of EMT markers and immunosuppressive proteins.  $\alpha$ 5-siRNA- or ns-RNA-transfected cells were treated with 10  $\mu$ M nicotine for 48 h. The gene expression profiles were investigated using qPCR analysis. Bar graphs show the relative expression  $\pm$  SEM of each subunit normalized to the corresponding expression in non-treated cells (NT; set to a value of 1; red dashed line. Meaningful changes of  $1 \pm 0.2$  is represented as a green dashed line). Measurements were made using 3–5 independent experiments performed in triplicate. The results obtained for SW480 (**A**), MCF7 (**B**), and DU145 cells (**C**) are shown. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \*  $p \leq 0.05$ .

Next, proliferation in the absence or presence of 1  $\mu$ M nicotine was studied in  $\alpha$ 5-WTor  $\alpha$ 5-MUT-transfected DU145 cells. In  $\alpha$ 5-WT-transfected cells, 1  $\mu$ M nicotine increased proliferation compared to non-treated cells (Figure 8C). In  $\alpha$ 5-MUT-transfected cells, nicotine failed to affect proliferation. Notably, basal proliferation without nicotine was per se higher in cells transfected with  $\alpha$ 5-MUT than in those transfected with  $\alpha$ 5-WT (Figure 8C). Migration in cells transfected with the  $\alpha$ 5-WT construct increased when cells were incubated with 1  $\mu$ M nicotine (Figure 8D). In cells expressing the mutated  $\alpha$ 5 subunit D398N, nicotine failed to alter migration. Moreover, basal migration was higher in cells transfected with  $\alpha$ 5-MUT than in those transfected with  $\alpha$ 5-WT (Figure 8D).

In lung basal epithelial cells expressing the D398N mutation in  $\alpha$ 5, basal proliferation is increased through a mechanism involving adenylyl cyclase [32]. Thus, we sought to determine if the increased basal proliferation in DU145 cells expressing  $\alpha$ 5-MUT is also due to altered adenylyl cyclase activation. In addition, the D398N mutation is localized in a section of the subunit that interacts with heteromeric G proteins [50]. Therefore, we also evaluated if activation of G proteins was involved within the pathway stimulating proliferation. Thus, DU145 cells were transfected with either  $\alpha$ 5-WT or  $\alpha$ 5-MUT, and proliferation was detected in the presence or absence of the adenylyl cyclase inhibitor SQ22536, a non-selective inhibitor of AC (SQ, 10  $\mu$ M), or the G-protein inhibitor pertussis toxin (PTX, 0.1  $\mu$ M). Our results show that, as described above, the presence of the mutation increased basal proliferation. This increase in basal proliferation was not affected by SQ. Therefore, we reason that AC is not involved in the mechanism (Figure 8E). When PTX was present, the proliferation levels of cells expressing  $\alpha$ 5-MUT were comparable to those of cells transfected with  $\alpha$ 5-WT. This result shows that the increased basal proliferation caused by the mutation present in  $\alpha$ 5 involved a G-protein-mediated mechanism.



**Figure 8.**  $\alpha$ 5SNP enhanced in vitro basal cell proliferation and migration in a prostate cancer cell model. Cells were transfected with either the wild-type  $\alpha$ 5 construct ( $\alpha$ 5-WT) or with its mutated version  $\alpha$ 5D398N ( $\alpha$ 5-MUT), and functional consequences were studied. (**A**) Depiction of the expression of the  $\alpha$ 5 subunit in non-transfected DU145 cells (NT; native expression) and in cells transfected with  $\alpha$ 5-WT (native plus foreign expression) or  $\alpha$ 5-MUT (native plus foreign expression); mRNA levels were determined via qPCR, and data show the mean  $\pm$  SEM of three independent cell culture experiments. (**B**) Immunofluorescence images showing the successful overexpression of  $\alpha$ 5-WT (left)

and  $\alpha$ 5-MUT (right) constructs in DU145 cells; the V5-tagged subunit was detected using incubation with the anti-V5 primary antibody followed by the appropriate Alexa Fluor 488 secondary antibody. Objective 100×, scale bar 40 µm. (**C**,**D**) Cell proliferation index and migration index (respectively) of DU145 cells transfected with  $\alpha$ 5-WT or  $\alpha$ 5-MUT-containing plasmids; cells were treated with 1 µM nicotine. The bar diagram shows data (mean  $\pm$  SEM). (**E**) Cell proliferation index of DU145 cells transfected with  $\alpha$ 5-WT or  $\alpha$ 5-MUT-containing plasmids. In the cells transfected with the latter, proliferation was analyzed in the presence or absence of SQ22536 (non-selective inhibitor of ACs, SQ, 10 µM) or PTX (G-protein inhibitor, pertussis toxin, 0.1 µM). The bar diagram shows the data (mean  $\pm$  SEM). (**F**) Cell proliferation index of non-transfected DU145 cells treated with 1 µM nicotine in the presence or absence of the inhibitors SQ22536 (SQ, 10 µM) or PTX (0.1 µM). Measurements were made using 3 independent experiments performed in triplicate. Statistical significance was analyzed using the Kruskal–Wallis test for non-parametric data with the GraphPad Prism (GraphPad 9.1.1 Software) software. A *p*-value of <0.05 was considered significant. \* *p* ≤ 0.05, \*\* *p* ≤ 0.001, \*\*\* *p* ≤ 0.0001.

Next, we evaluated whether SQ or PTX affected nicotine-induced proliferation in nontransfected DU145 cells (Figure 8F). As shown above, nicotine increased the proliferation of DU145 cells at 24 h and 48 h. The presence of the adenylyl cyclase inhibitor SQ did not affect nicotine-induced proliferation. On the other hand, when PTX was present, nicotine failed to induce proliferation. When given without nicotine, inhibitors did not modify the basal proliferation of DU145 cells (Supplementary Figure S1). Taken together, our results demonstrate that nicotine induces proliferation through specific activation of the  $\alpha$ 5 subunit in a G-protein-mediated mechanism.

### 4. Discussion

Ultimately,  $\alpha 5$  is expressed in many different types of tissues. However, its physiological and pathophysiological roles are not fully understood. Recent findings have associated this subunit with attention [41], food reward [72], addiction [73], and dopamine release mechanisms [74]. In addition,  $\alpha 5$  is associated with nicotine consumption [43] and lung cancer [75]. Elevated  $\alpha$ 5 protein and mRNA levels are present in different types of human cancers, such as lung, prostate, liver, breast, and gastric cancer [19,27,53,63,64,76,77], and are correlated with smoking status [25,66]. At a cellular level,  $\alpha$ 5 expression is thought to play a role in nicotine's regulation of proliferation, migration, and diverse pro-oncogenic signaling pathways in the A549 lung cancer cell line [76], while in other types of cancer cells, its function is not completely understood. In this study, we found that, upon knockdown of  $\alpha$ 5, nicotine failed to induce proliferation and migration in MCF7, DU145, and SW480 cells. This effect is thought to occur due to loss of the  $\alpha$ 5 subunit and not due to downregulation of the expression of other subunits, as we detected no change in other nAChR transcripts. Additionally, previous reports have demonstrated that in synaptosomes, neurons, and HEK-transfected cells, the presence of the  $\alpha$ 5 subunit does not affect the overall membrane expression of the  $\alpha 4\beta 2$  or  $\alpha 3\beta 4$  receptors [39,71,78,79]. This result suggests that nicotine-induced proliferation is reduced upon  $\alpha 5$  knockdown because of impaired receptor function and not because of the aberrant trafficking of receptors. However, further experiments are needed to evaluate receptor traffic in these conditions. Furthermore, no change in basal proliferation or migration was observed when this subunit was absent, suggesting that it is not involved, per se, in these cellular functions. Surprisingly, we did not find  $\alpha$ 7 to play a role in any nicotine-induced effects, as the specific inhibitor MLA did not change the nicotine-mediated increase in proliferation or migration in MCF7, SW480, and DU145 cells. This result is in contrast with previous observations in lung cancer cell lines such as A549, H1299, PC9, and H1975. In these cell lines,  $\alpha$ 7 plays an important role in nicotine-induced invasion, migration, proliferation, and EMT [21,23]. Furthermore, in human lung cancer patient samples, this subunit seems to be overexpressed and is associated with smoking levels [19].

Our study also shows that nicotine can induce changes in the mRNA expression of EMT markers and immune regulatory proteins in all cell lines tested, specifically through

 $\alpha$ 5. EMT is a vital step for the facilitation of a malignant and invasive phenotype in cancer, a process characterized by a decrease in epithelial proteins such as E-cadherin, as well as an increase in mesenchymal molecules such as fibronectin and vimentin [20]. Nicotine exposure was previously shown to induce EMT transition in lung and breast cancer cell lines [20,66]. In this study, we demonstrated the contribution of nAChRs in colon and prostate cancer cell EMT transition. Thus, our results are in line with previous observations in lung cancer patients, in which tobacco exposure favors EMT cellular transition [80]. In the human PCa cell line DU145 and the CRC cell line SW480, exposure to nicotine caused an increase in vimentin and the epithelial marker E-cadherin. Previous reports have shown that a loss of expression of the epithelial cell adhesion molecule E-cadherin promotes metastasis and is essential for EMT [81,82]. Other studies have reported that a decrease in this protein is not fully necessary to undergo EMT, nor, consequently, for the pro-oncogenic activity of cancer cells [83–86]. Nicotine exposure additionally helps maintain immune tolerance and restrain cancer autoimmunity by increasing the mRNA expression of negative immunoregulatory proteins such as PDL1 and CD47. In agreement with previous findings, nicotine increased both protein transcripts in A549 cells and was dependent on nAChR activation. In the MCF7 cell line, only CD47 was increased, and in the DU145 cell line, PDL1 was augmented. However, no effect was observed in the SW480 cell line. Previously, smoking was found to be correlated with increased levels of PDL-1 in lung cancer [25] and CD47 in brain metastases of lung cancer patients [26]. Accordingly, nicotine exposure increased CD47 levels in lung cancer cell lines [26] and PDL1 in lung cancer and melanoma cell lines [27,28]. Furthermore, nicotine-derived nitrosamine ketone (NNK) was shown to induce PDL1 expression in human bronchial epithelial cells [29]. When  $\alpha$ 5 was silenced in the MCF7, SW480, and DU145 cell lines, nicotine failed to regulate the mRNA expression of all markers evaluated. This result shows that nicotine controls the mRNA expression levels of these proteins via  $\alpha$ 5-containing nAChRs. Yet additional mechanistic studies are still needed to reveal whether activation of the  $\alpha 5$  affects the activity of other signaling pathways/proteins known to be influenced by the expression of such nicotinic receptor subtype, e.g., the vascular endothelial growth factor (VEGF) and epidermal growth factor receptor (EGFR) [33,87].

The non-synonymous single nucleotide polymorphism D398N of the  $\alpha$ 5 subunit, α5SNP, identified as a candidate mutation for smoking and lung cancer [32,38], was also studied for its contribution to nicotine's pro-oncogenic effects. We found that overexpression of  $\alpha$ 5SNP in the prostate cancer cell line DU145 caused an increase in basal proliferation, which was also shown for mouse lung epithelial cells [32]. In addition, overexpression of  $\alpha$ 5SNP increased migration. Surprisingly, when this mutation was present, nicotine failed to induce its pro-proliferative and pro-migratory effects. This result is aligned with previous observations showing that the presence of this D398N mutation in  $\alpha 3\beta 4\alpha 5$  nAChRs of HEK293-transfected cells reduced the relative maximum Ca<sup>2+</sup> response to various nAChR agonists [71]. Moreover, the presence of the D398N mutation in mouse lung epithelial cells increased basal proliferation via a mechanism involving adenylyl cyclase activation [32]. Whilst the nAChRs are very well known to possess ionotropic properties, these receptors also induce metabotropic signaling events [13] and are known to regulate adenylyl cyclase activity [88–90]. In the present work, we found that the observed increase in proliferation in DU145 cells expressing the D398N mutation was dependent upon a G-protein-mediated mechanism. Here, when DU145 cells carrying the mutation were in the presence of pertussis toxin, cellular proliferation was halted, whereas incubation with an adenylyl cyclase inhibitor did not intervene with the proliferative activity. The D398N mutation is localized in a section of the subunit that interacts with heteromeric G proteins; more specifically, the intracellular loop region of the  $\alpha$ 5 subunit interacts with the  $G_{\alpha 0}$  and  $G_{\beta \gamma}$  dimers [50]. Interestingly, the amino acid sequence surrounding this mutation was found to be very conserved among species [38], which possibly indicates the presence of an important structural function. Previous findings suggested that the D398N mutation renders the  $\alpha 3\beta 4$  receptor more efficient in promoting Ca<sup>2+</sup> release from

intracellular stores [71] and may also play a role in preventing receptor desensitization, thereby maintaining nicotine signaling [40]. In this regard, in neurons and microglial cell lines,  $\alpha$ 7-mediated Ca<sup>2+</sup> signals are, in part, mediated by a G-protein-mediated mechanism, ultimately leading to the release of Ca<sup>2+</sup> from intracellular stores [91,92]. Furthermore, we found that in non-transfected DU145 cells, nicotine increased proliferation through activation of the G-protein pathway via  $\alpha$ 5. Conversely, a previous report showed that activation of the  $\alpha$ 4 $\beta$ 2 receptors in T-cells led to inhibition of G-protein-mediated signaling, resulting in an increase in proliferation [93]. This result suggests that the contribution of nAChRs in G-protein-mediated signaling depends on cell type and subunit expression patterns. Future studies could provide a more in-depth analysis of the role of nAChRs in G-protein-mediated signaling in this context. Though previous reports have shown that activation of nAChRs has a dual ionotropic-metabotropic functional component, independent of each other [94–97], further experiments would have to be conducted to determine the involvement of Ca<sup>2+</sup> permeability or membrane depolarization in our present model.

Whether the endogenous  $\alpha$ 5 subunit is expressed on intracellular compartments in addition to the plasma membrane in the cell lines studied has not been addressed. In addition, despite the fact that one report has shown its expression in mitochondria [98], there are no data supporting the idea that it has a functional role in that organelle. In contrast, the  $\alpha$ 7 subunit expressed in mitochondria has been shown to be responsible for regulating cytochrome C release and activating additional signaling pathways important for cell survival [99,100].

Our results point to  $\alpha$ 5-containing receptors as key signaling nodes of nicotine's pro-oncogenic effects in diverse cancer types. This discovery is novel since, to date, no other reports have fully determined the importance of  $\alpha$ 5 in mediating nicotine's oncogenic functions at the cellular level in breast, colon, or prostate cancer cells. Our findings show the need for  $\alpha$ 5 inhibitors/modulators to further investigate  $\alpha$ 5's role in mediating nicotine's pro-oncogenic effects in cancer and as a putative drug target. Yet further studies will have to be carried out in order to determine whether the expression of the  $\alpha$ 5SNP can be used in the future as a tumor-agnostic marker.

#### 5. Conclusions

Based on our findings, we show here that the  $\alpha$ 5 subunit is the key mediator of nicotine's pro-oncogenic effects, whereas the silencing of the subunit by siRNA rendered the cells studied irresponsive to this stimulus. The signaling events mediated by this subunit are carried through a G- protein mediated mechanism. Most remarkably, the introduction of the  $\alpha$ 5SNP rendered the cells (i) irresponsive to nicotine, an effect reminiscent of the one observed when the mutation is present in neurons or heterologously expressed receptors in HEK cells [39–42], and (ii) with an augmented basal proliferation rate, because of increased G-protein activation. This hints that further studies are needed to clarify the role of  $\alpha$ 5-containing receptors and its SNP in cancer in general, and more specifically the signaling events leading to increased G-protein activation.

**Supplementary Materials:** The following supporting information can be downloaded at: https: //www.mdpi.com/article/10.3390/cells12152000/s1, Figure S1: Cell proliferation index of SW480 and DU145 human cancer cell lines in the presence of diverse inhibitors; Figure S2: Cell migration index in SW480 and MCF7cells; Figure S3: Silencing of  $\alpha$ 5 in the human cancer cell lines (A) MCF7, (B) SW480, and (C) DU145 does not affect expression of other nAChR subunits at the mRNA level; Figure S4: Overexpression of the WT or MUT version of  $\alpha$ 5 nAChR subunit does not affect the expression of other nAChR subunits at the mRNA level.

Author Contributions: Conceptualization, M.C.M.; methodology, M.C.M., I.P., D.R.-K. and F.B.; formal analysis, M.C.M., I.P., D.R.-K. and F.B.; investigation, M.C.M.; resources, M.C.M.; writing—original draft preparation, M.C.M.; writing—review and editing, M.C.M. and C.P.; visualization, M.C.M.; supervision, M.C.M. and C.P.; project administration, M.C.M.; funding acquisition, M.C.M. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the H2020-MSCA-IF project nAChRs-CRC-PCa-IF-RI, No. 101027428 (to M.C.M.) and by the Swiss National Science Foundation (Schweizerischer National-fonds zur Förderung der Wissenschaftlichen Forschung), grant number 310030\_200380/1 (to C.P.).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

**Data Availability Statement:** The data used and analyzed during the current study are available from the corresponding author upon reasonable request.

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

- Millar, N.S.; Gotti, C. Diversity of vertebrate nicotinic acetylcholine receptors. *Neuropharmacology* 2009, 56, 237–246. [CrossRef] [PubMed]
- Le Novère, N.; Changeux, J.P. Molecular evolution of the nicotinic acetylcholine receptor: An example of multigene family in excitable cells. J. Mol. Evol. 1995, 40, 155–172. [CrossRef]
- 3. Changeux, J.P. Nicotinic receptors and nicotine addiction. Comptes Rendus Biol. 2009, 332, 421–425. [CrossRef] [PubMed]
- Corradi, J.; Bouzat, C. Understanding the Bases of Function and Modulation of α7 Nicotinic Receptors: Implications for Drug Discovery. *Mol. Pharmacol.* 2016, 90, 288–299. [CrossRef]
- 5. Wongsamitkul, N.; Maldifassi, M.C.; Simeone, X.; Baur, R.; Ernst, M.; Sigel, E. α subunits in GABAA receptors are dispensable for GABA and diazepam action. *Sci. Rep.* **2017**, *7*, 15498. [CrossRef]
- 6. Maldifassi, M.C.; Baur, R.; Pierce, D.; Nourmahnad, A.; Forman, S.A.; Sigel, E. Novel positive allosteric modulators of GABAA receptors with anesthetic activity. *Sci. Rep.* 2016, *6*, 25943. [CrossRef] [PubMed]
- Thany, S.H.; Tricoire-Leignel, H. Emerging Pharmacological Properties of Cholinergic Synaptic Transmission: Comparison between Mammalian and Insect Synaptic and Extrasynaptic Nicotinic Receptors. *Curr. Neuropharmacol.* 2011, 9, 706–714. [CrossRef] [PubMed]
- Nemecz, Á.; Prevost, M.S.; Menny, A.; Corringer, P.J. Emerging Molecular Mechanisms of Signal Transduction in Pentameric Ligand-Gated Ion Channels. *Neuron* 2016, 90, 452–470. [CrossRef]
- Zoli, M.; Pucci, S.; Vilella, A.; Gotti, C. Neuronal and Extraneuronal Nicotinic Acetylcholine Receptors. *Curr. Neuropharmacol.* 2018, 16, 338–349. [CrossRef]
- Dineley, K.T.; Pandya, A.A.; Yakel, J.L. Nicotinic ACh receptors as therapeutic targets in CNS disorders. *Trends Pharmacol. Sci.* 2015, *36*, 96–108. [CrossRef]
- 11. Arias, H.R. Positive and negative modulation of nicotinic receptors. Adv. Protein Chem. Struct. Biol. 2010, 80, 153–203. [CrossRef]
- 12. Wessler, I.; Kirkpatrick, C.J. Acetylcholine beyond neurons: The non-neuronal cholinergic system in humans. *Br. J. Pharmacol.* **2008**, *154*, 1558–1571. [CrossRef] [PubMed]
- 13. Kabbani, N.; Nichols, R.A. Beyond the Channel: Metabotropic Signaling by Nicotinic Receptors. *Trends Pharmacol. Sci.* **2018**, *39*, 354–366. [CrossRef] [PubMed]
- 14. Rosas-Ballina, M.; Tracey, K.J. The neurology of the immune system: Neural reflexes regulate immunity. *Neuron* **2009**, *64*, 28–32. [CrossRef]
- Maldifassi, M.C.; Atienza, G.; Arnalich, F.; López-Collazo, E.; Cedillo, J.L.; Martín-Sánchez, C.; Bordas, A.; Renart, J.; Montiel, C. A new IRAK-M-mediated mechanism implicated in the anti-inflammatory effect of nicotine via α7 nicotinic receptors in human macrophages. *PLoS ONE* 2014, *9*, e108397. [CrossRef] [PubMed]
- 16. Corringer, P.J.; Le Novère, N.; Changeux, J.P. Nicotinic receptors at the amino acid level. *Annu. Rev. Pharmacol. Toxicol.* **2000**, *40*, 431–458. [CrossRef]
- 17. Grutter, T.; Le Novère, N.; Changeux, J.P. Rational understanding of nicotinic receptors drug binding. *Curr. Top Med. Chem.* **2004**, 4, 645–650. [CrossRef]
- Schaal, C.; Chellappan, S.P. Nicotine-mediated cell proliferation and tumor progression in smoking-related cancers. *Mol. Cancer Res.* 2014, 12, 14–23. [CrossRef]
- Bordas, A.; Cedillo, J.L.; Arnalich, F.; Esteban-Rodriguez, I.; Guerra-Pastrián, L.; de Castro, J.; Martín-Sánchez, C.; Atienza, G.; Fernández-Capitan, C.; Rios, J.J.; et al. Expression patterns for nicotinic acetylcholine receptor subunit genes in smoking-related lung cancers. *Oncotarget* 2017, *8*, 67878–67890. [CrossRef]
- Dasgupta, P.; Rizwani, W.; Pillai, S.; Kinkade, R.; Kovacs, M.; Rastogi, S.; Banerjee, S.; Carless, M.; Kim, E.; Coppola, D.; et al. Nicotine induces cell proliferation, invasion and epithelial-mesenchymal transition in a variety of human cancer cell lines. *Int. J. Cancer* 2009, 124, 36–45. [CrossRef]
- Mucchietto, V.; Fasoli, F.; Pucci, S.; Moretti, M.; Benfante, R.; Maroli, A.; Di Lascio, S.; Bolchi, C.; Pallavicini, M.; Dowell, C.; et al. α9- and α7-containing receptors mediate the pro-proliferative effects of nicotine in the A549 adenocarcinoma cell line. *Br. J. Pharmacol.* 2018, 175, 1957–1972. [CrossRef] [PubMed]

- 22. West, K.A.; Brognard, J.; Clark, A.S.; Linnoila, I.R.; Yang, X.; Swain, S.M.; Harris, C.; Belinsky, S.; Dennis, P.A. Rapid Akt activation by nicotine and a tobacco carcinogen modulates the phenotype of normal human airway epithelial cells. *J. Clin. Investig.* 2003, *111*, 81–90. [CrossRef] [PubMed]
- Zhang, C.; Ding, X.P.; Zhao, Q.N.; Yang, X.J.; An, S.M.; Wang, H.; Xu, L.; Zhu, L.; Chen, H.Z. Role of α7-nicotinic acetylcholine receptor in nicotine-induced invasion and epithelial-to-mesenchymal transition in human non-small cell lung cancer cells. *Oncotarget* 2016, 7, 59199–59208. [CrossRef]
- 24. Lian, S.; Xie, R.; Ye, Y.; Lu, Y.; Cheng, Y.; Xie, X.; Li, S.; Jia, L. Dual blockage of both PD-L1 and CD47 enhances immunotherapy against circulating tumor cells. *Sci. Rep.* **2019**, *9*, 4532. [CrossRef]
- Zhang, Y.; Jia, Y.; Li, P.; Li, H.; Xiao, D.; Wang, Y.; Ma, X. Reciprocal activation of α5-nAChR and STAT3 in nicotine-induced human lung cancer cell proliferation. *J. Genet. Genom.* 2017, 44, 355–362. [CrossRef]
- Wu, S.Y.; Xing, F.; Sharma, S.; Wu, K.; Tyagi, A.; Liu, Y.; Zhao, D.; Deshpande, R.P.; Shiozawa, Y.; Ahmed, T.; et al. Nicotine promotes brain metastasis by polarizing microglia and suppressing innate immune function. *J. Exp. Med.* 2020, 217, e20191131. [CrossRef]
- Zhu, P.; Jin, Z.; Kang, G.; Jia, Y.; Liu, D.; Zhang, Q.; Guo, F.; Jia, Y.; Jiao, Y.; Li, J.; et al. Alpha5 nicotinic acetylcholine receptor mediated immune escape of lung adenocarcinoma via STAT3/Jab1-PD-L1 signalling. *Cell Commun. Signal.* 2022, 20, 121. [CrossRef]
- Nguyen, H.D.; Liao, Y.C.; Ho, Y.S.; Chen, L.C.; Chang, H.W.; Cheng, T.C.; Liu, D.; Lee, W.R.; Shen, S.C.; Wu, C.H.; et al. The α9 Nicotinic Acetylcholine Receptor Mediates Nicotine-Induced PD-L1 Expression and Regulates Melanoma Cell Proliferation and Migration. *Cancers* 2019, *11*, 1991. [CrossRef]
- Kwok, H.H.; Gao, B.; Chan, K.H.; Ip, M.S.; Minna, J.D.; Lam, D.C. Nicotinic Acetylcholine Receptor Subunit α7 Mediates Cigarette Smoke-Induced PD-L1 Expression in Human Bronchial Epithelial Cells. *Cancers* 2021, 13, 5345. [CrossRef] [PubMed]
- Amos, C.I.; Wu, X.; Broderick, P.; Gorlov, I.P.; Gu, J.; Eisen, T.; Dong, Q.; Zhang, Q.; Gu, X.; Vijayakrishnan, J.; et al. Genome-wide association scan of tag SNPs identifies a susceptibility locus for lung cancer at *15q25.1*. *Nat. Genet.* 2008, *40*, 616–622. [CrossRef] [PubMed]
- Hung, R.J.; McKay, J.D.; Gaborieau, V.; Boffetta, P.; Hashibe, M.; Zaridze, D.; Mukeria, A.; Szeszenia-Dabrowska, N.; Lissowska, J.; Rudnai, P.; et al. A susceptibility locus for lung cancer maps to nicotinic acetylcholine receptor subunit genes on 15q25. Nature 2008, 452, 633–637. [CrossRef]
- Routhier, J.; Pons, S.; Freidja, M.L.; Dalstein, V.; Cutrona, J.; Jonquet, A.; Lalun, N.; Mérol, J.C.; Lathrop, M.; Stitzel, J.A.; et al. An innate contribution of human nicotinic receptor polymorphisms to COPD-like lesions. *Nat. Commun.* 2021, 12, 6384. [CrossRef] [PubMed]
- Ma, X.; Jia, Y.; Zu, S.; Li, R.; Jia, Y.; Zhao, Y.; Xiao, D.; Dang, N.; Wang, Y. α5 Nicotinic acetylcholine receptor mediates nicotineinduced HIF-1α and VEGF expression in non-small cell lung cancer. *Toxicol. Appl. Pharmacol.* 2014, 278, 172–179. [CrossRef] [PubMed]
- Sun, H.; Ma, X. α5-nAChR modulates nicotine-induced cell migration and invasion in A549 lung cancer cells. *Exp. Toxicol. Pathol.* 2015, 67, 477–482. [CrossRef] [PubMed]
- Chen, X.; Jia, Y.; Zhang, Y.; Zhou, D.; Sun, H.; Ma, X. α5-nAChR contributes to epithelial-mesenchymal transition and metastasis by regulating Jab1/Csn5 signalling in lung cancer. J. Cell Mol. Med. 2020, 24, 2497–2506. [CrossRef]
- 36. Wang, F.; Gerzanich, V.; Wells, G.B.; Anand, R.; Peng, X.; Keyser, K.; Lindstrom, J. Assembly of human neuronal nicotinic receptor alpha5 subunits with alpha3, beta2, and beta4 subunits. *J. Biol. Chem.* **1996**, 271, 17656–17665. [CrossRef]
- Le Novère, N.; Corringer, P.J.; Changeux, J.P. The diversity of subunit composition in nAChRs: Evolutionary origins, physiologic and pharmacologic consequences. J. Neurobiol. 2002, 53, 447–456. [CrossRef]
- 38. Maskos, U. The nicotinic receptor alpha5 coding polymorphism rs16969968 as a major target in disease: Functional dissection and remaining challenges. *J. Neurochem.* 2020, 154, 241–250. [CrossRef]
- Scholze, P.; Huck, S. The α5 Nicotinic Acetylcholine Receptor Subunit Differentially Modulates α4β2\* and α3β4\* Receptors. Front. Synaptic. Neurosci. 2020, 12, 607959. [CrossRef]
- Ray, C.; Soderblom, E.J.; Bai, Y.; Carroll, F.I.; Caron, M.G.; Barak, L.S. Probing the Allosteric Role of α5 Subunit of α3β4α5 Nicotinic Acetylcholine Receptors by Functionally Selective Modulators and Ligands. ACS Chem. Biol. 2017, 12, 702–714. [CrossRef]
- 41. Bailey, C.D.; Alves, N.C.; Nashmi, R.; De Biasi, M.; Lambe, E.K. Nicotinic α5 subunits drive developmental changes in the activation and morphology of prefrontal cortex layer VI neurons. *Biol. Psychiatry* **2012**, *71*, 120–128. [CrossRef] [PubMed]
- 42. Gotti, C.; Zoli, M.; Clementi, F. Brain nicotinic acetylcholine receptors: Native subtypes and their relevance. *Trends Pharmacol. Sci.* **2006**, *27*, 482–491. [CrossRef]
- 43. Fowler, C.D.; Lu, Q.; Johnson, P.M.; Marks, M.J.; Kenny, P.J. Habenular α5 nicotinic receptor subunit signalling controls nicotine intake. *Nature* **2011**, *471*, 597–601. [CrossRef]
- 44. Besson, M.; Guiducci, S.; Granon, S.; Guilloux, J.P.; Guiard, B.; Repérant, C.; Faure, P.; Pons, S.; Cannazza, G.; Zoli, M.; et al. Alterations in alpha5\* nicotinic acetylcholine receptors result in midbrain- and hippocampus-dependent behavioural and neural impairments. *Psychopharmacology* **2016**, *233*, 3297–3314. [CrossRef] [PubMed]
- Wang, J.C.; Cruchaga, C.; Saccone, N.L.; Bertelsen, S.; Liu, P.; Budde, J.P.; Duan, W.; Fox, L.; Grucza, R.A.; Kern, J.; et al. Risk for nicotine dependence and lung cancer is conferred by mRNA expression levels and amino acid change in *CHRNA5. Hum. Mol. Genet.* 2009, *18*, 3125–3135. [CrossRef] [PubMed]

- 46. Young, R.P.; Hopkins, R.J.; Hay, B.A.; Epton, M.J.; Black, P.N.; Gamble, G.D. Lung cancer gene associated with COPD: Triple whammy or possible confounding effect? *Eur. Respir. J.* **2008**, *32*, 1158–1164. [CrossRef]
- Saccone, N.L.; Culverhouse, R.C.; Schwantes-An, T.H.; Cannon, D.S.; Chen, X.; Cichon, S.; Giegling, I.; Han, S.; Han, Y.; Keskitalo-Vuokko, K.; et al. Multiple independent loci at chromosome 15q25.1 affect smoking quantity: A meta-analysis and comparison with lung cancer and COPD. *PLoS Genet.* 2010, 6, e1001053. [CrossRef] [PubMed]
- Pillai, S.G.; Ge, D.; Zhu, G.; Kong, X.; Shianna, K.V.; Need, A.C.; Feng, S.; Hersh, C.P.; Bakke, P.; Gulsvik, A.; et al. A genome-wide association study in chronic obstructive pulmonary disease (COPD): Identification of two major susceptibility loci. *PLoS Genet.* 2009, 5, e1000421. [CrossRef]
- Wilk, J.B.; Shrine, N.R.; Loehr, L.R.; Loehr, L.R.; Zhao, J.H.; Manichaikul, A.; Lopez, L.M.; Smith, A.V.; Heckbert, S.R.; Smolonska, J.; et al. Genome-wide association studies identify CHRNA5/3 and HTR4 in the development of airflow obstruction. Am. J. Respir. Crit. Care Med. 2012, 186, 622–632. [CrossRef]
- 50. Fischer, H.; Liu, D.M.; Lee, A.; Harries, J.C.; Adams, D.J. Selective modulation of neuronal nicotinic acetylcholine receptor channel subunits by Go-protein subunits. *J. Neurosci.* 2005, 25, 3571–3577. [CrossRef]
- 51. De Nunzio, C.; Andriole, G.L.; Thompson, I.M., Jr.; Freedland, S.J. Smoking and Prostate Cancer: A Systematic Review. *Eur. Urol. Focus.* **2015**, *1*, 28–38. [CrossRef] [PubMed]
- 52. Huncharek, M.; Haddock, K.S.; Reid, R.; Kupelnick, B. Smoking as a risk factor for prostate cancer: A meta-analysis of 24 prospective cohort studies. *Am. J. Public Health* **2010**, *100*, 693–701. [CrossRef] [PubMed]
- 53. Qi, J.C.; Xue, W.Y.; Zhang, Y.P.; Qu, C.B.; Lu, B.S.; Yin, Y.W.; Liu, K.L.; Wang, D.B.; Li, W.; Zhao, Z.M. Cholinergic α5 nicotinic receptor is involved in the proliferation and invasion of human prostate cancer cells. *Oncol. Rep.* **2020**, *43*, 159–168. [CrossRef]
- 54. Schuller, H.M. Is cancer triggered by altered signalling of nicotinic acetylcholine receptors? *Nat. Rev. Cancer* 2009, *9*, 195–205. [CrossRef]
- Tyagi, A.; Sharma, S.; Wu, K.; Xing, F.; Liu, Y.; Zhao, D.; Deshpande, R.P.; D'Agostino, R.B., Jr.; Watabe, K. Nicotine promotes breast cancer metastasis by stimulating N2 neutrophils and generating pre-metastatic niche in lung. *Nat. Commun.* 2021, 12, 474. [CrossRef]
- 56. Wei, P.L.; Kuo, L.J.; Huang, M.T.; Ting, W.C.; Ho, Y.S.; Wang, W.; An, J.; Chang, Y.J. Nicotine enhances colon cancer cell migration by induction of fibronectin. *Ann. Surg. Oncol.* **2011**, *18*, 1782–1790. [CrossRef]
- 57. Wong, H.P.; Yu, L.; Lam, E.K.; Tai, E.K.; Wu, W.K.; Cho, C.H. Nicotine promotes cell proliferation via alpha7-nicotinic acetylcholine receptor and catecholamine-synthesizing enzymes-mediated pathway in human colon adenocarcinoma HT-29 cells. *Toxicol. Appl. Pharmacol.* **2007**, *221*, 261–267. [CrossRef]
- Borgström, A.; Hauert, B.; Kappel, S.; Zoni, E.; Kiener, M.; Stokłosa, P.; Baur, R.; Spahn, M.; Kruithof-de Julio, M.; Peinelt, C. Small Molecular Inhibitors Block TRPM4 Currents in Prostate Cancer Cells, with Limited Impact on Cancer Hallmark Functions. *J. Mol. Biol.* 2021, 433, 166665. [CrossRef] [PubMed]
- Stokłosa, P.; Borgström, A.; Hauert, B.; Baur, R.; Peinelt, C. Investigation of Novel Small Molecular TRPM4 Inhibitors in Colorectal Cancer Cells. *Cancers* 2021, 13, 5400. [CrossRef] [PubMed]
- 60. Human Protein Atlas. Available online: https://www.proteinatlas.org (accessed on 1 May 2023).
- Cancer Genom Atlas (TCGA). Available online: https://www.cancer.gov/about-nci/organiza-tion/ccg/research/structuralgenomics/tcga (accessed on 1 May 2023).
- 62. Siegel, R.L.; Miller, K.D.; Wagle, N.S.; Jemal, A. Cancer statistics, 2023. CA Cancer J. Clin. 2023, 73, 17–48. [CrossRef] [PubMed]
- 63. Fu, Y.; Ci, H.; Du, W.; Dong, Q.; Jia, H. *CHRNA5* Contributes to Hepatocellular Carcinoma Progression by Regulating YAP Activity. *Pharmaceutics* **2022**, *14*, 275. [CrossRef] [PubMed]
- 64. Jia, Y.; Sun, H.; Wu, H.; Zhang, H.; Zhang, X.; Xiao, D.; Ma, X.; Wang, Y. Nicotine Inhibits Cisplatin-Induced Apoptosis via Regulating α5-nAChR/AKT Signaling in Human Gastric Cancer Cells. *PLoS ONE* **2016**, *11*, e0149120. [CrossRef] [PubMed]
- 65. Pucci, S.; Fasoli, F.; Moretti, M.; Benfante, R.; Di Lascio, S.; Viani, P.; Daga, A.; Gordon, T.J.; McIntosh, M.; Zoli, M.; et al. Choline and nicotine increase glioblastoma cell proliferation by binding and activating α7- and α9- containing nicotinic receptors. *Pharmacol. Res.* **2021**, 163, 105336. [CrossRef] [PubMed]
- 66. Cedillo, J.L.; Bordas, A.; Arnalich, F.; Esteban-Rodríguez, I.; Martín-Sánchez, C.; Extremera, M.; Atienza, G.; Rios, J.J.; Arribas, R.L.; Montiel, C. Anti-tumoral activity of the human-specific duplicated form of α7-nicotinic receptor subunit in tobacco-induced lung cancer progression. *Lung Cancer* 2019, *128*, 134–144. [CrossRef]
- 67. Maldifassi, M.C.; Momboisse, F.; Guerra, M.J.; Vielma, A.H.; Maripillán, J.; Báez-Matus, X.; Flores-Muñoz, C.; Cádiz, B.; Schmachtenberg, O.; Martínez, A.D.; et al. The interplay between α7 nicotinic acetylcholine receptors, pannexin-1 channels and P2X7 receptors elicit exocytosis in chromaffin cells. *J. Neurochem.* **2021**, 157, 1789–1808. [CrossRef]
- Kappel, S.; Stokłosa, P.; Hauert, B.; Ross-Kaschitza, D.; Borgström, A.; Baur, R.; Galván, J.A.; Zlobec, I.; Peinelt, C. TRPM4 is highly expressed in human colorectal tumor buds and contributes to proliferation, cell cycle, and invasion of colorectal cancer cells. *Mol. Oncol.* 2019, *13*, 2393–2405. [CrossRef] [PubMed]
- Albuquerque, E.X.; Pereira, E.F.; Alkondon, M.; Rogers, S.W. Mammalian nicotinic acetylcholine receptors: From structure to function. *Physiol. Rev.* 2009, *89*, 73–120. [CrossRef] [PubMed]
- Dinicola, S.; Masiello, M.G.; Proietti, S.; Coluccia, P.; Fabrizi, G.; Catizone, A.; Ricci, G.; de Toma, G.; Bizzarri, M.; Cucina, A. Nicotine increases colon cancer cell migration and invasion through epithelial to mesenchymal transition (EMT): COX-2 involvement. *J. Cell Physiol.* 2018, 233, 4935–4948. [CrossRef]

- Tammimäki, A.; Herder, P.; Li, P.; Esch, C.; Laughlin, J.R.; Akk, G.; Stitzel, J.A. Impact of human D398N single nucleotide polymorphism on intracellular calcium response mediated by α3β4α5 nicotinic acetylcholine receptors. *Neuropharmacology* 2012, 63, 1002–1011. [CrossRef]
- 72. Breum, A.W.; Falk, S.; Svendsen, C.S.A.; Nicolaisen, T.S.; Mathiesen, C.V.; Maskos, U.; Clemmensen, C. Divergent Roles of α5 and β4 Nicotinic Receptor Subunits in Food Reward and Nicotine-induced Weight Loss in Male Mice. *Endocrinology* 2022, 163, bqac079. [CrossRef]
- 73. Forget, B.; Scholze, P.; Langa, F.; Morel, C.; Pons, S.; Mondoloni, S.; Besson, M.; Durand-de Cuttoli, R.; Hay, A.; Tricoire, L.; et al. A Human Polymorphism in *CHRNA5* Is Linked to Relapse to Nicotine Seeking in Transgenic Rats. *Curr. Biol.* 2018, 28, 3244–3253.e7. [CrossRef] [PubMed]
- 74. Exley, R.; McIntosh, J.M.; Marks, M.J.; Maskos, U.; Cragg, S.J. Striatal α5 nicotinic receptor subunit regulates dopamine transmission in dorsal striatum. *J. Neurosci.* **2012**, *32*, 2352–2356. [CrossRef] [PubMed]
- 75. Improgo, M.R.; Scofield, M.D.; Tapper, A.R.; Gardner, P.D. The nicotinic acetylcholine receptor *CHRNA5/A3/B4* gene cluster: Dual role in nicotine addiction and lung cancer. *Prog. Neurobiol.* **2010**, *92*, 212–226. [CrossRef] [PubMed]
- Hollenhorst, M.I.; Krasteva-Christ, G. Nicotinic Acetylcholine Receptors in the Respiratory Tract. *Molecules* 2021, 26, 6097. [CrossRef] [PubMed]
- 77. Shehwana, H.; Keskus, A.G.; Ozdemir, S.E.; Acikgöz, A.A.; Biyik-Sit, R.; Cagnan, I.; Gunes, D.; Jahja, E.; Cingir-Koker, S.; Olmezer, G.; et al. *CHRNA5* belongs to the secondary estrogen signalling network exhibiting prognostic significance in breast cancer. *Cell Oncol.* 2021, 44, 453–472. [CrossRef]
- Jackson, K.J.; Marks, M.J.; Vann, R.E.; Chen, X.; Gamage, T.F.; Warner, J.A.; Damaj, M.I. Role of alpha5 nicotinic acetylcholine receptors in pharmacological and behavioural effects of nicotine in mice. J. Pharmacol. Exp. Ther. 2010, 334, 137–146. [CrossRef]
- 79. Brown, R.W.; Collins, A.C.; Lindstrom, J.M.; Whiteaker, P. Nicotinic alpha5 subunit deletion locally reduces high-affinity agonist activation without altering nicotinic receptor numbers. *J. Neurochem.* **2007**, *103*, 204–215. [CrossRef]
- 80. Strathdee, G. Epigenetic versus genetic alterations in the inactivation of E-cadherin. *Semin. Cancer Biol.* **2002**, *12*, 373–379. [CrossRef]
- Bracke, M.E.; Van Roy, F.M.; Mareel, M.M. The E-cadherin/catenin complex in invasion and metastasis. *Curr. Top Microbiol. Immunol.* 1996, 213, 123–161. [CrossRef]
- 82. Kang, Y.; Massagué, J. Epithelial-mesenchymal transitions: Twist in development and metastasis. *Cell* **2004**, *118*, 277–279. [CrossRef]
- 83. Bukholm, I.K.; Nesland, J.M.; Børresen-Dale, A.L. Re-expression of E-cadherin, alpha-catenin and beta-catenin, but not of gamma-catenin, in metastatic tissue from breast cancer patients. *J. Pathol.* **2000**, *190*, 15–19. [CrossRef]
- Lou, Y.; Preobrazhenska, O.; auf dem Keller, U.; Sutcliffe, M.; Barclay, L.; McDonald, P.C.; Roskelley, C.; Overall, C.M.; Dedhar, S. Epithelial-mesenchymal transition (EMT) is not sufficient for spontaneous murine breast cancer metastasis. *Dev. Dyn.* 2008, 237, 2755–2768. [CrossRef] [PubMed]
- 85. Hollestelle, A.; Peeters, J.K.; Smid, M.; Timmermans, M.; Verhoog, L.C.; Westenend, P.J.; Heine, A.A.; Chan, A.; Sieuwerts, A.M.; Wiemer, E.A.; et al. Loss of E-cadherin is not a necessity for epithelial to mesenchymal transition in human breast cancer. *Breast Cancer Res. Treat.* **2013**, *138*, 47–57. [CrossRef] [PubMed]
- Querzoli, P.; Coradini, D.; Pedriali, M.; Boracchi, P.; Ambrogi, F.; Raimondi, E.; La Sorda, R.; Lattanzio, R.; Rinaldi, R.; Lunardi, M.; et al. An immunohistochemically positive E-cadherin status is not always predictive for a good prognosis in human breast cancer. *Br. J. Cancer* 2010, *103*, 1835–1839. [CrossRef]
- Wang, M.L.; Hsu, Y.F.; Liu, C.H.; Kuo, C.H.; Chen, Y.C.; Yeh, Y.C.; Ho, H.L.; Wu, Y.C.; Chou, T.Y.; Wu, C.W. Low-Dose Nicotine Activates EGFR Signaling via α5-nAChR and Promotes Lung Adenocarcinoma Progression. *Int. J. Mol. Sci.* 2020, 21, 6829. [CrossRef]
- 88. Oshikawa, J.; Toya, Y.; Fujita, T.; Egawa, M.; Kawabe, J.; Umemura, S.; Ishikawa, Y. Nicotinic acetylcholine receptor alpha 7 regulates cAMP signal within lipid rafts. *Am. J. Physiol. Cell Physiol.* **2003**, *285*, C567–C574. [CrossRef]
- Cheng, Q.; Yakel, J.L. Activation of α7 nicotinic acetylcholine receptors increases intracellular cAMP levels via activation of AC1 in hippocampal neurons. *Neuropharmacology* 2015, 95, 405–414. [CrossRef]
- Zhu, S.; Huang, S.; Xia, G.; Wu, J.; Shen, Y.; Wang, Y.; Ostrom, R.S.; Du, A.; Shen, C.; Xu, C. Anti-inflammatory effects of α7-nicotinic ACh receptors are exerted through interactions with adenylyl cyclase-6. *Br. J. Pharmacol.* 2021, 178, 2324–2338. [CrossRef]
- King, J.R.; Nordman, J.C.; Bridges, S.P.; Lin, M.K.; Kabbani, N. Identification and Characterization of a G Protein-binding Cluster in α7 Nicotinic Acetylcholine Receptors. J. Biol. Chem. 2015, 290, 20060–20070. [CrossRef]
- King, J.R.; Gillevet, T.C.; Kabbani, N. A G-protein-coupled α7 nicotinic receptor regulates signaling and TNF-α release in microglia. *FEBS Open Bio.* 2017, 7, 1350–1361. [CrossRef]
- Nordman, J.C.; Muldoon, P.; Clark, S.; Damaj, M.I.; Kabbani, N. The α4 nicotinic receptor promotes CD4+ T-cell proliferation and a helper T-cell immune response. *Mol. Pharmacol.* 2014, *85*, 50–61. [CrossRef] [PubMed]
- Chrestia, J.F.; Bruzzone, A.; Esandi, M.D.C.; Bouzat, C. Tyrosine phosphorylation differentially fine-tunes ionotropic and metabotropic responses of human α7 nicotinic acetylcholine receptor. *Cell. Mol. Life. Sci.* 2021, 78, 5381–5395. [CrossRef] [PubMed]

- Kabbani, N.; Nordman, J.C.; Corgiat, B.A.; Veltri, D.P.; Shehu, A.; Seymour, V.A.; Adams, D.J. Are nicotinic acetylcholine receptors coupled to G proteins? *Bioessays* 2013, 35, 1025–1034. [CrossRef] [PubMed]
- Razani-Boroujerdi, S.; Boyd, R.T.; Dávila-García, M.I.; Nandi, J.S.; Mishra, N.C.; Singh, S.P.; Pena-Philippides, J.C.; Langley, R.; Sopori, M.L. T cells express alpha7-nicotinic acetylcholine receptor subunits that require a functional TCR and leukocyte-specific protein tyrosine kinase for nicotine-induced Ca<sup>2+</sup> response. *J. Immunol.* 2007, 179, 2889–2898. [CrossRef]
- Skok, M.V. To channel or not to channel? Functioning of nicotinic acetylcholine receptors in leukocytes. J. Leukoc. Biol. 2009, 86, 1–3. [CrossRef]
- Chernyavsky, A.; Chen, Y.; Wang, P.H.; Grando, S.A. Pemphigus vulgaris antibodies target the mitochondrial nicotinic acetylcholine receptors that protect keratinocytes from apoptolysis. *Int. Immunopharmacol.* 2015, 29, 76–80. [CrossRef]
- 99. Gergalova, G.; Lykhmus, O.; Kalashnyk, O.; Koval, L.; Chernyshov, V.; Kryukova, E.; Tsetlin, V.; Komisarenko, S.; Skok, M. Mitochondria express α7 nicotinic acetylcholine receptors to regulate Ca2+ accumulation and cytochrome c release: Study on isolated mitochondria. *PLoS ONE* 2012, 7, e31361. [CrossRef]
- 100. Skok, M.; Gergalova, G.; Lykhmus, O.; Kalashnyk, O.; Koval, L.; Uspenska, K. Nicotinic acetylcholine receptors in mitochondria: Subunit composition, function and signaling. *Neurotransmitter* **2016**, *3*, e1290. [CrossRef]

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.