# Supplementary Material

Sex differences in the association between cumulative use of cannabis and cognitive function in middle age: The Coronary Artery Risk Development in Young Adults (CARDIA) Study

Baptiste Pasquier, MMED<sup>1</sup>, Kristine Yaffe, MD<sup>2</sup>, Deborah A. Levine, MD, MPH<sup>3</sup>, Jamal S. Rana, MD, PhD<sup>4</sup>, Mark J. Pletcher, MD, MPH<sup>2</sup>, Kali Tal, PhD<sup>1</sup>, Stephen Sidney, MD<sup>5</sup>, Reto Auer, MD, MAS<sup>1,6</sup> and Julian Jakob, MD,<sup>1,7</sup>

<sup>1</sup> Institute of Primary Health Care (BIHAM), University of Bern, Bern, Switzerland <sup>2</sup> University of California San Francisco, San Francisco, CA, USA, <sup>3</sup> University of Michigan, Department of Internal Medicine and Cognitive Health Services Research Program, Ann Arbor, MI, USA, <sup>4</sup> Kaiser Permanente Northern California, Department of Cardiology, Oakland, CA USA, <sup>5</sup> Kaiser Permanente Northern California, Division of Research, Oakland, CA, USA,

<sup>6</sup> University General Medicine and Public Health Centre, University of Lausanne, Lausanne, Switzerland.
<sup>7</sup>Department of Paediatrics, University Hospital Bern, Inselspital, Bern, Switzerland

#### **Corresponding author**

Julian Jakob, MD, MAS Institute of Primary Health Care (BIHAM) University of Bern, Switzerland Email: julian.jakob@biham.unibe.ch

# Content

eMe	ethods
N	1easurements
	Cannabis use5
	Cigarette smoking exposure
	Alcohol exposure
	Other illicit drug exposure
	Cardiovascular risk factors
	Physical activity7
	Psychosocial measures
	Socioeconomic hardship7
	Additional measures of the Rey Auditory Verbal Learning Test (RAVLT)7
	Mirror star tracing test
	Changes in categories of current cannabis use between Year 25 and Year 30
S	tatistical analysis
	Inverse probability of censoring weights (IPCWs)
	Changes in midlife cognitive function by categories of cannabis exposure
eRe	sults
	Association between changes in cannabis use and changes in midlife CF10
eTa	bles and eFigures
	<b>Appendix Table 1:</b> Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup> . 2698 CARDIA participants (excluding participants with cannabis use in the 30 days prior to the study visit). Results stratified by sex. 11
	Appendix Table 2: Characteristics of 169 CARDIA participants with cognitive function test results at Year 30 with cannabis use in the 24 hours prior to the study visit). Characteristics stratified by sex
	<b>Appendix Table 3:</b> Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup> . 3145 CARDIA participants (including participants with cannabis use in the 24 hours prior to study visit). Results stratified by sex 16
	<b>Appendix Table 4:</b> Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup> . 2912 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study visit and participants with history of stroke or TIA <sup>b</sup> ). Results stratified by sex
	<b>Appendix Table 5:</b> Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup> . 2974 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study visit and additionally adjusted for the mirror star tracing test). Results stratified by sex
	<b>Appendix Table 6</b> : Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup> . 2974 CARDIA participants (excluding

participants with cannabis use in the 24 hours prior to the study visit). Results stratified by sex, Appendix Table 7: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2866 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study visit). Results stratified by sex Appendix Table 8: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2922 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study visit and participants with Appendix Table 9: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2914 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study visit and participants with Appendix Table 10: Unadjusted and adjusted association between measures of immediate memory and learning and cumulative exposure to cannabis in 'cannabis-years'a. 2974 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study). Results Appendix Table 11: Characteristics of 2786 CARDIA with measures of cognitive function at year 25 and year 30 (including participants with use in 24 hours prior to study visit). Results stratified by sex and categories of current exposure to cannabis......25 **Appendix Table 12**: Distribution of cognitive function at Year 25 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup> among 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams (including use in 24 hours prior to study visit). Results Appendix Table 13: Distribution of cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'a among 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams (including use in 24 hours prior to study visit). Results Appendix Table 14. Unadjusted 5-year change in midlife cognitive function by categories of current cannabis use between Year 25 and Year 30 exam. 2786 CARDIA participants with Appendix Table 15. Minimally adjusted 5-year change in midlife cognitive function by categories

Appendix Table 16: Unadjusted and adjusted association between cognitive function decline of > 0.1 SD between Year 25 and Year 30 exams and categories of exposure to cannabis at year 25 and year 30 among participants with measures of cognitive function at Year 25 and Year 30 exams (2786 participants, including use in 24 hours prior to study visit). Results stratified by sex.

 **Appendix Figure 2:** Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'. 3145 CARDIA participants (including participants with cannabis use in the 24 hours prior to study visit). Results stratified by sex. .... 36

# eMethods

#### Measurements

#### Cannabis use

We present below an example of one participant included in the study illustrating the method for computing cannabis-years more fully and how we applied linear imputation (Box 1). This is the same method of imputation used in a previous publication by our research group.<sup>1</sup>

visit	mj30d_	mj30d_imp	domj	life_	yomj_max
0	15	15	300	100 to 499 times	0.82
1		15	482		1.32
2	20	20	725	100 to 499 times	1.99
3		20	968		2.65
4		4	1017		2.79
5	4	4	1066	100 to 499 times	2.92
6		4	1115		3.05
7	20	20	1358	100 to 499 times	3.72
8		20	1601		4.39
9		20	1844		5.05
10	20	20	2087	500 to 1000	5.72
				times	
11		20	2330		6.38
12		20	2573		7.05
13		10	2695		7.38
14		10	2817		7.72
15	10	10	2939	500 to 1000	8.05
				times	
16		10	3061		8.39
17	-	10	3183		8.72
18	-	1	3195		8.75
19	-	1	3207		8.79
20	1	1	3219	100 to 499 times	8.82

Roy	1	Evample	of	computing	cannahis_	voare f	or one	narticir	ant.
DUX	1.	Example	ΟI	computing	carmapis-	years in		particip	Jani.

Visit: visit year; mj30d\_: Self-reported days of using cannabis during the month before the visit ("During the last 30 days, on how many days did you use cannabis?"), mj30d\_imp: imputed mj30d\_ variable; domj: computed cumulative days of cannabis use, life\_: self-reported lifetime use of cannabis queried at each visit ("About how many times in your lifetime have you used marijuana?"), yomj\_max: computed cumulative years of cannabis use (domj/365).

According to example 1, the participant reported having used cannabis 100 to 499 times during the lifetime (categorical variable life\_) at the baseline examination (visit 0). This was used to estimate the exposure prior to the first examination (domj at visit 0=300, where domj signifies "days of cannabis").

At this baseline examination, the participant reported using cannabis 15 days per month (mj30d\_). Multiplied by 12.17 months (365/30), we estimated that this participant used cannabis 182 days in the first year after the first exam (15 \* 12.17). The number of days of cannabis use in the month before the baseline examination was imputed forward at year 1 (mj30d\_imp). At year 2, the participant reported using cannabis 20 days per month; at year 5, he/she reported 4 days per month. These numbers were imputed backwards and forwards; when there were an uneven number of intervals (preventing us from evenly splitting the

imputation based on the prior value as opposed to the post-value), the exposure at the prior interval was favoured arbitrarily. Participants then accrued lifetime days of cannabis use over follow-up. The cumulative number of cannabis-years over lifetime is presented in the last column (yomj\_max). For this participant, the cumulative number of cannabis years was 8.82, corresponding to 3,219 estimated days of cannabis use. We found no change in the estimates when using alternate methods for imputing missing values such as using the mean number of days of cannabis use between two examinations with data on this variable.

#### Cigarette smoking exposure

Detailed cigarette smoking behavior was evaluated during each in-person CARDIA visit by responses to an interviewer-administered questionnaire.<sup>2</sup> Participants were also asked the number of cigarettes smoked daily at yearly contact over the phone between CARDIA visits. Current smoking was defined as smoking one cigarette or more per day. At baseline, participants were asked the number of years they had smoked in total, the age at which they started smoking regularly ("How old were you when you started smoking cigarettes regularly?") and years since cessation ("How many years ago did you stop?"), if appropriate. These data were used to estimate cumulative lifetime exposure to cigarettes in terms of pack-years, with 1 pack-year of exposure equivalent to 7300 cigarettes (1 year × 365 days/y × 1 pack/d × 20 cigarettes/pack).<sup>1</sup>

#### Alcohol exposure

Alcohol consumption was measured during each CARDIA visits. Participants were asked, "Did you drink any alcoholic beverages in the past year?" and three follow-up questions regarding how many drinks of wine, beer, and liquor they usually consumed per week. Assuming that one drink of beer, wine, or liquor contains 16.7 ml, 17.0 ml, or 19.1 ml of ethanol, respectively (per CARDIA protocol), we estimated total ethanol consumption per week in milliliters of ethanol and divided it by 17.24 ml of ethanol per average drink to estimate the usual number of drinks per week that each participant reported at each visit. We estimated lifetime alcohol consumption in "drink-years," defining 1 drink-year as the amount of alcohol consumed in 1 year by a person consuming 1 drink/day (365 days/year ×17.24 ml of alcohol/day = 6,293 ml of alcohol). Binge drinking at the Year 20 visit was assessed directly by asking participants: "During the past 30 days, on how many days did you have five or more drinks on the same occasion?" For the other visits, we computed bingeing as follows: At each visit, participants were asked: "In the past month what is the largest number of drinks you had in one day?" At baseline, participants were additionally asked: "How many days in the past month did you have about (number of drinks answered in the previous question) drinks?" We used the number of days participants reported having 5 or more drinks for these visits. For binge drinking events at the Year 7 visit, we used the closest available information about the number of days patients reported having 5 or more drinks if they reported having such a use in one day within the last month at the Year 7 visit.

#### Other illicit drug exposure

Other illicit substances queried included cocaine (including other forms of cocaine such as crack, powder, free base), amphetamines (speed, uppers, methamphetamines) and heroin.<sup>5</sup> Participants were asked: "Have you ever used (substance)?"; "During the last 30 days, on how many days did you use (substance)?" and "How many times in your lifetime have you used (substance)?" The number of days on cocaine, crack, speed, methamphetamines and heroin over the study duration was computed using current exposure at each visit and replaced by lifetime exposure when the latter was higher.

#### Cardiovascular risk factors

Blood pressure was measured on the right arm with a Hawksley random zero sphygmomanometer (WA Baum Company, Copaigue, NY) by trained and certified

technicians using standardized methods after the participant had rested for 5 minutes at baseline and Year 7, and at Year 20, a digital blood pressure monitor (Omron HEM-907XL; Online Fitness, Santa Monica, CA) was used.

Three measurements were obtained at 1-minute intervals. The average of the second and third measurements was used in analyses. Fasting total cholesterol and triglycerides were measured enzymatically at baseline, Years 7, and 20 by the Northwest Lipid Research Laboratory at the University of Washington (Seattle, WA). For all visits, HDL cholesterol was determined by dextran sulfate –magnesium precipitation on the Abbot Spectrum, and LDL cholesterol was calculated using the Friedewald equation. At each visit, weight and height were measured, and body mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. We built a cumulative measure of some cardiovascular risk factor (blood cholesterol, blood triglycerides, blood pressure) by calculating the area under the curve from year 0 through year 30.

#### Physical activity

Physical activity was measured with the CARDIA Physical Activity History questionnaire, which queries the amount of time per week spent performing 13 categories of leisure, occupational, and household physical activities over the past 12 months.

#### Psychosocial measures

Depression, symptoms of psychosis and ADHD have been associated with cannabis use and cognitive impairment<sup>6</sup> In the main model we adjusted for the self-reported depression using the Center for Epidemiologic Studies Depression scale (CES-D). A score of  $\geq$ 16 was the cut-off for both sexes, indicating clinically significant depressive symptoms.

We tested our results to sensitivity excluding participants having ever used antipsychotic medications (1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation) and excluding participants having ever used a treatment for ADHD (methylphenidate, amphetamine, dextroamphetamine,

metamphetamine, lisdexamfetamine). Antipsychotics and ADHD treatment were recorded verbatim and coded through a centralized automated system and according to the Iowa Drug Information Services. We used the American Hospital Formulary Services (AHSF) and the Women's Health and Aging Study (WHAS) classifications to classify medications used for depression. We had information on use of these medications at every visit.

#### Socioeconomic hardship

Income data were collected at Year 30 for 3383 adults of the CARDIA cohort study. Participants were asked "Which of these categories best describe your total combined family income for the past 12 months? This should include income (before taxes) from all sources, wages, veteran's benefits, help from relatives, rent from properties, and so on". 4 Categories of annual income were created at each visit: Less than \$25,000, \$25,000 to \$49,900, \$50,000 to \$99,900, more than \$100,000. We built a cumulative measure of socioeconomic hardship by calculating the area under the curve from year 0 to 30 from these categories of annual income.

#### Additional measures of the Rey Auditory Verbal Learning Test (RAVLT)

The RAVLT consisted of 5 presentations of a 15-item word list (List A) with recall, one presentation of a 15-item Interference List (List B) with recall, a Short Delay Free Recall of List A (Trial 6) and a Long Delay Free Recall of list A (Trial 7).<sup>7</sup> The 15 words were read slowly to the subject, requesting him/her to repeat them after the reading, independently from the order they were said. The same procedure was repeated in steps A2 (i.e., second recall of list A), A3, A4 and A5, pointing out that the subject had to remember all the words, including those said previously. Then a second list of words (List B) was read as a distracter, and the subject was asked to recall these new words. After that, the words of the first list were asked (Trial 6, Recall after Interference), without being exposed, in a task of immediate

recall from episodic memory. Ten minutes after this stage the subject was asked to recall the words of the first list (Trial 7, Delayed Recall) to assess the delayed recall of episodic memory.

The immediate recall after Trial 1 provided the score for **Immediate Recall**, the total words recalled across the five learning trials (A1 to A5) the **Total Encoding**, subtracting the words recalled in Trial V from the words recalled in Trial I (Trial V – I) the **amount learned in five trials**, the score of trial 6 provided the score for **Recall after Interference** and the score of Trial 7 provided the score for **Delayed Recall**.

#### Mirror star tracing test

At Year 2, the mirror star tracing test was conducted to elicit reactive blood pressure. In the mirror star-tracing test, participants had to trace the outline of a star from a reversed image displayed in a mirror while staying within narrow limits. Study participants were instructed to draw stars as quickly as possible with the fewest possible errors. If they moved out of the limits of the star, an error was scored. Total stars completed and total numbers of errors over three minutes were recorded. Although initially intended as a stressor to measure blood pressure reactivity and not as a cognitive test, some have suggested that the mirror star tracing test measures aspects of executive function.<sup>3</sup>

Changes in categories of current cannabis use between Year 25 and Year 30 We included participants who completed cognitive testing at both Year 25 and Year 30. We then classified participants according to their pattern of cannabis use in never users, past users (any use before Year 25 and not at Year 25 or 30), recent quitters (any use in past month at Year 25 but not at Year 30), recent restarters (any use in past month before Year 25, no use at Year 25 and any use in past month at Year 30) and continuous users (any use in past month at Year 25 and 30). No participant reported using cannabis for the first time within this five-year period. We did not have measures of acute use at Year 25 visit. To avoid comparing acute users (use in the 24 hours prior to the study visit) at Year 25 visit with nonacute users at Year 30 visit, we did not exclude users at Year 30 visit from this analysis.

#### Statistical analysis

#### Inverse probability of censoring weights (IPCWs)

To reduce the potential for informative censoring, we computed IPCWs.<sup>8</sup> Covariables included in the pooled logistic regression model used to estimate the IPCWs were: fixed covariables: race, sex, study center and education; time-dependent covariables were lagged values of: age, study visits, pack-years of cigarette smoking, current smoking, drink-years of alcohol use, binge drinking events, cumulative exposure to cocaine, crack, amphetamines and heroin. Drink-years of alcohol, binge-drinking episodes, tobacco pack-years, BMI, physical activity and CVRF were all modelled as 3-knot restricted cubic splines.

Changes in midlife cognitive function by categories of cannabis exposure We used descriptive statistics to compare participants with different cannabis use patterns separately for women and men. We estimated the 5-year change in the three cognitive test results *z* scores assessed at Year 25 and Year 30, for the five possible combinations of current cannabis use. We performed linear and logistic regression models. Linear regression analyses included an unadjusted, a minimally adjusted (age, race, study center and years of education) and a multivariable adjusted model. In the logistic regression analysis, we defined a lower cognitive function of -0.1 SD or more from the mean change between Year 25 and Year 30 as "cognitive decline". This was based on two studies. Auer et al. found that for every 5 cannabis-years, verbal memory was 0.15 standardized units lower, and Meier et al. reported a 0.6 SD difference in verbal memory and a 1.0 SD difference in informant-reported memory between never users and persistent cannabis users over 20 years.<sup>3,9</sup> We then calculated the odds of developing a 0.1 SD cognitive decline according to current cannabis use category compared to participants that never reported cannabis use. We tested this in unadjusted and adjusted logistic regression models, with covariables measured at Year 25. Models were adjusted to the same covariables as for the cross-sectional analysis. We also tested models with 0.5 SD defined as cut-off.

### eResults

Association between changes in cannabis use and changes in midlife CF 2,786 participants had CF scores at Year 25 and at Year 30. We describe participants' characteristics at baseline (see Appendix Table 11). After 5 years of follow-up, RAVLT scores changed by  $+0.2 \pm 2.6$  points for men and  $+0.1 \pm 2.6$  points for women, DSST scores by  $-0.2 \pm 8.1$  points for men and  $-0.3 \pm 10.2$  points for women, and Stroop scores  $-0.3 \pm 9.1$  points for men and  $-0.1 \pm 9.3$  points for women across all categories of cannabis use (see Appendix Table 12 and Table 13). Results of the linear regression models (unadjusted and minimally adjusted) are presented in Appendix Table 14 and 15. The chance of developing a 0.1 SD larger cognitive decline in delayed recall, DSST scores or Stroop scores over 5 years was not associated with changes in cannabis use during the same periods, even after we extensively adjusted for covariables at Year 25 (see Appendix Table 15, Figure 4).

# eTables and eFigures

**Appendix Table 1:** Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years' <sup>a</sup>. 2698 CARDIA participants (<u>excluding</u> participants with cannabis <u>use in the 30 days</u> prior to the study visit). Results stratified by sex.

		Standardized difference in each CF measures (95% CI) <sup>c</sup>							
		Men		Women					
<b>Cognitive Function Measure</b> - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center Adjusted for age, cardiovascular risk factors, depression and marital status, with IPCW <sup>d</sup>		Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>d</sup>			
Rey Auditory Verbal Learning-									
Test, delayed recall (RAVLT)									
- Number of participants	1102	1100	1078	1592	1592	1555			
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)			
- 1 day to < 0.5 cannabis years	-0.20 (-0.37 to -0.03)	-0.19 (-0.34 to -0.03)	-0.25 (-0.43 to -0.07)	0.08 (-0.05 to 0.21)	-0.01 (-0.13 to 0.11)	0.02 (-0.12 to 0.15)			
<ul> <li>- 0.5 to &lt; 2 cannabis years</li> </ul>	-0.30 (-0.48 to -0.12)	-0.19 (-0.37 to -0.02)	-0.33 (-0.57 to -0.09)	0.09 (-0.07 to 0.25)	-0.05 (-0.20 to 0.10)	0.04 (-0.14 to 0.21)			
- 2 to < 5 cannabis years	-0.31 (-0.57 to -0.04)	-0.05 (-0.28 to 0.19)	-0.22 (-0.51 to 0.08)	-0.46 (-0.75 to -0.17)	-0.30 (-0.57 to -0.05)	-0.20 (-0.48 to 0.08)			
- > 5 cannabis years	-0.45 (-0.79 to -0.12)	-0.12 (-0.46 to 0.19)	-0.36 (-0.69 to -0.03)	-0.27 (-0.65 to 0.10)	-0.19 (-0.52 to 0.15)	0.07 (-0.44 to 0.31)			
P-value for trend <sup>e</sup>	0.007	0.12	0.05	0.007	0.13	0.46			

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users with use within the last 30-day excluded (N=447).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>c</sup>Linear regression models determined the association between CF scores and self-reported cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring

**Appendix Table 2:** Characteristics of 169 CARDIA participants with cognitive function test results at Year 30 with cannabis <u>use in the</u> <u>24 hours</u> prior to the study visit). Characteristics stratified by sex.

	Cumulative exposure to cannabis <sup>a</sup>									
		М	en			Wo	men			
Variable	1 day to	0.5 to < 2	2 to < 5	> 5	1 day to	0.5 to < 2	2 to < 5	> 5		
	<0.5	cannabis	cannabis	cannabis	<0.5	cannabis	cannabis	cannabis		
	cannabis	years	years	years	cannabis	years	years	years	p-va	alue <sup>b</sup>
	years				years					
N (%)	0 (0)	8(8)	14(14)	80 (78)	1(2)	7(10)	6(9)	53 (79)		
Demographics										
Age, median (Q1; Q3)	- (-; -)	55 (51;58)	59 (56;59)	54 (52;57)	50 (50;50)	52 (50;59)	57 (49;60)	56 (51;59)	0.014	0.617
Race, N (Col. %) <sup>c</sup>									0.264	0.369
- Black	- (-; -)	3 (38)	10 (71)	42 (53)	1 (100)	4 (57)	2 (33)	35 (66)		
- White	- (-; -)	5 (63)	4 (29)	38 (48)	0 (0)	3 (43)	4 (67)	18 (34)		
Education, median (Q1;	- (-; -)	16 (14;20)	15 (13;16)	14 (12;16)	16 (16;16)	14 (13;16)	14 (11;17)	15 (12;16)	0.114	0.848
Q3), years										
Study center, N (Col. %) do									0.566	0.274
you need this detail>										
- Birmingham, AL	- (-; -)	1 (13)	1 (7)	9 (11)	0 (0)	2 (29)	0 (0)	3 (6)		
- Chicago, IL	- (-; -)	1 (13)	4 (29)	7 (9)	0 (0)	2 (29)	0 (0)	5 (9)		
- Minneapolis, MI	- (-; -)	2 (25)	3 (29)	26 (33)	0 (0)	2 (29)	2 (33)	14 (26)		
- Oakland, CA	- (-; -)	4 (50)	6 (43)	38 (48)	1 (100)	1 (14)	4 (67)	31 (58)		
Substance use exposure										
Cannabis use category, N									0.003	< 0.001
(Col. %) <sup>u</sup>										
- No current use	- (-; -)	0 (0)	0 (0)	1 (36)	0 (0)	0 (0)	0 (0)	0 (0)		
- 1 to 10 days per month	- (-; -)	5 (63)	5 (36)	9 (44)	0 (0)	7 (100)	1 (17)	5 (9)		
- 11 to 29 days per month	- (-; -)	3 (38)	7 (50)	32 (15)	0 (0)	0 (0)	5 (83)	23 (43)		
- 30 days per month	- (-; -)	0 (0)	2 (14)	38 (5)	1 (100)	0 (100)	0 (0)	25 (47)		
(everyday)										

Tobacco smoking, N (Col. %)									0.748	0.426
- Never smoker	- (-; -)	3 (38)	2 (14)	24 (30)	1 (100)	2 (29)	1 (17)	9 (17)		
- Former smoker	- (-; -)	3 (38)	6 (43)	29 (36)	0 (0)	4 (57)	3 (50)	25 (47)		
- Current smoker	- (-; -)	2 (25)	6 (43)	29 (34)	0 (0)	1 (14)	2 (33)	19 (36)		
Age started smoking among ever tobacco smokers, median (Q1: Q3)	- (-; -)	17 (15; 19)	16 (13;18)	18 (16; 20)	- (-; -)	18 (15; 31)	16 (16; 20)	17 (15; 23)	0.281	0.777
Cumulative tobacco exposure among ever smokers, median (Q1; Q3), pack-years <sup>e</sup>	- (-; -)	2 (0; 9)	12 (2; 21)	6 (0; 18)	- (-; -)	2 (0; 11)	4 (1; 16)	6 (1; 24)	0.362	0.327
Alcohol use										
<ul> <li>Cumulative alcohol use among ever drinkers, median (Q1, Q3), drink- years<sup>f</sup></li> </ul>	- (-; -)	29 (5; 15)	38 (16;58)	32 (16;75)	1 (1; 1)	18 (2; 44)	22 (3; 38)	16 (7; 36)	0.451	0.432
Binge drinking days,										
cumulative use, N (Col.%) <sup>®</sup>	()	1 (12)	2 (21)	10 (12)	1 (100)	2 (20)	2 (22)	10 (24)	0.857	0.908
- 11ever reported billgeling	- (-; -)	1 (13)	3 (21) 4 (20)	10 (13)	1 (100)	2 (29)	2 (33)	16 (34)		
- > 250 days	- (-; -) - (-; -)	2 (23) 5 (63)	7 (50)	52 (65)	0(0)	3 (43)	2 (33)	19 (36)		
Illicit drug use Current use <sup>h</sup>									< 0.001	0.842
Cocaine, crack, speed or methamphetamine, N (Col. %)	- (-; -)	0 (0)	1 (7)	6 (8)	0 (0)	0 (0)	0 (0)	3 (6)		
Heroin, N (Col. %)	- (-; -)	0 (0)	0 (0)	1 (7)	0 (0)	1 (14)	0 (0)	0 (0)	0.2	0.034
Physical activity										
Physical activity score,	-	584	436	415	574	217	301	241	0.870	0.509
median (Q1; Q3) <sup>i</sup>	(-; -)	(140; 970)	(152; 955)	(193; 690)	(574; 574)	(105; 247)	(133;492)	(94; 361)		
Anthropomorphic variable										
BMI, mean (SD) <sup>i</sup>	- (-; -)	27 (2.6)	26.7 (4.0)	29.8 (5.0)	48.3	32.4 (5.6)	28.6 (5.0)	29.8 (7.2)	0.035	0.06

Cardiovascular risk factors										
Systolic blood pressure,	· .									
mean (SD), mmHg	- (-; -)	123 (16)	122 (14)	123 (14)	108	118 (10)	119 (8)	123 (17)	0.986	0.709
Diastolic blood pressure, mean (SD), mmHg	- (-; -)	78 (11)	75 (10)	76 (11)	71	74 (8)	73 (13)	75 (13)	0.850	0.949
LDL-Cholesterol, mean (SD), mg/dl	- (-; -)	108 (25)	113 (36)	104 (29)	74	91 (39)	85 (31)	109 (35)	0.575	0.210
HDL-Cholesterol, mean (SD), mg/dl	- (-; -)	48 (11)	59 (16)	52 (13)	54	70 (21)	65 (25)	66 (22)	0.119	0.905
Triglycerides, mean (SD),	-	126	75	118	52	99	156	86	0.004	0.143
mg/dl	(-; -)	(68; 138)	(62; 92)	(82; 157)	(52; 52)	(86; 104)	(66; 357)	(60; 114)		
Diabetes mellitus, N (%)	- (-; -)	1 (13)	2 (14)	11 (14)	0 (0)	1 (14)	3 (50)	4 (8)	0.993	0.024
Psychological variables										
Depression, current CES-D >=16/30, N (%) <sup>k</sup>	- (-; -)	0 (0)	2 (14)	16 (20)	0 (0)	1 (14)	1 (17)	11 (21)	0.337	0.930
Socioeconomical variable										
Currently married, N (%)	- (-; -)	2 (25)	4 (29)	34 (43)	0 (0)	4 (57)	3 (50)	19 (36)	0.381	0.544
Rey Auditory Verbal										
Learning-Test, delayed										
recall (RAVLT) at Year 30										
- Raw mean (SD)	- (-; -)	6.1 (3.1)	7.2 (3.0)	6.3 (3.2)	9.0	8.3 (3.5)	6.2 (3.7)	7.5 (3.1)	0.592	0.643
Rey Auditory Verbal										
Learning-Test, delayed										
recall (RAVLT) at Year 25										
- Raw mean (SD)	- (-; -)	7.0 (3.5)	6.8 (3.7)	6.2 (3.1)	10.0	8.6 (3.4)	6.2 (2.5)	7.8 (3.3)	0.709	0.500
Digit Symbol Substitution										
Test (DSST) at Year 30										
- Raw mean (SD)	- (-; -)	67.1(8.8)	60.3(10.4)	61.4(11)	78.0	65.3 (19)	71.7 (12)	67 (17)	0.332	0.816
Digit Symbol Substitution										
Test (DSST) at Year 25	- (-; -)	70.5 (11)	64.3 (11)	62.2 (11)	78.0	68.7 (15)	74.8 (12)	68.8 (15)	0.168	0.730
- Raw mean (SD)										

Stroop Interference Test at Year 30										
- Raw mean (SD)	- (-; -)	-22.9(7.3)	-31.6(11)	-22.9(12)	-23.0	-25.6 (5)	-16.5 (11)	-25 (11)	0.037	0.331
Stroop Interference at Year 25 <sup>1</sup>										
- Raw mean (SD)	- (-; -)	-25.0(9.9)	-31.3(17)	-24.6(12)	-23.0	-24.7 (8.2)	-21.3 (6.4)	-24.8 (11)	0.234	0.885
Letter Fluency Test at Year										
30	- (-; -)	48.3 (3.0)	47.1 (14)	42.3 (12)	43.0	44.1 (19)	49.7 (12)	45.2 (12)	0.214	0.862
- Raw mean (SD)										
Category Fluency Test at Year 30	- (-; -)	22.4 (5.8)	19.8 (6.1)	20.0 (4.9)	15.0	19.9 (2.9)	19.7 (5.8)	20.0 (4.8)	0.445	0.774
- Raw mean (SD)										
MOCA Test at Year 30										
- Raw mean (SD)	- (-; -)	22.3 (3.0)	23.5 (4.5)	22.9 (3.1)	18.0	23.6 (2.2)	22.7 (3.9)	22.7 (3.7)	0.707	0.556

BMI = body mass index; CARDIA = Coronary Artery Risk Development in Young Adults study; Col. % = column percentage; LDL = low density lipoprotein (LDL); HDL = high-density lipoprotein; n = number of participants; Q1, Q3 = 1<sup>st</sup> and 3<sup>rd</sup> quartile (percentiles 25 and 75); SD = standard deviation

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y).

<sup>b</sup>P-values are from Kruskal-Wallis rank test for age, years of education, pack-years, drink-years, age started smoking, triglycerides, BMI and physical activity and from a χ2 test for race, study site, current smoking status, CES-D, cannabis use categories, cumulative binge drinking categories, illicit drug use categories, and marital status. P-values are from 1-way analyses of variance for the cognitive functions.

<sup>c</sup>By design, the CARDIA study sampled self-identified white men, white women, black men and black women in roughly equal numbers for participation in the study.

<sup>d</sup>Categories based on the answer to the question: "During the last 30 days, on how many days did you use marijuana?"

eSelf-reported cumulative exposure to cigarettes in pack-years: 1 pack-year of exposure = 7300 cigarettes (1 year × 365 days/y × 1 pack/d × 20 cigarettes/pack).

<sup>f</sup>Cumulative alcohol use in drink-years: 1 drink-year is the total amount of ethanol consumed by a person who had 1 alcoholic drink per day for 1 year (1 drink-year = 17.24 ml of ethanol/drink x 1 drink/d x 365 days/y = 6292.6 ml of ethanol).

<sup>g</sup>Binge drinking days, defined as ≥5 drinks per day. If bingeing were to be constant over 25 years in one individual, 250 binge drinking days would correspond to 10 days of bingeing each year for 25 years.

<sup>h</sup>Current use, defined as any use within the last 30 days. We computed the number of days on the illicit drug over the study duration based on current exposure at each visit, which we replaced with cumulative exposure when the latter was higher. Cocaine included all forms of cocaine, like crack, powder, free base; amphetamines included speed, uppers, and methamphetamines.

Physical activity, measured with the CARDIA Physical Activity History questionnaire, which queries the amount of time per week spent performing 13 categories of leisure, occupational, and household physical activities over the past 12 months.

<sup>j</sup>Calculated as weight in kilograms divided by height in meters squared.

<sup>k</sup>Self-reported depression was measured every five years, starting at the Year 5 visit, on the Center for Epidemiologic Studies Depression scale (CES-D).18 A score of ≥16 was the cut-off for both sexes, indicating clinically significant depressive symptoms.

We used the inverse of the Stroop score so we could interpret worse CF with negative standardized scores for all six CF tests.

**Appendix Table 3:** Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 3145 CARDIA participants (including participants with cannabis <u>use in the 24 hours prior</u> to study visit). Results stratified by sex.

		Standardized difference in each CF measures (95% CI) <sup>c</sup>								
		Men		Women						
Cognitive Function Measure - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>d</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>d</sup>				
Rey Auditory Verbal Learning-										
Test, delayed recall (RAVLT)										
- Number of participants	1350	1350	1320	1789	1786	1749				
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)				
- 1 day to < 0.5 cannabis years	-0.20 (-0.37 to -0.03)	-0.18 (-0.34 to -0.02)	-0.25 (-0.42 to -0.07)	0.08 (-0.05 to 0.21)	-0.01 (-0.13 to 0.11)	0.00 (-0.13 to 0.13)				
- 0.5 to < 2 cannabis years	-0.32 (-0.50 to -0.15)	-0.23 (-0.39 to -0.06)	-0.38 (-0.61 to -0.14)	0.06 (-0.10 to 0.21)	-0.06 (-0.21 to 0.08)	0.03 (-0.14 to 0.19)				
- 2 to < 5 cannabis years	-0.31 (-0.54 to -0.09)	-0.06 (-0.28 to 0.15)	-0.25 (-0.51 to -0.02)	-0.35 (-0.58 to -0.11)	-0.26 (-0.46 to -0.05)	-0.17 (-0.41 to 0.06)				
- > 5 cannabis years	-0.53 (-0.73 to -0.32)	-0.31 (-0.50 to -0.11)	-0.54 (-0.79 to -0.29)	-0.31 (-0.51 to 0.10)	-0.19 (-0.38 to 0.01)	-0.05 (-0.27 to 0.17)				
P-value for trend <sup>e</sup>	< 0.001	0.01	< 0.001	< 0.001	0.03	0.37				

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>c</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring

**Appendix Table 4:** Unadjusted and adjusted association between RAVLT at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2912 CARDIA participants (<u>excluding</u> participants with cannabis use in the 24 hours prior to the study visit and participants with <u>history of stroke or TIA<sup>b</sup></u>). Results stratified by sex.

		Standardized difference in each CF measures (95% CI) <sup>d</sup>								
		Men		Women						
<b>Cognitive Function Measure</b> - Self-reported cumulative exposure in cannabis-years <sup>c</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>e</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>e</sup>				
Rey Auditory Verbal Learning-										
lest, delayed recall (RAVLI)										
- Number of participants	1225	1225	1200	1695	1692	1658				
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)				
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.21 (-0.38 to -0.04)	-0.19 (-0.35 to -0.03)	-0.23 (-0.41 to -0.05)	0.09 (-0.04 to 0.23)	-0.00 (-0.11 to 0.13)	0.03 (-0.10 to 0.16)				
- 0.5 to < 2 cannabis years	-0.32 (-0.50 to -0.14)	-0.22 (-0.39 to -0.05)	-0.34 (-0.58 to -0.10)	0.10 (-0.05 to 0.26)	-0.03 (-0.17 to 0.12)	0.06 (-0.10 to 0.24)				
- 2 to < 5 cannabis years	-0.30 (-0.53 to -0.06)	-0.07 (-0.29 to 0.15)	-0.21 (-0.48 to 0.06)	-0.29 (-0.53 to -0.06)	-0.22 (-0.43 to -0.04)	-0.12 (-0.36 to 0.12)				
- > 5 cannabis years	-0.49 (-0.73 to -0.26)	-0.28 (-0.50 to -0.06)	-0.48 (-0.74 to -0.20)	-0.17 (-0.42 to 0.09)	-0.10 (-0.33 to 0.13)	0.04 (-0.23 to 0.32)				
P-value for trend <sup>f</sup>	< 0.001	0.04	0.008	<0.01	0.24	0.46				

CARDIA = Coronary Artery Risk Development in Young Adults study, TIA = Transient ischemic attack

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>Cannabis users with history of Stroke (N=50) or TIA (N=12).

°Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>d</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>e</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 5:** Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2974 CARDIA participants (<u>excluding participants</u> with cannabis use in the 24 hours prior to the study visit and additionally adjusted for the <u>mirror star tracing test</u>). Results stratified by sex.

		Standardized difference in each CF measures (95% CI) <sup>c</sup>							
		Men	Women						
Cognitive Function Measure - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression marital status and mirror star tracing test, with IPCW <sup>d</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression, marital status and mirror star tracing test, with IPCW <sup>d</sup>			
Rey Auditory Verbal Learning- Test, delayed recall (RAVIT)									
- Number of participants	1248	1248	1134	1722	1722	1683			
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)			
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.20 (-0.37 to -0.03)	-0.19 (-0.34 to -0.03)	-0.2 (-0.39 to -0.014)	0.08 (-0.05 to 0.21)	-0.01 (-0.13 to 0.11)	-0.14 (-0.09 to 0.17)			
- 0.5 to < 2 cannabis years	-0.31 (-0.49 to -0.14)	-0.21 (-0.37 to -0.05)	-0.31 (-0.56 to -0.7)	0.06 (-0.09 to 0.22)	-0.06 (-0.20 to 0.08)	-0.15 (-0.11 to 0.36)			
- 2 to < 5 cannabis years	-0.31 (-0.54 to -0.07)	-0.07 (-0.28 to 0.14)	-0.22 (-0.50 to 0.06)	-0.31 (-0.54 to -0.07)	-0.22 (-0.43 to -0.01)	-0.72 (-0.37 to 0.04)			
- > 5 cannabis years	-0.48 (-0.71 to -0.24)	-0.25 (-0.47 to -0.04)	-0.38 (-0.66 to -0.09)	-0.19 (-0.44 to 0.07)	-0.12 (-0.35 to 0.11)	0.30 (-0.50 to 1.11)			
P-value for trend <sup>e</sup>	< 0.001	0.05	0.08	<0.01	0.20	0.42			

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>c</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 6**: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2974 CARDIA participants (<u>excluding participants</u> with cannabis use in the 24 hours prior to the study visit). Results stratified by sex, <u>not adjusting for marital status</u>.

		Standardized difference in each CF measures (95% CI) <sup>c</sup>								
		Men		Women						
Cognitive Function Measure - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors and depression with IPCW <sup>d</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, and depression with IPCW <sup>d</sup>				
Rey Auditory Verbal Learning-										
Test, delayed recall (RAVLT)										
- Number of participants	1248	1248	1225	1722	1722	1688				
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0(Ref)				
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.20 (-0.37 to -0.03)	-0.18 (-0.34 to -0.03)	-0.24 (-0.42 to -0.06)	0.08 (-0.05 to 0.21)	-0.01 (-0.13 to 0.11)	0.0 (-0.13 to 0.13)				
- 0.5 to < 2 cannabis years	-0.32 (-0.50 to -0.14)	-0.20 (-0.37 to -0.04)	-0.36 (-0.59 to -0.12)	0.06 (-0.09 to 0.22)	-0.06 (-0.20 to 0.08)	0.03 (-0.15 to 0.19)				
- 2 to < 5 cannabis years	-0.31 (-0.54 to -0.07)	-0.07 (-0.28 to 0.15)	-0.28 (-0.54 to -0.01)	-0.31 (-0.54 to -0.07)	-0.22 (-0.43 to -0.01)	-0.15 (-0.39 to 0.09)				
- > 5 cannabis years	-0.48 (-0.71 to -0.24)	-0.27 (-0.48 to -0.05)	-0.48 (-0.75 to -0.21)	-0.19 (-0.44 to 0.07)	-0.12 (-0.35 to 0.11)	-0.01 (-0.28 to 0.27)				
P-value for trend <sup>e</sup>	< 0.001	0.05	0.009	0.002	0.19	0.60				

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

°Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 7**: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2866 CARDIA participants (<u>excluding</u> participants with cannabis use in the 24 hours prior to the study visit). Results stratified by sex and adjusting for socioeconomic hardship.

	Standardized difference in each CF measures (95% CI) <sup>c</sup>									
		Men			Women					
<b>Cognitive Function Measure</b> - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center, and socioeconomic hardship	Additionally adjusted for substance use, cardiovascular risk factors, depression, marital status and socioeconomic hardship, with IPCW <sup>d</sup>	Unadjusted model	Adjusted for age, race, education, study center, and socioeconomic hardship	Additionally adjusted for substance use, cardiovascular risk factors, depression, marital status and socioeconomic hardship, with IPCW <sup>d</sup>				
Rey Auditory Verbal Learning-										
Test, delayed recall (RAVLT)										
- Number of participants	1248	1242	1216	1722	1713	1676				
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0(Ref)				
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.20 (-0.37 to -0.03)	-0.19 (-0.35 to -0.03)	-0.26 (-0.44 to -0.08)	0.08 (-0.05 to 0.21)	-0.03 (-0.15 to 0.09)	-0.01 (-0.14 to 0.13)				
- 0.5 to < 2 cannabis years	-0.32 (-0.50 to -0.14)	-0.21 (-0.38 to -0.05)	-0.39 (-0.62 to -0.15)	0.06 (-0.09 to 0.22)	-0.06 (-0.21 to 0.08)	0.03 (-0.14 to 0.20)				
- 2 to < 5 cannabis years	-0.31 (-0.54 to -0.07)	-0.04 (-0.26 to 0.15)	-0.26 (-0.53 to -0.01)	-0.31 (-0.54 to -0.07)	-0.19 (-0.40 to 0.02)	-0.13 (-0.37 to 0.12)				
- > 5 cannabis years	-0.48 (-0.71 to -0.24)	-0.24 (-0.47 to -0.03)	-0.52 (-0.80 to -0.25)	-0.19 (-0.44 to 0.07)	-0.11 (-0.34 to 0.12)	-0.03 (-0.25 to 0.30)				
P-value for trend <sup>e</sup>	< 0.001	0.04	0.003	<0.01	0.43	0.68				

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>c</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 8**: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2922 CARDIA participants (<u>excluding participants</u> with cannabis use in the 24 hours prior to the study visit and participants with history of use of antipsychotic medications<sup>b</sup>). Results stratified by sex.

	Standardized difference in each CF measures (95% CI) <sup>d</sup>										
		Men			Women						
<b>Cognitive Function Measure</b> - Self-reported cumulative exposure in cannabis-years <sup>c</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>e</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW					
Rey Auditory Verbal Learning-											
Test, delayed recall (RAVLT)											
- Number of participants	1232	1223	1207	1694	1694	1655					
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	O(Ref)					
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.19 (-0.35 to -0.02)	-0.21 (-0.36 to -0.05)	-0.25 (-0.42 to -0.07)	0.10 (-0.03 to 0.23)	0.01 (-0.11 to 0.13)	0.03 (-0.11 to 0.16)					
- 0.5 to < 2 cannabis years	-0.30 (-0.48 to -0.12)	-0.21 (-0.38 to -0.05)	-0.41 (-0.63 to -0.18)	0.09 (-0.06 to 0.25)	-0.03 (-0.18 to 0.11)	0.05 (-0.12 to 0.21)					
- 2 to < 5 cannabis years	-0.29 (-0.52 to -0.05)	-0.06 (-0.27 to 0.17)	-0.27 (-0.53 to -0.01)	-0.30 (-0.54 to -0.07)	-0.21 (-0.42 to 0.01)	-0.15 (-0.39 to 0.09)					
- > 5 cannabis years	-0.46 (-0.69 to -0.23)	-0.24 (-0.46 to -0.03)	-0.50 (-0.77 to -0.24)	-0.19 (-0.44 to 0.07)	-0.11 (-0.34 to 0.12)	0.01 (-0.28 to 0.28)					
P-value for trend <sup>f</sup>	< 0.001	0.07	0.002	0.002	0.18	0.45					

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>(1st, 2nd and 3rd generation). 52 participants excluded.

°Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>d</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>e</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 9**: Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2914 CARDIA participants (<u>excluding participants</u> with cannabis use in the 24 hours prior to the study visit and participants with history of stimulant drug use<sup>b</sup>). Results stratified by sex.

	Standardized difference in each CF measures (95% CI) <sup>d</sup>										
		Men			Women						
<b>Cognitive Function Measure</b> - Self-reported cumulative exposure in cannabis-years <sup>c</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>e</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW					
Rey Auditory Verbal Learning-											
Test, delayed recall (RAVLT)											
- Number of participants	1223	1223	1197	1687	1687	1648					
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	O(Ref)					
- 1 day to < 0.5 cannabis years	-0.23 (-0.40 to -0.06)	-0.21 (-0.36 to -0.05)	-0.27 (-0.44 to -0.09)	0.08 (-0.05 to 0.21)	-0.01 (-0.13 to 0.10)	0.01 (-0.13 to 0.14)					
- 0.5 to < 2 cannabis years	-0.33 (-0.51 to -0.15)	-0.21 (-0.38 to -0.05)	-0.37 (-0.61 to -0.13)	0.05 (-0.11 to 0.21)	-0.07 (-0.21 to 0.08)	0.04 (-0.13 to 0.21)					
- 2 to < 5 cannabis years	-0.31 (-0.55 to -0.08)	-0.07 (-0.29 to 0.14)	-0.28 (-0.55 to -0.01)	-0.31 (-0.55 to -0.07)	-0.22 (-0.44 to -0.01)	-0.14 (-0.38 to 0.11)					
- > 5 cannabis years	-0.50 (-0.73 to -0.26)	-0.26 (-0.48 to -0.04)	-0.51 (-0.78 to -0.25)	-0.19 (-0.44 to 0.07)	-0.12 (-0.34 to 0.14)	0.03 (-0.25 to 0.30)					
P-value for trend <sup>f</sup>	< 0.001	0.04	0.004	0.002	0.18	0.60					

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>methylphenidate, amphetamine, dextroamphetamine, metamphetamine, lisdexamfetamine). 60 participants excluded.

°Years of cannabis exposure was modeled as a 5-level categorical predictor.

<sup>d</sup>Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>e</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

**Appendix Table 10:** Unadjusted and adjusted association between measures of immediate memory and learning and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup>. 2974 CARDIA participants (excluding participants with cannabis use in the 24 hours prior to the study). Results stratified by sex.

	Standardized difference in each CF measures (95% CI) <sup>c</sup>											
		Men			Women							
Cognitive Function Measure - Self-reported cumulative exposure in cannabis-years <sup>b</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression, and marital status, with IPCW <sup>d</sup>	Unadjusted model	Adjusted for age, race, education, study center,	Additionally adjusted for substance use, cardiovascular risk factors, depression, and marital status, with IPCW <sup>d</sup>						
Rey Auditory Verbal Learning-												
Test, Immediate Recall												
- Number of participants	1247	1247	1221	1721	1721	1681						
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)						
- 1 day to < 0.5 cannabis years	-0.11 (-0.28 to 0.06)	-0.11 (-0.27 to 0.05)	-0.07 (-0.26 to 0.12)	0.06 (-0.07 to 0.19)	-0.02 (-0.14 to 0.10)	-0.02 (-0.16 to 0.11)						
- 0.5 to < 2 cannabis years	-0.23 (-0.40 to -0.05)	-0.15 (-0.32 to 0.03)	-0.04 (-0.28 to 0.21)	0.04 (-0.11 to 0.20)	-0.07 (-0.22 to 0.09)	-0.02 (-0.21 to 0.17)						
- 2 to < 5 cannabis years	-0.30 (-0.53 to -0.06)	-0.12 (-0.34 to 0.11)	-0.05 (-0.34 to 0.25)	-0.17 (-0.41 to 0.06)	-0.12 (-0.34 to 0.11)	-0.05 (-0.27 to 0.18)						
- > 5 cannabis years	-0.46 (-0.69 to -0.22)	-0.28 (-0.50 to -0.05)	-0.05 (-0.34 to 0.23)	-0.17 (-0.43 to 0.09)	-0.12 (-0.37 to 0.12)	-0.05 (-0.33 to 0.22)						
P-value for trend <sup>e</sup>	< 0.001	0.19	0.96	0.09	0.67	0.99						
Rey Auditory Verbal Learning-												
Test, Total Encoding												
- Number of participants	1239	1239	1213	1713	1713	1685						
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)						
<ul> <li>1 day to &lt; 0.5 cannabis years</li> </ul>	-0.15 (-0.32 to 0.02)	-0.15 (-0.30 to 0.00)	-0.14 (-0.31 to 0.02)	0.11 (-0.02 to 0.25)	0.03 (-0.09 to 0.15)	0.06 (-0.07 to 0.19)						
- 0.5 to < 2 cannabis years	-0.30 (-0.47 to -0.12)	-0.17 (-0.33 to 0.00)	-0.15 (-0.38 to 0.09)	0.01 (-0.15 to 0.17)	-0.11(-0.25 to 0.04)	-0.03 (-0.21 to 0.14)						
- 2 to < 5 cannabis years	-0.33 (-0.56 to -0.09)	-0.07 (-0.28 to 0.14)	-0.05 (-0.33 to 0.23)	-0.36 (-0.59 to -0.12)	-0.25 (-0.45 to -0.04)	-0.17 (-0.40 to 0.07)						
- > 5 cannabis years	-0.53 (-0.76 to -0.30)	-0.28 (-0.49 to -0.07)	-0.20 (-0.45 to 0.06)	-0.19 (-0.45 to 0.06)	-0.10 (-0.33 to 0.13)	0.01 (-0.27 to 0.28)						
P-value for trend	< 0.001	0.08	0.37	< 0.001	0.01	0.13						

Rey Auditory Verbal Learning- Test, Amount learned in five						
trials						
- Number of participants	1241	1241	1215	1713	1713	1676
- Never used cannabis	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)	0 (Ref)
- 1 day to < 0.5 cannabis years	-0.13 (-0.30 to 0.04)	-0.13 (-0.30 to 0.04)	-0.21 (-0.41 to -0.02)	0.06 (-0.07 to 0.19)	0.04 (-0.09 to 0.18)	0.07 (-0.08 to 0.21)
- 0.5 to < 2 cannabis years	-0.20 (-0.38 to -0.03)	-0.14 (-0.31 to 0.04)	-0.26 (-0.51 to 0.02)	-0.05 (-0.20 to 0.11)	-0.08(-0.24 to 0.08)	-0.04 (-0.24 to 0.16)
- 2 to < 5 cannabis years	-0.18 (-0.41 to 0.06)	-0.03 (-0.27 to 0.20)	-0.16 (-0.46 to 0.13)	-0.09 (-0.32 to 0.15)	-0.03 (-0.27 to 0.21)	0.01 (-0.24 to 0.26)
- > 5 cannabis years	-0.17 (-0.40 to 0.07)	-0.03 (-0.27 to 0.21)	-0.27 (-0.56 to 0.03)	-0.07 (-0.33 to 0.18)	-0.03 (-0.27 to 0.23)	0.04 (-0.25 to 0.32)
P-value for trend	0.27	0.44	0.20	0.30	0.48	0.58

CARDIA = Coronary Artery Risk Development in Young Adults study

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169).

<sup>b</sup>Years of cannabis exposure was modeled as a 5-level categorical predictor.

\*Linear regression models determined the association between CF scores and cumulative exposure to cannabis use. Negative standardized scores indicate worse CF.

<sup>d</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring.

<sup>e</sup>Tests of statistical significance were 2-tailed, with an alpha level set at 0.05.

Cannabis use categories<sup>a</sup> at year 30 based on cannabis use status between year 25 and year 30 Men Women Never Users Past users Recent Never Past users Recent Recent Recent Variable (any use auitters restarters Users (any use quitters restarters before year (any use at (Any use before year (any use at (Any use Continuous Continuous 25 visit and before year 25 visit and before year year 25 year 25 users (any users (any not at visit and not at 25 visit. No and not at 25 visit. No not at visit use at year use at year use at year use at year 25 or 30) year 30) 25 and any 25 or 30) year 30) 25 and any 25 and any 25 and any use at year use at year use at year use at year 30) 30) 30) 30) Ν N = 168 N = 759 N = 55 N = 81 N = 261N = 1077 N = 69 N = 79 N = 135 N = 102 Demographics Age, median (Q1; Q3) 55 (52; 58) 55 (52; 58) 55 (51; 58) 56 (53; 58) 56 (52; 58) 56 (53; 58) 55 (52; 58) 56 (53; 58) 55 (52; 58) 56 (52; 58) Race, N (Col. %)<sup>b</sup> - Black 66 (39) 306 (40) 28 (51) 40 (49) 62 (46) 155 (59) 477(44) 42 (61) 43 (54) 57 (56) 453 (60) 73 (54) 27 (39) - White 102 (61) 27 (49) 41 (51) 106 (41) 600(56) 36 (46) 45 (44) Education, median (Q1; Q3), 16 (14; 20) 16 (14; 18) 15 (13; 17) 15 (13; 16) 15 (13; 16) 16 (14; 18) 16 (14;18) 15 (14;16) 15 (13;16) 15 (13; 16) years Study center, N (Col. %) 6 (7) 8 (8) - Birmingham, AL 74 (44) 190 (25) 18 (33) 17 (13) 119 (46) 194 (18) 16 (23) 15 (19) - Chicago, IL 45 (27) 168 (22) 14 (25) 17 (21) 23 (17) 64 (25) 247 (23) 17 (25) 14 (18) 18 (18) - Minneapolis, MI 28 (17) 50 (19) 200 (26) 9 (16) 30 (37) 44 (33) 284 (26) 13 (19) 20 (25) 29 (28) - Oakland, CA 21 (13) 201 (26) 14 (25) 28 (35) 51 (38) 28 (11) 30 (38) 47 (46) 352 (33) 23 (33) Substance use exposure 0 (0) Cumulative cannabis use, mean 3.5 (4.7) 3.2 (3.3) 12.1 (7.4) 0 (0) 0.6 (1.2) 2.5 (4.34) 3.3 (3.7) 9.9 (7.3) 1.1 (2.0) (SD)<sup>c</sup> Cannabis use category, N (Col. %)<sup>d</sup> - No current use 0 (0) 0 (0) 0 (0) 168 (100) 759 (100) 55 (100) 0 (0) 261 (100) 1077 (100) 69 (100)

**Appendix Table 11**: Characteristics of 2786 CARDIA with measures of cognitive function at year 25 and year 30 (including participants with use in 24 hours prior to study visit). Results stratified by sex and categories of current exposure to cannabis.

- 1 to 10 days per month	0 (0)	0 (0)	0 (0)	58(72)	60 (44)	0 (0)	0 (0)	0 (0)	69 (87)	48 (47)
- 11 to 29 days per month	0 (0)	0 (0)	0 (0)	20 (25)	40 (30)	0 (0)	0 (0)	0 (0)	9 (11)	29 (28)
- 30 days per month (everyday)	0 (0)	0 (0)	0 (0)	3 (4)	34(25)	0 (0)	0 (0)	0 (0)	1 (1)	24 (23)
Cannabis use within the last 24										
hours, N (Col. %)										
- Yes	0 (0)	0 (0)	0 (0)	19 (27)	65 (48)	0 (0)	0 (0)	0 (0)	17 /22)	42 (41)
Tobacco smoking,										
N (Col. %)										
- Never smoker	149 (89)	399 (53)	18 (33)	24 (30)	42 (31)	222 (85)	539 (50)	25 (36)	15 (19)	19 (19)
- Former smoker	14 (8)	256 (34)	21 (38)	35 (43)	56 (41)	34 (13)	420 (39)	27 (39)	39 (49)	54 (53)
- Current smoker	5 (3)	104 (14)	16 (29)	22 (27)	37 (27)	5 (2)	118 (11)	17 (25)	25 (32)	29 (28)
Age started smoking among	20 (17; 33)	17 (15; 21)	19 (17; 24)	17 (15; 19)	18 (16; 20)	25 (18; 34)	17 (15; 20)	17 (15; 18)	18 (16;22)	17 (15;21)
ever tobacco smokers, median										
(Q1: Q3)										
Cumulative tobacco exposure										
among ever smokers, median	0 (0; 0)	0 (0; 7)	2 (0; 17)	2 (0; 16)	4 (0; 18)	0 (0;0)	0 (0;6)	4 (0; 11)	8 (0; 17)	4 (0; 16)
(Q1; Q3), packyears <sup>e</sup>										
Alcohol use										
- Cumulative alcohol use among	2 (0; 12)	17 (6; 37)	26 (10; 67)	32 (13; 51)	35 (20; 70)	0 (0;3)	7(1; 20)	8 (2; 21)	17 (3; 28)	16 (7; 39)
ever drinkers, median (Q1, Q3),										
drink-years'										
Binge drinking days, cumulative										
use, N (Col. %) <sup>s</sup>		222 (22)	44 (20)		12 (12)	222 (22)		22 (12)	25 (22)	26 (25)
- never reported bingeing	114 (68)	223 (29)	11 (20)	14 (17)	13 (10)	232 (89)	584 (54)	29 (42)	25 (32)	36 (35)
- ≤ 250 days	28 (17)	211 (28)	14 (25)	19 (23)	32 (24)	24 (9)	322 (30)	22 (32)	27 (34)	25 (25)
- > 250 days	26 (15)	325 (43)	30 (55)	48 (59)	90 (67)	5 (2)	171 (16)	18 (26)	27 (34)	41 (40)
Illicit drug use <sup>h</sup>										
Current use										
Cocaine, crack, speed or	0 (0)	9 (1)	3 (5)	4 (5)	12 (9)	1 (0)	10 (1)	0 (0)	6 (8)	7 (7)
methamphetamine, N (Col. %)										
Heroin, N (Col. %)	0 (0)	4 (1)	0 (0)	3 (4)	1 (1)	0 (0)	1 (0)	0 (0)	1 (1)	0 (0)
Cumulative use										

Cocaine, crack, speed or methamphetamine N (Col. %)										
- Never reported using	165 (98)	288 (38)	17 (31)	8 (10)	7 (5)	253 (97)	494 (46)	25 (36)	11 (14)	18 (18)
- 1 to 25 days	2 (1)	178 (23)	8 (15)	23 (28)	23 (17)	6 (2)	263 (24)	12 (17)	18 (23)	13 (13)
- 25 to 250 days	0 (0)	130 (17)	10 (18)	26 (32)	51 (38)	1 (0)	215 (20)	16 (23)	24 (30)	33 (32)
- 250 days	1 (1)	163 (21)	20 (36)	24 (30)	54 (40)	1 (0)	105 (10)	16 (23)	26 (33)	38 (37)
Heroin, N (Col. %)										
- Never reported using	164 (98)	629 (83)	38 (69)	51 (63)	96 (71)	254 (97)	947 (88)	59 (86)	63 (80)	84 (82)
- 1-25 days	3 (2)	66 (9)	8 (15)	11 (14)	24 (18)	7 (3)	92 (9)	5 (7)	9 (11)	15 (15)
- >25 days	1 (1)	64 (8)	9 (16)	19 (23)	15 (11)	0 (0)	38 (4)	5 (7)	7 (9)	3 (3)
Physical activity										
Physical activity score, median	331.8 ±	385.2 ±	316.1 ±	420.5 ±	453.5 ±	220.8 ±	285.4 ±	260.6 ±	286.2 ±	279.6 ±
(Q1; Q3) <sup>i</sup>	262.0	293.1	237.0	359.0	306.8	210.9	239.3	232.5	234.2	228.9
Anthropomorphic variable										
BMI, mean (SD) <sup>j</sup>	31.5 ± 6.3	29.7 ± 6.1	29.8 ± 5.7	29.2 ± 5.3	28.3 ± 4.6	32.1 ± 8.2	30.6 ± 8.0	31.1 ± 7.6	31.6 ± 8.0	30.2 ± 7.6
Cardiovascular risk factors										
Systolic blood pressure, mean	121.2 ±	121.4 ±	125.2 ±	124.7 ±	123.4 ±	119.7 ±	118.6 ±	125.2 ±	119.3 ±	122.6 ±
(SD), mmHg	13.0	14.9	15.3	13.5	15.3	18.9	17.2	21.2	16.8	17.5
Diastolic blood pressure, mean										
(SD), mmHg	74.8 ± 10.0	74.2 ± 10.6	77.3 ± 11.5	76.9 ± 10.5	75.4 ± 10.5	73.1 ± 11.3	72.5 ± 10.9	76.9 ± 12.7	73.3 ± 11.4	75.6 ± 11.8
LDL-Cholesterol, mean (SD),	111.0 ±	110.0 ±	110.6 ±	104.5 ±	101.1 ±	111.9 ±	112.1 ±	106.9 ±	107.0 ±	115.5 ±
mg/dl	35.4	33.2	27.1	36.0	30.2	31.1	32.8	32.8	32.9	32.5
HDL-Cholesterol, mean (SD),	48.4 ±12.0	51.3 ± 14.5	54.6 ± 15.9	53.3 ± 16.4	54.1 ± 14.6	63.5 ± 15.8	67.2 ± 19.9	65.1 ± 20.0	62.7 ± 21.4	68.2 ± 20.7
mg/dl										
Triglycerides, mean (SD), mg/dl	121.0 ±	121.4 ±	117.4 ±	125.8 ±	136.7 ±	92.0 ± 49.5	95.4 ± 59.6	99.1 ± 43.0	115.4 ±	99.1 ± 60.7
	66.2	141.7	70.7	91.5	208.9				88.6	
Diabetes mellitus, N (%)	23 (14)	105 (14)	6 (11)	13 (16)	22 (16)	34 (13)	149 (14)	12 (17)	14 (18)	10 (10)
Psychological variables										
Depression, current CES-D	10 (6)	78 (10)	10 (18)	19 (23)	26 (19)	38 (15)	150 (14)	13 (19)	19 (24)	19 (19)
>=16/30, N (%) <sup>k</sup>										
Socioeconomical variable										
Currently married, N (%)	112 (67)	470 (62)	21 (38)	32 (40)	66 (49)	148 (57)	544 (51)	29 (42)	31 (39)	38 (37)
Past cardiovascular Diseases										

Non-fatal Stroke, N (%)	5 (3)	16 (2)	1 (2)	2 (2)	2 (1)	3 (1)	22 (2)	2 (3)	3 (4)	2 (2)
Non-fatal TIA, N (%)	0 (0)	2 (0)	1 (2)	0 (0)	1 (1)	1 (0)	6 (1)	1 (1)	0 (0)	0 (0)

BMI = body mass index; CARDIA = Coronary Artery Risk Development in Young Adults study; Col. % = column percentage; LDL = low density lipoprotein (LDL); HDL = high-density lipoprotein; n = number of participants; Q1, Q3 = 1<sup>st</sup> and 3<sup>rd</sup> quartile (percentiles 25 and 75); SD = standard deviation, TIA = Transient ischemic attack

<sup>a</sup>Categories based on the answer to the questions: "During the last 30 days, on how many days did you use marijuana?"

<sup>b</sup>By design, the CARDIA study sampled self-identified white men, white women, Black men and Black women in roughly equal numbers for participation in the study.

<sup>c</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabis use (1 year × 365 days/y).

<sup>d</sup>Categories based on the answer to the question: "During the last 30 days, on how many days did you use marijuana?"

eSelf-reported cumulative exposure to cigarettes in pack-years: 1 pack-year of exposure = 7300 cigarettes (1 year × 365 days/y × 1 pack/d × 20 cigarettes/pack).

<sup>1</sup>Cumulative alcohol use in drink-years: 1 drink-year is the total amount of ethanol consumed by a person who had 1 alcoholic drink per day for 1 year (1 drink-year = 17.24 ml of ethanol/drink x 1 drink/d x 365 days/y = 6292.6 ml of ethanol)

<sup>9</sup>Binge drinking days, defined as ≥5 drinks per day. If bingeing were constant over 25 years in one individual, 250 binge drinking days would correspond to 10 days of bingeing each year for 25 years. <sup>h</sup>Current use, defined as any use within the last 30 days. We computed the number of days on the illicit drug over the study duration based on current exposure at each visit, which we replaced with cumulative exposure when the latter was higher. Cocaine included all forms of cocaine. like crack, powder, free base: amphetamines included speed, uppers, and methamphetamines.

Physical activity, measured with the CARDIA Physical Activity History questionnaire, which queries the amount of time per week spent performing 13 categories of leisure, occupational, and household physical activities over the past 12 months

<sup>i</sup>Calculated as weight in kilograms divided by height in meters squared.

<sup>k</sup>Self-reported depression was measured every five years, starting at the Year 5 visit, on the Center for Epidemiologic Studies Depression scale (CES-D).18 A score of ≥16 was the cut-off for both sexes, indicating clinically significant depressive symptoms.

**Appendix Table 12**: Distribution of cognitive function at Year 25 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup> among 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams (including use in 24 hours prior to study visit). Results stratified by sex and categories of current exposure to cannabis.

	Cannabis use categories <sup>b</sup> at year 25 based on cannabis use status between year 25 and year 30											
			Men					Women				
Variable	Never Users	Past users (any use before year 25 visit and not at visit 25 or 30)	Recent quitters (any use at year 25 and not at year 30)	Recent restarters (Any use before year 25 visit. No use at year 25 and any use at year 30)	Continuous users (any use at year 25 and any use at year 30)	Never Users	Past users (any use before year 25 visit and not at visit 25 or 30)	Recent quitters (any use at year 25 and not at year 30)	Recent restarters (Any use before year 25 visit. No use at year 25 and any use at year 30)	Continuous users (any use at year 25 and any use at year 30)	p-v	alue <sup>c</sup>
Rey Auditory Verbal Learning-												
Number of participants	1.60	750			405	264	4077	60	70	400		.0.001
- Number of participants	168	759	55	81	135	261	10//	69	79	102	<0.00	<0.001
- Raw mean (SD)	8.4 (3.3)	7.6; (3.1)	7.2 (3.1)	7.0 (3.5)	6.7 (3.4)	9.1 (3.2)	9.4 (3.0)	8.4 (3.0)	8.5 (3.0)	8.1 (3.5)		
	0.27	0.01	-0.09	-0.15	-0.26	-0.04	0.07	-0.23	-0.21	-0.35	<u> </u>	
Digit Symbol Substitution Test (DSST)												
- Number of participants	166	753	55	81	136	258	1079	69	77	102		
- Raw mean (SD)	68.5 (15.2)	66.4 (15.4)	62.2 (15.1)	63.5 (16.7)	65.5 (14.1)	71.8(15.8)	75.7(15.0)	69.7(14.8)	70.6(15.7)	69.7(16.2)	0.03	<0.01
- Standardized mean	0.15	0.02	-0.29	-0.16	-0.04	-0.15	0.10	-0.29	-0.23	-0.29		
Stroop Interference Test <sup>e</sup>												
- Number of participants	168	747	53	81	131	258	1071	70	78	101		
- Raw mean (SD)	-23.3(12.7)	-22.4(10.1)	-23.8(10.5)	-24.1(12.3)	-22.4(10.3)	-22.7(10.3)	-21.6(10.4)	-24.7(12.5)	-25.2(16.0)	-24.3(11.0)	0.56	<0.01
- Standardized mean	-0.03	0.03	-0.12	-0.12	0.01	-0.03	0.06	0.22	-0.27	-0.19		

CARDIA = Coronary Artery Risk Development in Young Adults study, SD = standard deviation.

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabs use (1 year × 365 days/y).

<sup>b</sup>Categories based on the answer to the question: "During the last 30 days, on how many days did you use marijuana?"

°P-values are from 1-way analyses of variance. All P values two sided.

<sup>d</sup>Each CF measure was standardized by subtracting the mean and then dividing the score by the within-CARDIA sex-specific standard deviation.

"We used the inverse of the Stroop score so we could interpret worse CF with negative standardized scores for all three CF tests.

**Appendix Table 13**: Distribution of cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'<sup>a</sup> among 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams (including use in 24 hours prior to study visit). Results stratified by sex and categories of current exposure to cannabis.

	Cannabis use categories <sup>b</sup> at year 30 based on cannabis use status between year 25 and year 30											
			Men					Women				
Variable	Never Users	Past users (any use before year 25 visit and not at visit 25 or 30)	Recent quitters (any use at year 25 and not at year 30)	Recent restarters (Any use before year 25 visit. No use at year 25 and any use at year 30)	Continuous users (any use at year 25 and any use at year 30)	Never Users	Past users (any use before year 25 visit and not at visit 25 or 30)	Recent quitters (any use at year 25 and not at year 30)	Recent restarters (Any use before year 25 visit. No use at year 25 and any use at year 30)	Continuous users (any use at year 25 and any use at year 30)	p-v	alue <sup>c</sup>
Rey Auditory Verbal Learning-												
- Number of participants	168	759	55	81	135	261	1077	69	79	102	<0.00	<0.001
- Raw mean (SD)	8.5 (3.5)	7.8: (3.3)	7.4 (3.1)	7.0 (3.7)	6.8 (3.3)	9.3 (3.2)	9.5 (3.3)	8.4 (2.9)	8.3 (3.4)	8.3 (3.4)	1	.0.001
- Standardized mean <sup>d</sup>	0.25	0.02	-0.09	-0.21	-0.27	-0.01	0.07	-0.27	-0.30	-0.30		
Digit Symbol Substitution Test (DSST)												
- Number of participants	166	753	55	81	136	258	1079	69	77	102		
- Raw mean (SD)	66.1 (15.4)	63.8 (16.1)	59.7 (16.9)	61.5 (15.6)	63.4 (13.7)	69.0(17.0)	72.9(16.1)	68.6(14.5)	66.9(17.3)	68.6(17.1)	0.06	<0.01
- Standardized mean	0.15	0.01	-0.25	-0.14	-0.02	-0.15	0.08	-0.17	-0.28	-0.17		
Stroop Interference Test <sup>e</sup>												
- Number of participants	168	747	53	81	131	258	1071	70	78	101		
- Raw mean (SD)	-23.5(15.8)	-23.0(12.6)	-23.9(10.5)	-23.6(13.4)	-21.8(10.2)	-23.3(11.1)	-21.7(10.4)	-22.4(10.9)	-23.8(12.3)	-24.8(11.8)	0.76	0.02
- Standardized mean	-0.04	0.01	-0.07	-0.05	0.09	-0.09	0.05	-0.01	-0.13	-0.23		

CARDIA = Coronary Artery Risk Development in Young Adults study, SD = standard deviation.

<sup>a</sup>Self-reported cumulative exposure to cannabis joints in cannabis-years; 1 cannabis-year of exposure = 365 days of cannabs use (1 year × 365 days/y).

<sup>b</sup>Categories based on the answer to the question: "During the last 30 days, on how many days did you use marijuana?"

°P-values are from 1-way analyses of variance. All P values two sided.

<sup>d</sup>Each CF measure was standardized by subtracting the mean and then dividing the score by the within-CARDIA sex-specific standard deviation.

"We used the inverse of the Stroop score so we could interpret worse CF with negative standardized scores for all three CF tests.

Test results			Men					Women			p-value <sup>b</sup>	
across	Never	Past users	Recent	Recent	Continuous	Never	Past users	Recent	Recent	Continuous		
categories of	users		quitters	restarters	users	users		quitters	restarters	users		
current			-									
exposure to												
cannabis <sup>a</sup>												•
Rey Auditory	0.00 (-0.11	0.04 (-0.02	0.02 (-0.18	-0.03 (-0.20	0.02 (-0.11	0.02 (-0.08	-0.01 (-	-0.04 (-0.22	-0.08 (-0.25	0.05 (-0.10	0.94	0.79
Verbal	to 0.12)	to 0.10)	to 0.23)	to 0.14)	to 0.15)	to 0.11)	0.05 to	to 0.14)	to 0.09)	to 0.20)		
Learning-Test,							0.04)					
delayed recall												
(RAVLT)												
Digit Symbol	0.01 (-0.07	0.00 (-0.03	0.02 (-0.11	0.05 (-0.06	0.05 (-0.04	-0.02 (-	-0.04 (-	0.08 (-0.06	-0.08 (-0.21	0.08 (-0.04	0.86	0.17
Substitution	to 0.08)	to 0.04)	to 0.15)	to 0.15)	to 0.13)	0.10 to	0.07 to -	to 0.23)	to 0.06)	to 0.20)		
Test (DSST)						0.05)	0.00)					
Stroop	0.01 (-0.11	-0.03 (-	0.02 (-0.20	0.06 (-0.11	0.07 (-0.06	-0.03 (-	0.01 (-0.04	0.23 (0.04	0.16 (-0.02	-0.01 (-0.17	0.64	0.08
Interference	$t_0 (0.13)$	0.09 to	$t_0 (0.24)$	to 0.24)	to 0.21)	0.13 to	to 0.06)	to 0.43)	to 0.35)	$t_0 0.15$		
Test		0.03)				0.07)			10 0.00)			
1030		0.007				0.077						

**Appendix Table 14**. Unadjusted 5-year change in midlife cognitive function by categories of current cannabis use between Year 25 and Year 30 exams. 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams

<sup>a</sup>Never users: Never used cannabis; **Past users:** Any use before Year 25 visit and no current use at Year 25 or Year 30 visits; **Recent quitters:** Current use at Year 25 visit and no current use at Year 25 visit. **Recent restarters:** Any use before Year 25 visit. No current use at Year 25 visit and current use at Year 30 visit. **Continuous users:** Current use at Year 25 and at Year 30 visit. **RAVLT** - Rey Auditory Verbal Learning Test; DSST – Digit Symbol Substitution Test

<sup>b</sup>Tests of statistical significance were 2-tailed, with an alpha level set at 0.05

**Appendix Table 15**. Minimally adjusted 5-year change in midlife cognitive function by categories of current cannabis use between Year 25 and Year 30 exams. 2786 CARDIA participants with measures of cognitive function at Year 25 and Year 30 exams

Test results			Men					Women			p-valu	ue <sup>b</sup>
across	Never	Past users	Recent	Recent	Continuous	Never	Past users	Recent	Recent	Continuous		
categories of	users		quitters	restarters	users	users		quitters	restarters	users		
current												
exposure to												
cannabis"	0.07/0.05	0.04/0.02	0.05 ( 0.45	0.00 ( 0.05	0.02/0.46	0.05 ( 0.05	0.02 (	0.02 ( 0.20	0.00 ( 0.00	0.05 ( 0.40	0.50	0.00
Rey Auditory	0.07(-0.05)	0.04 (-0.02	0.05 (-0.15	-0.09 (-0.25	-0.02 (-0.16	0.05(-0.05)	-0.02 (-	-0.02 (-0.20	-0.06 (-0.23	0.05 (-0.10	0.58	0.69
Verbal	(0.19)	10 0.09)	to 0.25)	to 0.08)	to 0.11)	10 0.14)	0.06 to	10 0.16)	to 0.10)	to 0.20)		
Learning-Test,							0.03)					
delayed recall												
(KAVLI)												
Digit Symbol	0.03 (-0.05	0.00 (-0.03	0.04 (-0.09	0.03 (-0.08	0.03 (-0.05	-0.01.(-	-0.04 (-	0.09 (-0.05	-0.08(-0.22	0.08 (-0.04	0.91	0.15
Substitution	to 0 11)	$t_0 = 0.04$	$t_0 (0.03)$	to 0 13)	to 0 11)	0 09 to	0.08 to -	to 0 24)	to 0.06)	to 0 20)	0.51	0.15
Test (DSST)	(0 0.11)	10 0.04)	(0 0.17)	(0 0.15)	(0 0.11)	0.07)	0.00	10 0.24)	10 0.00)	10 0.207		
1030 (0331)						0.077	0.007					
Churchen	0.02 ( 0.10	0.02 (	0.02/0.20	0.06 ( 0.12	0.07/0.07	0.01 (	0.01 ( 0.04	0.21 (0.02	0.12 ( 0.06	0.07/0.22	0.65	0.17
Stroop	0.03 (-0.10	-0.03 (-	0.02 (-0.20	0.06 (-0.12	0.07 (-0.07	-0.01 (-	0.01 (-0.04	0.21 (0.02	0.12 (-0.06	-0.07 (-0.23	0.05	0.17
Interference	10 0.15)	0.09 (0	10 0.24)	10 0.23)	10 0.20)	0.11 (0	10 0.06)	(0 0.40)	(0.30)	(0 0.10)		
lest		0.03)				0.10)						
						1			1			1

<sup>a</sup>Never users: Never used cannabis; **Past users**: Any use before Year 25 visit and no current use at Year 25 or Year 30 visits; **Recent quitters**: Current use at Year 25 visit and no current use at Year 25 or Year 30 visits; **Recent quitters**: Current use at Year 25 visit and no current use at Year 30 visit. **Recent restarters**: Any use before Year 25 visit. No current use at Year 25 visit and current use at Year 30 visit. **Continuous users**: Current use at Year 25 and at Year 30 visit. Results from minimally adjusted linear regression models, adjusted for age, race, study site, education. RAVLT - Rey Auditory Verbal Learning Test; DSST – Digit Symbol Substitution Test <sup>b</sup>Tests of statistical significance were 2-tailed, with an alpha level set at 0.05

**Appendix Table 16**: Unadjusted and adjusted association between cognitive function decline of > 0.1 SD between Year 25 and Year 30 exams and categories of exposure to cannabis at year 25 and year 30 among participants with measures of cognitive function at Year 25 and Year 30 exams (2786 participants, including use in 24 hours prior to study visit). Results stratified by sex.

	Odds Ratio (95% CI) <sup>b</sup>							
Test results across categories of current exposure to cannabis <sup>a</sup>	Men			Women				
	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>c</sup>	Unadjusted model	Adjusted for age, race, education, study center	Additionally adjusted for substance use, cardiovascular risk factors, depression and marital status, with IPCW <sup>c</sup>		
Rey Auditory Verbal Learning-								
Test, delayed recall (RAVLT)								
- Never users	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)		
- Past users	0.99 (0.71 - 1.40)	1.10 (0.77 - 1.56)	1.12 (0.73 - 1.73)	0.91 (0.69 - 1.20)	1.01 (0.75 - 1.36)	0.99 (0.70 - 1.40)		
- Recent quitters	1.29 (0.70 - 2.38)	1.34 (0.72 - 2.50)	1.45 (0.68 - 3.10)	1.09 (0.64 - 1.87)	1.11 (0.64 - 1.93)	0.94 (0.50 - 1.77)		
- Recent restarters	1.37 (0.80 - 2.33)	1.66 (0.95 - 2.89)	1.71 (0.89 - 3.25)	1.14 (0.69 - 1.90)	1.18 (0.69 - 2.00)	0.94 (0.51 - 1.75)		
- Continuous users	1.00 (0.63 - 1.59)	1.18 (0.72 - 1.92)	1.33 (0.71 – 2.49)	1.06 (0.66 - 1.68)	1.15 (0.70 - 1.87)	0.93 (0.52 - 1.67)		
P-value for trend <sup>d</sup>	0.63	0.42	0.47	0.75	0.94	0.99		
Digit Symbol Substitution Test (DSST)								
- Never users	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)		
- Past users	1.01 (0.72 - 1.42)	1.13 (0.79 - 1.60)	1.26 (0.83 - 1.92)	1.10 (0.84 - 1.45)	1.22 (0.91 - 1.63)	1.13 (0.81 - 1.59)		
- Recent quitters	0.94 (0.50 - 1.74)	0.96 (0.51 - 1.80)	1.09 (0.51 - 2.33)	0.69 (0.40 - 1.21)	0.69 (0.39 - 1.22)	0.59 (0.32 - 1.09)		
- Recent restarters	1.02 (0.59 – 1.74)	1.21 (0.68 - 2.11)	1.38 (0.71 – 2.66)	1.29 (0.77 - 2.14)	1.39 (0.82 - 2.38)	1.11 (0.62 - 2.01)		
- Continuous users	0.93 (0.58 - 1.47)	1.07 (0.66 - 1.74)	1.30 (0.71 - 2.40)	1.05 (0.66 - 1.67)	1.14 (0.70 – 1.84)	0.99 (0.54 – 1.80)		
P-value for trend	0.85	0.93	0.83	0.40	0.17	0.20		
Stroop Interference Test								
- Never users	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)		
- Past users	1.16 (0.82 - 1.65)	1.21 (0.85 - 1.73)	1.12 (0.71 - 1.75)	0.99 (0.75 - 1.30)	1.03 (0.78 - 1.37)	1.13 (0.81 - 1.56)		
- Recent quitters	1.09 (0.58 - 2.06)	1.11 (0.58 - 2.12)	0.92 (0.44 – 1.94)	0.73 (0.43 - 1.26)	0.77 (0.44 - 1.32)	0.80 (0.44 - 1.46)		
- Recent restarters	1.06 (0.61 - 1.84)	1.11 (0.63 - 1.95)	1.17 (0.58 - 2.35)	0.90 (0.54 - 1.50)	0.96 (0.57 - 1.62)	0.95 (0.52 - 1.76)		

- Continuous users	1.08 (0.67 - 1.73)	1.15 (0.70 - 1.88)	1.11 (0.59 – 2.10)	1.15 (0.73 - 1.81)	1.25 (0.78 – 2.01)	1.37 (0.77 – 2.40)
P-value for trend	0.93	0.88	0.96	0.70	0.64	0.51

<sup>a</sup>Never users: Never used cannabis; Past users: Any use before Year 25 visit and no current use at Year 25 or Year 30 visits; Recent quitters: Current use at Year 25 visit and no current use at Year 30 visit. Recent restarters: Any use before Year 25 visit. No current use at Year 25 visit and current use at Year 30 visit. Continuous users: Current use at Year 25 and at Year 30 visit. <sup>b</sup>Cognitive decline defined as sex-specific decline > 0.1 standard deviation (SD) from the mean change. Results are shown as odds ratio compared with never users (baseline). <sup>c</sup>Analyses weighted by the inverse probability of censoring (IPCW) to address potential bias by informative censoring

<sup>d</sup>Tests of statistical significance were 2-tailed, with an alpha level set at 0.05

**Appendix Figure 1:** Adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'. 2698 CARDIA participants (<u>excluding</u> participants with cannabis use in <u>the 30 days</u> prior to the study visit). Results stratified by sex.



CARDIA = Coronary Artery Risk Development in Young Adults study

Self-reported cumulative exposure to cannabis joints in cannabis-years; 1-cannabis-year of exposure = 365 days of cannabis use (1 year x 365 days/y). Cannabis users within the last 30 days of the Year 30 visit excluded (N=447). Model results adjusted for age, race, study site, education, cigarette smoking (current, cumulative, age starting smoking), alcohol, illicit drug use, cardiovascular risk factors, physical activity, depression, marital status and compared with never users (baseline). All test results standardized: a 1-unit negative deviation indicates a standard deviation worse CF than the mean. We used the inverse of the Stroop score so we could interpret worse CF with negative standardized scores for all six CF tests. RAVLT - Rey Auditory Verbal Learning Test; DSST – Digit Symbol Substitution Test; MOCA - Montreal cognitive assessment **Appendix Figure 2:** Unadjusted and adjusted association between cognitive function at Year 30 and cumulative exposure to cannabis in 'cannabis-years'. 3145 CARDIA participants (including participants with cannabis use in <u>the 24 hours</u> prior to study visit). Results stratified by sex.



CARDIA = Coronary Artery Risk Development in Young Adults study

Self-reported cumulative exposure to cannabis joints in cannabis-years; 1-cannabis-year of exposure = 365 days of cannabis use (1 year x 365 days/y). Model results adjusted for age, race, study site, education, cigarette smoking (current, cumulative, age starting smoking), alcohol, illicit drug use, cardiovascular risk factors, physical activity, depression, marital status and compared with never users (baseline). All test results standardized: a 1-unit negative deviation indicates a standard deviation worse CF than the mean. We used the inverse of the Stroop score so we could interpret worse CF with negative standardized scores for all six CF tests. RAVLT - Rey Auditory Verbal Learning Test; DSST – Digit Symbol Substitution Test; MOCA - Montreal cognitive assessment **Appendix Figure 3:** Adjusted association between RAVLT and cumulative exposure to cannabis in 'cannabis-years'. 2974 CARDIA participants (excluding participants with cannabis use in the <u>24 hours</u> prior to the study). Results stratified by sex and <u>race</u>.



CARDIA = Coronary Artery Risk Development in Young Adults study

Self-reported cumulative exposure to cannabis joints in cannabis-years; 1-cannabis-year of exposure = 365 days of cannabis use (1 year x 365 days/y). Cannabis users within the 24 hours prior of the Year 30 visit excluded (N=169). Model results adjusted for age, study site, education, cigarette smoking (current, cumulative, age started smoking), alcohol, illicit drug use, cardiovascular risk factors, physical activity, depression, marital status and compared with never users (baseline). All test results standardized: a 1-unit negative deviation indicates a standard deviation worse CF than the mean. RAVLT - Rey Auditory Verbal Learning Test **Appendix Figure 4:** Adjusted association between cognitive function decline of > 0.1 SD between Year 25 and Year 30 exams and categories of exposure to cannabis at Year 25 and Year 30 among participants with measures of cognitive function at Year 25 and 30 exams (2786 participants, including use in 24 hours prior to study visit). Results stratified by sex.



Accelerated cognitive decline defined as sex-specific decline > 0.1 standard deviation (SD) from the mean change. Model results shown as odds ratio compared with never users (baseline).

Never users: Never used cannabis; Past users: Any use before Year 25 visit and no current use at Year 25 or Year 30 visits; Recent quitters: Current use at Year 25 visit and no current use at Year 30 visit. Recent restarters: Any use before Year 25 visit. No current use at Year 25 visit and current use at Year 30 visit. Continuous users: Current use at Year 25 and at Year 30 visit.

Results from multivariable logistic regression models, adjusted for age, race, study site, education, cigarette smoking (current, cumulative, age starting smoking), alcohol, illicit drug use, cardiovascular risk factors, physical activity, depression, marital status. RAVLT - Rey Auditory Verbal Learning Test; DSST – Digit Symbol Substitution Test

## eReferences

 Pletcher MJ, Vittinghoff E, Kalhan R, et al. Association between marijuana exposure and pulmonary function over 20 years. Jama 2012;307(2):173-81, doi:10.1001/jama.2011.1961
 Wagenknecht LE, Perkins LL, Cutter GR, et al. Cigarette smoking behavior is strongly related to educational status: the CARDIA study. Prev Med 1990;19(2):158-69, doi:10.1016/0091-7435(90)90017-e

3. Auer R, Vittinghoff E, Yaffe K, et al. Association Between Lifetime Marijuana Use and Cognitive Function in Middle Age: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. JAMA Intern Med 2016;176(3):352-61, doi:10.1001/jamainternmed.2015.7841

Alcoholism NIoAAa. Helping Patients Who Drink Too Much: A Clinician's Guide. 2005;
 Kertesz SG, Pletcher MJ, Safford M, et al. Illicit drug use in young adults and subsequent decline in general health: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. Drug Alcohol Depend 2007;88(2-3):224-33, doi:10.1016/j.drugalcdep.2006.10.017

6. Leadbeater BJ, Ames ME, Linden-Carmichael AN. Age-varying effects of cannabis use frequency and disorder on symptoms of psychosis, depression and anxiety in adolescents and adults. Addiction 2019;114(2):278-293, doi:10.1111/add.14459

7. Strauss E SE, Spreen O. A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary. . Oxford University Press; 2006;

8. Hernán MA BB, Robins JM. Marginal structural models to estimate the causal effect of zidovudine on the survival of HIV-positive men. Epidemiology 2000;Sep;11(5):561-70.):

9. Meier MH, Caspi A, A RK, et al. Long-Term Cannabis Use and Cognitive Reserves and Hippocampal Volume in Midlife. Am J Psychiatry 2022;179(5):362-374, doi:10.1176/appi.ajp.2021.21060664