

Swiss recommendations for the diagnosis, management and follow-up of post-COVID condition in primary care medicine (2023)

Mayssam Nehme^a, Lara Diem^b, Claudio L. A. Bassetti^b, Idris Guessous^{ac}, Working group Swiss post-COVID recommendations for primary care physicians

^a Division of Primary Care Medicine, Geneva University Hospitals, Geneva, Switzerland

^b Department of Neurology, Inselspital, University Hospital Bern, Bern, Switzerland

^c Faculty of Medicine, University of Geneva, Geneva, Switzerland

The COVID-19 pandemic and the post-infectious sequelae of SARS-CoV-2 have been major health concerns. The diagnosis of post-COVID condition is based on the WHO definition: “Post COVID-19 condition occurs in individuals with a history of probable or confirmed SARS-CoV-2 infection, usually 3 months from the onset of COVID-19 with symptoms and that last for at least 2 months and cannot be explained by an alternative diagnosis” [1]. Post-COVID condition affects approximately 9–22% of people after a SARS-CoV-2 infection, irrespective of the severity of the acute phase of the infection [2]. As of the year 2022, 3,022,992 people have been infected with SARS-CoV-2 in Switzerland [3]. Between 272,000 and 634,000 people in Switzerland are estimated to have been affected by post-COVID condition in 2022 alone.

Despite the large number of affected people and the recognition of post-infection syndromes in the literature [4, 5], knowledge of the underlying mechanisms, prognosis and management of these conditions is still limited. The pathogenesis of post-COVID condition is still unclear, but studies suggest possible immune dysfunction and inflammation, endothelial dysfunction leading to microthrombosis or persistence of viral particles [6]. The clinical picture of post-COVID condition is heterogeneous, including fatigue, difficulty concentrating, headaches, sleep disorders, autonomic dysfunction, pain, and respiratory, cardiac and gastrointestinal symptoms [2, 7]. The three most frequent symptoms are fatigue, difficulty concentrating and sleep disorders [7, 8]. In post-COVID condition, fatigue is characterized by post-exertional malaise (PEM) [9]. PEM refers to the worsening of symptoms after physical, mental or emotional exertion [10] and was reported in 8.2% of SARS-CoV-2–infected individuals 15 months after the initial infection [9]. Patients’ quality of life and functional capacity are impacted by these symptoms, leading to an increased cost to the individual and to society in general [11].

In 2022, the Federal Office of Public Health (FOPH), in coordination with the Federation of Swiss Doctors (FMH), introduced an initiative to draft recommendations for primary care physicians for the management of post-COVID condition. Primary care physicians are the cornerstone of management and follow-up of post-COVID condition, and

these recommendations aimed to help them in the diagnosis, management and follow-up of this disease. The recommendations for primary care physicians were based on an initial effort by the Division of Primary Care Medicine at the Geneva University Hospitals (HUG), in collaboration with several specialists at HUG, that was published in February 2021 and updated in November 2021 [12]. The authors recognized a need for national recommendations to ensure a more coordinated approach in the management of post-COVID condition. This was particularly important in the context of a condition that was not fully understood, and physicians had to avoid harm to the patient. Hence was born a collaborative effort to share experiences and lessons learned by Swiss hospitals and medical [8] and scientific authorities since the start of the pandemic in March 2020, leading to the creation of post-COVID recommendations for primary care physicians.

The Division of Primary Care Medicine at HUG and the Department of Neurology at Inselspital, the University Hospital of Bern, steered the sounding board and working groups and led the work on post-COVID recommendations for primary care physicians. Representatives from different fields of medicine were invited to participate. Fields included general internal medicine, neurology, pulmonary medicine, cardiology, psychiatry, infectious diseases, otorhinolaryngology, rheumatology, dermatology, gastroenterology, neuropsychology, physical therapy, occupational therapy (ergotherapy), Swiss Insurance Medicine and patient associations. Working groups helped define the content of the post-COVID recommendations and discuss different approaches applied in Switzerland. Working groups helped to generate a consensus on the recommendations, which were then discussed and validated by the sounding board. The sounding board gathered the representatives from the major scientific societies in Switzerland and from patient organizations. The final document has been validated by the major scientific societies whose disciplines are linked to the diagnosis and management of post-COVID condition.

To date, the management of post-COVID condition relies on the management of daily energy levels, the prevention of PEM and an interdisciplinary approach to help patients

Mayssam Nehme, MD
Service de médecine de
premier recours
Hôpitaux universitaires de
Genève (HUG)
Consultation Post-COVID
Rue Gabrielle Perret
Gentil 4
CH-1205 Genève
Mayssam.Nehme[at]
hcuge.ch

cope with their symptoms and find, when possible, ways to return and adapt to daily life. The role of primary care physicians is critical in the identification, diagnosis and follow-up of post-COVID condition. This document aimed to help physicians in understanding which symptoms could be due to post-COVID condition, to provide them with an approach to diagnosis after excluding other causes and to recommend strategies for the management of persistent symptoms and their impact on daily life. Primary care physicians can use this document, as well as its abridged version, to orient patients in their care pathway, to consider the suggested validated scales to identify and follow up on symptoms and their severity and to use the suggested approaches to treat the symptoms when possible, or at least to reduce their impact on daily life. Offering patients self-management tools, as well as tools to understand and cope with their disease, is also an essential component of management. Patients and physicians can find this information, as well as other resources for understanding and managing their symptoms, on www.rafael-postcovid.ch, an interactive platform with a medical chatbot, verified information and webinars for patients and healthcare professionals created by HUG in partnership with several hospitals, medical networks and national and international organizations. Information is also available from the Altea network (www.altea-network.ch), which is managed by an interdisciplinary team consisting of members of different fields of medicine, communication, social marketing, design and law. Patients can also benefit from patient-to-patient support and updated information on ongoing activities for post-COVID patients via the Long Covid Schweiz association (www.long-covid-info.ch).

Finally, these recommendations were possible thanks to the commitment and collaboration of all the experts who contributed to this work and to the support of the FOPH, the FMH and the scientific societies. While the future will likely bring more solutions for this condition, through ongoing collaborations [13], registers [14] and randomized controlled trials in Switzerland [15–17] and abroad [18], this work was essential for providing guidance given the current state of knowledge and hopefully will benefit a great number of primary care physicians, healthcare professionals and patients.

Acknowledgments

Working group Swiss post-COVID recommendations for primary care physicians: Allali Gilles, Antonini Pietro, Assal Frederic, Bassetti Claudio L.A., Baudet Corinne, Benzakour Lamyae, Bollag Yvonne, Britt Chantal, Brugger Silvio, Busche Philipp, Chmiel Corinne, Diem Lara, Di Gallo Alain, Eckerle Isabella, Finckh Axel, Frei Linda, Fretz Gregory, Frossard Jean-Louis, Funke-Chambour Manuela, Garzoni Christian, Guerreiro Ivan, Guessous Idris, Haller Dagmar M., Hersche Ruth, Kaiser Laurent, Lador Frederic, Landis Basile, Lauper Kim, L'huillier Arnaud G., McGuire Francis, Menouret Emmanuel, Meyer Philippe, Moreth Jens, Najjar Iris, Nehme Mayssam, Penner Iris-Katharina, Peron Julie, Perrin Anne, Posfay-Barbe Klara, Quinto Carlos, Sandor Peter, Schäffler Hilde, Schlunegger Michael, Schmidt-Leuenberger Joachim, Streit Sven, Spillman Nicole, Toutous-Trellu Laurence, Tschudi Andri, Vetter Pauline, Weber Pascal, Weil Barbara, Weise Andrea

Financial disclosure

This this work was funded by the Federal Office of Public Health.

Potential competing interests

All authors have completed and submitted the International Committee of Medical Journal Editors form for disclosure of potential conflicts of interest. No potential conflict of interest related to the content of this manuscript was disclosed.

References

- World Health Organization. A clinical case definition of post COVID-19 condition by a Delphi consensus. Last updated October 6, 2021 <https://apps.who.int/iris/bitstream/handle/10665/345824/WHO-2019-nCoV-Post-COVID-19-condition-Clinical-case-definition-2021.1-eng.pdf> [Access October 10, 2021].
- Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. *Nat Rev Microbiol*. 2023 Mar;21(3):133–46. <http://dx.doi.org/10.1038/s41579-022-00846-2>. PubMed. 1740-1534
- Bundesamt für Gesundheit: Coronavirus: Situation Schweiz; [Available from: <https://www.bag.admin.ch/bag/de/home/krankheiten/ausbrueche-epidemien-pandemien/aktuelle-ausbrueche-epidemien/novel-cov/situation-schweiz-und-international.html>]
- Stefano GB. Historical Insight into Infections and Disorders Associated with Neurological and Psychiatric Sequelae Similar to Long COVID. *Med Sci Monit*. 2021 Feb;27:e931447. <http://dx.doi.org/10.12659/MSM.931447>. PubMed. 1643-3750
- Choutka J, Jansari V, Hornig M, Iwasaki A. Unexplained post-acute infection syndromes [Erratum in: *Nat Med*. 2022 Aug;28] [8] [1723. PMID: 35585196]. *Nat Med*. 2022 May;28(5):911–23. <http://dx.doi.org/10.1038/s41591-022-01810-6>. PubMed. 1546-170X
- Nehme M, Ducrot A, Salmon D, Guessous I. Post-Covid: nouveautés 2022 et prochaines étapes [Post-Covid: 2022 updates and next steps]. *Rev Med Suisse*. 2023 Feb 1;19(812):160-166. French. doi: <http://dx.doi.org/10.53738/REVME.2023.19.812.160>. PMID: 36723639.
- Nehme M, Vetter P, Chappuis F, Kaiser L, Covicare Study Team, Guessous I. Prevalence of post-COVID Condition 12 Weeks after Omicron Infection Compared to Negative Controls and Association with Vaccination Status [in publication].
- Corinne C. MediX long COVID recommendations <https://www.medix.ch/wissen/guidelines/long-covid-fact-sheet/> [Last updated April 2023].
- Nehme M, Braillard O, Chappuis F, Covicare study team, Guessous I. The chronification of post-COVID condition associated with neurocognitive symptoms, functional impairment and increased healthcare utilization. *Scientific Reports* [Accepted, in publication].
- Centers for Disease Control and Prevention. Post-exertional malaise (PEM) [Available from: <https://www.cdc.gov/me-cfs/healthcare-providers/clinical-care-patients-mecfs/treating-most-disruptive-symptoms.html>]
- Bach K. New data shows long Covid is keeping as many as 4 million people out of work. *Brookings Institute*. Aug 24, 2022.
- Hôpitaux Universitaires de Genève. Prise en charge des patients souffrant de séquelles à long terme d'une infection au SARS-CoV-2. Février 2021 - mise à jour Novembre 2021 <https://www.hug.ch/sites/interhug/files/structures/coronavirus/guidelines-postcovid-29112021.pdf>
- Bassetti CL, Helbok R, Adorjan K, Falkai P. European Psychiatric Association-European Academy of Neurology statement on post-COVID syndrome. *Eur J Neurol*. 2023 Jan;30(1):294–5. <http://dx.doi.org/10.1111/ene.15572>. PubMed. 1468-1331
- Beghi E, Helbok R, Crean M, Chou SH, McNett M, Moro E, et al.; EAN Neuro-COVID Task Force. The European Academy of Neurology COVID-19 registry (ENERGY): an international instrument for surveillance of neurological complications in patients with COVID-19. *Eur J Neurol*. 2021 Oct;28(10):3303–23. <http://dx.doi.org/10.1111/ene.14652>. PubMed. 1468-1331
- Clinicaltrials.gov NCT05497089 trial. Temelimab as a Disease Modifying Therapy in Patients With Neuropsychiatric Symptoms in Post-COVID 19 or PASC Syndrome <https://clinicaltrials.gov/ct2/show/NCT05497089> [Last accessed January 20, 2023].
- Clinicaltrials.gov NCT05118711 Sequelae of COVID-19 With Focus on Exercise Capacity and Underlying Mechanisms (COR-PHYS) <https://clinicaltrials.gov/study/NCT05118711> [Last accessed July 11, 2023].
- Clinicaltrials.gov NCT05890534 Pycnogenol® in Post-COVID-19 Condition (PYCNOVID) <https://classic.clinicaltrials.gov/ct2/show/NCT05890534> [Last accessed July 11, 2023].
- Ducrot A, Nehme M, Sum Yu W. RAFAEL team, Schneider F, Posday-Barbe K, Guessous I. Summary of clinical trials for post-COVID condition <https://www.rafael-postcovid.ch/sites/default/files/inline-files/ClinicalTrials-Version%20professionnels.pdf>