


Journal Pre-proof

Extracellular sodium regulates fibroblast growth factor 23 (FGF23) formation

Zsuzsa Radvanyi, Eun Jin Yoo, Palanivel Kandasamy, Adrian Salas-Bastos, Sophie Monnerat, Julie Refardt, Mirjam Christ-Crain, Himeka Hayashi, Yasuhiko Kondo, Jonathan Jantsch, Isabel Rubio-Aliaga, Lukas Sommer, Carsten A. Wagner, Matthias A. Hediger, Hyug Moo Kwon, Johannes Loffing, Ganesh Pathare



PII: S0021-9258(23)02508-5

DOI: <https://doi.org/10.1016/j.jbc.2023.105480>

Reference: JBC 105480

To appear in: *Journal of Biological Chemistry*

Received Date: 4 July 2023

Revised Date: 30 October 2023

Accepted Date: 16 November 2023

Please cite this article as: Radvanyi Z, Yoo EJ, Kandasamy P, Salas-Bastos A, Monnerat S, Refardt J, Christ-Crain M, Hayashi H, Kondo Y, Jantsch J, Rubio-Aliaga I, Sommer L, Wagner CA, Hediger MA, Kwon HM, Loffing J, Pathare G, Extracellular sodium regulates fibroblast growth factor 23 (FGF23) formation, *Journal of Biological Chemistry* (2023), doi: <https://doi.org/10.1016/j.jbc.2023.105480>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 THE AUTHORS. Published by Elsevier Inc on behalf of American Society for Biochemistry and Molecular Biology.

CRedit author statement

Ganesh Pathare: Conceptualization, Supervision and Funding acquisition; **Zsuzsa Radvanyi, Eun Jin Yoo, Palanivel Kandasamy, Sophie Monnerat, Julie Refardt, Mirjam Christ-Crain** and **Ganesh Pathare:** Methodology; **Adrian Salas-Bastos:** RNA-seq data analysis; **Himeka Hayashi, Yasuhiko Kondo, Jonathan Jantsch** and **Isabel Rubio-Aliaga:** Resources; **Zsuzsa Radvanyi, Eun Jin, Lukas Sommer, Carsten A. Wagner, Matthias A. Hediger, Hyug Moo Kwon** and **Johannes Loffing** and **Ganesh Pathare:** Formal analysis; **Zsuzsa Radvanyi, Eun Jin** and **Ganesh Pathare:** Writing - Original Draft; **Lukas Sommer, Carsten A. Wagner, Matthias A. Hediger, Hyug Moo Kwon** and **Johannes Loffing:** Writing- Reviewing and Editing.

Extracellular sodium regulates fibroblast growth factor 23 (FGF23) formation.

Zsuzsa Radvanyi^{1,2†}, Eun Jin Yoo^{3†}, Palanivel Kandasamy⁴, Adrian Salas-Bastos¹, Sophie Monnerat^{5,6}, Julie Refardt^{5,6}, Mirjam Christ-Crain^{5,6}, Himeka Hayashi⁷, Yasuhiko Kondo⁷, Jonathan Jantsch^{8,9}, Isabel Rubio-Aliaga^{10,2}, Lukas Sommer¹, Carsten A. Wagner^{10,2}, Matthias A. Hediger⁴, Hyug Moo Kwon³, Johannes Loffing^{1,2}, Ganesh Pathare^{1,2*}

¹Institute of Anatomy, University of Zurich, Zurich, Switzerland; ²Swiss National Centre of Competence in Research “Kidney Control of Homeostasis”, Switzerland; ³Department of Biological Sciences, Ulsan National Institute of Science and Technology, Ulsan, Republic of Korea; ⁴Membrane Transport Discovery Lab, Department of Nephrology and Hypertension and Department of Biomedical Research, Inselspital, University of Bern, Bern, Switzerland; ⁵Department of Endocrinology, Diabetology and Metabolism, University Hospital Basel, Basel, Switzerland; ⁶Department of Clinical Research, University of Basel, Basel, Switzerland; ⁷Department of Animal Sciences, Teikyo University of Science, Yamanashi, Japan; ⁸Institute of Clinical Microbiology and Hygiene, University Hospital of Regensburg and University of Regensburg, Regensburg, Germany; ⁹Institute for Medical Microbiology, Immunology, and Hygiene, and Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany; ¹⁰Institute of Physiology, University of Zurich, Zurich, Switzerland

†contributed equally to this work

***Correspondence author:** Ganesh Pathare, Institute of Anatomy, University of Zurich, Winterthurerstrasse 190, 8057, CH, Email: ganesh.pathare@anatomy.uzh.ch

Keywords: FGF23, extracellular-sodium, hyponatremia, NFAT5, bone and kidney

Competing Interests

The authors declare no potential conflicts of interest.

ABSTRACT

The bone-derived hormone fibroblast growth factor-23 (FGF23) has recently received much attention due to its association with chronic kidney disease and cardiovascular disease progression. Extracellular sodium concentration ($[Na^+]$) plays a significant role in bone metabolism. Hyponatremia (lower serum $[Na^+]$) has recently been shown to be independently associated with FGF23 levels in patients with chronic systolic heart failure. However, nothing is known about the direct impact of $[Na^+]$ on FGF23 production. Here, we show that an elevated $[Na^+]$ (+20 mM) suppressed FGF23 formation, whereas low $[Na^+]$ (-20 mM) increased FGF23 synthesis in the osteoblast-like cell lines UMR-106 and MC3T3-E1. Similar bidirectional changes in FGF23 abundance were observed when osmolality was altered by mannitol but not by urea, suggesting a role of tonicity in FGF23 formation. Moreover, these changes in FGF23 were inversely proportional to the expression of NFAT5 (nuclear factor of activated T cells-5), a transcription factor responsible for tonicity-mediated cellular adaptations. Furthermore arginine vasopressin (AVP), which is often responsible for hyponatremia, did not affect FGF23 production. Next, we performed a comprehensive and unbiased RNA-seq analysis of UMR-106 cells exposed to low vs. high $[Na^+]$, which revealed several novel genes involved in cellular adaptation to altered tonicity. Additional analysis of cells with Crisp-Cas9 mediated NFAT5 deletion indicated that NFAT5 controls numerous genes associated with FGF23 synthesis, thereby confirming its role in $[Na^+]$ -mediated FGF23 regulation. In line with these in vitro observations, we found that hyponatremia patients have higher FGF23 levels. Our results suggest that $[Na^+]$ is a critical regulator of FGF23 synthesis.

SIGNIFICANCE STATEMENT

Our study emphasizes the critical involvement of extracellular-sodium-NFAT5 signaling in the regulation of FGF23 production. Elevated FGF23 levels are correlated with the development of left ventricular hypertrophy, while FGF23 deficiency in murine models simulates age-related diseases. Moreover, FGF23 exhibits potential as a biomarker for conditions such as chronic kidney disease and cardiovascular disorders, although its exact role in disease progression remains elusive. Consequently, investigating the regulation of FGF23 production holds significant importance

INTRODUCTION

Fibroblast growth factor-23 (FGF23) was discovered as markedly elevated 'phosphatonin' in patients with autosomal dominant hypophosphataemic rickets (ADHR)¹. High levels of FGF23 were also found in X-linked hypophosphatemia (XLH), which is caused by inactivating mutations in the *PheX* gene^{2,3}. Eventually, it was found that FGF23 regulates phosphate and vitamin D homeostasis by inhibiting renal sodium phosphate cotransporters and suppressing the process of vitamin D biosynthesis, respectively⁴. As a result, disruption of the FGF23 endocrine axis plays a key role in the pathophysiology of renal and bone disorders as well as aging⁴. In recent years, FGF23 has gained significant interest due to its strong association with poor prognosis in chronic kidney disease (CKD) and cardiovascular disease (reviewed in⁵). In mice, high FGF23 induces left ventricular hypertrophy⁶. FGF23 is mainly produced by osteoblasts and osteocytes. Upon secretion, it undergoes cleavage, leading to the presence of both intact (iFGF23) and C-terminal fragments (cFGF23) in the circulation⁷. Dietary phosphate^{8,9}, vitamin D⁸, insulin¹⁰, volume-regulation^{11,12}, aldosterone^{13,14}, iron status¹⁵ and inflammation¹⁶ have been identified as endogenous regulators of FGF23 in bone. Dietary salt intake, which does not impact plasma Na⁺ levels, is thought to regulate FGF23 through aldosterone^{13,14,17}. Furthermore, mice lacking Klotho (a co-receptor essential for FGF23 actions) exhibited an extended lifespan when they were fed a high-salt diet, while reduced salt intake had a modest effect on survival¹⁸. However, nothing is known regarding the direct role of extracellular sodium ion concentration ([Na⁺]) or osmolality in regulating FGF23 synthesis.

Both hyponatremia (serum [Na⁺] <135 mM) and hypernatremia (serum [Na⁺] >145 mM) can affect bone remodeling. Several studies have provided compelling evidence linking hyponatremia to bone loss, osteoporosis, and heightened bone fragility¹⁹⁻²⁴. Most hyponatremia cases are due to the syndrome of inappropriate antidiuretic hormone secretion (SIAD), which is characterized by high arginine vasopressin (AVP) levels (reviewed in²⁵). On a mechanistic level, both high AVP^{26,27} and low [Na⁺] levels^{19,28} are postulated to promote osteoclastogenesis and inhibit osteoblastogenesis, leading to bone loss in hyponatremia. Correction of hyponatremia in hospitalized patients has a positive impact on osteoblast function^{29,30}. On the other hand, high [Na⁺] may also enhance bone loss through an increase in osteoclastic resorption³¹. When cells were exposed to high [Na⁺], the expression of the RANKL decoy receptor

osteoprotegerin (*Opg*) increased in both osteoclast-precursor cells and osteoblasts³², suggesting a direct effect of elevated $[Na^+]$ on bone.

The intracellular milieu is almost immediately affected by changes in $[Na^+]$ or osmolality. Within cells, osmoregulation is mainly governed by the tonicity-responsive transcription factor tonicity-responsive enhancer-binding protein (TonEBP), also called the nuclear factor of activated T cells 5 (NFAT5)³³. Hypertonic conditions induce the upregulation of NFAT5, leading to the transcription of numerous NFAT5 target genes associated with adapting to high $[Na^+]$ ³³. Nevertheless, NFAT5 remains active even under isotonic conditions and can be either upregulated or downregulated in response to changes in tonicity³⁴. A recent genome-wide association study (GWAS) on plasma $[Na^+]$ concentration identified genetic variants in NFAT5³⁵. This suggests that NFAT5 may participate in the regulation of systemic $[Na^+]$ / water balance.

A recent study has shown that hyponatremia is independently associated with FGF23 levels in patients with chronic systolic heart failure³⁶. Given the important role of $[Na^+]$ in osteoblast functions, we hypothesized that the altered $[Na^+]$ levels may regulate the production of FGF23 through NFAT5. By manipulating culture media, we studied whether altered $[Na^+]$ mediated tonicity or osmolality, or both were responsible for FGF23 secretion by osteoblasts. Additionally, we studied the regulation of FGF23 by high AVP, which is often observed in hyponatremic patients.

RESULTS

Elevated $[Na^+]$ mediated hypertonicity suppresses FGF23 formation.

NaCl, as an impermeable solute, creates a hypertonic environment when its extracellular levels increase. NFAT5 is the key transcription factor involved in the adaptation to hypertonicity³³. As demonstrated in Fig. 1A/B/C, +NaCl (20 mM) and mannitol (40 mM), significantly increased NFAT5 protein and mRNA expression in the rat osteoblast-like cell line UMR-106. However, 40 mM urea did not affect NFAT5 protein/mRNA expression since it elevates osmolality rather than tonicity via cell membrane permeability. The osmolality of normal DMEM cell culture media was 300.7 ± 2.0 mOsm/Kg. There was an almost similar increase in osmolality after the addition of +NaCl, mannitol, and urea (Table S1). Importantly, the total

FGF23 (cFGF23) in the cell supernatant was markedly suppressed by +NaCl and mannitol, but not by urea (Fig. 1D). Similarly, in Fig. 1E, *Fgf23* mRNA in cells was also significantly suppressed by +NaCl and mannitol, but not by urea. iFGF23 was not detected due to its very low levels in the cell supernatant (not shown). Consistent with previous reports³², we confirmed that +NaCl concentration up to ~80 mM does not affect cell survival (Fig. 1F/S4). The +NaCl effect on both *Fgf23* and *Nfat5* mRNA was concentration-dependent, with as little as 10 mM +NaCl significantly suppressing *Fgf23* mRNA (Fig. 1G). Time-course experiments showed that *Nfat5* mRNA peaked at 8 h of high-NaCl treatment, while the nadir of *Fgf23* mRNA levels was observed at 24 h (Fig. 1H). The impact of high NaCl on FGF23 was further replicated in MC3T3-E1 cells. After 6 days of differentiation in osteogenic media, these cells were subjected to a 24-hour treatment with NaCl, mannitol, and urea. High NaCl, mannitol, and urea treatments induced similar changes in *Fgf23* and *Nfat5* mRNA levels as observed in UMR-106 cells (Fig. S1A/B). As a negative control, cells cultured in non-osteogenic media did not exhibit any *Fgf23* mRNA expression.

Hypotonicity increases FGF23 production.

We explored the possibility that hyponatremia might have a direct effect on FGF23 formation. To mimic hyponatremia, cells were cultured in low $[Na^+]$ media. To generate cell culture media with low $[Na^+]$ concentrations, custom-made NaCl-free media was reconstituted with specific amounts of NaCl to obtain media in which $[Na^+]$ were -5, -10, -15, and -20 mM lower than in control media. As shown in Fig. 2A, we found that reducing $[Na^+]$ in the culture media increased *Fgf23* mRNA levels in a dose-dependent manner. To understand the underlying mechanisms by which low $[Na^+]$ stimulates FGF23 formation, we manipulated the low $[Na^+]$ media (-20 mM) with and without correction of osmolality by the addition of 40 mM mannitol or 40 mM urea. As shown in Table S2, the osmolality of low $[Na^+]$ medium was 260.7 ± 1.5 mOsm/kg. There was an almost similar correction in osmolality after the addition of mannitol and urea (~302 mOsm/Kg). Interestingly, the addition of mannitol to the low $[Na^+]$ medium reversed the increase in FGF23; however, the addition of urea did not affect FGF23 formation. This effect was observed at both mRNA and protein levels measured from cells and cell supernatants, respectively (Fig 2B/C). These findings suggested that low tonicity rather than low osmolality was responsible for the increase in FGF23 upon reducing $[Na^+]$ concentrations. Expectedly, upon reducing $[Na^+]$ concentration, *Nfat5* mRNA was significantly lower, the

effect, again, was reversed by mannitol but not urea (Fig. 2D). Similar changes were observed in MC3T3-E1 cells (Fig. S2).

No evidence for FGF23 regulation by AVP.

Elevated AVP levels in hyponatremia could potentially affect osteoblast function through its receptors. AVP exerts major pathophysiological actions through three distinct receptor isoforms designated V_{1a} , V_{1b} , and V_2 . Both V_{1a} and V_2 receptors have been found in murine osteoblasts and osteoclasts, where they may take part in bone remodelling^{26,27}. To this end, we checked the expression pattern of AVP receptors in the rat osteoblast-like UMR-106 cell line. Endpoint PCR was performed on RNA isolated from rat liver (positive control for V_{1a} receptor), kidney (positive control for V_2 receptor), and UMR-106 cells. As shown in Fig. 3A, V_{1a} receptors were very abundant in the liver, while also expressed in kidney tissue as well as UMR-106 cells. However, V_2 receptors were mostly detected in kidney tissue with only slight expression in liver and UMR-106 cells. To further confirm that V_{1a} is the major AVP receptor expressed in UMR-106 cells, qRT-PCR was performed. As shown in Fig. 3B, V_{1a} mRNA levels were ~7 times higher than V_2 mRNA in UMR-106 cells. As expected, the pituitary specific- V_{1b} receptor³⁷ was undetectable in UMR-106 cells. Next, we treated UMR-106 cells with AVP (1-1000 nM) for 24 hours (Fig. 3C). However, this did not result in any significant change in *Fgf23* mRNA expression. Moreover, AVP (100 nM) applied for various time points between 3-24 h also did not affect *Fgf23* mRNA levels (Fig. 3D). We found that the main AVP receptor expressed in UMR-106 cells is V_{1a} ; however, AVP binds more strongly to V_2 than V_{1a} receptors in rats ($K_d=1.7$ nm for the V_{1a} ; $K_d=0.4$ nm for the V_2)³⁷. Therefore, next, we explored the effect of specific V_{1a} agonist ([Phe²]OVT, [Phe²,Orn⁸]vasotocin)³⁸ and V_2 agonist (ddAVP) on *Fgf23* synthesis. However, both V_{1a} agonist and V_2 agonist, at low and high concentrations, did not affect *Fgf23* production significantly, when treated for a duration of 24h (Fig. 3E/F). To investigate the impact of hyponatremia and high AVP together on FGF23 production, we exposed cells to a low-NaCl medium with a V_{1a} agonist or AVP. However, this combined treatment did not result in a significant alteration in FGF23 formation when compared to cells exposed to low-NaCl alone (Fig. 3G). Consistent with these *in-vitro* data, the lack of V_{1a} receptors in mice had no effect on FGF23 levels in the serum (Fig. 3H). Overall, based on these findings, we concluded that

although V_{1a} receptors are adequately expressed in UMR-106 cells, AVP does not modulate FGF23 formation.

Tonicity-mediated FGF23 regulation is NFAT5-dependent.

Our results show that elevated $[Na^+]$ levels (+20 mM) suppressed FGF23 formation, whereas low $[Na^+]$ levels (-20 mM) led to an increase in FGF23 synthesis. These bidirectional changes in FGF23 were inversely proportional to NFAT5 activity. In mice, *Fgf23* is expressed only in limited tissues such as the calvaria, spleen, and thymus, whereas *Nfat5* is almost ubiquitously expressed, including in the organs that produce *Fgf23* (Fig. S3). Therefore, NFAT5 may regulate FGF23 production. To test whether $[Na^+]$ mediated FGF23 changes were mediated by NFAT5 or not, we used the Crispr-Cas9 technology to knockout NFAT5 in UMR-106 cells (*NFAT5^{KO}*). A single-cell derived clone of *NFAT5^{KO}* was generated with a complete knockout, confirmed by immunoblotting (Fig. 4A) and qRT-PCR (Fig. 4B). Unlike in control cells, +NaCl treatment did not affect NFAT5 protein or mRNA expressions in *NFAT5^{KO}* cells (Fig 4A/B). Moreover, cellular damage measured by LDH activity was more prominent in *NFAT5^{KO}* cells than control cells, when treated with very high NaCl levels (Fig 4C). These observations confirmed and validated successful NFAT5 deletion in UMR-106 cells. Importantly, *Fgf23* mRNA measured by qRT-PCR showed blunted response to tonicity in *NFAT5^{KO}* cells compared with control cells (Fig. 4D). However, the tonicity response to *Fgf23* was not completely abolished in *NFAT5^{KO}* cells.

Next, we employed RNA-Seq to generate a comprehensive, unbiased survey of genes regulated by low vs. high $[Na^+]$ in UMR-106 cells in control and *NFAT5^{KO}* cells. The RNA-seq of control vs. *NFAT5^{KO}* cells revealed differential gene regulation upon -NaCl vs. +NaCl treatment as depicted in Venn diagrams (Fig. 4E). This suggested that transcriptionally active NFAT5 alters the expression of numerous genes in response to +NaCl (SI dataset 1/2). In the same line, in response to +NaCl treatment, control cells exhibited significant changes in a total of 5360 genes ($p < 0.05$), whereas *NFAT5^{KO}* cells showed alterations in 4097 genes (Fig. S5). Block 1/4/6 of the heatmap (Fig. 4F) summarizes the genes that exhibit significant up or downregulation in response to +NaCl treatment in control cells, while only slight or no alterations in *NFAT5^{KO}* cells. Therefore, we focused only on these unique sets of potential NFAT5 targets (SI dataset 3).

These NFAT5- target genes (n=2615) are further summarized in a volcano plot (Fig. 4G). *Slc14a2*, *Fxyd2*, *Sgk1*, *Pnpla6*, *Igf1bp7*, and *Itga10* are known NFAT5 targets³⁹, and our study confirmed this. Importantly, only control cells showed statistically significant downregulation of *Fgf23* after +NaCl treatment. The gene ontology (GO) analysis of the upregulated potential NFAT5 targets indicated the activation of ‘ossification’ and ‘bone-mineralization’ signaling pathways (Fig. 4H, SI dataset 4). After manually screening the genes involved in these pathways, we found that many of the identified genes regulate FGF23 and/or bone mineralization (Fig. 4I). Genes like *Phex*, *Mmp3*, *Nfkb1a*, *Foxo1*, and *Akt1* are known to regulate FGF23 formation⁴⁰. As proof of concept, we analyzed *Phex*, a known suppressor of FGF23, and found that *Phex* mRNA is significantly upregulated by the treatment of +NaCl and mannitol but not by urea (Fig. 4J). Interestingly NaCl-mediated upregulation of *Phex*, as confirmed by qRT-PCR, was not observed in *NFAT5^{KO}* cells (Fig. S6) confirming it's a tonicity or NFAT5 target. However, other known regulators of FGF23 such as *Ibsp*, *Enpp1*, *Ankh*, *Pth1r*, *Dmp1*, *Fgfr1*, and *Sost*, were altered in both control and *NFAT5^{KO}* cells (Fig. S7), suggesting NFAT5-independent mechanism in FGF23 regulation by [Na⁺]. Fig. S8 summarized potential NFAT5-target genes, which were previously postulated in [Na⁺] sensing and [Na⁺] homeostasis. Fig. S9 and S10 illustrate the analysis of all the KEGG and GO pathways, respectively.

Hyponatremia patients have higher FGF23 levels.

To test for the potential relevance of our findings *in vitro* for the situation *in vivo*, we investigated in a pilot study the levels of FGF23 in hyponatremic patients (with GFR>60 ml/min) and compared them to the values of healthy subjects. The description of matching healthy controls and patients, including the etiology of hyponatremia, is provided in Table S3 and S4 respectively. Serum [Na⁺] levels in hyponatremic patients were significantly lower than in the control group (Fig. 5A). Serum cFGF23 and iFGF23 isoforms were measured by ELISAs. As shown in Figure 5B, hyponatremic patients exhibited a significant increase in serum cFGF23 levels. Although there was a trend of increased serum iFGF23 in hyponatremic patients, the difference was not statistically significant (Fig. 5C). To evaluate the impact of altered FGF23 levels on phosphate homeostasis in these patients, we measured serum phosphate (Pi) levels. However, as shown in Fig. 5D, no significant difference was observed in serum phosphate levels between the control and hyponatremic patients. To perform correlation analysis, we plotted serum [Na⁺] against serum cFGF23 and

iFGF23. Both cFGF23 and iFGF23 demonstrated a negative correlation with serum $[\text{Na}^+]$, but only cFGF23 exhibited a statistically significant correlation with $[\text{Na}^+]$ (Fig. 5E). Prior studies suggested that some of the hyponatremia patients may have higher inflammatory cytokines (reviewed in⁴¹). At least in theory, this could potentially account for elevated FGF23 levels. To this end, we measured $\text{TNF}\alpha$ and IL6 levels in hyponatremia patients (Fig. S11). Nonetheless, there were no significant differences in these cytokines between hyponatremic patients and healthy subjects.

DISCUSSION

Our experiments demonstrate that pathophysiologically relevant changes in $[\text{Na}^+]$ directly regulate FGF23 formation by osteoblasts. By manipulating the culture media using $+\text{NaCl}$, mannitol, or urea we showed that the cellular adaptation to tonicity, rather than osmolality, plays a significant role in regulating FGF23 levels upon change in $[\text{Na}^+]$. Additionally, comprehensive RNA-seq analysis and NFAT5 knockdown experiments defined the role of the osmo-sensitive transcription factor NFAT5 in FGF23 regulation. We found no evidence that AVP, which increases in hyponatremia, regulates FGF23 production in the osteoblast cell line UMR-106.

We found that FGF23 regulation by tonicity is bidirectional, i.e. high $[\text{Na}^+]$ suppresses FGF23 while low $[\text{Na}^+]$ elevates FGF23 production, which is dependent upon NFAT5 activity. Under isotonic conditions, NFAT5 is both cytosolic and nuclear^{34,42}. With an increase in tonicity, it translocates to the nucleus, initiating the activation of a group of genes responsible for protecting cells against osmotic stress. This activation occurs through NFAT5 binding to a specific regulatory element called the osmotic response element or the tonicity-responsive enhancer, which is located in the promoter region of these genes⁴³⁻⁴⁵. As a result, a tonicity-dependent increase in nuclear NFAT5 is proposed as the key event in driving its target genes. While hypertonic stress promotes nuclear translocation and activates enhancer activity of NFAT5, hypotonicity induces its nuclear export supporting bidirectional regulation in response to tonicity changes^{34,42,46}. The c-terminus of NFAT5 also contains a transactivation domain (TAD) and its activity varies directly with $[\text{Na}^+]$ concentration (low to high)⁴⁶. NFAT5 regulates numerous gene expressions through its nuclear translocation during hypertonicity and nucleocytoplasmic export during hypotonicity. The $+\text{NaCl}$ treatment

in UMR-106 cells likely triggered a transcriptomics response via NFAT5, potentially impacting FGF23 regulation. Our study identified previously known genes that regulate FGF23 and bone mineralization as potential novel targets of NFAT5. For instance, *Phex* was one of the earliest genes identified to inhibit FGF23 formation^{2,3}. Now, we found that *Phex* is a NFAT5 target and gets remarkably upregulated by tonicity. Although our results demonstrated that FGF23 formation depends upon tonicity and NFAT5, the exact mechanism responsible for this phenomenon is unclear. Moreover, the tonicity response to *Fgf23* was not completely abolished in *NFAT5^{KO}* cells, suggesting that NFAT5-independent mechanisms do also contribute to the regulation of *Fgf23* by $[Na^+]$.

Osmoregulation and $[Na^+]$ sensing mechanisms were mainly studied for cells in the hypothalamus (reviewed in⁴⁷). In line with studies from others^{29,30,32}, we found that osteoblasts can sense osmotically relevant changes in $[Na^+]$ and respond with an altered expression of several genes including *Fgf23*. Studies in RAW264.7 pre-osteoclastic cells showed increased osteoclast formation and resorptive activity in response to low $[Na^+]$ concentration^{20,21}. The molecular mechanism by which $[Na^+]$ is detected by bone cells is unknown. The expression of the osmoprotective transcription factor NFAT5 is regulated by local $[Na^+]$ content in many organs^{48,49}. Here, we demonstrate that the expression of this transcription factor in UMR-106 osteoblast-like cells is regulated by $[Na^+]$. Previous studies focused on the effect of major changes in $[Na^+]$ (e.g. 50-100 mM NaCl hypo- and hypertonic conditions) on NFAT5 regulation^{33,34,42}. However, such drastic changes of $[Na^+]$ occur only in the renal medulla, but not in bone. Now, our findings suggest that even differences of 20-40 mM NaCl significantly regulate NFAT5 activity in osteoblasts and osteoclasts, which is consistent with recent other studies^{21,32}. Serum/glucocorticoid regulated kinase 1 (*Sgk1*)⁵⁰, transient receptor potential vanilloid type 4 (*Trpv4*)⁵¹, Na^+/K^+ /ATPase (*Atp1a1*) and its associated protein such as sucrose non-fermenting-1-related serine/threonine kinase-SIK1 (*Snrk*)⁵² were previously postulated to take part in $[Na^+]$ sensing in various cells. Secondly, Na^+/H^+ exchanger activity⁵³ (*Slc9a1*), and Na^+/Ca^{2+} exchanger activity⁵⁴ (*Slc8a1*) have been implicated in $[Na^+]$ homeostasis in bone cells. Our RNA-seq data show that, except for TRPV4, all these genes were significantly upregulated in control cells but not in *NFAT5^{KO}* cells upon +NaCl treatment, indicating that these are NFAT5 targets (Fig. S8). Based on our findings and considering the role of NFAT5 as a master regulator of tonicity, it is tempting to

speculate that changes in NFAT5 activation may be one of the earliest events in osteoblast adaptation to $[\text{Na}^+]$ levels. Therefore, NFAT5 likely plays a crucial role in $[\text{Na}^+]$ sensing in osteoblasts.

Our findings revealed that AVP does not take part in FGF23 formation. Most patients with chronic hyponatremia exhibit elevated levels of AVP that are inconsistent with the osmotic balance, even when SIAD is not the underlying cause of hyponatremia^{55,56} (reviewed in⁵⁷). The research group of Zaidi conducted seminal studies that have established a primary role for AVP signaling in bone mass regulation. Mice injected with AVP exhibited reduced osteoblast formation and increased osteoclast formation via the V_{1a} receptor and ERK signaling^{26,27} (reviewed in^{58,59}). Conversely, mice injected with the V_{1a} antagonist SR49059 or genetic V_{1a} deficiency showed enhanced bone mass²⁶. In contrast, the V_2 receptor did not play a significant role in bone²⁷. In line with these studies, we observed a significant expression of V_{1a} receptors but only a minimal expression of V_2 receptors in UMR-106 cells. However, there was no evidence that AVP, V_{1a} agonist and V_2 agonist regulate FGF23 production in UMR-106 cells. Consistently, serum FGF23 levels in V_{1a} KO mice were unchanged when compared to wild-type mice. These findings support a direct effect of $[\text{Na}^+]$ on FGF23 formation in hyponatremia. For instance, Verbalis *et.al.*^{19,60} employed the SIAD model, where chronic hyponatremia induced by ddAVP and water loading promoted water retention. When water loading was not performed, the group of animals did not develop hyponatremia, and their bone mass did not show a significant reduction compared with the hyponatremia group. Moreover, cultured cells subjected to low $[\text{Na}^+]$ concentration exhibit increased bone resorption pathways and elicited gene expression changes driving osteoclast differentiation and functions^{20,21}, implying a direct effect of low $[\text{Na}^+]$ in bone loss. Overall, it seems that increased AVP levels play a significant role in bone loss among certain SIAD patients, but it may not take part in regulating FGF23 production.

We found a positive association between elevated FGF23 levels and hyponatremia in a pilot study conducted in humans. However, it remains unclear whether the increased FGF23 levels in hyponatremic patients are primarily influenced by low $[\text{Na}^+]$ and/or by the underlying cause of hyponatremia. While elevated inflammatory cytokines in hyponatremia patients⁴¹ could potentially increase FGF23, we did not detect significantly higher cytokine levels in the patient samples. Previously it was shown that in multivariate

linear regression analysis, hyponatremia is independently associated with FGF23 level in patients with chronic systolic heart failure³⁶. This further suggests that FGF23 elevation in hyponatremic patients might be due to low $[Na^+]$ levels.

A strength of our study is unfolding the novel role of $[Na^+]$ and NFAT5 in FGF23 production using two osteoblast cell lines. Nonetheless, we would like to acknowledge the limitations. Firstly, the precise mechanism by which NFAT5 governs the regulation of FGF23 is currently unclear. Secondly, the human study is constrained by the limited number of patients available for investigating the impact of hyponatremia on FGF23 levels. This may have precluded observing statistically significant changes in iFGF23. A larger sample size with hyponatremic patients will be necessary to confirm and unravel the possible relationship between $[Na^+]$ and FGF23 levels in humans. In conclusion, our study identified a crucial signaling between $[Na^+]$, NFAT5, and FGF23 formation, warranting further investigation to confirm its clinical significance.

EXPERIMENTAL PROCEDURES

All the chemicals were purchased from Sigma Aldrich unless otherwise stated.

Culture of UMR106 cells.

A rat osteoblastic cell line UMR-106 was originally purchased from ATCC (CRL-1661) and cultured at 5% CO_2 in a growth medium consisting of Dulbecco's Modified Eagle's Medium DMEM, low glucose, GlutaMAX™ (gibco; Cat. #21885-025) supplemented with 10% heat-inactivated FBS (Amimed; Cat. #2-01F30-I; lot nr: F07156P) and 100 U/ml penicillin, and 100 μ g/ml streptomycin (gibco; Cat. #15140122). A single lot of FBS was utilized throughout the entire study. Cells between passages 4 and 10 were used in this study. When the cells reached near-confluency on 6-well plates, the cell medium was replaced with the experimental medium (2 ml/well) consisting of DMEM containing 1 nM of 1,25-dihydroxyvitamin D3 (Tocris; Cat. #2551), as described previously⁶¹. The cells were further treated with or without +NaCl, mannitol, or urea for specified concentrations and durations. For low-NaCl treatments, custom-made NaCl-free cell culture media was ordered commercially (Biotechne, Cat. #CUST07ATLB). The media formulation was developed to ensure that all the other components of the media were exactly similar to DMEM (gibco; Cat. # 21885-025), with the sole exception of the NaCl content. Thereafter NaCl was added to the medium to

prepare a hypotonic medium to study the effect of low $[\text{Na}^+]$ concentration. The osmolality of all the culture media were measured on a Vapro 5600 (Wescor, Logan, UT) vapor pressure osmometer⁶². AVP (Tocris Cat. #2935) and ddAVP (Tocris; Cat. #3396) were freshly prepared by dissolving them in water and treated for a specified duration and concentration. The V_{1a} agonist ($[\text{Phe}^2]\text{OVT}$, $[\text{Phe}^2, \text{Orn}^8]\text{vasotocin}$) was a kind gift from Prof. Maurice Manning, The University of Toledo, USA. Its pharmacological properties in rat bioassay are reported earlier in a review³⁸ by Manning *et.al*.

For FGF23 measurement in cell media, cells were cultured in a 100 mm petri dish. After 24 h of treatment with +NaCl, mannitol, and urea, cell media was concentrated using Pierce™ Protein Concentrator PES, 10K MWCO (Thermo Fisher; Cat. #88527). The resultant concentrate (150 μl) was used to measure c-terminal FGF23 by ELISA (Quidel; Cat. #60-6300).

Culture of MC3T3-E1 cells

MC3T3-E1 Subclone 4 mouse pre-osteoblast cells were maintained in α -Minimum Essential Medium (α -MEM) supplemented with nucleosides, 2 mM L-glutamine (Thermo Fisher, Cat. #22571038), and 10% FBS, with 100 U/ml penicillin and 100 $\mu\text{g}/\text{ml}$ streptomycin. Cells at passages 4 to 10 were seeded onto 12-well plates and cultured for 24 hours. Osteogenic differentiation was initiated by culturing the cells in a medium containing 50 $\mu\text{g}/\text{ml}$ ascorbic acid and 10 mM β -glycerophosphate for 6 days. Cells were then subjected to treatment with high NaCl, mannitol, or urea for another 24 hours in the presence of calcitriol (10 nM). In experiments pertaining to low-NaCl treatment, following a 6-day differentiation in α -MEM, the cells were subjected to 24 hr treatment with sodium-free DMEM (with ascorbic acid, β -glycerophosphate and calcitriol) containing the specified amounts of NaCl, mannitol, or urea, as detailed in the results section.

Animal experiments

To check *Fgf23* and *Nfat5* mRNA expression, organs were harvested from C57BL/6J wildtype mice (male, 8-12 weeks old) according to Swiss law and were approved by the veterinary administration of the Canton of Zurich (Kantonales Veterinäramt), Switzerland. The V_{1a} receptor KO mice were generated as previously reported⁶³. The experiments with V_{1a} KO mice were conducted at the animal husbandry facility of the Teikyo

University of Science. Around 400 μ l of blood was drawn from 8-9 weeks old WT and V_{1a} receptor KO male littermates. The intact-FGF23 and c-terminal FGF23 were measured by ELISAs (Quidel, Cat. #60-6800, Quidel, Cat. #60-6300 respectively) according to the manufacturer's instructions.

LDH and MTT assay:

UMR-106 cells (control and $NFAT5^{KO}$ cells) were plated in a 12-well plate and cultured for 24 hr. Next, different concentrations of +NaCl were added to the culture media for a further 24 h. LDH was measured on aliquots of the +NaCl-treated cellular supernatant using the CyQUANT™ LDH Cytotoxicity Assay Kit (Thermo Fisher; Cat. #C20300). LDH activity was measured by measuring absorbance at 490 nm. The LDH release in +NaCl-treated cells is expressed in arbitrary units normalized to untreated cells. For MTT assay, UMR-106 were plated in 96-well plate for 24 hr followed by 24 hr NaCl treatment. The CyQUANT™ MTT cell viability assay (Thermo Fisher, Cat. #V13154) was performed according to the manufacturer's instructions. Absorbance measurements at 570 nm were made using a microplate reader.

Immunoblotting

Cells were homogenized in ice-cold RIPA lysis buffer (Thermo Fisher, Cat. #89900). Then cell lysates were centrifuged for 10 min at 2000 g and supernatants were used for immunoblotting. Protein concentration was measured by Bradford assay (CooAssay Protein Dosage Reagent; Uptima, Cat. #UPF86421). Equal amounts of protein (30 μ g) were loaded in Laemmli buffer (pH 6.8) on 8% polyacrylamide gels. Electrophoretically separated proteins were blotted to nitrocellulose membranes at 100 V for 2 hours. Next, membranes were blocked in Odyssey blocking buffer (LI-COR; Cat. # 927-70001) for 1 hour and incubated with the diluted NFAT5 primary antibody (Thermo Fisher; Cat. #PA1-023) and β -actin (Santa Cruz; Cat. #sc-47778X) overnight at 4°C. On the following day, secondary antibodies -goat-anti-rabbit IRDye 800 (LI-COR; Cat. # 926-32211), goat-anti-mouse IRDye 680 (LI-COR; Cat. # 926-68070) were added to the membranes in Casein Blocking solution in deionized water (1:10) and incubated for 1 hour at room temperature. Then membranes were repeatedly washed with phosphate-buffered saline containing 0.1% Tween 20. The fluorescent signal was visualized using Odyssey IR imaging system (LI-COR Biosciences). Optical densities were quantified with ImageJ S5 and were normalized to β -actin.

qRT-PCR.

The RNA was extracted from UMR-106/MC3T3-E1 cells and mouse tissues using a Nucleospin RNA isolation kit (Macherry Nagel; Cat. #740955). Rat tissue RNA was isolated as described in our previous study⁶⁴. The RNA was transcribed using a High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher; Cat. #4374966) and subjected to qPCR with SybrGreen Master Mix (Roche; Cat. #4707516001) using primers (Table 1). The relative quantification of gene expression based on double-delta Ct (threshold cycle) analysis was performed after normalization to *Tbp* expression.

Generation NFAT5 knockout in UMR106 cells by CRISPR/Cas9

Single guide RNAs (sgRNAs) specifically targeting *Rattus norvegicus* NFAT5 at exon 1-4 were designed using the CRISPR designing tool (<https://www.benchling.com/crispr>). All sgRNA primers were synthesized by Microsynth AG, Balgach, Switzerland. sgRNA primers were (Table 2) cloned into BbsI-HF linearized pU6-(BbsI)_CBh-Cas9-T2A-mCherry (Addgene, Plasmid Cat. #64324) or pX330-U6-Chimeric_BB-CBh-hSpCas9 (Addgene, Plasmid Cat. #42230). Sanger sequencing (Microsynth AG) was used to verify the correct cloning of shRNA into the vectors. UMR106 cells were plated at 4×10^5 per well in a 6-well plate and transfected with 2 μ g of NFAT5 gRNAs per well using the FuGENE[®] HD Transfection Reagent (Promega, USA). 24 h post-transfection, cells were trypsinized and single cells were sorted into a 96-well plate by FACS (BD BioSciences). A total of five clones derived from single-cell were propagated. NFAT5 knockout was confirmed by immunoblotting and qPCR. Following primers were used to confirm *Nfat5* deletion in *NFAT5^{KO}* cells by qPCR: *Nfat5-KO* Fwd: GCCCTCGGACTTCATCTCAT; *Nfat5-KO* Rev: ACAGATTCTTCCAATAGTCCAGC

RNA-sequencing and data analysis

Control and *NFAT5^{KO}* cells were cultured as explained earlier. Cells were treated with low-NaCl and high-NaCl (difference of 40 mM NaCl) for 24 h. RNA was isolated using a Nucleospin kit as explained earlier. RNA-seq was performed commercially by Novogene (UK) Company Limited. In brief, amplified cDNA samples were subjected to different quality control standards. RNA sample was used for library preparation using NEB Next[®] Ultra RNA Library Prep Kit for Illumina[®]. Indices were included to multiplex multiple

samples. Briefly, mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. After fragmentation, the first strand cDNA was synthesized using random hexamer primers followed by the second strand cDNA synthesis. The library was ready after end repair, A-tailing, adapter ligation, and size selection. After amplification and purification, the insert size of the library was validated on an Agilent 2100 and quantified using quantitative qPCR. Libraries were then sequenced on Illumina NovaSeq 6000 S4 flow cell with PE150 according to results from library quality control and expected data volume.

We included 12 datasets in the RNA-Seq data analysis: three biological replicates in each group (control with -NaCl, control with +NaCl, *NFAT5^{KO}* with -NaCl, and *NFAT5^{KO}* with +NaCl). The fold changes were calculated by dividing the arithmetic mean of the normalized read counts by the number of replicates. The statistical significance of differential gene expression in RNA-Seq data was determined with *p* value <0.05. The supporting datasets S1 and S2 contain all of the genes identified upon +NaCl treatment in control and *NFAT5^{KO}* cells respectively. While supporting datasets S3 contains unique genes, which were either significantly upregulated or downregulated in control cells but unchanged in *NFAT5^{KO}* cells upon +NaCl treatment. Venn diagrams for the sets of genes up- or down-regulated were made by intersecting the lists of gene names between control and *NFAT5^{KO}* cells (R package VennDiagram version 1.71). The heatmap of DEGs was made with the R package ComplexHeatmap (version 2.4.3) using as input the full set of genes differentially expressed between the normal and +NaCl conditions (*p* value < 0.05) for both control and *NFAT5^{KO}* cells. The input expression matrix was normalized by row (gene) by computing a z-score. Both experimental groups (columns) and gene expression profiles (rows) were clustered using the euclidean distance and the hierarchical clustering algorithm. Gene set groups were obtained by cutting the dendrogram in eight slices with the cutree default implementation of the Heatmap function. Functional over-representation analysis of gene ontology biological processes was performed with the R package clusterProfiler (version 3.16.1) using as input the genes uniquely Up-regulated in the control cells upon +NaCl treatment (*p* value < 0.05) and considering a significance *q*-value cutoff of 0.05.

Clinical study

Samples of patients with hyponatremia were collected during a prospective multicentric observational study (the Co-MED study, NCT01456533)⁶⁵ conducted at the University Hospital of Basel Switzerland, and the Medical University Clinic Aarau, Switzerland, from June 2011 to August 2013. The human study reported in this report abide by the Declaration of Helsinki principles. Samples of healthy controls were collected during a prospective double-blind, placebo-controlled randomized crossover study (the DIVE Study, NCT02729766)⁶⁶ conducted at the University Hospital Basel, Switzerland, from March to June 2016. Venous blood samples were collected in the morning hours, between 8 am and 10 am. The local ethics committee (Ethic Committee of north-west Switzerland (EKNZ)) approved the study protocols and written informed consent was obtained from all study participants, including informed consent to further use of biologic material. Patients and healthy controls were matched based on age and BMI (n=6 each group). Detailed characteristics of the healthy controls and hyponatremia patients can be found in table S3 and S4, respectively. The serum electrolyte analysis [Na⁺] was performed immediately after collection on a ABL800FLEX blood gas analyzer (Radiometer, Thalwil, Switzerland). The serum was separated, and samples were stored at -80 °C until use. Commercial ELISA kits were employed to measure cFGF23 (Quidel, Cat. #60-6100), iFGF23 (Quidel, Cat. #60-6600), TNF α (Thermo Fisher, Cat. #900-K25) and IL6 (Thermo Fisher, Cat. #900-K16) in the serum according to the manufacturer's instructions. Serum phosphate was measured by QuantiChrom™ Phosphate Assay Kit (Bioassay systems Cat. #DIPI-500).

STATISTICS

All the gene or protein expression values are expressed as arithmetic means \pm SEM, where n represents number of independent experiments (biological replicates). An unpaired Student's t-test, one-way or two-way ANOVA was used for comparisons between the groups using GraphPad Prism. In cases where the p value is not mentioned, the following applies: ns (not significant) $p > 0.05$, * $p \leq 0.05$, ** $p < 0.01$, and *** $p < 0.001$.

ACKNOWLEDGMENTS

This work is supported by the Swiss National Science Foundation through the National Center of Competence in Research NCCR Kidney.CH. [grant number N-403-03-55 (to Ganesh Pathare)]. The authors sincerely thank Klaudia Kopper and Monique Carrel for their excellent technical assistance. Additionally, we express our gratitude to Prof. Maurice Manning from The University of Toledo, USA, for generously providing the V_{1a} agonist used in this study. Furthermore, the authors express their gratitude to Prof. Franz Weber, Centre of Dental Medicine, University of Zurich for providing the MC3T3-E1 cells.

AUTHORS CONTRIBUTIONS

GP conceptualized the study; ZR, EJ and GP designed the research; ZR, EJ, PK, SM, JR, MCC and GP performed the research. ASB performed RNA-seq data analysis; HH, YK, JJ, and IRA contributed new reagents/analytic tools; ZR, EJ, LS, CAW, MAH, HMK, JL and GP analyzed data; ZR, EJ, and GP wrote the paper; LS, CAW, MAH, HMK, JL reviewed and edited the manuscript; GP was responsible for supervision; and GP was responsible for funding acquisition.

DATA AVAILABILITY

All the relevant data associated with the current study can be found in the main text or supplementary material. The raw RNA-seq data used in this publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through GEO Series accession number GSE235694 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE235694>). All data are available from the authors upon request.

REFERENCES

1. White, K. E., Evans, W. E., O'Riordan, J. L. H., Speer, M. C., Econs, M. J., Lorenz-Depiereux, B., Grabowski, M., Meitinger, T. & Strom, T. M. Autosomal dominant hypophosphataemic rickets is associated with mutations in FGF23. *Nat. Genet.* **26**, 345–348 (2000).
2. Liu, S., Guo, R., Simpson, L. G., Xiao, Z. S., Burnham, C. E. & Quarles, L. D. Regulation of fibroblastic growth factor 23 expression but not degradation by PHEX. *J. Biol. Chem.* **278**, 37419–37426 (2003).
3. Yuan, B., Takaiwa, M., Clemens, T. L., Feng, J. Q., Kumar, R., Rowe, P. S., Xie, Y. & Drezner, M. K. Aberrant Phex function in osteoblasts and osteocytes alone underlies murine X-linked hypophosphatemia. *J. Clin. Invest.* **118**, 722–734 (2008).
4. Shimada, T., Kakitani, M., Yamazaki, Y., Hasegawa, H., Takeuchi, Y., Fujita, T., Fukumoto, S., Tomizuka, K. & Yamashita, T. Targeted ablation of Fgf23 demonstrates an essential physiological role of FGF23 in phosphate and vitamin D metabolism. *J. Clin. Invest.* **113**, 561–568 (2004).
5. Vervloet, M. Renal and extrarenal effects of fibroblast growth factor 23. *Nature Reviews Nephrology* (2019) doi:10.1038/s41581-018-0087-2.
6. Grabner, A., Amaral, A. P., Schramm, K., Singh, S., Sloan, A., Yanucil, C., Li, J., Shehadeh, L. A., Hare, J. M., David, V., Martin, A., Fornoni, A., Di Marco, G. S., Kentrup, D., Reuter, S., Mayer, A. B., Pavenstädt, H., Stypmann, J., Kuhn, C., *et al.* Activation of cardiac fibroblast growth factor receptor 4 causes left ventricular hypertrophy. *Cell Metab* **22**, 1020–1032 (2015).
7. Sharma, S., Katz, R., Bullen, A. L., Chaves, P. H. M., De Leeuw, P. W., Kroon, A. A., Houben, A. J. H. M., Shlipak, M. G. & Ix, J. H. Intact and c-terminal FGF23 assays-do kidney function, inflammation, and low iron influence relationships with outcomes? *J. Clin. Endocrinol. Metab.* **105**, e4875 (2020).
8. Saito, H., Maeda, A., Ohtomo, S. I., Hirata, M., Kusano, K., Kato, S., Ogata, E., Segawa, H., Miyamoto, K. I. & Fukushima, N. Circulating FGF-23 is regulated by 1 α ,25-dihydroxyvitamin D 3 and phosphorus in vivo. *J. Biol. Chem.* **280**, 2543–2549 (2005).

9. Perwad, F., Azam, N., Zhang, M. Y. H., Yamashita, T., Tenenhouse, H. S. & Portale, A. A. Dietary and serum phosphorus regulate fibroblast growth factor 23 expression and 1,25-dihydroxyvitamin D metabolism in mice. *Endocrinology* **146**, 5358–5364 (2005).
10. Bär, L., Feger, M., Fajol, A., Klotz, L. O., Zeng, S., Lang, F., Hocher, B. & Föller, M. Insulin suppresses the production of fibroblast growth factor 23 (FGF23). *Proc. Natl. Acad. Sci. U. S. A.* **115**, 5804–5809 (2018).
11. Pathare, G., Föller, M., Daryadel, A., Mutig, K., Bogatikov, E., Fajol, A., Almilaji, A., Michael, D., Stange, G., Voelkl, J., Wagner, C. A., Bachmann, S. & Lang, F. OSR1-sensitive renal tubular phosphate reabsorption. *Kidney Blood Press. Res.* **36**, (2012).
12. Pathare, G., Föller, M., Michael, D., Walker, B., Hierlmeier, M., Mannheim, J. G., Pichler, B. J. & Lang, F. Enhanced FGF23 serum concentrations and phosphaturia in gene targeted mice expressing WNK-resistant spak. *Kidney Blood Press. Res.* (2012) doi:10.1159/000343393.
13. Zhang, B., Umbach, A. T., Chen, H., Yan, J., Fakhri, H., Fajol, A., Salker, M. S., Spichtig, D., Daryadel, A., Wagner, C. A., Föller, M. & Lang, F. Up-regulation of FGF23 release by aldosterone. *Biochem. Biophys. Res. Commun.* (2016) doi:10.1016/j.bbrc.2016.01.034.
14. Pathare, G., Anderegg, M., Albano, G., Lang, F. & Fuster, D. G. Elevated FGF23 Levels in Mice Lacking the Thiazide-Sensitive NaCl cotransporter (NCC). *Sci. Rep.* (2018) doi:10.1038/s41598-018-22041-1.
15. Farrow, E. G., Yu, X., Summers, L. J., Davis, S. I., Fleet, J. C., Allen, M. R., Robling, A. G., Stayrook, K. R., Jideonwo, V., Magers, M. J., Garringer, H. J., Vidal, R., Chan, R. J., Goodwin, C. B., Hui, S. L., Peacock, M. & White, K. E. Iron deficiency drives an autosomal dominant hypophosphatemic rickets (ADHR) phenotype in fibroblast growth factor-23 (Fgf23) knock-in mice. *Proc. Natl. Acad. Sci. U. S. A.* **108**, (2011).
16. Masuda, Y., Ohta, H., Morita, Y., Nakayama, Y., Miyake, A., Itoh, N. & Konishi, M. Expression of Fgf23 in Activated Dendritic Cells and Macrophages in Response to Immunological Stimuli in Mice. *Biol. Pharm. Bull.* **38**, 687–693 (2015).

17. Humalda, J. K., Yeung, S. M. H., Geleijnse, J. M., Gijssbers, L., Riphagen, I. J., Hoorn, E. J., Rotmans, J. I., Vogt, L., Navis, G., Bakker, S. J. L. & de Borst, M. H. Effects of Potassium or Sodium Supplementation on Mineral Homeostasis: A Controlled Dietary Intervention Study. *J. Clin. Endocrinol. Metab.* **105**, e3246–e3256 (2020).
18. Fischer, S. S., Kempe, D. S., Leibrock, C. B., Rexhepaj, R., Siraskar, B., Boini, K. M., Ackermann, T. F., Föller, M., Hoher, B., Rosenblatt, K. P., Kuro-o, M. & Lang, F. Hyperaldosteronism in Klotho-deficient mice. *Am. J. Physiol. Renal Physiol.* **299**, (2010).
19. Verbalis, J. G., Barsony, J., Sugimura, Y., Tian, Y., Adams, D. J., Carter, E. A. & Resnick, H. E. Hyponatremia-induced osteoporosis. *J. Bone Miner. Res.* **25**, 554–563 (2010).
20. Barsony, J., Sugimura, Y. & Verbalis, J. G. Osteoclast response to low extracellular sodium and the mechanism of hyponatremia-induced bone loss. *J. Biol. Chem.* **286**, 10864–10875 (2011).
21. Barsony, J., Xu, Q. & Verbalis, J. G. Hyponatremia elicits gene expression changes driving osteoclast differentiation and functions. *Mol. Cell. Endocrinol.* **554**, 111724 (2022).
22. Usala, R. L., Fernandez, S. J., Mete, M., Cowen, L., Shara, N. M., Barsony, J. & Verbalis, J. G. Hyponatremia Is Associated With Increased Osteoporosis and Bone Fractures in a Large US Health System Population. *J. Clin. Endocrinol. Metab.* **100**, 3021–3031 (2015).
23. Hoorn, E. J., Rivadeneira, F., Van Meurs, J. B. J., Ziere, G., Stricker, B. H. C., Hofman, A., Pols, H. A. P., Zietse, R., Uitterlinden, A. G. & Zillikens, M. C. Mild hyponatremia as a risk factor for fractures: the Rotterdam Study. *J. Bone Miner. Res.* **26**, 1822–1828 (2011).
24. Jamal, S. A., Arampatzis, S., Harrison, S. L., Bucur, R. C., Ensrud, K., Orwoll, E. S. & Bauer, D. C. Hyponatremia and Fractures: Findings From the MrOS Study. *J. Bone Miner. Res.* **30**, 970–975 (2015).
25. Cowen, L. E., Hodak, S. P. & Verbalis, J. G. Age-Associated Abnormalities of Water Homeostasis. *Endocrinol. Metab. Clin. North Am.* **42**, 349–370 (2013).
26. Tamma, R., Sun, L., Cuscito, C., Lu, P., Corcelli, M., Li, J., Colaianni, G., Moonga, S. S., Di

- Benedetto, A., Grano, M., Colucci, S., Yuen, T., New, M. I., Zallone, A. & Zaidi, M. Regulation of bone remodeling by vasopressin explains the bone loss in hyponatremia. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 18644–18649 (2013).
27. Sun, L., Tamma, R., Yuen, T., Colaianni, G., Ji, Y., Cuscito, C., Bailey, J., Dhawan, S., Lu, P., Calvano, C. D., Zhu, L. L., Zambonin, C. G., Di Benedetto, A., Stachnik, A., Liu, P., Grano, M., Colucci, S., Davies, T. F., New, M. I., *et al.* Functions of vasopressin and oxytocin in bone mass regulation. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 164–169 (2016).
28. Barsony, J., Manigrasso, M. B., Xu, Q., Tam, H. & Verbalis, J. G. Chronic hyponatremia exacerbates multiple manifestations of senescence in male rats. *Age (Omaha)*. **35**, 271 (2013).
29. Monnerat, S., Refardt, J., Potasso, L., Meier, C. & Christ-Crain, M. An Increase in Plasma Sodium Levels Is Associated With an Increase in Osteoblast Function in Chronic SIAD. *J. Clin. Endocrinol. Metab.* (2023) doi:10.1210/clinem/dgad238.
30. Potasso, L., Refardt, J., Meier, C. & Christ-Crain, M. Effect of hyponatremia normalization on osteoblast function in patients with SIAD. *Eur. J. Endocrinol.* **186**, 1–8 (2021).
31. Wu, L., Luthringer, B. J. C., Feyerabend, F., Zhang, Z., Machens, H. G., Maeda, M., Taipaleenmäki, H., Hesse, E., Willumeit-Römer, R. & Schilling, A. F. Increased levels of sodium chloride directly increase osteoclastic differentiation and resorption in mice and men. *Osteoporos. Int.* **28**, 3215 (2017).
32. Schroder, A., Neubert, P., Titze, J., Bozec, A., Neuhofer, W., Proff, P., Kirschneck, C. & Jantsch, J. Osteoprotective action of low-salt diet requires myeloid cell–derived NFAT5. *JCI Insight* **4**, (2019).
33. Miyakawa, H., Woo, S. K., Dahl, S. C., Handler, J. S. & Kwon, H. M. Tonicity-responsive enhancer binding protein, a Rel-like protein that stimulates transcription in response to hypertonicity. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 2538–2542 (1999).
34. Woo, S. K., Dahl, S. C., Handler, J. S. & Kwon, H. M. Bidirectional regulation of tonicity-responsive

- enhancer binding protein in response to changes in tonicity. *Am. J. Physiol. Renal Physiol.* **278**, (2000).
35. Böger, C. A., Gorski, M., McMahon, G. M., Xu, H., Chang, Y. P. C., Van Der Most, P. J., Navis, G., Nolte, I. M., De Borst, M. H., Zhang, W., Lehne, B., Loh, M., Tan, S. T., Boerwinkle, E., Grams, M. E., Sekula, P., Li, M., Wilmot, B., Moon, J. G., *et al.* NFAT5 and SLC4A10 Loci Associate with Plasma Osmolality. *J. Am. Soc. Nephrol.* **28**, 2311–2321 (2017).
36. Wohlfahrt, P., Melenovsky, V., Kotrc, M., Benes, J., Jabor, A., Franekova, J., Lemaire, S., Kautzner, J. & Jarolim, P. Association of Fibroblast Growth Factor-23 Levels and Angiotensin-Converting Enzyme Inhibition in Chronic Systolic Heart Failure. *JACC Hear. Fail.* **3**, 829–839 (2015).
37. Gouzènes, L., Sabatier, N., Richard, P., Moos, F. C. & Dayanithi, G. V1a- and V2-type vasopressin receptors mediate vasopressin-induced Ca²⁺ responses in isolated rat supraoptic neurones. *J. Physiol.* **517**, 771 (1999).
38. Manning, M., Misicka, A., Olma, A., Bankowski, K., Stoev, S., Chini, B., Durroux, T., Mouillac, B., Corbani, M. & Guillon, G. Oxytocin and Vasopressin Agonists and Antagonists as Research Tools and Potential Therapeutics. *J. Neuroendocrinol.* **24**, 609 (2012).
39. Izumi, Y., Yang, W., Zhu, J., Burg, M. B. & Ferraris, J. D. RNA-Seq analysis of high NaCl-induced gene expression. *Physiol. Genomics* **47**, 500–513 (2015).
40. Bär, L., Stournaras, C., Lang, F. & Föller, M. Regulation of fibroblast growth factor 23 (FGF23) in health and disease. *FEBS Letters* vol. 593 1879–1900 (2019).
41. Park, S. J. & Shin, J. II. Inflammation and hyponatremia: An underrecognized condition? *Korean Journal of Pediatrics* vol. 56 519–522 (2013).
42. Xu, S., Wong, C. C. L., Tong, E. H. Y., Chung, S. S. M., John R. Yates, I., Yin, Y. & Ko, B. C. B. Phosphorylation by Casein Kinase 1 Regulates Tonicity-induced Osmotic Response Element-binding Protein/Tonicity Enhancer-binding Protein Nucleocytoplasmic Trafficking. *J. Biol. Chem.*

- 283**, 17624 (2008).
43. Takenaka, M., Preston, A. S., Kwon, H. M. & Handler, J. S. The tonicity-sensitive element that mediates increased transcription of the betaine transporter gene in response to hypertonic stress. *J. Biol. Chem.* **269**, 29379–29381 (1994).
 44. Ko, B. C. B., Ruepp, B., Bohren, K. M., Gabbay, K. H. & Chung, S. S. M. Identification and characterization of multiple osmotic response sequences in the human aldose reductase gene. *J. Biol. Chem.* **272**, 16431–16437 (1997).
 45. Rim, J. S., Atta, M. G., Dahl, S. C., Berry, G. T., Handler, J. S. & Kwon, H. M. Transcription of the sodium/myo-inositol cotransporter gene is regulated by multiple tonicity-responsive enhancers spread over 50 kilobase pairs in the 5'-flanking region. *J. Biol. Chem.* **273**, 20615–20621 (1998).
 46. Ferraris, J. D., Williams, C. K., Persaud, P., Zhang, Z., Chen, Y. & Burg, M. B. Activity of the TonEBP/OREBP transactivation domain varies directly with extracellular NaCl concentration. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 739–744 (2002).
 47. Denton, D. A., Mckinley, M. J. & Weisinger, R. S. Hypothalamic integration of body fluid regulation. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 7397 (1996).
 48. Wiig, H., Schröder, A., Neuhofer, W., Jantsch, J., Kopp, C., Karlsen, T. V., Boschmann, M., Goss, J., Bry, M., Rakova, N., Dahlmann, A., Brenner, S., Tenstad, O., Nurmi, H., Mervaala, E., Wagner, H., Beck, F. X., Müller, D. N., Kerjaschki, D., *et al.* Immune cells control skin lymphatic electrolyte homeostasis and blood pressure. *J. Clin. Invest.* **123**, 2803–2815 (2013).
 49. Machnik, A., Neuhofer, W., Jantsch, J., Dahlmann, A., Tammela, T., MacHura, K., Park, J. K., Beck, F. X., Müller, D. N., Derer, W., Goss, J., Ziomber, A., Dietsch, P., Wagner, H., Van Rooijen, N., Kurtz, A., Hilgers, K. F., Alitalo, K., Eckardt, K. U., *et al.* Macrophages regulate salt-dependent volume and blood pressure by a vascular endothelial growth factor-C-dependent buffering mechanism. *Nat. Med.* **15**, 545–552 (2009).
 50. Wu, C., Yosef, N., Thalhamer, T., Zhu, C., Xiao, S., Kishi, Y., Regev, A. & Kuchroo, V. K. Induction

- of pathogenic TH 17 cells by inducible salt-sensing kinase SGK1. *Nature* **496**, 513–517 (2013).
51. Tian, W., Fu, Y., Garcia-Elias, A., Fernández-Fernández, J. M., Vicente, R., Kramer, P. L., Klein, R. F., Hitzemann, R., Orwoll, E. S., Wilmot, B., McWeeney, S., Valverde, M. A. & Cohen, D. M. A loss-of-function nonsynonymous polymorphism in the osmoregulatory TRPV4 gene is associated with human hyponatremia. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 14034–14039 (2009).
52. Sjöström, M., Stenström, K., Eneling, K., Zwiller, J., Katz, A. I., Takemori, H. & Bertorello, A. M. SIK1 is part of a cell sodium-sensing network that regulates active sodium transport through a calcium-dependent process. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 16922–16927 (2007).
53. Ravesloot, J. H., Eisen, T., Baron, R., Boron, W. F. & Boron, W. F. Role of Na-H Exchangers and Vacuolar H⁺ Pumps in Intracellular pH Regulation in Neonatal Rat Osteoclasts. *J. Gen. Physiol.* **105**, 177–208 (1995).
54. Moonga, B. S., Davidson, R., Sun, L., Adebajo, O. A., Moser, J., Abedin, M., Zaidi, N., Huang, C. L. H. & Zaidi, M. Identification and characterization of a sodium/calcium exchanger, NCX-1, in osteoclasts and its role in bone resorption. *Biochem. Biophys. Res. Commun.* **283**, 770–775 (2001).
55. Schrier, R. W. & Fassett, R. G. Pathogenesis of sodium and water retention in cardiac failure. *Ren. Fail.* **20**, 773–781 (1998).
56. Cárdenas, A. & Arroyo, V. Mechanisms of water and sodium retention in cirrhosis and the pathogenesis of ascites. *Best Pract. Res. Clin. Endocrinol. Metab.* **17**, 607–622 (2003).
57. Hannon, M. J. & Verbalis, J. G. Sodium homeostasis and bone. *Curr. Opin. Nephrol. Hypertens.* **23**, 370–376 (2014).
58. Zaidi, M., Sun, L., Liu, P., Davies, T. F., New, M., Zallone, A. & Yuen, T. Pituitary-bone connection in skeletal regulation. *Horm. Mol. Biol. Clin. Investig.* **28**, 85–94 (2016).
59. Zaidi, M., New, M. I., Blair, H. C., Zallone, A., Baliram, R., Davies, T. F., Cardozo, C., Iqbal, J., Sun, L., Rosen, C. J. & Yuen, T. Actions of pituitary hormones beyond traditional targets. *J.*

- Endocrinol.* **237**, R83–R98 (2018).
60. Verbalis, J. G. & Drutarosky, M. D. Adaptation to chronic hypoosmolality in rats. *Kidney Int.* **34**, 351–360 (1988).
61. Akiyama, K. ichi, Miura, Y., Hayashi, H., Sakata, A., Matsumura, Y., Kojima, M., Tsuchiya, K., Nitta, K., Shiizaki, K., Kurosu, H. & Kuro-o, M. Calciprotein particles regulate fibroblast growth factor-23 expression in osteoblasts. *Kidney Int.* **97**, 702–712 (2020).
62. Pathare, G., Dhayat, N. A., Mohebbi, N., Wagner, C. A., Bobulescu, I. A., Moe, O. W. & Fuster, D. G. Changes in V-ATPase subunits of human urinary exosomes reflect the renal response to acute acid/alkali loading and the defects in distal renal tubular acidosis. *Kidney Int.* **93**, 871–880 (2018).
63. Koshimizu, T. A., Nasa, Y., Tanoue, A., Oikawa, R., Kawahara, Y., Kiyono, Y., Adachi, T., Tanaka, T., Kuwaki, T., Mori, T., Takeo, S., Okamura, H. & Tsujimoto, G. V1a vasopressin receptors maintain normal blood pressure by regulating circulating blood volume and baroreflex sensitivity. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 7807–7812 (2006).
64. Verouti, S. N., Lambert, D., Mathis, D., Pathare, G., Escher, G., Vogt, B. & Fuster, D. G. Solute carrier SLC16A12 is critical for creatine and guanidinoacetate handling in the kidney. *Am. J. Physiol. - Ren. Physiol.* (2021) doi:10.1152/AJPRENAL.00475.2020.
65. Nigro, N., Winzeler, B., Suter-Widmer, I., Schuetz, P., Arici, B., Bally, M., Blum, C. A., Nickel, C. H., Bingisser, R., Bock, A., Huber, A., Müller, B. & Christ-Crain, M. Evaluation of copeptin and commonly used laboratory parameters for the differential diagnosis of profound hyponatraemia in hospitalized patients: ‘The Co-MED Study’. *Clin. Endocrinol. (Oxf)*. **86**, 456–462 (2017).
66. Refardt, J., Winzeler, B., Meienberg, F., Vogt, D. R. & Christ-Crain, M. Empagliflozin Increases Short-Term Urinary Volume Output in Artificially Induced Syndrome of Inappropriate Antidiuresis. *Int. J. Endocrinol.* **2017**, (2017).

FIGURE LEGENDS

Fig. 1. High NaCl suppresses FGF23 formation in UMR-106 cells. A) Representative immunoblots of NFAT5 and β -actin upon addition of 20 mM NaCl, 40 mM mannitol, and 40 mM urea treatment for 16 h. B) Quantification of NFAT5 protein normalized with β -actin (n=3). C) Fold change *Nfat5* mRNA levels; D) cFGF23 levels in cell media; E) Fold change *Fgf23* mRNA levels, after treating 20 mM NaCl, 40 mM mannitol, and 40 mM urea for 24 h (n=3-4). F) Lactate dehydrogenase (LDH) release in the cells following 0, 40, 80, 120, and 140 mM +NaCl treatment for 24 h (n=3). G) Fold change *Fgf23* and *Nfat5* mRNA levels following 0, 2, 5, 10, 20, and 30 mM +NaCl treatment for 24 h (n=4). One-way ANOVA summary for *Nfat5*: $F=13.39$; $p<0.0001$; $R\ sq.=0.79$, and for *Fgf23*: $F=56.19$; $p<0.0001$; $R\ sq.=0.94$. H) Fold change *Fgf23* and *Nfat5* mRNA levels after 20 mM +NaCl treatment over 0-48 h (n=3). p value (using an unpaired t-test) indicates statistically significant difference between *Nfat5* and *Fgf23* mRNA, when compared to 0 h (normalized to one, for each time-point). All the values are expressed in arithmetic means \pm SEM.

Fig. 2. Hypotonicity elevates FGF23 formation in UMR-106 cells. A) Fold change *Fgf23* mRNA levels upon treating NaCl-deficient culture media by -5, -10, -15, and -20 mM NaCl for 24 h (n=3). B) Fold change *Fgf23* mRNA; C) cFGF23 levels in media; D) *Nfat5* mRNA, after treating cells with -20 mM NaCl-deficient culture media. The osmolality was corrected by adding mannitol (40 mM) or urea (40 mM) for 24 h (n=3). All the values are expressed in arithmetic means \pm SEM. ns (not significant) $p > 0.05$

Fig. 3. No evidence for FGF23 regulation by AVP. A) Agarose gel electrophoresis for detection of rat V_{1a} and V_2 receptor mRNA in rat liver, kidney, and UMR-106 cells. The cDNA (2 μ g) obtained by RT was amplified for 34 cycles by PCR. B) V_{1a} , V_2 , and V_{1b} mRNA expressions quantified by qRT-PCR in UMR-106 cells (n=3). ND: not detected. C) Fold change *Fgf23* mRNA abundance relative to *Tbp* in UMR-106 cells treated without (Ctrl) or with the indicated concentrations of AVP for 24 h. (n=3). D) Fold change *Fgf23* levels in UMR-106 cells treated without (Ctrl) or with 100 nM AVP for indicated durations (n=3). E) Fold change *Fgf23* levels in UMR-106 cells treated without (Ctrl) or with the indicated concentrations of V_{1a} agonist, ([Phe²]OVT, [Phe²,Orn⁸]vasotocin) for 24 h. (n=3). F) Fold change *Fgf23* levels in UMR-106 cells treated without (Ctrl) or with the indicated concentrations of V_2 agonist, ddAVP for 24 h (n=3). G) Fold

change *Fgf23* mRNA in UMR-106 cells treated with -20 mM NaCl with or without AVP and V_{1a} agonist (each 10 nM) for 24h (n=3). H) Serum cFGF23 and iFGF23 levels in wild-type and V_{1a} KO mice (n=4-5, each group).

Fig. 4. The $[Na^+]$ mediated regulation of FGF23 in UMR-106 cells requires NFAT5. A) Original immunoblot of NFAT5 and β -actin after -NaCl (-20 mM) and +NaCl (+20 mM) treatment for 24 h in control and $NFAT5^{KO}$ cells. B) *Nfat5* mRNA levels in control and $NFAT5^{KO}$ cells after -NaCl and +NaCl treatment for 24 h (n=3). *Nfat5* mRNA levels in $NFAT5^{KO}$ cells were not detected. C) LDH release in cell supernatants treated with different NaCl concentrations (n=3). *p* value (using an unpaired t-test) indicates statistically significant difference between LDH levels in control and $NFAT5^{KO}$ cell supernatant at that particular +NaCl concentration. The two-way ANOVA reveals a $p < 0.0001$ for the treatment effect and $p = 0.0028$ for the interaction. D) *Fgf23* mRNA levels measured by qRT-PCR in control and $NFAT5^{KO}$ UMR-106 cells after -NaCl vs. +NaCl treatment for 24h (n=3, each group). E) Venn diagrams showing a number of upregulated/downregulated genes in -NaCl vs. +NaCl treatment for 24 h. F) Heatmap of gene expression levels in control and $NFAT5^{KO}$ cells upon -NaCl vs. +NaCl treatment. G) Volcano plot of upregulated/downregulated genes in control cells upon -NaCl vs. +NaCl treatment. These genes were differentially expressed in control, but not in $NFAT5^{KO}$ cells upon -NaCl vs. +NaCl treatment. H) GO analysis of the upregulated genes in control cells that remain unchanged in $NFAT5^{KO}$ cells upon -NaCl vs. +NaCl treatment. I) FGF23-regulating genes that are statistically significantly ($p < 0.05$) upregulated/downregulated in control cells but remain unchanged in $NFAT5^{KO}$ cells upon -NaCl vs. +NaCl treatment. J) *Phex* mRNA levels measured by qRT-PCR upon -NaCl vs. +NaCl for 24 h in UMR-106 cells. Osmolality was corrected to +NaCl by adding mannitol or urea (n=4).

Fig. 5. Hyponatremia patients have higher FGF23 levels. Serum $[Na^+]$ (A), cFGF23 (B), iFGF23 (C), and phosphate (D) in healthy matching controls and hyponatremic patients (n=6, each group). E) Correlation analysis between serum $[Na^+]$ and serum cFGF23/iFGF23. $r = -0.6138$, *p* value (two-tailed) = 0.0337 for $[Na^+]$ vs cFGF23; $r = -0.4830$, *p* value (two-tailed) = 0.1117 for $[Na^+]$ vs iFGF23.

Main tables

Extracellular sodium regulates fibroblast growth factor 23 (FGF23) formation.

Zsuzsa Radvanyi^{1,2†}, Eun Jin Yoo^{3†}, Palanivel Kandasamy⁴, Adrian Salas-Bastos¹, Sophie Monnerat^{5,6}, Julie Refardt^{5,6}, Mirjam Christ-Crain^{5,6}, Himeka Hayashi⁷, Yasuhiko Kondo⁷, Jonathan Jantsch^{8,9}, Isabel Rubio-Aliaga^{10,2}, Lukas Sommer¹, Carsten A. Wagner^{10,2}, Matthias A. Hediger⁴, Hyug Moo Kwon³, Johannes Loffing^{1,2}, Ganesh Pathare^{1,2*}

¹Institute of Anatomy, University of Zurich, Zurich, Switzerland; ²Swiss National Centre of Competence in Research “Kidney Control of Homeostasis”, Switzerland; ³Department of Biological Sciences, Ulsan National Institute of Science and Technology, Ulsan, Republic of Korea; ⁴Membrane Transport Discovery Lab, Department of Nephrology and Hypertension and Department of Biomedical Research, Inselspital, University of Bern, Bern, Switzerland; ⁵Department of Endocrinology, Diabetology and Metabolism, University Hospital Basel, Basel, Switzerland; ⁶Department of Clinical Research, University of Basel, Basel, Switzerland; ⁷Department of Animal Sciences, Teikyo University of Science, Yamanashi, Japan; ⁸Institute of Clinical Microbiology and Hygiene, University Hospital of Regensburg and University of Regensburg, Regensburg, Germany; ⁹Institute for Medical Microbiology, Immunology, and Hygiene, and Center for Molecular Medicine Cologne (CMMC), University of Cologne, Cologne, Germany; ¹⁰Institute of Physiology, University of Zurich, Zurich, Switzerland

†contributed equally to this work

***Correspondence author:** Ganesh Pathare, Institute of Anatomy, University of Zurich, Winterthurerstrasse 190, 8057, CH, Email: ganesh.pathare@anatomy.uzh.ch

Keywords: FGF23, extracellular-sodium, hyponatremia, NFAT5, bone and kidney

This word file includes:

Main tables

Main tables

Gene name	Primer sequence
<i>Fgf23 (Rat)</i>	Forward- TTGGATCGTATCACTTCAGC
	Reverse- TGCTTCGGTGACAGGTAG
<i>Nfat5 (Rat)</i>	Forward- TCCTGGCTC ATCTCAGCAGAC
	Reverse- ACCGAGGAGGCCAGTGGGGC
<i>Phex (Rat)</i>	Forward- CAGGCATCACATTCACCAAC
	Reverse- GGAGGACTGTGAGCACCAAT
<i>Tbp (Rat)</i>	Forward- TGGGATTGTACCACAGCTCCA
	Reverse- CTCATGATGACTGCAGCAAACC
<i>V1a (Rat)</i>	Forward- AACGAACAGCACTGGGATGTG
	Reverse- GGAATGAATCTGATGGATTTGGAA
<i>V2 (Rat)</i>	Forward- TAGCATACCGCCATGGA
	Reverse- CAAAGATGAAGAGCTGAGGCA
<i>V1b (Rat)</i>	Forward- AGATTCTACCAATGTGGCTTTC
	Reverse- ATGGTGGCTCAAGGAACG
<i>Gapdh (Rat)</i>	Forward- AGGTCGGTGTGAACGGATTTG
	Reverse- CATTCTCAGCCTTGACTGTGCC
<i>Fgf23 (Mouse)</i>	Forward- TTTCCAGGTTCTGTCTAGG
	Reverse- CTCGCAGGTGACTCTCAG
<i>Nfat5 (Mouse)</i>	Forward- TGTTCAGCCATTTACGTACTCC
	Reverse- ACATTCAAAGCACCAGCTGCT
<i>Tbp (Mouse)</i>	Forward- ACCCTTCACCAATGACTCCTATG
	Reverse- TGACTGCAGCAAATCGCTTGG

Table 1: List of primers used for qPCR

GuideRNA name	Primer's sequence
NFAT5-Exon 1-guideRNA1	Forward- CACCGAGCAATGAGATGAAGTCCGA
	Reverse- AAACCTCGGACTTCATCTCATTGCTC
NFAT5-Exon 3-guideRNA2	Forward- CACCGCATCAATGAGTCAGACAAG
	Reverse- AAACCTTGTCTGACTCATTGATGC
NFAT5-Exon 4-guideRNA3	Forward- CACCGCGTGGGGGTAAGTAACAGAG
	Reverse- AAACCTCTGTTACTTACCCCCACGC
NFAT5-Exon 4-guideRNA4	Forward- CACCGTTCCTACAATGATAACACTG
	Reverse- AAACCAGTGTTATCATTGTAGGAAC

Table 2: List of GuideRNA used for the generation of CRISPR/Cas9 knockout of NFAT5.

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof