











REVIEW ARTICLE

Apnoeic oxygenation during paediatric tracheal intubation: a systematic review and meta-analysis

Alexander Fuchs^{1,2,*}, Gabriela Koepf^{1,†}, Markus Huber¹, Jonas Aebli¹, Arash Afshari³, Rachele Bonfiglio², Robert Greif^{4,5}, Andrea C. Lusardi², Carolina S. Romero⁶, Marc von Gernler⁷, Nicola Disma^{2,‡} and Thomas Riva^{1,‡}

¹Department of Anaesthesiology and Pain Medicine, Bern University Hospital, Inselspital, University of Bern, Bern, Switzerland, ²Unit for Research in Anaesthesia, IRCCS Istituto Giannina Gaslini, Genoa, Italy, ³Department of Paediatric And Obstetric Anesthesia, Juliane Marie Centre, Rigshospitalet, University of Copenhagen, Institute of Clinical Medicine, Copenhagen, Denmark, ⁴School of Medicine, Sigmund Freud University Vienna, Vienna, Austria, ⁵University of Bern, Bern, Switzerland, ⁶Anesthesia, Critical Care and Pain Department, Hospital General Universitario De Valencia, Research Methods Department, Universidad Europea de Valencia, Valencia, Spain and ⁷Medical Library, University of Bern, Bern, Switzerland

*Corresponding author. E-mail: alexander.fuchs@insel.ch

†Contributed equally and share the first authorship.

‡Contributed equally and share the last authorship.

Abstract

Background: Supplemental oxygen administration by apnoeic oxygenation during laryngoscopy for tracheal intubation is intended to prolong safe apnoea time, reduce the risk of hypoxaemia, and increase the success rate of first-attempt tracheal intubation under general anaesthesia. This systematic review examined the efficacy and effectiveness of apnoeic oxygenation during tracheal intubation in children.

Methods: This systematic review and meta-analysis included randomised controlled trials and non-randomised studies in paediatric patients requiring tracheal intubation, evaluating apnoeic oxygenation by any method compared with patients without apnoeic oxygenation. Searched databases were MEDLINE, Embase, Cochrane Library, CINAHL, [ClinicalTrials.gov](https://clinicaltrials.gov), International Clinical Trials Registry Platform (ICTRP), Scopus, and Web of Science from inception to March 22, 2023. Data extraction and risk of bias assessment followed the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) recommendation.

Results: After initial selection of 40 708 articles, 15 studies summarising 9802 children were included (10 randomised controlled trials, four pre-post studies, one prospective observational study) published between 1988 and 2023. Eight randomised controlled trials were included for meta-analysis ($n=1070$ children; 803 from operating theatres, 267 from neonatal intensive care units). Apnoeic oxygenation increased intubation first-pass success with no physiological instability (risk ratio [RR] 1.27, 95% confidence interval [CI] 1.03–1.57, $P=0.04$, $I^2=0$), higher oxygen saturation during intubation (mean difference 3.6%, 95% CI 0.8–6.5%, $P=0.02$, $I^2=63\%$), and decreased incidence of hypoxaemia (RR 0.24, 95% CI 0.17–0.33, $P<0.01$, $I^2=51\%$) compared with no supplementary oxygen administration.

Conclusion: This systematic review with meta-analysis confirms that apnoeic oxygenation during tracheal intubation of children significantly increases first-pass intubation success rate. Furthermore, apnoeic oxygenation enables stable physiological conditions by maintaining oxygen saturation within the normal range.

Received: 18 August 2023; Accepted: 24 October 2023

© 2023 The Authors. Published by Elsevier Ltd on behalf of British Journal of Anaesthesia. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

For Permissions, please email: permissions@elsevier.com

Clinical trial registration: Protocol registered prospectively on PROSPERO (registration number: CRD42022369000) on December 2, 2022.

Keywords: airway; apnoea; apnoeic oxygenation; paediatric anaesthesia; supplemental oxygen; tracheal intubation

Editor's key points

- Supplemental oxygen administration by apnoeic oxygenation during laryngoscopy for tracheal intubation could prolong safe apnoea time, reduce the risk of hypoxaemia, and increase the success rate of first attempt tracheal intubation under general anaesthesia.
- This systematic review examined the efficacy of apnoeic oxygenation in facilitating tracheal intubation in children.
- Apnoeic oxygenation increased intubation first-pass success, increased oxygen saturation during intubation, and decreased incidence of hypoxaemia compared with no supplementary oxygen.
- However, the included studies were heterogeneous, and more high-quality randomised controlled trials are warranted with well-defined outcome variables.

Tracheal intubation aims to establish a patent airway to ensure ventilation of the lungs during surgery, procedures, or respiratory insufficiency. It is lifesaving for children with severe acute respiratory failure. The number of tracheal intubation attempts is associated with an increased incidence of severe complications.^{1,2} Interventions to improve first-attempt intubation success rate and increase safety are pivotal.³ Tracheal intubation in the operating theatre is associated with an incidence of difficult airway of ~0.9% in children up to 16 yr old,⁴ but alarmingly this is 5.8% in neonates,¹ and is mostly unanticipated. Furthermore, the incidence of haemoglobin desaturation appears to be as high as 40%, potentially leading to severe adverse events in neonates.¹ Prolonged severe desaturation can lead to hypoxic encephalopathy,⁵ cardiac arrest,^{6,7} or death.⁸

Facemask preoxygenation in children can be difficult because of lack of cooperation leading to improper mask seal,⁹ and is relatively ineffective, especially in infants, considering hypoxaemia occurs within seconds after cessation of spontaneous or assisted ventilation.¹⁰ Safe apnoea time is generally described as the time from cessation of breathing or ventilation until a patient attains a critical level of peripheral arterial oxygen saturation.¹¹ Beyond this point, oxygenation decreases rapidly to critically low blood and tissue oxygen levels endangering vital functions.¹²

Apnoeic oxygenation to reduce the risk of hypoxaemia and extend safe apnoea time was initially described in the early 1900s.^{13,14} It consists of administering a constant stream of oxygen 100% while the patient is not breathing. The physiological requirements for adequate apnoeic oxygenation are a patent upper and lower airway, diffusion of highly concentrated oxygen into the alveoli, minimal pulmonary shunting, and presence of cardiac function. Low-flow oxygen with different delivery techniques achieves this objective, and

mounting evidence supports the possible advantages of high-flow nasal oxygen administration.¹⁵ However, apnoeic oxygenation is currently not a routine practice during tracheal intubation in most institutions.^{2,16,17} This systematic review evaluates the efficacy of apnoeic oxygenation on first-attempt intubation success, oxygen saturation, and adverse effects during paediatric tracheal intubation.

Methods

The review protocol for this systematic review and meta-analysis was registered with PROSPERO (registration number: CRD42022369000) on December 2, 2022. We report our findings according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses guidelines (PRISMA).¹⁸

Eligibility criteria

We included peer-reviewed randomised controlled trials (RCTs) and non-randomised studies (non-RCTs, interrupted time series, controlled before-and-after studies, and cohort studies) in paediatric patients (age <16 yr) requiring tracheal intubation. Included studies compared apnoeic oxygenation by any method or device with a control group without apnoeic oxygenation. Apnoeic oxygenation was defined as any passive insufflation of oxygen of any flow into the nose or mouth without ventilation. Unpublished studies, case series, conference abstracts, trial protocols, duplicates, and unretrievable articles were excluded.

Searched databases and search strategy

A literature search strategy was devised for the following databases: MEDLINE, Embase, Cochrane Library, CINAHL, Web of Science Core Collection, Scopus, [ClinicalTrials.gov](https://www.clinicaltrials.gov), and International Clinical Trials Registry Platform (ICTRP). A medical information specialist (MvG) developed an initial search strategy in MEDLINE with a test against a list of core references to ensure that key publications were included. After refinement, the information specialist set up the search strategy for each information source based on database-specific index terms and free text. The free text search included synonyms, acronyms, and similar terms. No database-provided limits have been applied in any sources considering study types, languages, publication years, or any other formal criteria. The search was finalised on March 22, 2023. The detailed final search strategies were published,¹⁹ and additional information about the systematic search is available in the [Supplementary material](#).

Study selection and assessment

After identifying relevant publications, all were imported into EndNote (EndNote20, Clarivate, Philadelphia, PA, USA). We used Deduklick (Risklick AG, Bern, Switzerland) for

deduplication (MvG). An equal number of titles and abstracts were distributed to four groups with two study researchers (ND and ACL, AF and GK, TR and JA, RB and CSR). The researchers independently screened all titles and abstracts using the blinded mode in Rayyan²⁰ for systematic reviews. Disagreements were resolved through discussion or consulting a third senior researcher (RG). All available data were extracted (GK, CSR), including study characteristics, design, interventions, populations, study methods, and outcomes of significance to the review question and specific objectives (Table 1). Any discrepancies were resolved through discussion or consultation with a senior researcher (RG).

The risk of bias was assessed by five authors (AA, AF, CSR, GK, RB) using Version 2 of the Cochrane risk-of-bias tool for randomised trials (RoB 2) for randomised trials,²¹ Risk Of Bias In Non-randomised Studies - of Interventions (ROBINS-I) for observational studies,²² and the checklist from the National Institutes of Health (NIH) quality assessment tool for before-after studies.²³ Disagreements were resolved by consensus or discussion with a senior researcher (RG). If information from included studies was not available to answer our primary and secondary outcomes or assess the overall quality of the studies, the corresponding authors were approached.

Three authors (AF, GK, MH) independently assessed the certainty of evidence with Grading of Recommendations Assessment, Development, and Evaluation (GRADE).²⁴ Any disagreements that arose were resolved through consensus. Six clinically relevant outcomes were defined and included in the GRADE table of certainty of evidence (Table 2).

Outcomes

The primary outcome was the incidence of successful first-pass tracheal intubation. Secondary outcomes were: number of intubation attempts; incidence of life-threatening events during tracheal intubation (i.e. cardiac arrest, failed intubation, failed ventilation, or both, emergency front of neck access, severe hypoxaemia (SpO₂ <80%), severe bradycardia (<60 bpm), or death); barotrauma (i.e., pneumothorax or pneumomediastinum confirmed with ultrasound or chest radiography); hypoxaemia (SpO₂ <90%), bradycardia (<80 beats min⁻¹ or <100 beats min⁻¹), injury of the upper airway (e.g. bleeding or swelling), or unplanned admission to the intensive care unit secondary to tracheal intubation attempt.

Statistical methods

Only RCTs were considered for meta-analysis. Effect sizes (risk ratios [RRs] for binary outcomes and mean differences for continuous outcomes) were calculated when at least two RCTs reported data for an outcome.

As considerable between-study heterogeneity was detected, a random-effects model was used for analysis of effect sizes. We chose to apply the inverse variance method for continuous outcomes, and Mantel-Haenszel for binary outcomes. When median and inter-quartile ranges were reported as summary statistics for continuous outcomes, a quantile estimation method was used to estimate the mean and standard deviation (SD). The between-study heterogeneity was assessed with Higgins and Thompson's I^2 statistic. Statistical tests for funnel plot asymmetry were not performed given the small number of studies per outcome. All statistical computations were performed with R Version 4.0.5.²⁵

Results

We identified 27,084 articles after deduplication published between 1988 and 2023 (Fig. 1); 15 studies (including 9802 patients) met inclusion criteria. Among these, 10 were RCTs,^{26–35} four pre-post studies,^{36–39} and one a prospective observational study.⁴⁰ Ten RCTs were considered eligible for effect size calculation; eight RCTs had an intervention and a control group, one study described two intervention arms and a control group (Steiner and colleagues³⁴), and one study reported the outcomes for two subgroups only (Foran and colleagues³⁰). The two intervention arms in Steiner and colleagues³⁴ were analysed as separate and independent group comparisons. Similarly, the two subgroups in Foran and colleagues³⁰ were analysed as two independent intervention arms compared with the control group. The limitation of this approach is outlined in the Limitations section. The meta-analysis of RCTs involved 1070 children, 803 from the operating theatre and 267 from the neonatal intensive care unit. Study characteristics are outlined in Table 1.

Risk of bias assessment

Figure 2 summarises the risk of bias assessment (RoB2, ROBINS-I, pre-post-studies). The overall risk of bias from RCTs was considered to be low in four trials^{26,31,33,35} and to have some concerns in six trials.^{27–30,32,34} Across the trials, the risk of deviations from intended interventions and missing outcome data was deemed low. Some concerns were noted for two trials^{27,28} mainly because of inadequate reporting of the randomisation procedure (allocation concealment), concerns over the validity of outcome measurements were noted in three trials^{30,32,34} as the assessors were aware of the interventions received by study participants, while in three trials^{27,29,30} we were unable to assess adherence to a prespecified study protocol because of unavailability of the latter (outcome-reporting bias). Despite our efforts to contact the authors, we failed to obtain additional clarifying information. The quality rating for the before-after studies assessment was good in one study³⁶ and fair in three studies.^{37–39}

Meta-analyses

The forest plots of the meta-analyses appear in Figures 3 and 4.

First-pass success rate of tracheal intubation

Three RCTs^{28,30,33} with a total of 374 patients reported the success rate of first attempt at tracheal intubation, which was achieved in 66.8% (N=102/155) of the patients with apnoeic oxygenation and 51.5% (N=87/169) in the control group. The overall pooled analysis (Fig. 3a) showed a higher likelihood of first-pass successful tracheal intubation in the apnoeic oxygenation group compared with the control group (RR 1.27, 95% CI 1.03–1.57, P=0.04; $I^2=0$). However, a single study³³ largely dominates the pooled estimate, contributing 84.8% to the pooled effect size. The certainty of evidence was graded low given the small sample size, and as the optimal information size was not met.

Number of tracheal intubation attempts

Three RCTs^{28,30,33} with a total of 311 patients met inclusion criteria for assessment of number of tracheal intubation

Table 1 Summary of the findings for the included studies and for the observational and pre-post trials. aOR, adjusted odds ratio; ETT, ; int., intervention group; NHF, nasal high flow; PICU, paediatric intensive care unit; RR, risk ratio; Risk of bias methodological quality tools used.* RoB 2: A revised Cochrane risk-of-bias tool for randomised trials. † ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. ‡ checklist from the NIH Quality Assessment Tool for Before-After (Pre-Post) Studies.

Author year (country)	Study design	Total n	Setting	Age	Intervention	Control	Primary outcome	Secondary outcomes	Conclusions in the study	Comment	Risk of bias
Randomised controlled trials											
Ledbetter and colleagues 1988 (USA) ²⁷	RCT Two arms	20	Operating room (single centre)	1–24 Months	n=10 Flow: 4 L min ⁻¹ Device: Miller #1 Oxyscope	n=10 Flow: 0 L min ⁻¹ Device: Miller #1 Oxyscope	Maintenance of transcutaneous oxygen tension during tracheal intubation in children mean (SD) int. 24.2 (7.7) 95% CI (18.7–29.6) vs control 15.6 (6.3) CI (11.2–20.1); P-value 0.014	None	Oxygen insufflation during laryngoscopy and intubation of spontaneously breathing anaesthetised infants effectively minimises the decrease in transcutaneous oxygen tension (TcPo ₂) from pre-laryngoscopy levels	This study is not directly comparable with the other RCTs since TcPo ₂ was used and this measure specifically reflects the oxygen tension in the tissues and not in blood	Some concerns*
Windpassinger and colleagues 2016 (Austria) ²⁶	RCT Two arms	48	Operating room (single centre)	0–2 Yr	n=24 Flow: 4 L min ⁻¹ Device: Air Traq laryngoscope size 0–1	n=24 Flow: 0 L min ⁻¹ Device: Air Traq laryngoscope size 0–1	Time to SpO ₂ 95% after intubation with or without posterior pharynx oxygen insufflation mean (SD) int. 166 (47) s vs control 131 (39) s	Time to intubation	Oxygen insufflation prolongs the period of adequate oxygen saturation in infants and small children by an amount that is potentially clinically important	The duration of safe apnoea after intubation was longer with apnoeic oxygenation in small children than without oxygen	Low risk*
Humphreys and colleagues 2017 (Australia) ³⁵	RCT Two arms	48	Operating room (single centre)	0–10 Yr	n=24 Flow: flow rates 0–15 kg, 2 L kg ⁻¹ min ⁻¹ 15–30 kg, 35 L min ⁻¹ 30–50 kg, 40 L min ⁻¹ >50 kg, 50 L min ⁻¹ Device: Optiflow	n=24 Flow: 0 L min ⁻¹ Device: none	Time to SpO ₂ 92% in seconds intervention vs control 0–6 months: 192 vs 109 (95% CI 28.8), 7–24 months: 237 vs 147 (95% CI 18.9) 2–5 yr: 320 vs 191 (95% CI 15.3) 6–10 yr: 430 vs 261 (95% CI 37.5) Lowest SpO ₂ Mean (SD) int. 97.55% (2.93%) vs control 95.9% (5.75%) P-value 0.049	Changes in transcutaneous oxygen saturation Changes in transcutaneous CO ₂ tension	Transnasal humidified rapid-insufflation ventilatory exchange (THRIVE) prolongs the safe apnoea time in healthy children but has no effect to improve CO ₂ clearance	Safe apnoea time was with different oxygen flow rates (weight and age dependent) significantly longer than without supplementary oxygen	Low risk*
Dias and colleagues 2017 (India) ²⁹	RCT Two arms	95	Operating room (single centre)	<6 Months	n=47 Flow: 4 L min ⁻¹ Device: Oxiport	n=48 Flow: 0 L min ⁻¹ Device: Miller Blade	Lowest SpO ₂ Mean (SD) int. 97.55% (2.93%) vs control 95.9% (5.75%) P-value 0.049	Incidence of SpO ₂ <90% SpO ₂ 85–89% SpO ₂ <85% Correlation between time to intubation and SpO ₂	Apnoeic laryngeal oxygen insufflation with Oxiport laryngoscope blade decreases the incidence of severe desaturation during neonatal and infant intubations	The use of apnoeic oxygenation mostly in neonates requiring general anaesthesia shows a reduction in the incidence of desaturation	Some concerns*
Bruckner and colleagues 2021 (Austria) ²⁸	RCT Two arms	16	Neonatology (single centre)	<1 Month	n=7 Flow: flow rates 6 L min ⁻¹ <1 kg 7 L min ⁻¹ 1–2 kg 8 L min ⁻¹ >2 kg Device: oxygen connected direct to the tracheal tube	n=9 Flow: 0 L min ⁻¹ Device: Miller Blade	Number of intubation attempts median (IQR) int. 1 (1–2) vs control 4 (2–5) P-value 0.056	First-pass success rate Duration until successful intubation	Continuous gas flow through the endotracheal tube during intubation might result in fewer intubation attempts and a higher rate of successful intubation on the first attempt	With different oxygen flow rates during intubation occurred less aborted intubations, because of desaturation, bradycardia, or both. Multiple intubation attempts were required without supplementary oxygen and the cumulative time to successful intubation was clearly prolonged	Some concerns*
Foran and colleagues	RCT Two arms	50	Neonatology (single centre)	<1 Month	n=22 (<34 weeks 15/ ≥34 weeks 7)	n=28 (<34 weeks 18/ ≥34 weeks 10)	Duration of SpO ₂ <75% <34 weeks' gestation Median int. 29 s (0–126	Duration of SpO ₂ >85% and 65% Lowest oxygen	No significant differences were noted in duration of oxygen saturation of	The duration and not the incidence of hypoxia was defined as primary	Some concerns*

Continued

Table 1 Continued

Author year (country)	Study design	Total n	Setting	Age	Intervention	Control	Primary outcome	Secondary outcomes	Conclusions in the study	Comment	Risk of bias
2023 (Ireland) ³⁰	Stratified by gestational age				Flow: 6 L min ⁻¹ Device: Optiflow	Flow: 0 L min ⁻¹ Device: none	s) vs control 43 s (0 -132 s) (P=0.78) ≥34 weeks' gestation Median int. 0 (0-32 s) vs control 0 (0-20 s) (P-value 0.9)	saturation Bradycardia Number of intubation attempts Time to successful intubation Mortality rate	<75% between groups. The rate of desaturation appeared to decrease with use of NIF therapy in the preterm infant and postulated that greater success may be related to the increased time allowed for intubation	outcome. NIF therapy increases the rate of intubation without desaturation or bradycardia in newborns	
Gandhi and colleagues 2021 (India) ³¹	RCT Two arms	80	Operating room (single centre)	<1 Yr	n=40 Flow: 2 L min ⁻¹ Device: Oxiport	n=40 Flow: 0 L min ⁻¹ Device: Miller Blade	Lowest SpO ₂ Mean (6) int. 97.77 (2.81) vs control 92.42 (3.71) with a P-value <0.001	Time for 1% desaturation from baseline Desaturation rate Time for intubation Heart rate	Apnoeic laryngeal oxygen insufflation with Oxiport Blade decreases the incidence and severity of desaturation during intubation in neonates and infants	Significantly higher saturation was maintained with oxygen insufflation in neonates and infants compared with without oxygen	Low risk*
Olayan and colleagues 2018 (Saudi Arabia) ³²	RCT Two arms	30	Operating room (single centre)	1-8 Yr	n=15 Flow: 3 L min ⁻¹ Device: nasal cannula	n=15 Flow: 0 L min ⁻¹ Device: nasal cannula	Apnoea time until SpO ₂ 92% or time to successful intubation Median (IQR) int. 39 (19 -64) vs control 134 (21 -55) P=0.870.	SpO ₂ <95% SpO ₂ <92%	There was no difference in the duration of safe apnoea between the apnoeic oxygenation group and the standard care group.	This study defined safe apnoea duration as time to first event. The measure represents the time to successful secure the airway in both groups	Some concerns*
Steiner and colleagues 2016 (USA) ³³	RCT Three arms	457	Operating room (single centre)	1-17 Yr	Intervention 1 n=153 Flow: flow rates 2 L min ⁻¹ , blade sizes 0 -1 3 L min ⁻¹ , blade sizes 2 -3 Device: Truview Intervention 2 n=145 Flow: flow rates 2 L min ⁻¹ , blade sizes 1 -2 3 L min ⁻¹ , blade sizes 1 -2 3 L min ⁻¹ , blade size >3 Device: oxygen cannula attached to the side of standard laryngoscope	n=159 Flow: 0 L min ⁻¹ Device: standard laryngoscope	Time for 1% desaturation from baseline 25th percentile (95% CI) 75 s (37-122 s) vs int. 2.67 s (35-149) vs control 30 s (24-39 s)	SpO ₂ <90%	Laryngeal oxygen insufflation increases the time to 1% desaturation and reduces the overall rate of desaturation during laryngoscopy in children	Regardless the delivery method, supplementary oxygen reduces the rate of desaturation in apnoeic children	Some concerns*
Hodgson and colleagues 2022 (Australia) ³³	RCT Two arms	251	Delivery room or neonatal intensive care (multicentre)	<1 Month	n=124 Flow: 8 L min ⁻¹ Device: Optiflow	n=127 Flow: 0 L min ⁻¹ Device: Optiflow	First-pass success rate without physiological instability Int. 50% (62/124) control 31.5% (69/127) Adjusted risk difference 17.6 (95% CI 6.0-29.2)	Median SpO ₂ Time to and duration of desaturation Duration and number of intubation attempts	The likelihood of successful intubation on the first attempt without physiological instability in the infant was higher with high-flow therapy than with standard care	NIF therapy maintained the physiological stability and prolonged the time available to intubation	Low risk*
Soneru and colleagues 2019 (USA) ⁴⁰	Prospective cohort study	356	Operating room (single centre)	<8 Yr	n=160 Flow: 5 L min ⁻¹ Device: nasal cannula	n=196 Flow: 0 L min ⁻¹ Device: nasal cannula	SpO ₂ <95% RR before intubation P-value 0.0001	SpO ₂ <95% SpO ₂ <90% Lowest SpO ₂ Intervention by	Apnoeic nasal oxygenation during intubation of paediatric surgical patients prolongs	Especially in trainees, extended safe apnoea time is relevant for the	Serious ¹

Continued

Table 1 Continued

Author year (country)	Study design	Total n	Setting	Age	Intervention	Control	Primary outcome	Secondary outcomes	Conclusions in the study	Comment	Risk of bias
Pre-Post Studies											
Overmann and colleagues 2019 (USA) ³⁸	Observational cohort study (before and after intervention)	305	Emergency department (single centre)	<18 Yr	Postintervention n=227 Flow: flow rates <3 Yr 2 L min ⁻¹ 3-8 Yr 4 L min ⁻¹ >8 years 6 L min ⁻¹ Device: nasal cannula	Preintervention n=78 Flow: 0 L min ⁻¹ Device: none	SpO ₂ <90% post 14% (95% CI 7-24%) vs pre 22% (95% CI 17-28%); P-value 0.05	Time to desaturation Duration of desaturation Lowest SpO ₂	Apnoeic oxygenation was not associated with a lower risk of oxyhaemoglobin desaturation during RSI [†]	As a part of an intervention bundle in an emergency department, apnoeic oxygenation with low flow rates was not associated with lower desaturation during rapid sequence intubation in children	Fair [†]
Napolitano and colleagues 2019 (USA) ³⁶	Observational cohort study (before and after intervention)	1373	Paediatric intensive care unit (single centre)	<18 Yr	Postintervention n=712 Flow: flow rates <12 months 5 L min ⁻¹ 1-7 Yr 10 L min ⁻¹ >8 Yr 15 L min ⁻¹ Device: nasal cannula	Preintervention n=661 Flow: 0 L min ⁻¹ Device: none	SpO ₂ <80% post 11.8% vs pre 15.4%; P-value 0.049	SpO ₂ <70% Adverse events	Apnoeic oxygenation in PICU was associated with significant reduction in moderate and severe oxygen desaturation	The implementation of apnoeic oxygenation in a PICU was feasible and associated with less desaturation	Good [†]
Napolitano and colleagues 2023 (USA) ³⁷	Observational cohort study (before and after intervention)	6549	Paediatric intensive care unit (multicentre)	<18 Yr	Postintervention n=3318 Flow: flow rates <12 months 5 L min ⁻¹ 1-7 yr 10 L min ⁻¹ >8 Yr 15 L min ⁻¹ Device: nasal cannula	Preintervention n=3231 Flow: 0 L min ⁻¹ Device: none	Tracheal intubation associated events (TIAEs) aOR 0.75 (95% CI 0.58-0.98); P-value 0.03	SpO ₂ <80%	Apnoeic oxygenation utilisation was associated with a lower incidence of adverse TIAEs in children who required tracheal intubation in the PICU	The implementation of apnoeic oxygenation in 14 PICUs was associated with a lower occurrence of adverse events	Fair [†]
Vukovic and colleagues 2019 (USA) ³⁹	Observational cohort study (before and after intervention)	149	Emergency department (single centre)	<18 Yr	Postintervention n=90 Flow: flow rates <2 Yr 4 L min ⁻¹ 2-12 Yr 6 L min ⁻¹ >12 Yr 8 L min ⁻¹ Device: nasal cannula	Preintervention n=59 Flow: 0 L min ⁻¹ Device: none	Lowest SpO ₂ Median [IQR] post 100 [95-100] vs pre 93 [69-99]; P-value 0.001	Number of attempts	Apnoeic oxygenation is associated with decreases in hypoxaemia during paediatric endotracheal intubation	The use of apnoeic oxygenation and video laryngoscope in a paediatric emergency department showed a reduction in the incidence of hypoxaemia	Fair [†]

Table 2 Grading of Recommendations Assessment, Development, and Evaluation (GRADE) table of certainty of evidence for the main outcomes. The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI). CI, confidence interval; RR, risk ratio. *Optimal information size is not met - when the dominant study (in terms of sample size) is removed as sensitivity analysis. †Wide variance of point estimates across studies, minimal overlap of confidence intervals, high I-squared statistic (>50%) ‡Optimal information size not met - in particular when the dominant study (in terms of sample size) is removed as sensitivity analysis. §Wide variance of point estimates across studies, minimal overlap of confidence intervals, high I-squared statistic (60%) ¶Very high I-squared statistic (96%), however, this heterogeneity is largely dominated by a single study.

Patient or population: tracheal intubation in children aged 0–16 yr requiring elective or emergency tracheal intubation.

Setting: in-hospital patients.

Intervention: apnoeic oxygenation with either low- or high-flow oxygen.

Comparison: no oxygen.

Outcomes	No. of participants (studies) Follow-up	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects	
				Risk with no oxygen	Risk difference with apnoeic oxygenation with either low- or high-flow oxygen
First-pass success rate (%)	322 Bruckner and colleagues (2021) ²⁸ Foran and colleagues (2023; pre-term group) ³⁰ Foran and colleagues (2023; term group) ³⁰ Hodgson and colleagues (2022) ³³	⊕⊕○○ Low*	RR 1.27 (1.03–1.57)	51 Per 100	14 More per 100 (2 more to 29 more)
Incidence of hypoxaemia (SpO2 <90%)	769 Windpassinger and colleagues (2016) ²⁶ Dias and colleagues (2017) ²⁹ Steiner and colleagues (2016; VL-O ₂ group) ³⁴ Steiner and colleagues (2016; DL-O ₂ group) ³⁴	⊕⊕○○ Low ^{†,‡}	RR 0.24 (0.17–0.33)	40 Per 100	31 Fewer per 100 (33 fewer to 27 fewer)
Lowest SpO2 (%)	500 Dias and colleagues (2017) ²⁹ Foran and colleagues (2022; Pre-term group) Foran and colleagues (2023; term group) ³⁰ Gandhi and colleagues (2021) ³¹ Olayan and colleagues (2018) ³² Hodgson and colleagues (2022) ³³	⊕⊕⊕⊕ High	—	The mean lowest SpO2 (%) ranged from 44.3 to 99.0%	Mean difference 3.6% higher (0.8 higher to 6.5 higher)
Incidence of bradycardia (<80 beats min ⁻¹ or <100 beats min ⁻¹)	306 Foran and colleagues (2023; pre-term group) ³⁰ Foran and colleagues (2023; term group) ³⁰	⊕⊕○○ Low	RR 0.76 (0.37–1.53)	10 Per 100	2 Fewer per 100 (6 fewer to 5 more)

Continued

Table 2 Continued

Number of attempts (-)	311	Hodgson and colleagues (2022) ³³	⊕⊕⊕○	—	The mean number of attempts ranged from 1.5 to 3.7	Mean difference 0.6 lower (2.4 lower to 1.6 higher)
		Bruckner and colleagues (2021) ²⁸	Moderate [†]			
First intubation success time (s)	1117	Foran and colleagues (2023; pre-term group) ³⁰			The mean first intubation success time ranged from 19.1 to 74.0 s	Mean difference 10.5 s higher (4.4 lower to 25.4 higher)
		Foran and colleagues (2023; term group) ³⁰				
		Hodgson and colleagues (2022) ³³	⊕⊕⊕○	—		
		Ledbetter and colleagues (1988) ²⁷	Moderate [§]			
		Windpassinger and colleagues (2016) ²⁶				
		Dias and colleagues (2017) ²⁹				
		Gandhi and colleagues (2021) ³¹				
	Steiner and colleagues (2016; VL-O ₂ group) ³⁴					
	Steiner and colleagues (2016; DL-O ₂ group) ³⁴					
	Hodgson and colleagues (2022) ³³					


GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.



attempts (153 neonates with apnoeic oxygenation and 158 in the control group; Fig. 4c). No statistical difference in the number of intubation attempts was detected (mean difference -0.6 , 95% CI -2.4 to 1.2 , $P=0.35$). Between-study heterogeneity was substantial ($I^2=60\%$), and the certainty of the evidence was graded moderate because of a wide variance of point estimates across studies and minimal overlap of confidence intervals.

Lowest oxygen saturation

Five RCTs^{29–33} with six comparisons reported the lowest haemoglobin oxygen saturation during tracheal intubation for 500 children, 248 children in the apnoeic oxygenation group and 252 in the control group (Fig. 3b). Patients in the apnoeic oxygenation group had a higher oxygen saturation than those in the control group (mean difference 3.6%, 95% CI 0.8–6.5%, $P=0.02$). Substantial between-study heterogeneity ($I^2=63\%$) was detected. The certainty of evidence was graded high.

Incidence of hypoxaemia

Three RCTs^{26,29,34} with 769 children reported the incidence of hypoxaemia during tracheal intubation (379 children in the apnoeic oxygenation group and 390 children in the control group; Fig. 4a). Overall pooled estimate showed a significant reduction in hypoxaemia incidence (RR 0.24, 95% CI 0.17–0.33, $P<0.01$). Between-study heterogeneity was substantial ($I^2=51\%$). The certainty of evidence was graded very low because of a wide variance of point estimates across studies, a minimal overlap of confidence intervals, a high I^2 statistic, and as optimal information sizes were not met. More importantly, 99.4% of the pooled effect size is derived from the two group comparisons of a single study.³⁴

Incidence of bradycardia

Two RCTs^{30,33} reported the incidence of bradycardia with 306 children (146 in the apnoeic oxygenation group and 160 in the control group; Fig. 4b). No reduction was detected in the




Fig 2. Risk of bias assessment. NIH, National Institutes of Health.

incidence bradycardia when applying apnoeic oxygenation (RR 0.76, 95% CI 0.37–1.53, $P=0.44$; $I^2=0$). The study by Hodgson and colleagues³³ contributed the entire weight (100%) to the pooled estimates. The certainty of evidence was graded low.

Time to successful intubation

Six RCTs^{26,27,29,31,33,34} with seven group comparisons reported the time to successful intubation with a total of 1117 children




Fig 3. Forest plots for the outcomes (a) first-pass success rate, (b) lowest SpO₂, and (c) apnoea times during paediatric tracheal intubation. CI, confidence interval; df, degrees of freedom; MH, Mantel-Haenszel; sd, standard deviation.

(553 children with apnoeic oxygenation and 564 in the control group; Fig. 4d). The pooled estimate showed no difference in the intubation success time (mean difference 10.5 s, 95% CI –4.4 to 25.4 s, $P=0.14$). Between-study heterogeneity was very high ($I^2=96\%$), and the certainty of evidence was graded moderate as the heterogeneity was largely dominated by a single study.³³

Apnoea time

Two RCTs^{26,33} reported apnoea times during paediatric tracheal intubation with a total of 296 patients (148 children with apnoeic oxygenation and 148 in the control group; Fig. 3c). No difference was detected in apnoea times (mean difference 19 s, 95% CI –143 to 182 s, $P=0.38$). Between-study heterogeneity was substantial ($I^2=60\%$), and the random effects models demonstrated large uncertainties in the pooled estimates as the two studies differed in their mean estimates and sd by a factor of almost 4. The evidence was not graded as the two studies used different apnoea time definitions. The first study³³ reported on duration of apnoea for intubation, whereas in the other study²⁶ children were first intubated and

then left apnoeic until the onset of desaturation with or without oxygenation.

Adverse events

Two RCTs^{28,33} reported adverse events during the tracheal intubation. Hodgson and colleagues³³ reported serious adverse events such as chest compressions or epinephrine administration within 1 h after tracheal intubation in 0 of the 124 patients with apnoeic oxygenation vs 1.6% ($N=2/127$) in the control group. Pneumothorax was reported in 1.6% ($N=2/124$) patients in the apnoeic oxygenation group and 4.7% ($N=6/127$) in the control group. Death within 72 h after tracheal intubation in 0.8% ($N=1/124$) of patients with apnoeic oxygenation and 2.4% ($N=3/127$) in the control group. Bruckner and colleagues²⁸ reported aborted tracheal intubation because of desaturation, bradycardia, or both in 33% ($N=3/10$) of intubation attempts with apnoeic oxygenation and in 69% ($N=20/29$) when intubation was attempted without oxygen. Because of variability in methods and reporting across these studies, no meta-analysis was conducted.




Fig 4. Forest plots for the outcomes (a) incidence of hypoxaemia, (b) bradycardia, (c) number of intubation attempts, and (d) first pass intubation success time. CI, confidence interval; df, degrees of freedom; MH, Mantel-Haenszel; sd, standard deviation.

Discussion

This systematic review and meta-analysis investigated the effectiveness of apnoeic oxygenation during tracheal intubation in children under 16 yr of age. Despite the low certainty of evidence according to GRADE, apnoeic oxygenation was

associated with a higher probability of first-pass tracheal intubation success and a reduced number of intubation attempts for each patient. Regardless of the method of administration, apnoeic oxygenation reduced the incidence of hypoxaemia when compared with no oxygen administration. Apnoeic oxygen administered during tracheal intubation

improved and stabilised respiratory and haemodynamic variables and facilitated overall airway management.

First-attempt success is critical in paediatric airway management, as adverse events are directly associated with the overall number of intubation attempts.^{2,41,42} Our findings indicate that administration of oxygen reduces the overall number of attempts by improving first attempt success rate. This is highly important for neonates, small infants, and children with limited cardiopulmonary reserve, as they are more prone to rapid oxygen desaturation. These patients often have higher oxygen consumption, lower closing capacity, low functional residual capacity, and increased risk of airway collapse compared with older children.⁴³

The main obstacle of tracheal intubation in children is often the short apnoea time before severe arterial oxygen desaturation, which is more severe in neonates, infants, and those with severe comorbidities.¹² Thus, methods to extend the safe apnoea time are highly desirable. Across the studies included in this systematic review, oxygen delivery techniques, flow rates, and significant hypoxaemia levels were heterogeneous. Nevertheless, in eight RCTs^{26–29,31,33–35} apnoeic oxygenation during intubation prolonged the safe apnoea time, decreased the incidence of hypoxaemia, and reduced the number of intubation attempts or increased first-pass success without physiological instability. These findings are supported by five observational or pre-post studies^{36–40} that were not included in our meta-analyses but showed a benefit of apnoeic oxygenation during tracheal intubation with fewer adverse events, better intubation conditions, and reduction of hypoxaemia. Maintaining adequate oxygenation during tracheal intubation is crucial in neonates and infants to prevent hypoxaemia and the associated complications.^{10,12,44,45}

On the one hand, using oxygen at high concentrations in premature babies and infants can lead to complications triggered by oxidative stress, including bronchopulmonary dysplasia⁴⁶ or severe retinopathy of prematurity.⁴⁷ Administering supplemental oxygen to children with a cyanotic congenital heart disease might lead to perfusion mismatch, resulting in haemodynamic instability.⁴⁸ However, hypoxaemia is also harmful, and haemoglobin oxygen saturation values of 85–89% in premature babies increase the risk of death before hospital discharge.⁴⁹ In addition to potential harm with oxygen, there is also the risk of barotrauma. Hodgson and colleagues³³ diagnosed pneumothorax more often in patients without apnoeic oxygenation and Napolitano and colleagues³⁷ showed fewer adverse events, including pneumothorax, in an observational study. Causality remains unclear, but rescue facemask ventilation during intubation attempts might pose a risk of barotrauma rather than apnoeic oxygenation. High-flow nasal oxygen administration was reported to reduce the risk of pneumothorax compared with continuous positive airway pressure for respiratory support in preterm infants.⁵⁰ In summary, the relatively short period of potential hyperoxia during intubation with apnoeic oxygenation appears to outweigh the benefits of avoiding hypoxaemia

Limitations

The significant heterogeneity in study designs of the included trials constitutes a major limitation together with the limited number of included patients and studies. A further limitation is our assumption of the independence of different treatment arms³⁴ or subgroups.³⁰ It is likely that the true uncertainty of

the pooled effect sizes (i.e. the confidence intervals), might be wider than reported in this study.

Avoiding hypoxaemia and keeping a normal oxygen saturation is the first step of a cascade of events that might lead to serious adverse events such as bradycardia or cardiac arrest. The aggregated evidence from this systematic review confirms that apnoeic oxygenation during airway management reduces the incidence of hypoxaemia, but the strength of evidence remains low. This is partly as a result of the exclusion of several studies investigating apnoeic oxygenation because of the lack of a control group with no intervention.^{15,51,52} Moreover, only two RCTs^{28,33} reported data on adverse events, both of which reported a higher incidence of adverse events in children with no apnoeic oxygen administration. However, data were insufficient and not consistently reported to perform a meta-analysis. This is also supported by the findings of the observational studies by Napolitano and colleagues.^{36,37} They found an association between oxygen administration and a lower incidence of severe adverse events during tracheal intubation but not a reduction in tracheal intubation attempts or severe peri-intubation hypoxaemia.

Unanswered questions and future research

This systematic review illustrates a high variability in the choice of outcome measures across the included studies. Future study designs on paediatric airway management should standardise the choice of outcomes with actual clinical relevance, allowing better comparison and meta-analysis of studies. First-pass success rate of tracheal intubation and adverse events should be included in any future research as they represent the clinical outcomes of highest relevance. The optimal technique for oxygen delivery during intubation, oxygen flow rate, and concentration of oxygen to be administered during oxygenation have yet to be determined. As financial resources in healthcare and environmental consciousness are increasingly important, cost-effectiveness and environmental impact analysis should be included in future studies.

Severe hypoxaemia, severe bradycardia, cannot intubate and cannot ventilate, emergency front of neck access, cardiac arrest, and death were rarely reported in the included studies, given the relatively small sample sizes of the included individual studies.

Conclusions

This systematic review provides evidence for improved first-attempt intubation success rate during tracheal paediatric intubation with apnoeic oxygenation. Apnoeic oxygenation during tracheal intubation in children decreases the risk of oxygen desaturation. However, as the included data were heterogeneous, more high-quality randomised controlled trials are warranted with a well-defined core set of outcome variables.

Authors' contributions

Wrote the protocol and manuscript: all authors

Study statistician: MH

Medical librarian: MvG

Articles were extracted by: GK, JA, RB, ND, ACL, MvG, TR, AF
Methodologists: AA, CSR

Were involved in the data analysis and interpretation and read and approved the final manuscript: all authors

Declaration of interests

The authors declare that they have no conflicts of interest.

Funding

This study was investigator-initiated and financed solely with departmental grants. The departments had no influence in the conduct of the study or interpretation of the data. We thank the departments for their support of protected research time.

Data sharing

The data presented in this manuscript are available.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bja.2023.10.039>.

References

- Disma N, Virag K, Riva T, et al. Difficult tracheal intubation in neonates and infants. NEonate and Children audit of Anaesthesia pRactice IN Europe (NECTARINE): a prospective European multicentre observational study. *Br J Anaesth* 2021; **126**: 1173–81
- Fiadjo JE, Nishisaki A, Jagannathan N, et al. Airway management complications in children with difficult tracheal intubation from the Pediatric Difficult Intubation (PeDI) registry: a prospective cohort analysis. *Lancet Respir Med* 2016; **4**: 37–48
- Foglia EE, Ades A, Sawyer T, et al. Neonatal intubation practice and outcomes: an international registry study. *Pediatrics* 2019; **143**: e20180902
- Engelhardt T, Virag K, Veyckemans F, Habre W, Network AGotESoACT. Airway management in paediatric anaesthesia in Europe-insights from APRICOT (Anaesthesia Practice in Children Observational Trial): a prospective multicentre observational study in 261 hospitals in Europe. *Br J Anaesth* 2018; **121**: 66–75
- Nishisaki A, Ferry S, Colborn S, et al. Characterization of tracheal intubation process of care and safety outcomes in a tertiary pediatric intensive care unit. *Pediatr Crit Care Med* 2012; **13**: e5–10
- Stinson HR, Srinivasan V, Topjian AA, et al. Failure of invasive airway placement on the first attempt is associated with progression to cardiac arrest in pediatric acute respiratory compromise. *Pediatr Crit Care Med* 2018; **19**: 9–16
- Shiima Y, Berg RA, Bogner HR, et al. Cardiac arrests associated with tracheal intubations in PICUs: a multicenter cohort study. *Crit Care Med* 2016; **44**: 1675–82
- Schwartz DE, Matthay MA, Cohen NH. Death and other complications of emergency airway management in critically ill adults. A prospective investigation of 297 tracheal intubations. *Anesthesiology* 1995; **82**: 367–76
- Holm-Knudsen RJ, Carlin JB, McKenzie IM. Distress at induction of anaesthesia in children. A survey of incidence, associated factors and recovery characteristics. *Pediatr Anesth* 1998; **8**: 383–92
- Kothari R, Hodgson KA, Davis PG, Thio M, Manley BJ, O'Curra E. Time to desaturation in preterm infants undergoing endotracheal intubation. *Arch Dis Child Fetal Neonatal Ed* 2021; **106**: 603–7
- Bhatia PK, Bhandari SC, Tulsiani KL, Kumar Y. End-tidal oxygraphy and safe duration of apnoea in young adults and elderly patients. *Anaesthesia* 1997; **52**: 169–78
- Patel R, Lenczyk M, Hannallah RS, McGill WA. Age and the onset of desaturation in apnoeic children. *Can J Anaesth* 1994; **41**: 771–4
- Volhard F. Ueber kuenstliche Atmung durch Ventilation der Trachea und eine einfache Vorrichtung zur rhythmischen kuenstlichen Atmung. *Munch Med Wochenschr* 1908; **55**: 209–11
- Meltzer S, Auer J. Continuous respiration without respiratory movements. *J Exp Med* 1909; **11**: 622–5
- Riva T, Pedersen TH, Seiler S, et al. Transnasal humidified rapid insufflation ventilatory exchange for oxygenation of children during apnoea: a prospective randomised controlled trial. *Br J Anaesth* 2018; **120**: 592–9
- Else SDN, Kovatsis PG. A narrative review of oxygenation during pediatric intubation and airway procedures. *Anesth Analg* 2020; **130**: 831–40
- King MR, Jagannathan N. Best practice recommendations for difficult airway management in children-is it time for an update? *Br J Anaesth* 2018; **121**: 4–7
- Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021; **88**, 105906
- von Gernler M, Riva T, Fuchs A, Romero CS, Greif R. Apneic oxygenation during pediatric tracheal intubation [Ovid MEDLINE]. *searchRxiv* 2023: 2023. <https://doi.org/10.1079/searchRxiv.2022.00081>
- Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan-a web and mobile app for systematic reviews. *Syst Rev* 2016; **5**: 210
- Sterne JAC, Savovic J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019; **366**: l4898
- Sterne JA, Hernan MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016; **355**: i4919
- National Heart L, Institute B. *Study quality assessment tools: for observational cohort and cross-sectional studies, for before-after (pre-post) studies with no control group, for case-control studies*. Maryland: USA Im Internet; 2019. Available from: www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools. [Accessed 23 November 2023]
- Atkins D, Best D, Briss PA, et al. GRADE Working Group. Grading quality of evidence and strength of recommendations. *BMJ* 2004; **328**: 1490–4
- R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2022
- Windpassinger M, Plattner O, Gemeiner J, et al. Pharyngeal oxygen insufflation during airtraq laryngoscopy slows arterial desaturation in infants and small children. *Anesth Analg* 2016; **122**: 1153–7
- Ledbetter JL, Rasch DK, Pollard TG, Helsen P, Smith RB. Reducing the risks of laryngoscopy in anaesthetised infants. *Anaesthesia* 1988; **43**: 151–3
- Bruckner M, Morris NM, Pichler G, et al. In newborn infants a new intubation method may reduce the number of intubation attempts: a randomized pilot study. *Children (Basel)* 2021; **8**: 553

29. Dias R, Dave N, Chhabria R, Shah H, Garasia M. A randomised comparative study of Miller laryngoscope blade versus Oxiport((R)) Miller laryngoscope blade for neonatal and infant intubations. *Indian J Anaesth* 2017; **61**: 404–9
30. Foran J, Moore CM, Ni Chathasaigh CM, Moore S, Purna JR, Curley A. Nasal high-flow therapy to Optimise Stability during Intubation: the NOSI pilot trial. *Arch Dis Child Fetal Neonatal Ed* 2023; **108**: 244–9
31. Gandhi N, Bhatt S, Goswami A. Comparative study of oxiport laryngoscope blade versus miller laryngoscope blade for intubation in neonates and infants during general anesthesia, a prospective randomized controlled study. *Indian J Clin Anaesth* 2021; **8**: 356–60
32. Olayan L, Alatassi A, Patel J, Milton S. Apnoeic oxygenation by nasal cannula during airway management in children undergoing general anaesthesia: a pilot randomised controlled trial. *Perioper Med (Lond)* 2018; **7**: 3
33. Hodgson KA, Owen LS, Kamlin COF, et al. Nasal high-flow therapy during neonatal endotracheal intubation. *N Engl J Med* 2022; **386**: 1627–37
34. Steiner JW, Sessler DI, Makarova N, et al. Use of deep laryngeal oxygen insufflation during laryngoscopy in children: a randomized clinical trial. *Br J Anaesth* 2016; **117**: 350–7
35. Humphreys S, Lee-Archer P, Reyne G, Long D, Williams T, Schibler A. Transnasal humidified rapid-insufflation ventilatory exchange (THRIVE) in children: a randomized controlled trial. *Br J Anaesth* 2017; **118**: 232–8
36. Napolitano N, Laverriere EK, Craig N, et al. Apneic oxygenation as a quality improvement intervention in an academic PICU. *Pediatr Crit Care Med* 2019; **20**: e531–7
37. Napolitano N, Polikoff L, Edwards L, et al. Effect of apneic oxygenation with intubation to reduce severe desaturation and adverse tracheal intubation-associated events in critically ill children. *Crit Care* 2023; **27**: 26
38. Overmann KM, Boyd SD, Zhang Y, Kerrey BT. Apneic oxygenation to prevent oxyhemoglobin desaturation during rapid sequence intubation in a pediatric emergency department. *Am J Emerg Med* 2019; **37**: 1416–21
39. Vukovic AA, Hanson HR, Murphy SL, Mercurio D, Sheedy CA, Arnold DH. Apneic oxygenation reduces hypoxemia during endotracheal intubation in the pediatric emergency department. *Am J Emerg Med* 2019; **37**: 27–32
40. Soneru CN, Hurt HF, Petersen TR, Davis DD, Braude DA, Falcon RJ. Apneic nasal oxygenation and safe apnea time during pediatric intubations by learners. *Paediatr Anaesth* 2019; **29**: 628–34
41. Garcia-Marcinkiewicz AG, Kovatsis PG, Hunyady AI, et al. First-attempt success rate of video laryngoscopy in small infants (VISI): a multicentre, randomised controlled trial. *Lancet* 2020; **396**: 1905–13
42. Riva T, Engelhardt T, Basciani R, et al. Direct versus video laryngoscopy with standard blades for neonatal and infant tracheal intubation with supplemental oxygen: a multicentre, non-inferiority, randomised controlled trial. *Lancet Child Adolesc Health* 2022; **7**: 101–11
43. Hardman JG, Wills JS. The development of hypoxaemia during apnoea in children: a computational modelling investigation. *Br J Anaesth* 2006; **97**: 564–70
44. Hatch LD, Grubb PH, Lea AS, et al. Endotracheal intubation in neonates: a prospective study of adverse safety events in 162 infants. *J Pediatr* 2016; **168**: 62–66.e6
45. Humphreys S, Pham TM, Stocker C, Schibler A. The effect of induction of anesthesia and intubation on end-expiratory lung level and regional ventilation distribution in cardiac children. *Paediatr Anaesth* 2011; **21**: 887–93
46. Kaplan E, Bar-Yishay E, Prais D, et al. Encouraging pulmonary outcome for surviving, neurologically intact, extremely premature infants in the postsurfactant era. *Chest* 2012; **142**: 725–33
47. Hellstrom A, Smith LE, Dammann O. Retinopathy of prematurity. *Lancet* 2013; **382**: 1445–57
48. Marino BS, Bird GL, Wernovsky G. Diagnosis and management of the newborn with suspected congenital heart disease. *Clin Perinatol* 2001; **28**: 91–136
49. Sola A, Golombek SG, Montes Bueno MT, et al. Safe oxygen saturation targeting and monitoring in preterm infants: can we avoid hypoxia and hyperoxia? *Acta Paediatr* 2014; **103**: 1009–18
50. Hong H, Li XX, Li J, Zhang ZQ. High-flow nasal cannula versus nasal continuous positive airway pressure for respiratory support in preterm infants: a meta-analysis of randomized controlled trials. *J Matern Fetal Neonatal Med* 2021; **34**: 259–66
51. Riva T, Preel N, Theiler L, et al. Evaluating the ventilatory effect of transnasal humidified rapid insufflation ventilatory exchange in apnoeic small children with two different oxygen flow rates: a randomised controlled trial. *Anaesthesia* 2021; **76**: 924–32
52. Riva T, Seiler S, Stucki F, Greif R, Theiler L. High-flow nasal cannula therapy and apnea time in laryngeal surgery. *Paediatr Anaesth* 2016; **26**: 1206–8

Handling Editor: Hugh C Hemmings Jr