Published for SISSA by O Springer

RECEIVED: November 25, 2022 ACCEPTED: April 18, 2023 PUBLISHED: April 28, 2023

3 definitions of BF theory on homology 3-spheres

Matthias Blau,^a Mbambu Kakona^{b,c} and George Thompson^c

^aAlbert Einstein Center for Fundamental Physics,

Institute for Theoretical Physics, University of Bern,

Sidlerstrasse 5, CH-3012 Bern, Switzerland

^bEast African Institute for Fundamental Research (EAIFR), University of Rwanda,

KN 7 Ave, Kigali, Rwanda

^cAbdus Salam International Centre for Theoretical Physics, Strada Costiera 11 Trieste, Italy

E-mail: blau@itp.unibe.ch, kakona@eaifr.org, thompson@ictp.it

ABSTRACT: 3-dimensional BF theory with gauge group G (= Chern-Simons theory with non-compact gauge group TG) is a deceptively simple yet subtle topological gauge theory. Formally, its partition function is a sum/integral over the moduli space \mathcal{M} of flat connections, weighted by the Ray-Singer torsion. In practice, however, this formal expression is almost invariably singular and ill-defined.

In order to improve upon this, we perform a direct evaluation of the path integral for certain classes of 3-manifolds (namely integral and rational Seifert homology spheres). By a suitable choice of gauge, we sidestep the issue of having to integrate over \mathcal{M} and reduce the partition function to a finite-dimensional Abelian matrix integral which, however, itself requires a definition. We offer 3 definitions of this integral, firstly via residues, and then via a large k limit of the corresponding $G \times G$ or $G_{\mathbb{C}}$ Chern-Simons matrix integrals (obtained previously). We then check and discuss to which extent the results capture the expected sum/integral over all flat connections.

KEYWORDS: Chern-Simons Theories, Topological Field Theories

ARXIV EPRINT: 2211.10136

source: https://doi.org/10.48350/189691 | downloaded: 8.5.2024



Contents

1	Introduction			
2	 BF-theory on a 3-manifold and path integrals 2.1 BF theory as a TG Chern-Simons theory 2.2 Path integral for BF theory: formal considerations 2.3 Path integral for BF theory and massive Ray-Singer torsion 2.4 Expectations for the partition function on Seifert manifolds 	6 6 7 9 10		
3	 F-theory on a Seifert fibred 3-manifold Seifert 3-manifolds briefly BF action and gauge transformations on a Seifert manifold Gauge fixing 1: temporal gauge for A and B Gauge fixing 2: Abelianisation and emergence of non-trivial line bundles Reduction of the partition function to a finite-dimensional integral 			
4	Definition 1: direct evaluation of the <i>BF</i> (path) integral 4.1 Symmetries and other properties of the finite dimensional integrals 4.2 The ϕ_C contribution to \mathbb{Z} homology spheres 4.3 The ϕ_C contribution to \mathbb{Q} homology spheres 4.4 A residue formula for SU(2) 4.5 SU(2) <i>BF</i> theory on Brieskorn spheres $\Sigma(a_1, a_2, a_3)$ 4.6 A residue formula for higher rank and outlook	19 19 20 22 24 26 27		
5	Large k asymptotics of $G \times G$ and $G_{\mathbb{C}}$ Chern-Simons theory 5.1 Definition 2: BF and the large k limit of $G_k \times G_{-k}$ Chern-Simons theory 5.2 Definition 3: BF and the large k or s limit of $G_{\mathbb{C}}$ Chern-Simons theory			
Α	 The tangent bundle group TG and its Lie algebra tg A.1 Basic properties of TG and tg A.2 Diagonalisation / Abelianisation in TG and tg 			
в	 A symmetry of complex Chern-Simons theory on Seifert 3-manifolds B.1 Discrete symmetries B.2 Holomorphic factorisation 	39 40 42		

1 Introduction

Gauge theories with non-compact gauge groups are notoriously difficult to make sense of, issues including questions of convergence and propagators with the wrong signs which make unitarity and hence the physical meaning of the theory far from clear. One set of theories for which one may make some headway are topological field theories where a particle interpretation is not required. Our first investigation in that direction where the partition function can be evaluated exactly [1] was to consider the Schwarz type theories [2–4] formally representing the Ray-Singer Torsion [5, 6] which have non-compact symmetries. These symmetries though are essentially Abelian and we would like to consider now a class of theories with non-Abelian non-compact gauge groups. In particular we have in mind the topological BF theories [7–10] based on a compact gauge group G. In two dimensions such a theory is the zero coupling limit of Yang-Mills theory with gauge group G and as such does not have an associated non-compact symmetry. In dimension greater than or equal to 3, however, BF theories possess additional non-compact shift symmetries. We will focus on 3-dimensional BF theory here, where the combined gauge group turns out to be the non-compact group TG, the tangent bundle group of G.

Witten [11] first introduced 3-dimensional BF theories as a variant of Chern-Simons theory [12] for non-compact gauge groups (in fact, it can be regarded as a Chern-Simons theory for the non-compact gauge group TG). The action is

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B \wedge F_{A} \right) \tag{1.1}$$

where A is a connection on a G-bundle over M and B is a Lie algebra valued 1-form. The action enjoys the usual compact G gauge symmetry, as well as a non-compact shift gauge symmetry $B \to B + d_A \Lambda$. This theory is deceptively simple. The path integral over B yields a delta function constraint on the curvature 2-form F_A , so that formally the path integral for the partition fuction of BF theory is

$$Z_{BF}[M,G] = \int DADB \, \exp\left(i \int_{M} \operatorname{Tr}\left(B \wedge F_{A}\right)\right) \simeq \int DA\delta\left(F_{A}\right) \tag{1.2}$$

This seems to suggest that the path integral simply reduces to an integral over the moduli space of flat connections, with some measure. In general, however, this is not correct. Rather, the complete classical equations of motion are

$$F_A = 0, \quad d_A B = 0 \tag{1.3}$$

which taken together (and modulo gauge transformations) can formally be viewed as describing the tangent bundle of the moduli space $\mathfrak{M}[M,G]$ of flat G connections on the 3-manifold M, $T\mathfrak{M}[M,G]$. The partition function is then an integral over this space (and ghost zero modes etc.), again with some measure to be determined. However, in general this space can be very singular and the finite dimensional integral is ill-defined. The various types of singularities that lurk inside the path integral associated with the action (1.1) include issues with reducible connections (manifested as ghost zero modes) as well as the non-compactness of the moduli space itself (due to B zero modes).

One situation in which it is pretty clear what sort of contribution one should find from the path integral is when one expands the path integral around an isolated irreducible flat connection. In that case, there are no non-trivial solutions to the equations of motion for Band no ghost zero modes, and following Witten [11], the contribution of such an *acyclic* connection ω on a 3-manifold M can be shown to be precisely the Ray-Singer analytic torsion $\tau_M(\omega)$ [5, 6] of that connection. Thus on manifolds on which the moduli space of flat connections includes such acyclic flat connections, one should perhaps expect the partition function to take the form

$$Z_{BF}[M,G] = \sum_{\text{acyclic}} \tau_M(\omega_{(\alpha)}) + \dots$$
(1.4)

where the ellipses indicate the contributions from the non-acyclic flat connections. If M is an Integral Homology Sphere (\mathbb{Z} Homology Sphere, $H_1(M, \mathbb{Z}) = 0$), then for G = SU(2) the trivial connection is the only reducible flat connection, and there are indeed \mathbb{Z} Homology Spheres for which all the flat connections are isolated. In such a situation one should expect the BF partition function to be largely captured by the above expression. One of the aims of this paper will be to check to which extent this expectation is borne out by actual explicit evaluations of the path integral of BF theory.

Moreover, by using the Massive Ray-Singer Torsion $\tau_M(\omega, \mu)$ introduced in [1] (among other reasons precisely with its application to BF theory in mind), one can extend the considerations of Witten to the case that the flat connections are not necessarily irreducible and isolated, and thus obtain a formal expression for the regularised partition function of BF theory (section 2), which takes the form (2.20)

$$Z_{BF}[M,G;\mu] = \sum \tau_M(\omega_{(\alpha)},\mu)$$
(1.5)

of a sum / integral over the moduli space of flat connections. This expression can in turn be expanded in the mass-parameter μ to reproduce different kinds of contributions to the partition function [1]. However, this approach outlined above to defining the partition function of BF theory is rather formal: one would need to determine by hand the spaces of flat connections and zero modes associated with them, calculate the (massive) Ray-Singer torsion by some means, etc. So this does not all by itself lead to an *evaluation* of the BFpartition function via path integral methods.

In order to be able to verify to which extent an actual evaluation of the BF path integral reproduces the above formal expectations, one needs a class of manifolds for which explicit path integral calculations can be performed. To that end we will concentrate on Seifert fibred 3-manifolds. These spaces allow for a significant simplification in the calculations via suitable gauge choices. The first of these is that the 'time' (fibre) components of both A and B (the components of A and B along the fibres of the defining Seifert fibrations) can be taken to be constant along the fibre. This is a generalisation of the usual temporal gauge $\partial_0 A_0 = 0$ of a gauge field on a circle, and allows one to partially push the calculation down a dimension, to the base of the Seifert fibration. And now further simplification comes from the approach [13-18] to evaluating the path integral through Abelianisation. For Chern-Simons theory with gauge group G the end result is a finite dimensional integral over the Cartan subalgebra t of the Lie algebra \mathfrak{g} of G. One of the characteristic features of this approach is that the reduction to a finite-dimensional integral (over a linear space) completely by passes the need to define (and integrate over) the moduli space of flat connections. Nevertheless, this approach can be shown to reproduce the results of perturbative or localisation calculations (when available), which in principle require an exact evaluation of the latter.

Returning to BF theory, the main technical result of this paper is to show that this calculational method can be extended to BF theory with its non-compact gauge group TG. The result is a finite-dimensional integral over the space $\mathfrak{t} \oplus \mathfrak{t}$, which plays the role of the Cartan subalgebra of the Lie algebra of TG. This once again bypasses the need to integrate over the moduli space of flat connections. The finite-dimensional integrals that we find are similar to those for Chern-Simons theory, but they are more singular than their Chern-Simons counterparts, having poles on the integration contour an integral over λ which is a distribution, and hence require a definition. For example, the partition function for SU(2) and M a \mathbb{Z} Homology Sphere devolves to

$$Z_{BF}[M, \mathrm{SU}(2)] \simeq \sum_{n_0} \int_{\mathbb{R} \times \mathbb{R}} \tau_M(\phi) \, . \, \exp\left(i\lambda \left(\phi/P + 2\pi i n_0\right)\right) \tag{1.6}$$

(see the body of the paper for all the definitions). The important point to note here is that the Ray-Singer Torsion $\tau_M(\phi)$ has poles at $\phi = m\pi$ $m \neq sP$ and zero when m is proportional to P. The λ integral sets $\phi = Pn_0$ which formally avoids the poles. Still we can only conclude what the integral should be after it is properly regularised (defined).

In the case of Chern-Simons theory the integrals that we find there also have poles, however, there we could argue that the contributions at the poles should not be included (lying on the walls of the Weyl chamber) and we introduced a mass regulator that essentially allowed us to avoid them. Unfortunately, in the present setting of BF theory, we do not have a guiding principle and so we offer 3 different possible definitions of BF theory and discuss their advantages as well as their shortcomings:

1. Direct Definition via Residues

The first, and the one we spend most time on (section 4), is through the direct definition of the finite-dimensional integral (1.6). In order to get a handle on the poles of the Ray-Singer Torsion on $t \times t$, one defines the theory in such a way that it is given by the residue of all the possible poles including delta function contributions. We find that, rather remarkably, this does reproduce the expected contributions of *reducible* flat connections which we can follow by making use of the Massive Ray-Singer Torsion. In particular we show in detail how this gives the expected results, as a sum over contributions from the isolated non-trivial flat connections, for Lens spaces. Indeed, this definition appears to capture the essence of the contribution of reducible connections in general. However, this approach does not reproduce the expected result (1.4) for connections that are isolated and irreducible. We note that, suprisingly, some of the poles on the contour correspond to irreducible flat connections, while others correspond to 'complex' flat connections. We offer a brief attempt at an explanation (related to gauge fixing) for why the irreducible connections arise in this way.

In view of this, we approach the problem from a different point of view in section 5, based on the fact that the gauge group TG can be regarded as a contraction of either the compact group $G \times G$ or the complex group $G_{\mathbb{C}}$. Correspondingly, the other 2 definitions

that we consider are to regard BF theory as arising either in the large k limit of $G \times G$ Chern-Simons theory at levels k and -k, or in certain limits of the coupling t = k + is of $G_{\mathbb{C}}$ Chern-Simons theory. This regularises the theory to some extent and also side-steps the gauge-fixing issue alluded to above.

2. Definition via a large k Limit of $G \times G$ Compact Chern-Simons Theory

The advantage of using the compact $G \times G$ theory is that the Hilbert space of states is finite dimensional and only goes over to the infinite dimensional Hilbert space of states of BF theory in the limit. This therefore acts as a natural regulator. A disadvantage is that certain conditions must be met by the connections so that the correspondence that we need exists. We offer examples of Brieskorn \mathbb{Z} Homology Spheres (Seifert manifolds with 3 exceptional fibres) [19] where the conditions are met and one obtains the partition function of BF theory in the expected form (1.4). These conditions are not met by all manifolds, however. For example, the equality between BF theory and the large k limit of the $G \times G$ theory fails for certain Lens spaces. As the Lens spaces are \mathbb{Q} Homology Spheres one would like to conjecture that the correspondence holds for isolated flat connections of \mathbb{Z} Homology Spheres in general.

3. Definition via a large k Limit of $G_{\mathbb{C}}$ Complex Chern-Simons Theory

As for $G_{\mathbb{C}}$ Chern-Simons theory [20–22] with complex level k+is, there are many formal correspondences with BF theory. The most obvious is to set k = 0 and let $s \to \infty$, which leads directly to the BF action (1.1) and its non-compact symmetries (2.2). Given the difficulties we face with this action, this is not the limit that we consider. Rather, we take s = 0 and $k \to \infty$. An advantage of this approach is that the finite dimensional integrals that arise are slight variants of those discussed by Lawrence and Rozansky [24] for SU(2) and for general G by Mariño [25] in the context of G Chern-Simons theory. In principle then the same strategies that apply there can be used here, though we leave that for the future.

To the extent that the perturbative large k evaluation of Chern-Simons theory reduces to a sum over contributions from flat connections, this is then also true for the partition function of BF theory. Overall, the definition of BF theory via $G_{\mathbb{C}}$ Chern-Simons theory appears to be the most complete (however, one needs to ensure that one does not count complex flat connections that cannot be conjugated into flat G connections).

One approach not followed here but which is certainly of interest is resurgence [26]. Gukov, Mariño and Putrov [27] show that for SU(2) one can start with the Abelian contribution to the Chern-Simons partition function in the large k limit and Borel resum in order to obtain contributions from non-Abelian flat connections. Once in the complex space of connections there is the possibility that $SL(2, \mathbb{C})$ connections which are not (conjugate to) SU(2) flat connections will contribute. They show that miraculously this does not happen. The importance of this for us is that, as we will see, we appear to obtain contributions from connections in our first approach that would, from this point of view be considered to be 'complex'.

This paper is organised as follows. Section 2 follows Witten's [11] approach to obtaining the partition function for isolated irreducible flat connections. There we also consider what formally happens when the connections are either not isolated or are reducible. In these cases we can profitably make use of the massive Ray-Singer Torsion introduced in [1]. This allows us to express our expectations for the form of the partition function in concrete terms.

In section 3 we formulate BF theory on Seifert 3-manifolds which are either \mathbb{Z} or \mathbb{Q} homology spheres. Particular attention is paid to issues with gauge fixing of ϕ and λ , the components of A and B along the fibre respectively. We will shows that ϕ and (with an assumption about ϕ) λ can be chosen to be constant along the fibre (temporal gauge), and to simultaneously take values in a fixed Cartan subalgebra $\mathfrak{t} \oplus \mathfrak{t}$ of TG (Abelianisation in TG). We then give a brief outline of how to pass from the functional integral to a finite dimensional integral over $\mathfrak{t} \oplus \mathfrak{t}$.

As already mentioned, the integral in question has poles on the integration contour and so requires a definition. In section 4 we thus give our first definition of BF theory, as a residue of a particular function related to the integrand of the finite dimensional integral, and analyse its consequences. Given the shortcomings of that definition, the alternative Definitions 2 and 3 described above are explored in section 5.

Certain technical details are relegated to an appendix (A for information about the group TG and its Lie algebra, and B for details about a certain useful discrete symmetry of complex Chern-Simons theory on Seifert manifolds).

2 BF-theory on a 3-manifold and path integrals

In this section we give a brief review of 3-dimensional BF theory and Witten's formula for the path integral which holds when there are no zero modes. This is followed by a discussion of a generalisation which takes zero modes into account.

2.1 BF theory as a TG Chern-Simons theory

As recalled in the Introduction, 3-dimensional BF theory on a 3-manifold M is defined by the action (1.1)

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B \wedge F_{A} \right) \tag{2.1}$$

with A a connection on a G-bundle over M and $B \in \Omega^1(M, \mathfrak{g})$. The gauge symmetries of this action are the usual G gauge symmetry acting on A and B, as well as a shift gauge symmetry for the field B,

$$(A, B) \to (g^{-1}Ag + g^{-1}dg, g^{-1}(B + d_A\Lambda)g)$$
 (2.2)

This shift gauge symmetry is non-compact, parametrised by $\Lambda \in \Omega^0(M, \mathfrak{g})$, and thus, even for G a compact Lie group, 3-dimensional BF theory is an example of a gauge theory with a non-compact gauge group (and this is the main reason we are interested in this theory here). In fact, BF-theory is not just some non-compact analogue of Chern-Simons theory: in a precise sense it is a Chern-Simons theory for the non-compact gauge group TG, the tangent bundle group of G. As it will frequently be useful to have this perspective in the back of one's mind in the following, here (and in more detail in appendix A) we quickly review the relevant facts regarding the group TG, its Lie algebra \mathfrak{tg} , and Chern-Simons theory based on it.

Thus, consider a compact semi-simple Lie group G (throughout we will also assume that G is connected and simply-connected) with Lie algebra \mathfrak{g} , a basis of generators j_a and commutation relations $[j_a, j_b] = f_{ab}^c j_c$. Then the Lie algebra \mathfrak{tg} of the Lie group TG has generators (j_a, p_a) and commutation relations (A.8)

$$[j_a, j_b] = f_{ab}^c j_c, \quad [j_a, p_b] = [p_a, j_b] = f_{ab}^c p_c, \quad [p_a, p_b] = 0.$$
(2.3)

For the considerations of section 5 it will be useful to keep in mind, that this algebra can be obtained as a contraction of the Lie algebra of $G \times G$ or $G_{\mathbb{C}}$ (see (A.12)).

Given an invariant non-degenerate scalar product $g_{ab} = \text{Tr } j_a j_b$ on \mathfrak{g} , an invariant and non-degenerate scalar product on \mathfrak{tg} is (A.18)

$$\ll j_a, j_b \gg = \ll p_a, p_b \gg = 0, \quad \ll j_a, p_b \gg = g_{ab}.$$
 (2.4)

A TG-gauge field can be expanded as

$$C = A^a j_a + B^a p_a \,, \tag{2.5}$$

with field strength

$$F_C = dC + \frac{1}{2}[C, C] = F_A^a j_a + d_A B^a p_a \,. \tag{2.6}$$

The Chern-Simons action for C with respect to the above scalar product is (with a for present purposes convenient choice of normalisation, and with an integration by parts)

$$I_{CS} \equiv \frac{1}{2} \int_M \ll C, dC + \frac{1}{3} [C, C] \gg = \int_M \operatorname{Tr} B \wedge F_A = I_{BF}$$
(2.7)

The equations of motion (1.3) of *BF*-theory are then evidently simply the flatness conditions for the connection C,

$$F_C = 0 \quad \Leftrightarrow \quad F_A = 0, \quad d_A B = 0 \tag{2.8}$$

Moreover, the gauge symmetries (2.2) of *BF*-theory are precisely the *TG* gauge symmetries of the *TG*-connection *C*.

2.2 Path integral for BF theory: formal considerations

Evidently (and as recalled in the Introduction), formally, the path integral for the partition function of BF theory is

$$Z_{BF}[M,G] = \int DADB \exp\left(i \int_{M} \operatorname{Tr} \left(B \wedge F_{A}\right)\right)$$
$$\simeq \int DA\delta\left(F_{A}\right)$$
(2.9)

so that we may expand about flat connections to give a more complete evaluation. Following Witten [11], we use the standard Faddeev-Popov covariant gauge-fixing procedure around

these flat connections. More details about gauge fixing Chern-Simons theory with noncompact gauge groups can be found in [28]. The action and symmetries about a flat background connection $\omega_{(\alpha)}$ are

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B \wedge F_{A+\omega_{(\alpha)}} \right)$$
(2.10)

$$(A + \omega_{(\alpha)}, B) \to (g^{-1}(A + \omega_{(\alpha)})g + g^{-1}dg, g^{-1}\left(B + d_{A + \omega_{(\alpha)}}\Lambda\right)g)$$
(2.11)

while the covariant background field gauge fixing conditions

$$d_{\omega_{(\alpha)}} * A = 0. \quad d_{\omega_{(\alpha)}} * B = 0$$
 (2.12)

require us to make a choice of metric on M. Then the ghost and gauge fixing action is

$$I_{GF} = \int_{M} \operatorname{Tr} \left(u d_{\omega_{(\alpha)}} * B + v d_{\omega_{(\alpha)}} * A + \overline{f} d_{\omega_{(\alpha)}} * d_{(\omega_{(\alpha)} + A)} f + \overline{g} d_{\omega_{(\alpha)}} * d_{(\omega_{(\alpha)} + A)} g \right)$$
(2.13)

If the space of flat connections is not made up of isolated points then one would need to integrate over them which one could do through the introduction of collective coordinates. One should also take into account the possible zero modes of both B and of the ghosts. Those zero modes form non-compact spaces $\mathrm{H}^{1}_{\omega_{\alpha}}(M,\mathfrak{g})$, and $\mathrm{H}^{0}_{\omega_{\alpha}}(M,\mathfrak{g})$ respectively. Taking the above caveats into account the path integral formally becomes

$$Z_{BF}[M,G] = \sum_{\alpha} \int DADB \exp\left(iI_{BF} + iI_{GF}\right)$$
$$= \sum_{T\mathfrak{M}_{\alpha}} \tau_M(\omega_{(\alpha)})$$
(2.14)

Here the sum and integral symbol over $T\mathfrak{M}_{\alpha}$ is meant to indicate that one sums over distinct components of the moduli spaces of classical solutions and integrates over the moduli of continuous families of solutions. These spaces include the zero modes of A and B as well as of those of the ghosts f and g (the multiplier field and anti-ghost zero modes canonically cancel each other). The space $T\mathfrak{M}_{\alpha}$ space can be very singular and so (2.14) as it stands is rather symbolic in general.

On the 3-manifolds of interest in this paper, the moduli space includes acyclic flat connections, i.e. flat connections which are isolated and irreducible, so that there are no zero modes at all at these solutions. The path integral around such a connection simplifies by using the following scaling argument. About such an isolated and irreducible connection send

$$A \to tA, \quad B \to t^{-1}B, \quad t \in \mathbb{R}_+$$
 (2.15)

(with compensating scaling for the multiplier fields). This transformation has trivial Jacobian, and the action remains well defined in the limit $t \to 0$. Indeed, in this limit,

$$I_{BF} \to \int_{M} Bd_{\omega_{(\alpha)}} A$$
$$I_{GF} \to \int_{M} \operatorname{Tr} \left(ud_{\omega_{(\alpha)}} * B + vd_{\omega_{(\alpha)}} * A + \overline{f} * \Delta_{\omega_{(\alpha)}} f + \overline{g} * \Delta_{\omega_{(\alpha)}} g \right)$$
(2.16)

and the partition function around this connection is simply a standard path integral representation [3, 9] of the Ray-Singer torsion [5] $\tau_M(\omega_{(\alpha)})$ of the flat connection $\omega_{(\alpha)}$,

$$\tau_M(\omega_{(\alpha)}) = \left(\operatorname{Det} \Delta^0_{\omega_{(\alpha)}}\right)^{3/2} \cdot \left(\operatorname{Det} \Delta^1_{\omega_{(\alpha)}}\right)^{-1/2}$$
(2.17)

(the superscripts on the twisted Laplacians indicating the degrees of the forms on which they act). Thus on such 3-manifolds, the partition function now becomes

$$Z_{BF}[M,G] = \sum_{\text{acyclic}} \tau_M(\omega_{(\alpha)}) + \dots$$
(2.18)

where the ellipses indicate the rest of the contributions to the path integral. As we have already explained, there may be zero modes to deal with in the extra terms (2.18) which manifest themselves as zeros of the determinants in (2.17) and which invalidate the simplifications that we were allowed to make for the isolated irreducible connections.

One way to proceed is to adopt the prescription of Ray and Singer [6] by first excising the zero modes and then adding a correction term to ensure metric independence. This 'extra' gauge fixing of the zero modes may be achieved by a BRST procedure [9] the details of which have been eplained in some detail in [1]. The advantage of this method is that then the torsion is a natural measure for the finite dimensional integral that appears in (2.14). While this defines the integrand, one is still confronted with the problem of determining and defining the space over which this is to be integrated. Thus, while formally this appears to be a good definition, at a practical level it seems somewhat intractable at the moment.

2.3 Path integral for BF theory and massive Ray-Singer torsion

An alternative method for regularising such zero modes, and for keeping track of the ellipses in the formula (2.18) was advocated in [1]. The idea is to add a mass to the Laplacians that appear in (2.17), thus lifting all the zero modes and side-stepping (or at least initially bypassing) the problem of having to deal with them directly. As a first step, what this means is that instead of using the twisted Laplacian Δ_{ω} in the definition of the Ray-Singer Torsion we instead use the massive Laplacian $\Delta_{\omega} + \mu^2$ which now manifestly has a positive definite spectrum. The Massive Ray-Singer Torsion, on a 3-manifold, for a flat connection ω , is then defined to be

$$\tau_M(\omega,\mu) = \left(\text{Det} \left(\Delta_{\omega}^0 + \mu^2 \right) \right)^{3/2} \, . \, \left(\text{Det} \left(\Delta_{\omega}^1 + \mu^2 \right) \right)^{-1/2} \tag{2.19}$$

One can then define a regularised partition function by

$$Z_{BF}[M,G;\mu] = \sum \tau_M(\omega_{(\alpha)},\mu)$$
(2.20)

where we do not necessarily attempt to integrate over the tangent space as those modes have been lifted. Here a caveat is required: in [1] we showed that for manifolds which are mapping tori it is possible to maintain complete gauge invariance even with the introduction of a mass. However, a mass term cannot be introduced in a gauge invariant way on a general 3-manifold. We accept that the gauge symmetry is broken and may be reinstated if required by some renormalisation. The way in which one obtains the actual Ray-Singer Torsion from the massive Ray-Singer torsion is to take a limit

$$\tau_M(\omega) \equiv \lim_{\mu \to 0} \mu^{-3\dim \mathcal{H}^0_\omega + \dim \mathcal{H}^1_\omega} \tau_M(\omega, \mu)$$
(2.21)

In this way the BF partition function will then take the form

$$Z_{BF}[M,G,\mu] = \sum \mu^{3\dim \mathrm{H}^{0}_{\omega} - \dim \mathrm{H}^{1}_{\omega}} \left(\tau_{M}(\omega) + \dots\right)$$
(2.22)

and we will need to specify which terms we are interested in. The ellipses in this formula refer to terms higher order in μ than the zero-th order Torsion (essentially constants and derivatives of the Ray-Singer Torsion).

At a formal level this is, perhaps, as far as one may go. It must be said that this situation is not very satisfying and has calculational drawbacks. The formula (2.20) is very implicit and requires knowledge outside of the path integral in order to be used. The flat connections need to be found, the cohomology groups about the flat connections must be determined and a formula for the Massive Ray-Singer Torsion must be given.

It is therefore of considerable interest to consider 3-manifolds for which the BF partition function can also be calculated directly and explicitly, and where the result can be compared with the formal expectations for the partition function outlined above. One such class of manifolds is Seifert manifolds.

2.4 Expectations for the partition function on Seifert manifolds

The Ray-Singer torsion for Seifert 3-manifolds is known (through its equivalence to Reidemeister Torsion [29]). About an acyclic flat connection, i.e. with trivial twisted cohomology, it is

$$\tau_M(\omega) = \tau_{S^1}(\phi)^{2-N} \prod_{i=1}^N \tau_{S^1}(\phi/a_i)$$
(2.23)

where ϕ is the component of the connection in the direction of the fibre of the Seifert 3-manifold and N (the number of orbifold points) and a_i (the order of the isotropy group at the *i*'th orbifold point) are integers that are part of the defining data of a Seifert 3-manifold (see section 3.1). We should also specify the representation of the group G for which we are evaluating the Torsion; however, as throughout we fix on the adjoint representation, we do not need to indicate it in the notation.

Including the contributions from the non-acyclic connections, one expects the partition function to take the form (2.18)

$$Z_{BF}[M,G] \simeq \sum_{\text{acyclic}} \tau_{S^1}(\phi)^{2-N} \prod_{i=1}^N \tau_{S^1}(\phi/a_i) + \dots$$
 (2.24)

and, as in the general situation above, one way to keep track of the ellipses in the above formula is to use the massive Ray-Singer torsion. In the case of Seifert manifolds all we need is the massive Ray-Singer torsion on the circle, which is well understood [1]. Thus a suitably regularised version of (2.24) is

$$Z_{BF}[M,G;\mu] = \sum \tau_{S^1}(\phi,\mu)^{2-N} \prod_{i=1}^N \tau_{S^1}(\phi/a_i,\mu) + \dots$$
(2.25)

One can now use a mass expansion to identify the individual contributions to the regularised partition function.

The analysis that we have presented thus far, in the covariant gauges, is an extension of that of Witten to those flat connections which are not necessarily flat and isolated. One shortcoming with this approach, as we have already explained, is that with a background field approach one must, by independent means, find the flat connections and determine the cohomology groups about them. One would prefer an approach that evaluates the path integral directly and does not require the split between the classical and quantum fields. The rest of the paper is devoted to producing just such an approach.

In particular, we will show in section 3 that the procedure developed in [13–18] to reduce the partition function of Chern-Simons theory for a compact gauge group G to a finite-dimensional integral (over the Cartan subalgebra \mathfrak{t} of \mathfrak{g}) can be extended to the case at hand, namely BF theory, or Chern-Simons theory with gauge group TG.

The finite dimensional integrals have poles on the contour of integration and so it then remains to define and give a meaning to them. As we will see in section 4, a direct calculation of this finite-dimensional integral reproduces the expected form of the partition function only partially. In particular it captures the contributions of reducible flat connections as expected but surprisingly does not give the expected results for the isolated and irreducible flat connections.

This prompts us to investigate two other possible definitions in section 5 which do lead to the expected form of the partition function discussed in this section.

3 BF-theory on a Seifert fibred 3-manifold

In this section we specialise to Seifert fibred 3-manifolds. The extra structure afforded by the S^1 principal bundle structure (over an orbifold base) allows for convenient choices of gauge as well as for regularising the Ray-Singer Torsion with the introduction of a mass term [1]. We are very brief with the background material on Seifert manifolds as it has appeared before. We spend some time on the gauge fixing as this is quite new for the TGtheories, while the evaluation of the determinants is so close to that of the determinants evaluated in [17] that we borrow liberally from there.

3.1 Seifert 3-manifolds briefly

From now on M denotes a Seifert 3-manifold. Then M is a 3-manifold which is a circle V-bundle over the 2 dimensional orbifold Σ_V of genus g with N orbifold points. A general Seifert manifold is written as $M[\deg \mathcal{L}_M, g, (a_1, b_1), \ldots, (a_N, b_N)]$ where the a_i are the isotropies of the orbifold points, the b_i are the weights of the line V-bundle at the orbifold

points and deg \mathcal{L}_M is the degree of that line bundle. As the Seifert 3-manifold is the circle V-bundle of the line V-bundle \mathcal{L}_M it is also designated by $S(\mathcal{L}_M)$.

Let \mathcal{L}_0 be a topological line bundle at some smooth point on Σ_V with degree 1 and \mathcal{L}_i be the line V-bundles at the *i*-th orbifold points with degree $1/a_i$ then

$$\mathcal{L}_M = \mathcal{L}_0^{b_0} \otimes \mathcal{L}_1^{b_1} \otimes \dots \otimes \mathcal{L}_N^{b_N}$$
(3.1)

and M is the circle bundle of \mathcal{L}_M [30] (we use normalised Seifert data so that $1 \leq b_i < a_i$).

The Seifert 3-manifold is smooth iff $gcd(a_i, b_i) = 1$ for each *i*. It is a \mathbb{Z} Homology Sphere $(H_1(M, \mathbb{Z}) = 0)$ iff the line bundle \mathcal{L}_M that defines it satisfies

$$g = 0, \quad c_1(\mathcal{L}_M) = b_0 + \sum_{i=1}^N \frac{b_i}{a_i} = \pm \frac{1}{a_1 \dots a_N} \equiv \pm \frac{1}{P}$$
 (3.2)

(here we have introduced the notation $P = a_1 \dots a_N$). One consequence of these conditions is that $gcd(a_i, a_j) = 1$ for $i \neq j$. If one takes a tensor power of this line V-bundle, $\mathcal{L}_{M_d} = \mathcal{L}_M^{\otimes d}$, then the Seifert manifold $M_d = M/\mathbb{Z}_d$ is a \mathbb{Q} Homology Sphere $(H_1(M_d, \mathbb{Q}) = 0)$ with

$$g = 0, \quad c_1(\mathcal{L}_{M_d}) = c_1(\mathcal{L}_M^{\otimes d}) = \pm \frac{d}{a_1 \dots a_N} = \pm \frac{d}{P}$$
(3.3)

and

$$d| = |\mathbf{H}_1(M_d, \mathbb{Z})| \tag{3.4}$$

These are the 3-manifolds that we will exclusively concentrate on. The reason for this choice of M is two fold.

Firstly, if M is neither a \mathbb{Z} nor a \mathbb{Q} homology sphere, then it will necessarily have a non-zero dimensional moduli space of Abelian flat connections which in turn means that there will be Abelian B zero modes and hence non-compact directions to deal with.¹ We do not want to have to worry about such a situation here, as it is somewhat tangential to the other issues that we wish to address.

Far from all \mathbb{Z} or \mathbb{Q} homology spheres are Seifert. The reason for choosing M to be more specifically a Seifert \mathbb{Z} or \mathbb{Q} homology sphere is a more pragmatic one. Having M a Seifert manifold means that we have Fourier mode decomposition of all the fields, so their components can be ultimately viewed as living on Σ_V , and, as shown in [13–18], there are specific gauge choices available that allow one to significantly simplify the evaluation of the partition function.

3.2 BF action and gauge transformations on a Seifert manifold

As the Seifert fibred 3-manifold M is a principal U(1) bundle, it also comes equipped with the fundamental vector field ξ which generates the U(1) action. We also equip M with a (nowhere vanishing) connection 1-form κ , i.e.

$$\iota_{\xi}\kappa = 1, \quad L_{\xi}\kappa = \iota_{\xi}d\kappa = 0, \qquad (3.5)$$

¹Even if M is a \mathbb{Z} or \mathbb{Q} homology sphere, there may be non-zero dimensional components of the moduli space of flat connections, but there are many M for which the moduli space of flat connections is made up of a finite number of points.

and

$$\int_{M} \kappa \wedge d\kappa = c_1 \left(\mathcal{L}_M \right) \,. \tag{3.6}$$

Note that, acting on a form of any degree, one has

$$\iota_{\xi}\kappa + \kappa\iota_{\xi} = 1 \tag{3.7}$$

so that $\iota_{\xi}\kappa$ and $\kappa\iota_{\xi}$ are projection operators onto the space of horizontal and vertical forms of fixed degree respectively.

Given a trivial G-bundle on a Seifert 3-manifold M, we correspondingly decompose the connection A and 1-form B of BF theory as

$$A = A_{\Sigma} + \kappa \phi, \quad B = B_{\Sigma} + \kappa \lambda \tag{3.8}$$

where

$$A_{\Sigma} = \iota_{\xi} \kappa A, \quad \phi = \iota_{\xi} A, \quad B_{\Sigma} = \iota_{\xi} \kappa B, \quad \lambda = \iota_{\xi} B \tag{3.9}$$

We may also decompose the exterior derivative as

$$d = \iota_{\xi} \kappa d + \kappa \iota_{\xi} d = d^{\Sigma} + \kappa \iota_{\xi} d \tag{3.10}$$

with twisted versions

$$d_A = \iota_\xi \kappa d_A + \kappa \iota_\xi d_A \tag{3.11}$$

which acts on 0-forms as

$$d_A\beta = d_A^\Sigma\beta + \kappa D_\phi\beta \tag{3.12}$$

In terms of this decomposition, the gauge transformations (2.2) take the form, with t a local fibre coordinate

$$A_{\Sigma} \to g^{-1} d_A^{\Sigma} g, \quad \phi \to g^{-1} (\partial_t + \phi) g$$

$$B_{\Sigma} \to g^{-1} (B_{\Sigma} + d_A^{\Sigma} \Lambda) g, \quad \lambda \to g^{-1} (\lambda + D_{\phi} \Lambda) g$$
(3.13)

and the action (2.1) becomes

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B_{\Sigma} \wedge \kappa D_{\phi} A_{\Sigma} + \kappa_{\Sigma} B \wedge d^{\Sigma} \phi + \lambda \kappa \wedge F_{A_{\Sigma}} + \kappa \wedge d\kappa \phi \lambda \right)$$
(3.14)

with horizontal curvature

$$F_{A_{\Sigma}} = d^{\Sigma} A_{\Sigma} + A_{\Sigma}^2 \tag{3.15}$$

3.3 Gauge fixing 1: temporal gauge for A and B

One of the great benefits of considering a Seifert manifold is that there are very useful non-covariant gauges available. In particular, on a Seifert manifold we can always choose the "temporal gauge" that the fibre-component ϕ of A is constant along the fibre,

$$\iota_{\xi} d\phi = 0. \tag{3.16}$$

It turns out that (for generic ϕ) one can also impose the analogouus condition on λ ,

$$\iota_{\xi} d\lambda = 0. \tag{3.17}$$

In local coordinates with t a fibre coordinate, these conditions simply read

$$\frac{\partial}{\partial t}\phi = 0, \quad \text{and} \quad \frac{\partial}{\partial t}\lambda = 0$$
 (3.18)

The first of these can be achieved by having ϕ gauge equivalent to some 'time' independent field ϕ_0 ; i.e. we have to solve

$$g(t)^{-1}(\partial_t + \phi)g(t) = \phi_0 \quad \Leftrightarrow \quad \partial_t g(t) = g(t)\phi_0 - \phi(t)g(t) \tag{3.19}$$

This equation has the solution

$$g(t) = P \exp\left(\int_{t}^{0} \phi(s) ds\right) e^{t\phi_0}$$
(3.20)

(where we have without loss of generality chosen g(0) = 1). Periodicity g(1) = g(0) of the gauge parameter now determines ϕ_0 to be the average value of ϕ in the sense that

$$e^{\phi_0} = P \exp\left(\int_0^1 \phi(s) ds\right) \tag{3.21}$$

The second gauge condition, $\partial_t \lambda = 0$, is perhaps not familiar and as there is a hidden subtlety we will deal with it in some detail. Suppose we have already implemented the first gauge choice (thus in the following $\phi = \phi_0$ is taken to satisfy $\partial_t \phi = 0$). To arrive at λ constant along the fibre we need to solve

$$\lambda + D_{\phi}\Lambda = \lambda_0 \tag{3.22}$$

for some λ_0 constant in t to be determined. Writing

$$\partial_t \left(e^{t\phi} \Lambda e^{-t\phi} \right) = e^{t\phi} (D_{\phi} \Lambda) e^{-t\phi}$$
(3.23)

one sees that this equation has the solution.

$$h(t)\Lambda(t)h^{-1}(t) = \int_0^t e^{s\phi} \left(\lambda_0 - \lambda(s)\right) e^{-s\phi} ds + \Lambda(0)$$
(3.24)

where

$$h(t) = e^{t\phi}, \quad h = e^{\phi} \tag{3.25}$$

Periodicity in the gauge parameter $\Lambda(1) = \Lambda(0)$ gives us the equation

$$(\operatorname{Ad}(h) - 1)\Lambda(0) = \int_0^1 e^{s\phi} \left(\lambda_0 - \lambda(s)\right) e^{-s\phi} ds$$
(3.26)

Before going on we note that a shift of Λ by a constant (in t) Λ_0 simply amounts to a shift in λ_0 ,

$$\Lambda \to \Lambda + \Lambda_0 \quad \Rightarrow \quad \lambda_0 \to \lambda_0 + [\phi, \Lambda_0] \tag{3.27}$$

Without loss of generality we can therefore assume that $\Lambda(0) = 0$, so that the periodicity condition (3.26) implies

$$\int_{0}^{1} e^{s\phi} (\lambda_0 - \lambda(s)) e^{-s\phi} ds = 0.$$
 (3.28)

The integral involving λ_0 can be done explicitly, with the result

$$\frac{\mathrm{e}^{\operatorname{\mathsf{ad}}(\phi)} - 1}{\mathrm{\mathsf{ad}}(\phi)} \lambda_0 = \int_0^1 \mathrm{e}^{s \operatorname{\mathsf{ad}}(\phi)} \lambda(s) ds \,. \tag{3.29}$$

For generic values of ϕ , the operator on the left-hand side can be inverted to give

$$\lambda_0 = \operatorname{ad}(\phi)(\operatorname{e}^{\operatorname{ad}(\phi)} - 1)^{-1} \int_0^1 \operatorname{e}^{s \operatorname{ad}(\phi)} \lambda(s) ds$$
(3.30)

which tells us in which sense λ_0 can be regarded as a ϕ -weighted average of λ over the fibre.

However, we see that this solution fails when ϕ is not suitably generic, so that the operator $\exp \operatorname{ad}(\phi) - 1$ has zero-modes. This should not come as a suprise, as this is precisely the condition that the operator $D_{\phi} = \partial_t + \operatorname{ad}(\phi)$ has zero-modes, so that the gauge fixing condition (3.22) cannot be solved for all $\lambda(t)$. Indeed, for a zero mode ψ of D_{ϕ} one has (see also the discussion in section 2.3.4 of [1])

$$D_{\phi}\psi = 0 \quad \Rightarrow \quad \psi(t) = e^{t \operatorname{ad}(\phi)}\psi(0), \qquad (3.31)$$

and periodicity $\psi(1) = \psi(0)$ implies

$$\psi(1) = \psi(0) \quad \Rightarrow \quad \left(e^{\operatorname{\mathsf{ad}}(\phi)} - 1\right)\psi(0) = 0.$$
(3.32)

For ϕ taking values in the Cartan subalgebra t of \mathfrak{g} , this is simply the condition that ϕ has a component that is an element of the integral lattice I(G) of G.

Normally this kind of constraint arising only for highly non-generic field configurations would not be an issue as one is integrating over ϕ and the points where ϕ is integral have measure zero. Unfortunately, in *BF* theory we get delta function support onto particular values of ϕ . In particular we will see when we are considering \mathbb{Z} Homology Spheres that these integral points are the ones that are selected (cf. (4.6)).

3.4 Gauge fixing 2: Abelianisation and emergence of non-trivial line bundles

After having chosen the gauges (3.18), we still have gauge transformations available that are constant along the fibre. For a compact gauge group, one can locally use this residual gauge feedom to conjugate the field ϕ into the Cartan subalgebra t of \mathfrak{g} , i.e. one can impose the gauge condition

$$\phi^{\mathfrak{k}} = 0 \quad \Leftrightarrow \quad \phi = \phi^{\mathfrak{t}}, \tag{3.33}$$

where \mathfrak{k} is the orthogonal complement of the Cartan subalgebra \mathfrak{t} of \mathfrak{g} ,

$$\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{k} \,. \tag{3.34}$$

We will come back to the global issues involved in this choice of gauge below.

However, first of all we need to address the question if we can do something analogous in the case at hand, with gauge group TG. Since TG is neither compact nor semi-simple, this is not a priori obvious. However, we show in appendix A.2 that one can use the adjoint action of TG on its Lie algebra to conjugate any element into an element of $\mathfrak{t} \oplus \mathfrak{t}$, which plays the role of the Cartan subalgebra of the Lie algebra of TG. From one perspective, the reason we are able to do this is that TG arises as the Inönü-Wigner contraction of a group where this is certainly possible, namely $G \times G$.

Explicitly, this means that we can use the gauge transformation

$$(\phi, \lambda) \mapsto (g, \Lambda)^{-1} \cdot (\phi, \lambda) \cdot (g, \Lambda) = (g^{-1} \phi g, g^{-1} (\lambda + [\phi, \Lambda])g)$$
(3.35)

to (locally) impose simultaneously the Abelianising gauge conditions

$$\phi^{\mathfrak{k}} = 0, \quad \lambda^{\mathfrak{k}} = 0 \quad \Leftrightarrow \quad \phi = \phi^{\mathfrak{t}}, \quad \lambda = \lambda^{\mathfrak{t}}$$
(3.36)

Now let us turn to the global issues regarding this gauge choice, which implies that we will be dealing with an Abelian gauge theory on Σ_V . As has been described in detail in [14, 31], there are topological obstructions to imposing the gauge choice on ϕ globally; in particular, conjugating ϕ to the Cartan subalgebra \mathfrak{t} on Σ_V is not possible with smooth gauge transformations. However, if we insist on doing so anyway, the price to be paid is that we must sum over all available line bundles on Σ_V . The reason for the emergence of this sum has been explained a number of times [13–16, 18, 31].

Since TG is contractible to G (and there are no topological issues involved in the shift transformation $\lambda \to \lambda + [\phi, \Lambda]$ beyond those involving ϕ itself), there are no additional topological obstructions arising from diagonalisation in TG.

In practice there are two ways to introduce the non-trivial line bundles and the sum over them. They both have merits and in any case are equivalent. The first makes contact with flat connections rather straightforwardly while the other is computationally easier.

We begin by describing the method using background fields described in more detail in [18]. The available line V-bundles on Σ_V are

$$\mathcal{L} = \mathcal{L}_0^{\otimes n_0} \otimes \mathcal{L}_1^{\otimes n_1} \otimes \dots \otimes \mathcal{L}_N^{\otimes n_N}$$
(3.37)

and the basic idea is to allow for a non-trivial background connection on patches about each divisor, with the divisors being the singular points and with one divisor being a regular point of Σ_V . It was shown in [18] that in the tubular neighbourhood $V_{(a_i,b_i)}$ of the *i*-th singular point one could introduce a connection 1-form κ_i which would glue to a global one form κ on M. In terms of these local forms and the local surgery data

$$a_i s_i - b_i r_i = -1, \quad \text{with} \quad (a_i, b_i) = 1$$
(3.38)

the local orbifold data, the background connection takes the simple form²

$$\mathcal{A}_B = 2\pi i \left(\mathbf{n}_0 \kappa_0 + \sum_{i=1}^N \mathbf{n}_i r_i \kappa_i \right)$$
(3.39)

²A word on notation. Throughout we write connections in a basis of simple roots α_i . For semi-simple simply laced Lie algebras we have that the natural inner product $\langle \alpha_i, \alpha_j \rangle = 2\delta_{ij}$. The bold facing of integers indicates that we mean $\mathbf{m} = m^i \alpha_i$ and for each *i* we have $m^i \in \mathbb{Z}$. We will write that $\mathbf{m} \in I(G)$ where I(G)is the integral lattice of *G*. As *G* is simply connected we have identified the root lattice with I(G).

where the \mathbf{n} are Hermitian and lie in \mathfrak{t} . With an abuse of notation, we may write the background as

$$\kappa \phi_B = 2\pi i \left(\mathbf{n}_0 \kappa_0 + \sum_{i=1}^N \mathbf{n}_i r_i \kappa_i \right)$$
(3.40)

Each of the local forms κ_i extends outside of the tubular neighbourhood to $d\theta$ on the fibre. One also has that

$$\int_{V_{(a_i,b_i)}} \kappa_i d\kappa_i = \frac{b_i}{a_i} \tag{3.41}$$

with the regular point having weight $a_0 = 1$. Glued together the κ_i form a global, smooth 1-form $\kappa = \kappa_0 \cup \kappa_1 \cup \cdots \cup \kappa_N$ that defines a principal bundle structure on M with

$$c_1(\mathcal{L}_M) = \int_M \kappa \wedge d\kappa = b_0 + \sum_{i=1}^N \frac{b_i}{a_i}$$
(3.42)

With the background field in place, one makes the substitution

$$\kappa\phi \to \kappa\phi + \kappa\phi_B$$
 (3.43)

everywhere in the path integral, with the second term on the right given by (3.40).

Alternatively [13–16] one sets all the $\mathbf{n}_i \ i \neq 0$ to zero and $\kappa_0 \to \kappa$ (and this is the natural procedure from the point of view of obstruction theory outlined in [31] and extended here to the manifolds that we are considering). Otherwise one keeps the substitution (3.43) in the path integral. In appendix B we show that there is enough symmetry in the theory to pass between these different formulations.

3.5 Reduction of the partition function to a finite-dimensional integral

With all the preliminaries in place we can now outline how to perform the path integral for the partition function.

Firstly the action (3.14), taking into account the non-trivial bundles that arise on Abelianisation, goes over to

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B_{\Sigma} \wedge \kappa D_{\phi + \phi_{B}} A_{\Sigma} + \kappa B_{\Sigma} \wedge d^{\Sigma} \phi + \lambda \kappa \wedge F_{A_{\Sigma}} + \kappa \wedge d\kappa (\phi + \phi_{B}) \lambda \right)$$
(3.44)

with $\kappa \phi_B$ and $\kappa d\kappa \phi_B$ suitably understood. Indeed we note that the part of B_{Σ} that is constant along the fibre and takes values in the Cartan subalgebra t only appears through the term

$$\int_{M} \operatorname{Tr} \left(\kappa B_{\Sigma} \wedge d^{\Sigma} \phi \right). \tag{3.45}$$

Thus integrating out these modes of B_{Σ} imposes the condition that $d^{\Sigma}\phi = 0$. Given the gauge conditions $\partial_t \phi = 0$, this implies that ϕ is constant on M. Likewise the component of A_{Σ} constant along the fibre and in the Cartan subalgebra only appears in

$$\int_{M} \operatorname{Tr} \left(\kappa A_{\Sigma} \wedge d^{\Sigma} \lambda \right) \tag{3.46}$$

so integrating over that mode of A_{Σ} fixes $d^{\Sigma}\lambda = 0$ which, together with $\partial_t \lambda = 0$, tells us that λ is also constant on M. The other components of A_{Σ}^t and B_{Σ}^t are paired with each other and lead to an overall constant when integrated over.

We should also take care of any possible harmonic modes of the components of A and B that we have integrated over. If there any such harmonic modes of B, they would be non-compact directions and would lead to divergences. In order to avoid this we fix on the genus of Σ_V to be zero, as in (3.2) and (3.3) thus ensuring that there are no B_{Σ}^t zero modes to contend with. The stronger choice that we have made of having M be a \mathbb{Z} or \mathbb{Q} homology sphere is to ensure that there are no moduli of Abelian connections on M as then we would necessarily have Abelian B zero modes.

The form that we now have for the action is

$$I_{BF} = \int_{M} \operatorname{Tr} \left(B_{\Sigma}^{\mathfrak{k}} \wedge \kappa D_{\phi + \phi_{B}} A_{\Sigma}^{\mathfrak{k}} + \kappa \wedge \lambda A_{\Sigma}^{\mathfrak{k}} A_{\Sigma}^{\mathfrak{k}} \right) + \operatorname{Tr} \lambda \left(c_{1}(\mathcal{L}_{M})\phi + 2\pi i \widehat{\mathbf{q}} \right)$$
(3.47)

where

$$\widehat{\mathbf{q}} = \sum_{i=0}^{N} \frac{1}{a_i} \mathbf{n}_i \tag{3.48}$$

We are now ready to integrate out the fields $A_{\Sigma}^{\mathfrak{k}}$ and $B_{\Sigma}^{\mathfrak{k}}$ that appear in (3.47). There are also ghost terms of the same type to take into account. The functional determinant that we obtain and need to evaluate is

$$\operatorname{Det} \left(\begin{array}{c} \operatorname{ad}_{\lambda} & D_{\phi + \phi_B} \\ -D_{\phi + \phi_B} & 0 \end{array} \right) \Big|_{\Omega^1_H(M, \mathfrak{k}) \oplus \Omega^1_H(M, \mathfrak{k})}^{-1/2}$$
(3.49)

There is a similar determinant from the ghost terms. As usual we will expand in Fourier modes along the S^1 fibre and we will regulate with a ζ -function regularisation. What one finds is, even upon regularisation, that the ad_{λ} part of the determinant does not contribute. In this way what finally needs to be evaluated is

$$\left|\operatorname{Det}\left(D_{\phi+\phi_B}\right)\right|_{2\Omega^0_H(M,\mathfrak{k})\ominus\Omega^1_H(M,\mathfrak{k})}\tag{3.50}$$

For every Fourier mode $2\Omega_H^0(M, \mathfrak{k}) \oplus \Omega_H^1(M, \mathfrak{k})$ is roughly the Euler characteristic of Σ_V . This, almost, cancellation of all the modes in the functional determinant can be used to evaluate it. However, the field ϕ_B is not constant on Σ_V unlike ϕ . This issue was circumvented in [18] by using a density form of the index theorem originally found in [13]. The determinant in [18] is the square root of (3.50), so we may straightforwardly borrow the result from there. In this way we see that the absolute value (3.50) is

$$\tau_M(\phi; \mathbf{n}_i) = \tau_{S^1}(\phi + 2\pi i \mathbf{n}_0)^{2-2g-N} \cdot \prod_{i=1}^N \tau_{S^1}((\phi + 2\pi i r_i \mathbf{n}_i)/a_i)$$
(3.51)

where τ_M is the Ray-Singer torsion of M. The form of τ_M shows that it is given by circles that is in terms of the Ray-Singer torsion of S^{1} 's [29]. For completeness we note that

$$\tau_{S^1}(\varphi) = \det_{\mathfrak{k}} \left(I - \mathsf{Ad}(\exp\varphi) \right) \tag{3.52}$$

so that \mathbf{n}_0 drops out of (3.51) since it has integral (lattice) entries.

We regularise the Ray-Singer Torsion by introducing a mass [1]. In the present context the mass is introduced into the determinants (3.49). This is not consistent with the TGsymmetry but does preserve the Abelian part of it. One exchanges (3.52) with

$$\tau_{S^1}(\varphi,\mu) = \det_{\mathfrak{g}} \left(I - e^{-2\pi\mu} \operatorname{Ad}(\exp\varphi) \right)$$
(3.53)

and it is understood that we use the massive form for the Ray-Singer Torsion of the torsion on the circles on the right hand side of (3.51). For the details of how this comes about see Example 2.5 in [1]. We should note we have introduced extra factors of μ (namely by the factor $(1 - e^{-2\pi\mu})^{\dim T}$) by passing to (3.53) and hence we have also changed the normalisation of the path integral

Putting everthing together, we see that we have now managed to reduce the path integral for the BF partition function to a finite dimensional integral over the (by now constant) fields ϕ and λ ,

$$Z_{BF}[M,G] = \sum_{\mathbf{n}_0} \cdots \sum_{\mathbf{n}_N} \int_{\mathbf{t} \oplus \mathbf{t}} \tau_M(\phi; \mathbf{n}_i) \cdot \exp\left(i \operatorname{Tr} \lambda \left(c_1(\mathcal{L}_M)\phi + 2\pi i \widehat{\mathbf{q}}\right)\right)$$
(3.54)

This is the integral that we will look at in detail in the following section.

4 Definition 1: direct evaluation of the BF (path) integral

As we saw in the last section we are able to perform the functional integral representing the partition function Z_{BF} to the point that we are left with finite dimensional integrals. In this section we analyse the remaining integrals and notice that there are issues with them. Indeed this prompts us to search for a definition for Z_{BF} . This definition does not agree with the perturbative result of Witten [11] in the case of isolated and irreducible connections and we end the section by explaining where there could be possible issues with one of the gauge conditions that we have chosen. The formulae obtained capture those Abelian connections in \mathbb{Q} Homology Spheres and in these cases reproduce the expected result (2.25).

4.1 Symmetries and other properties of the finite dimensional integrals

At this point we have collected all the pieces and the BF partition function, for the 3-manifolds under consideration, is now a finite dimensional integral

$$Z_{BF}[M,G] = \sum_{\mathbf{n}_0} \cdots \sum_{\mathbf{n}_N} \int_{\mathbf{t} \oplus \mathbf{t}} \tau_M(\phi; \mathbf{n}_i) \cdot \exp\left(i \operatorname{Tr} \lambda \left(d\phi/P + 2\pi i \widehat{\mathbf{q}}\right)\right)$$
(4.1)

where the integral over $\mathbf{t} \oplus \mathbf{t}$ is that over the constant Abelian "fields" ϕ and λ and we have used the fact that $c_1(\mathcal{L}_M) = d/P$ for the 3-manifolds we are considering as given in (3.2) and (3.3) which in particular means that g = 0. There are no sums over the $\mathbf{n}_i \ i \neq 0$ if we have set them to zero, in which case we also have that $\hat{\mathbf{q}} = \mathbf{n}_0$. Note that as the gauge group G is compact connected and simply connected the G bundles over M are trivial and, on Abelianisation, the line V bundles at the smooth point are 'honest' line bundles so that $\mathbf{n}_0 \in \mathbb{Z}^{\mathbf{rk}}$. Consequently, in this case, \mathbf{n}_0 only appears in the action in (4.1) (for either formulation of background).

In order to understand and ultimately evaluate the integral (4.1), we note the following properties of the integrand:

1. both the action and the Ray-Singer torsion are invariant under the discrete symmetry

$$\phi \to \phi + 2\pi i P \mathbf{r}, \quad \mathbf{n}_0 \to \mathbf{n}_0 - d\mathbf{r} \quad \text{with} \quad \mathbf{r} \in \mathbb{Z}^{\mathbf{rk}(G)}, \quad P = \prod_{i=1}^N a_i$$
(4.2)

- 2. the Ray-Singer torsion with $N \ge 3$ has poles at $\alpha(\phi) = n\pi$ with $n \ne Pm$ (while for $N \le 2$ there are no poles)
- 3. the Ray-Singer torsion has zeros at $\alpha(\phi) = m\pi P$
- 4. the Ray-Singer torsion is an even function of its arguments, in the sense that

$$\tau_M(\phi, \mathbf{n}_i) = \tau_M(-\phi, a_i - \mathbf{n}_i) = \tau_M(-\phi, -\mathbf{n}_i)$$
(4.3)

Using the symmetry (4.2), we may either 'compactify' the field ϕ or limit the range of \mathbf{n}_0 . Either the way, the integral over λ leads to a delta function constraint on ϕ , namely

$$\phi = \phi_C \equiv -\frac{2\pi i P}{d} \widehat{\mathbf{q}} \tag{4.4}$$

and so it appears that only these values contribute to the path integral. These values correspond to reducible connections as we have seen in the past [17] and so one might conclude that the only flat connections that contribute to the BF partition function are those that are Abelian. However, because of the presence of poles, this conclusion is a bit quick, and we will take a closer look at this issue in section 4.4.

Before moving on, it is worthwhile reminding ourselves that there are restrictions on Mso the analysis we are performing holds. As a counter-example, presume that M is a product $\Sigma \times S^1$, or more generally a Seifert mapping torus, i.e. with $c_1(\mathcal{L}_M) = 0$ (so definitely not one of the manifolds that we are considering). Then there would be no symmetry such as (4.2) for the action and the integral over λ would not lead to (4.4).

4.2 The ϕ_C contribution to \mathbb{Z} homology spheres

Here we specialise to the ϕ_C contribution to the path integral for M either an \mathbb{Z} or a \mathbb{Q} Homology Sphere. We will use the symmetries available not to limit the range of ϕ but rather to set $\mathbf{n}_0 \in I(G)/dI(G)$ and we will keep the \mathbf{n}_i and so take the classical field contribution to be

$$\phi_C = -2\pi i \frac{P}{d} \sum_{i=0}^N \frac{\mathbf{n}_i}{a_i} \tag{4.5}$$

and any other choice leads to the same conclusions that we arrive at presently. As we only want the ϕ_C contribution we must ensure that τ_M has no poles and this is achieved by passing to the massive Ray-Singer torsion $\tau_M(\phi, \mu)$ which we do. If M is a Z homology sphere then d = 1 and the symmetry allows us to set $\mathbf{n}_0 = 0$ so that

$$\phi_C = -2\pi i P \sum_{i=1}^N \frac{\mathbf{n}_i}{a_i} \in I(G) \tag{4.6}$$

Note that these are precisely the (potentially problematic) integral values of ϕ discussed at the end of section 3.3. We will (have to) come back to this below. A tiny bit of gymnastics³ shows that

$$\phi_C + 2\pi i \mathbf{n}_j r_j = 0 \mod 2\pi i a_j I(G) \tag{4.7}$$

One immediately sees that the Ray-Singer Torsion (without a mass) on the circle vanishes and the overall ratio in (3.51) is zero. The poles and the zeros of the finite dimensional integral arise as the zeros of the (inverse) determinant of the D_{ϕ} operator. The poles are there when D_{ϕ} acts on 1-forms and the zeros arise when acting on 0-forms at the special values (4.4) of ϕ .

In order to understand the contributions we turn on the mass regulator, as in (3.53). Then we find

$$\tau_M(\phi,\mu) = \det_{\mathfrak{g}} \left(I - e^{-2\pi\mu} \right)^{2-N} \prod_{i=1}^N \det_{\mathfrak{g}} \left(I - e^{-2\pi\mu/a_i} \right)$$
(4.8)

The small mass limit gives us, as in (2.22),

$$\tau_M(\phi,\mu) = (2\pi\mu)^{2\dim G} (a_1 \dots a_N)^{-\dim G} + \dots$$
(4.9)

The exponent of μ is understood as follows: the torsion of M has been pushed to that on S^1 and as on S^1 the twisted cohomology groups satisfy $H^0_{\omega} = H^1_{\omega}$ the combination $3 \dim H^0_{\omega} - \dim H^1_{\omega} = 2 \dim H^0_{\omega}$ and $\dim H^0_{\omega} = \dim G$ for the trivial connection which is the maximally reducible connection.

This result is quite far from the known Reidemeister Torsion for regular elements of a Brieskorn Sphere $\Sigma(a_1, a_2, a_3)$ [32],

$$(a_1 a_2 a_3)^{-1} \prod_{i=1}^3 4 \sin^2 \left(\frac{n_i r_i \pi}{a_i}\right) \tag{4.10}$$

in the case of SU(2). The semi-classical analysis of Witten would have suggested

$$Z_{BF}[\Sigma(a_1, a_2, a_3) \operatorname{SU}(2)] = (a_1 a_2 a_3)^{-1} \prod_{i=1}^{3} \sum_{n_i=1}^{a_i-1} 4 \sin^2\left(\frac{n_i r_i \pi}{a_i}\right) + \text{reducible terms} \quad (4.11)$$

rather than the result that we obtained.

From (4.9) we understand that the ϕ_C contributions that we are looking at come from reducible connections, for these manifolds the trivial connection as G = SU(2), and so it appears that we see only the non acyclic terms in (4.11). For example for S^3 the only flat

³By (3.2) $Pb_j/a_j = 1 - a_j t_j$, $t_j \in \mathbb{Z}$ and $Pb_j r_j/a_j = r_j - r_j a_j t_j = (1 - a_j s_j) P/a_j$, $t_j \in \mathbb{Z}$ whence $r_j - P/a_j = 0 \mod a_j \mathbb{Z}$. Consequently, $\phi_C + 2\pi i \mathbf{n}_j r_j = -2\pi i P \sum_{i \neq j} \mathbf{n}_i/a_i + 2\pi i \mathbf{n}_j (r_j - P/a_j) = 0 \mod 2\pi i a_j I(G)$.

connection is the trivial connection and, as it has maximal reducibility, one sees that a formula of the type (4.9) correctly captures this fact. Indeed, for any (trivial) G bundle over a \mathbb{Z} Homology sphere the trivial connection is flat and isolated and would give a contribution as above. What are missing are the contributions of non-reducible or, for higher rank groups, less reducible connections.

4.3 The ϕ_C contribution to \mathbb{Q} homology spheres

The zeros that were faced when M is an \mathbb{Z} Homology Sphere are avoided to some extent when the manifold is a \mathbb{Q} Homology Sphere. The first advantage is that the points at which we wish to evaluate the torsion are no longer necessarily at integer values but rather at

$$\phi_C = -2\pi i \frac{P}{d} \sum_{i=0}^N \frac{\mathbf{n}_i}{a_i} \tag{4.12}$$

Even if ϕ_C does at some points take integer values, the mass regularised torsion remains well defined and that is what we will use. We will give an example of this later on. Recall that here we fix the symmetry (4.2) by insisting that $\mathbf{n}_0 \in I(G)/d.I(G)$.

The ϕ_C contribution to the partition function is, with the masses re-instated so that we may follow the reducibility of the connections,

$$Z_{BF}[M,G,\mu]|_{\phi_{C}} = \sum_{\mathbf{n}_{0}} \cdots \sum_{\mathbf{n}_{N}} \tau_{S^{1}} (-2\pi i (P/d) \widehat{\mathbf{q}} + 2\pi i \mathbf{n}_{0}, \mu)^{2-N} \cdot \prod_{i=1}^{N} \tau_{S^{1}} ((-2\pi i (P/d) \widehat{\mathbf{q}} + 2\pi i r_{i} \mathbf{n}_{i})/a_{i}, \mu/a_{i})$$

$$(4.13)$$

This result can be understood in terms of the homotopy representations for flat connections on the Seifert \mathbb{Q} Homology Sphere to which we turn shortly. Notice that the contribution from these connections is precisely of the form that was anticipated in (2.20). However, for present purposes the most useful form for us is where we have set the \mathbf{n}_i for $i \geq 1$ to zero and restrict $\mathbf{n}_0 \in I(G)/d.I(G)$

$$Z_{BF}[M,G,\mu]|_{\phi_C} = \sum_{\mathbf{n}_0 \in I(G)/d.I(G)} \tau_{S^1} (-2\pi i (P/d)\mathbf{n}_0,\mu)^{2-N} \cdot \prod_{i=1}^N \tau_{S^1} (-2\pi i (P/d)\mathbf{n}_0/a_i,\mu/a_i)$$
(4.14)

The classical examples of \mathbb{Q} Homology Spheres are of course the Lens spaces L(p,q). The simplest Seifert presentation of the Lens space L(p,q) is with N = 1, d = p and $a = q^*$ where $qq^* = 1 \mod p$, $p = b_0q^* + b_1$. Substituting these values into (4.14)

$$Z_{BF}[L(p,q),G,\mu] = \sum_{\mathbf{n}\in I(G)/p.I(G)} \tau_{S^1}(2\pi i q^* \mathbf{n}/p + \mu).\tau_{S^1}(2\pi i \mathbf{n}/p + \mu/q^*)$$
(4.15)

To specialise to G = SU(2) we note that the massive Ray-Singer Torsion (3.53) takes the form

$$\tau_{S^1}(\phi,\mu) = (1 - e^{-2\pi\mu}) \cdot (1 + e^{-4\pi\mu} - e^{-2\pi\mu} 2\cos 2\phi)$$
(4.16)

so that on specialising (4.15) to G = SU(2) one obtains

$$Z_{BF}[L(p,q), \mathrm{SU}(2), \mu] = \left(\frac{\mu^2}{q^*}\right)^3 + \dots + \left(\frac{\mu^2}{q^*}\right) \sum_{n=1}^{p-1} \sin^2\left(2\pi i q^* n/p\right) \cdot \sin^2\left(2\pi i n/p\right) + \dots$$
(4.17)

We note that happily (4.15) and (4.17) have the form that we expected from the covariant analysis in section 2 and in particular agree with the general form (2.22).

The first term in (4.17) corresponds to the trivial connection with reducibility of order 3 while the second term is the sum over the Abelian connections with reducibility of order 1 and the torsions agree with the formulae obtained by Ray [33] for Lens spaces. This should also be compared with the Chern-Simons large k limit formula (5.7) in [34] and we see that there is complete agreement with the expression for the Ray-Singer Torsion of the Lens spaces for the Abelian connections (the only difference with Jeffrey's formula is that we are summing over the Ray-Singer Torsion not its square root).

We note that ϕ_C in this case is the complete contribution to the integral (4.1) for, as noted in item 2 after (4.2), when N = 1 there are no poles in the integrand and the integral is well defined as is.

Now consider quotients of the Poincare sphere $\Sigma(2,3,5) = M[-1,g = 0,(2,1),$ (3,1), (5,1)]. The fundamental group of $\Sigma(2,3,5)$ is $\pi_1(\Sigma(2,3,5)) = I^*$ the binary icosahedral group of order 120 while the quotient manifold $\Sigma(2,3,5)_d = S(\mathcal{L}_{\Sigma(2,3,5)}^{\otimes d})$ has fundamental group $\mathbb{Z}_d \times I^*$ (Theorem 2 (v) page 112 in [35]). This means that $\Sigma(2,3,5)_d$ comes equipped with the non-Abelian flat connections of $\Sigma(2,3,5)$ through the I^* factor of the fundamental group and with Abelian connections through the \mathbb{Z}_d factor of π_1 . In this case we have

$$Z_{BF}[\Sigma(2,3,5)_d,G,\mu]|_{\phi_C} \simeq \sum_{0 \le \mathbf{n} \le d-1} \tau_{S^1} (60\pi i \mathbf{n}/d + \mu)^{-1} \cdot \tau_{S^1} (30\pi i \mathbf{n}/d + \mu/2) \cdot \tau_{S^1} (20\pi i \mathbf{n}/d + \mu/3) \cdot \tau_{S^1} (12\pi i \mathbf{n}/d + \mu/5)$$

$$(4.18)$$

Apart from having the standard form we note that we can expect, at least in the case of SU(2) that this formula correctly captures the contributions of the reducible connections.

Indeed all the so called small Seifert manifolds with genus zero and orientable base have finite fundamental groups with cyclic factors. Hence quite generally the space of flat connections for these manifolds naturally splits into the non-Abelian and Abelian parts. The cyclic group is generated by the fibre generator h. For large Seifert manifolds the fibre generator also generates the cyclic group as a normal subgroup of the infinite order fundamental group. Then $\mathbb{Z}_d \subset \pi_1(M_d)$ and $\mathbb{Z}_d = (\pi_1/[\pi_1, \pi_1])(M_d)$.

The interpretation of the formulae at ϕ_C that we arrived at above in terms of homotopy representations has been given in [17]. At these reducible connections the Chern-Simons invariant agrees with that given by Kirk and Klassen in [36], Auckly [37] and Nishi [38] and is consistent with the identification of the connections being reducible and flat. Also the known relationship between the Ray-Singer Torsion for flat connections and Reidemeister torsion corroborates the identification of the contributions that we find.

4.4 A residue formula for SU(2)

As we saw in section 4.3 that Abelian reducible connections are accounted for by ϕ_C we now take another and closer look at the finite-dimensional integral to see where the contributions from non-Abelian (ir)reducibles could come from.

The integrand of (4.1) has poles and zeros and so we must give meaning to the integral. A similar situation arises in Chern-Simons theory [13] where one explicitly removes the poles with the addition of a mass term. There is a good reason for this in Chern-Simons theory as, say on $\Sigma \times S^1$, one is counting the finite number of states in the theory. Here, if we use a mass regularisation then there is the same effect, namely the poles do not contribute. However, this then excludes important contributions to the *BF* partition function. Also, it should be said, that for *BF* theory one does not have a finite dimensional Hilbert space of states and so there is no 'directive' to ensure that the poles should not be counted in some way.

One could rotate the ϕ integral to say $\exp(-i\epsilon) \times \mathfrak{t}$ which would ensure that one avoids the poles altogether. However, one would need to simultaneously require a different path of integration for λ to ensure convergence of the integrals. Passing to the massive Ray-Singer Torsion is another way of avoiding the poles as the mass pushes the poles off the real \mathfrak{t} axis and into the complex plane $\mathbb{C} \times \mathfrak{t}$.

Short of a guiding principle we use the first symmetry in the Properties of the Integrand to restrict the range of each component of the ϕ field to lie in $(-\pi P, \pi P)$. In this way we do not have to worry about convergence issues with respect to integration over ϕ . However, there are still the poles to contend with and the integration over λ .

In order to be concrete and to fix ideas we now focus on G = SU(2). From the outset we are tasked with having to make sense of

$$Z_{BF}[M, \mathrm{SU}(2)] \sim \sum_{n \in \mathbb{Z}} \int_{\mathbb{R} \times \mathbb{R}} \tau_M(\phi) \cdot \exp\left(i\lambda \left(\phi \frac{d}{P} - 2\pi n\right)\right)$$
(4.19)

We firstly sum over n to exchange λ with the integers and exchange the integral with the sum

$$Z_{BF}[M, \mathrm{SU}(2)] \sim \int_{\mathbb{R}} \tau_M(\phi) \cdot \sum_{n \in \mathbb{Z}} \exp\left(in\phi \frac{d}{P}\right)$$
(4.20)

As $\tau_M(\phi)$ is even in ϕ we have

$$Z_{BF}[M, \mathrm{SU}(2)] \sim \int_{-\pi P}^{\pi P} \tau_M(\phi) \cdot \frac{(1 + \exp\left(id\phi/P\right))}{(1 - \exp\left(id\phi/P\right))}$$
$$\sim \int_{-\pi P}^{\pi P} \tau_M(\phi) \cdot \frac{f(\phi)}{(1 - \exp\left(id\phi/P\right))}$$
(4.21)

and we have used the symmetries available to limit the range of ϕ to lie in the range $(-\pi P, \pi P)$. Here the function f is

$$f(\phi) = (1 + \exp\left(id\phi/P\right)) \tag{4.22}$$

There are now poles on the real axis coming both from the Ray-Singer Torsion as well as from the result of the geometric sum. We push all of these, in the complex plane, to lie above the real line (say by use of an $i\epsilon$ prescription).

The contour that we choose in the complex plane in order to evaluate the integral is as follows: travel along the real axis from $-\pi P$ to πP then straight up along the imaginary axis to $\pi P + iR$ followed by an arc to $-\pi P + iR$ then straight down to $-\pi P$ and finally take the $R \to \infty$ limit. For the integral over the arc to make sense one regularises the integrand by multiplying by $\exp(i\epsilon\phi^2)$ with $\epsilon > 0$. The integrals over the other parts of the contour are convergent as we will see below. The fact that the prescription for the poles on the interval $(-\pi P, \pi P)$ are now such that they all lie inside the contour (in the upper half plane) means that the contour integral is just given by the residue of all of those poles.

Denote the contribution of each segment of the integral by the start and end points as I(x, y) so that we have, with $R \to \infty$

$$I(-\pi P, \pi P) + I(\pi P, \pi P + iR) - I(-\pi P, -\pi P + iR) = 2\pi i \operatorname{Res} \left(\tau_M(\phi) \cdot \frac{(1 + \exp(id\phi/P))}{(1 - \exp(id\phi/P))} \right)$$
(4.23)

as the integral over the arc is zero. Had we preserved the symmetry $\phi \rightarrow \phi + 2\pi P$ with our regulator then we would have been able to declare that $I(\pi P, \pi P + iR) = I(-\pi P, -\pi P + iR)$. However as ϕ becomes large the integrand of $I(\pm \pi P, \pm \pi P + iR)$ goes as

$$\exp\left(i\epsilon(i\phi\pm\pi P)^2 - 2\phi\left(N-2-\sum_{i=1}^N 1/a_i\right)\right) + \dots$$
(4.24)

and for $N \geq 3$ (but not for $\Sigma(2,3,5)$ the Poincare sphere) there is convergence without the need for the Gaussian oscillatory behaviour so that one may take the $\epsilon \to 0$ limit and the integrals over the two vertical parts of the contour cancel each other in the limit. In short, we have that with this prescription for handling the poles that

$$Z_{BF}[M, \mathrm{SU}(2)] = 2\pi i \mathrm{Res}\left(\tau_M(\phi) \cdot \frac{(1 + \exp\left(id\phi/P\right))}{(1 - \exp\left(id\phi/P\right))}\right)$$
(4.25)

One would also like to understand a little better what the residue formula implies. To that end let

$$\tau_M(\phi) = \prod_{\alpha>0} (\sin(\alpha(\phi)))^{4-2N} \cdot \prod_{\alpha>0} \prod_{i=1}^N (\sin(\alpha(\phi)/a_i))^2$$
$$= \prod_{\alpha>0} (\sin(\alpha(\phi)))^{4-2N} \cdot \tilde{\tau}_M(\phi)$$
(4.26)

The residue formula in the case of SU(2) takes the simple form

$$Z_{BF}[M, \mathrm{SU}(2)] = Z_{BF}[M, \mathrm{SU}(2)]|_{\phi_C} + \sum_{r=0}^{2N-5} {\binom{2N-5-r}{r}} \sum_{m=1-P}^{P-1} \left(f(m\pi) \cdot \frac{1}{1-\mathrm{e}^{im\pi d/P}}\right)^{(2N-5-r)} \tilde{\tau}_M(m\pi)^{(r)}$$
(4.27)

where the sum over m indicates the poles of $(\sin(\alpha(\phi)))^{4-2N}$. One sees that this is essentially a sum over derivatives of the Reidemeister torsion $\tilde{\tau}_M$ (up to normalisation).

4.5 SU(2) BF theory on Brieskorn spheres $\Sigma(a_1, a_2, a_3)$

We now look concretely at SU(2) BF Theory on Brieskorn Spheres $\Sigma(a_1, a_2, a_3)$, in order to see if the suggested contour reproduces the expectations from perturbation theory as derived by Witten [11]. Even though the result for the path integral is not exactly that predicted by Witten, we will see that this choice does have contributions from irreducible flat connections which is quite pleasing in its own right though, as we explain below, not completely unexpected.

With N = 3 and d = 1 the path integral becomes, apart from the ϕ_C contributions of the previous sections

$$Z_{BF}[\Sigma(a_1, a_2, a_3), SU(2)] \simeq \sum_{m=1}^{P-1} \left(\frac{1}{P} \csc^2(m\pi/2P) - 4\cot(m\pi/P) \sum_{i=1}^3 \frac{1}{a_i} \cot(m\pi/a_i) \right) \tilde{\tau}_{\Sigma(a_1, a_2, a_3)}(m\pi)$$
$$\simeq \frac{1}{a_1 a_2 a_3} \sum_{m=1}^{P-1} h(m) \tilde{\tau}_{\Sigma(a_1, a_2, a_3)}(m\pi)$$
(4.28)

To compare with (4.11) we note that the integer m can be written in various ways in particular

$$m = a_1\alpha_1 + n_1r_1 = a_2\alpha_2 + n_2r_2 = a_3\alpha_3 + n_3r_3, \quad \alpha_i, n_i \in \mathbb{Z}$$
(4.29)

to arrive at

$$Z_{BF}[\Sigma(a_1, a_2, a_3), \mathrm{SU}(2)] \simeq \frac{1}{a_1 a_2 a_3} \sum_{n_i} g(n_i) \prod_{i=1}^3 4 \sin^2\left(\frac{\pi n_i r_i}{a_i}\right)$$
(4.30)

which would agree with (4.11) if the function g is n_i independent.

As our first example (even though as we noted the derivation provided fails) consider the Poincaré homology sphere $\Sigma(2,3,5)$. The Ray-Singer-Torsion is the same for the values of (n_1, n_2, n_3) corresponding to (1, 1, r), (1, 1, 5 - r) and (1, 2, r). In this way we have, apart from the trivial connection, the two possible non-Abelian connections (1, 1, 1) and (1, 1, 2). By Proposition 2.8 of [39], this count is correct for the flat connections on $\Sigma(2, 3, 5)$.

As our next example let us fix on $\Sigma(2,3,7)$ for which there are 2 irreducible isolated flat connections. The Ray-Singer-Torsion is the same for the values of (n_1, n_2, n_3) corresponding to (1,1,r), (1,1,7-r) and (1,2,r). If we identify those contributions as corresponding to the same isolated and irreducible connection then we get 4 copies each of three different connections, say of (1,1,1), (1,1,2) and (1,1,3). In that case the sum in (4.30) is over the 3 different connections each of which corresponds to 4 different values of m as tabulated below (the values of g are approximate)

	m	(n_1, n_2, n_3)	$g/P = \sum_m h(m)/P$
	1, 13, 29, 41	(1, 1, 1)	-34.45
	5, 19, 23, 37	(1, 1, 2)	6.96
1	1, 17, 25, 31	(1, 1, 3)	3.35

Hence the partition function does not just give the sum of these connections with equal weight.

The situation is somewhat less clear than just indicated and rather more puzzling as the third contribution m = 11, 17, 25, 31 does not correspond to a flat SU(2) connection on $\Sigma(2,3,7)$. Rather this connection is one identified in [27] as a flat SL(2, \mathbb{C}) connection that is conjugate to SL(2, \mathbb{R}) and not to SU(2) (cf. also [40]).

The holonomies of the isolated irreducible flat connections are elements of the group and the holonomies can, independently, be conjugated into a preferred maximal torus (even though they are not reducible). The contribution to the Reidemeister Torsion of these holonomies is then determined by the integers **n** as described in [32] equation 2.8 for Brieskorn spheres. To apply this to an actual flat connection there are conditions on the integers arising from the presentation of the fundamental group. What we have in our situation is that all such integers arise as the poles of the Ray-Singer Torsion and, unfortunately, our residue formula (4.25) does not restrict only to the ones that correspond to honest flat SU(2) connections.

4.6 A residue formula for higher rank and outlook

Here we would like to briefly give a formula in terms of residues for the partition function (4.1) to use as a definition for general G.

A possible definition, is

$$Z_{BF}[M,G] \simeq 2\pi i \operatorname{Res}\left(f(\phi)\tau_M(\phi) \cdot \prod_{j=1}^{\dim \mathfrak{t}} \frac{1}{1 - e^{i\phi_j d/P}}\right)$$
(4.32)

where $\phi = \phi_j \alpha^j$ and the function f is such that $f(2\pi \mathbf{n} P/d) = 1$ for all $\mathbf{n} \in I(G)$,

$$f(\phi) = \sum_{\mathbf{n}\in\Lambda} e^{in^j \phi_j d/P} f_{\mathbf{n}}$$
(4.33)

The poles of $\prod_j (1 - \exp(i\phi_j d/P))^{-1}$ are at ϕ_C , (4.4), so summing over its residues at these poles is equivalent to performing the λ integral. The properties of $f(\phi)$ are so that at those poles f is unity (whence the contribution is exactly the same as performing the λ integral). The function f is obtained, as in the SU(2) case, by first performing the sum over \mathbf{n} to restrict the form of λ and symmetry properties within the integral

$$\sum_{\mathbf{n}} \int_{\mathfrak{t}} d\lambda \, \exp\left(i \operatorname{Tr} \lambda \left(d\phi/P + 2\pi i \mathbf{n}\right)\right) \to \prod_{j=1}^{\dim \mathfrak{t}} \frac{f(\phi)}{1 - \mathrm{e}^{i\phi_j d/P}} \tag{4.34}$$

It must also be remembered that the range of ϕ is constrained.

The poles of τ_M may also contribute to the residue. If one takes the attitude that one should use the massive Ray-Singer Torsion directly then there are no extra poles and (4.32) agrees with the naive evaluation of the path integral. Alternatively, (4.32) allows one to take the poles of τ_M into account after which one may reinstate the mass.

We have not shown that there is a contour that leads to (4.32) but one may reasonably hope that a generalisation of that used in the SU(2) case will be available.

Before ending this section we would like to suggest a possible apriori reason for why the definition adopted here has shortcomings. We start by observing that it is somewhat surprising that the possible isolated irreducible flat connections are determined by poles in the Ray-Singer Torsion. This is certainly counterintuitive as the poles of the Ray-Singer Torsion are understood to arise when one has moduli, i.e. 1-form zero modes. A possible reason for this is a shortcoming in the gauge-fixing choice that has been made. As already noted in order to gauge fix λ to be constant on the fibre we require that ϕ be generic 'enough'. To see that this may be the cause of concern, imagine that one wishes, as we did in section 2, to expand around an isolated irreducible flat connection. We do this but we still insist on gauge fixing the quantum fields as we did in section 3 so that λ is constant along the fibre. Then we would need to solve the equivalent of (3.22)

$$\lambda + D_{\phi_f + \phi} \Lambda = \lambda_0 \tag{4.35}$$

where ϕ_f is the fibre component of the background flat connection and ϕ is the fibre component of the quantum field. If one thinks of the quantum field as having a \hbar in front of it then to lowest order the equation to be solved is

$$\lambda + D_{\phi_f} \Lambda \simeq \lambda_0 \tag{4.36}$$

and the field ϕ_f is certainly not generic. The discussion in section 3 tells us that precisely for these ϕ_f , as explained around (3.31)–(3.32), the operator D_{ϕ_f} is not invertible and the gauge $\partial_t \lambda = 0$ cannot be achieved.

Given this question about the gauge choice and the appearance of a complex flat connection in the SU(2) theory one may wonder if resurgence is an approach that may demystify the situation. The BF theory, as we have approached it here, arises as one possible limit of $G_{\mathbb{C}}$ Chern-Simons theory (see the next section). In the SU(2) case the reducible connections are Abelian and one may hope that the resurgence programme will yield the correct non-Abelian connections as they do in [27] with the correct contribution to the partition function.

5 Large k asymptotics of $G \times G$ and $G_{\mathbb{C}}$ Chern-Simons theory

Our aim in this section is to see if embedding BF theory in a 'bigger' theory may act to give a suitable definition which can be used in case M is an \mathbb{Z} Homology Sphere. To that end we consider some naive large k limits of Chern-Simons theory with the compact gauge group $G \times G$ (with levels k and -k for the two factors) and the complexified group $G_{\mathbb{C}}$. The 'fattened' groups $G_{-k} \times G_k$ and $G_{\mathbb{C}}$, as has already been explained (see appendix A), have been chosen as in the limits to be discussed they both contract to TG.

The advantage of such an approach is that we avoid the possible issues we had with the generic nature (or not) of ϕ as the values of ϕ signalled out prior to taking the large k limit allow for the gauge where λ is constant along the fibre.

In the case of $G_{-k} \times G_k$ we find that the theory does not appear to 'decompactify' enough in the large k limit to capture all the features of the TG theory. However, in examples, we show that it does actually pick up those flat connections which are both isolated and irreducible correctly. The $G_{\mathbb{C}}$ Chern-Simons theory, on the other hand, has all the features of the TG theory from the outset. We show that the partition function has the formal properties that would allow an application of the large k analysis that have worked in the case of the compact groups.

5.1 Definition 2: BF and the large k limit of $G_k \times G_{-k}$ Chern-Simons theory

Here we consider the Chern-Simons action for two connections on two copies of the same bundle for the product gauge group $G \times G$ and for the following gymnastics we require that the bundle in question is trivial. The action under consideration is

$$CS(A_{+}, A_{-}) = \frac{1}{4\pi} \int_{M} \left[\operatorname{Tr} \left(A_{+} \wedge dA_{+} + \frac{2}{3} A_{+} \wedge A_{+} \wedge A_{+} \right) - \operatorname{Tr} \left(A_{-} \wedge dA_{-} + \frac{2}{3} A_{-} \wedge A_{-} \wedge A_{-} \right) \right]$$
(5.1)

The relative sign means that the two gauge theories have opposite levels and we denote such a theory, with action $k CS(A_+, A_-)$ as a $G_k \times G_{-k}$ Chern-Simons theory. If we denote the level k Chern-Simons theory partition function for group G by $Z_{CS}[M, G_k]$ then the partition function for the $G_k \times G_{-k}$ theory is

$$Z_{CS}[M, G_k \times G_{-k}] = Z_{CS}[M, G_k] . Z_{CS}[M, G_{-k}]$$
(5.2)

Now to show how this is related to BF theory make the substitutions

$$A_{\pm} = A \pm \frac{\pi}{k} B, \quad A = \frac{1}{2} (A_{+} + A_{-}), \quad B = \frac{k}{2\pi} (A_{+} - A_{-})$$
 (5.3)

to arrive at

$$CS(A_+, A_-) = \int_M \operatorname{Tr}\left(B \wedge F_A + \frac{\pi^2}{k^2}B \wedge B \wedge B\right)$$
(5.4)

There may be a Jacobian J, depending on k in passing to the new variables in the path integral, so the relationship that we expect is finally

$$Z_{BF}[M,G] = \lim_{k \to \infty} J(k) Z_{CS}[M,G_k \times G_{-k}]$$
(5.5)

where the limit as $k \to \infty$ formally ensures that the cubic term in B may be neglected. We allow J to also soak up any other factors of k that may be present due to zero modes and so on.

This formal result says that the $k \to \infty$ limit of a $G_k \times G_{-k}$ Chern-Simons theory is equivalent to a pure BF theory. However, this statement certainly requires some elaboration as adding and subtracting connections at will as in (5.3) is not usually an operation that makes sense in the theory of bundles. This is mirrored in the gauge transformations that one obtains for A and B which are also not what one might call 'standard'. To emphasise that the limit in question is by no means obvious consider the 3-manifold $\Sigma \times S^1$. Chern-Simons theory on such a manifold is perfectly sensible and the large k limit is well understood, the partition function having leading term $k^{n/2} \operatorname{Vol}(\mathfrak{M}[\Sigma, G])$ the symplectic volume of the moduli space of G connections on Σ where $n = \dim(\mathfrak{M}[\Sigma, G])$ [42]. The large k limit of the $G_k \times G_{-k}$ partition function is then proportional to $k^n \operatorname{Vol}(\mathfrak{M}[\Sigma, G])^2$ and not what one would expect from BF theory, say $k^n \operatorname{Vol}(T\mathfrak{M}[\Sigma, G])$ (even though this is ill defined).

So we need to better understand when the limit will indeed correspond to BF theory. At large k, the stationary phase approximation tells us that the functional integral [12] localises around the critical points of the action CS(A), which in this case are precisely the flat connections. We then have, for isolated flat connections, as $k \to \infty$,

$$Z_k[M] \sim \sum_{\alpha} e^{2\pi i k C S(\alpha)} e^{-\frac{i\pi}{2}I(\alpha)} \sqrt{\tau_M(\alpha)}$$
(5.6)

where α , $\tau_M(\alpha)$ and $I(\alpha)$ are the flat connections, the Ray-Singer torsion at the flat connection and the spectral flow to the flat connection respectively.

The $G_k \times G_{-k}$ theory then has the asymptotic form

$$Z_k[M]Z_{-k}[M] \sim \sum_{\alpha,\beta} e^{ik \left(CS(\alpha) - CS(\beta)\right)} f_{\alpha\beta} \sqrt{\tau_M(\alpha)} \sqrt{\tau_M(\beta)}$$
(5.7)

where

$$f_{\alpha\beta} = \exp\left(-\frac{i\pi}{2}[I(A_{\alpha}) - I(A_{\beta})]\right)$$
(5.8)

One would expect therefore, that in the limit as $k \to \infty$, that providing that flat connections α and β having the same Chern-Simons invariant implies they are the same flat connection,

 $CS(\alpha) = CS(\beta) \mod 2\pi\mathbb{Z} \implies \alpha = \beta$ (5.9)

then the oscillations in (5.7) would ensure that we would indeed only need to sum over $\alpha = \beta$. In such a situation then one could reasonably expect that (5.5) holds. If this condition does not hold then there may be 'non-diagonal' contributions to the sum (5.7) and so the relationship with *BF* theory becomes more tenuous.

Indeed for any connected component of non-isolated flat connections all the flat connections in that component have the same Chern-Simons invariant and fail our test (5.9). The flat connections on $\Sigma \times S^1$ that we discussed previously are of this type and this, to some extent, explains why they do not match the expectations for BF theory. The lesson here is that we must concentrate on isolated flat connections. Fortunately for us Fintushil and Stern [39] have shown that the Brieskorn spheres (the Seifert Z homology spheres with 3 exceptional fibres) have moduli spaces of flat SU(2) connections made up of a finite number of discrete points.

We shall now illustrate when (5.9) holds with a few examples in the literature. Consider firstly the Poincaré Homology Sphere, $M = \Sigma(2,3,5)$. The Poincaré Z homology sphere has three flat SU(2) connections, one of which is the reducible trivial connection and two which are non-Abelian. Freed and Gompf [19] determine the large k behaviour for the Chern-Simons partition function and find

$$Z_k[\Sigma(2,3,5)] \sim \sqrt{\frac{2}{5}} e^{-3\pi i/4} \left[\sin\left(\frac{\pi}{5}\right) e^{-\pi i(k+2)/60} + \sin\left(\frac{2\pi}{5}\right) e^{-49\pi i(k+2)/60} \right]$$
(5.10)

Taking the modulus square, and presuming the limit of the norm agrees with the norm of the limit, one gets

$$\lim_{k \to \infty} Z_k[\Sigma(2,3,5)] Z_{-k}[\Sigma(2,3,5)] \sim \frac{2}{5} \left[\sin^2\left(\frac{\pi}{5}\right) + \sin^2\left(\frac{2\pi}{5}\right) \right]$$
(5.11)

this corresponds to the sum of the Ray-Singer torsions of the irreducible connections for $\Sigma(2,3,5)$ up to a finite normalisation. Similarly [19] provides us with the example of $\Sigma(2,3,17)$ and, with the same caveat on exchanging limits and norms, we find that the norm square of the Chern-Simons partition function is the sum of the Ray-Singer Torsion over the 6 irreducible flat SU(2) connections.

There are manifolds, however, where you find two or more isolated flat connections giving the same Chern-Simons invariant so that (5.9) does not hold. Lens spaces yield examples of this phenomena. As we have seen for Lens spaces L(p,q) the Ray-Singer Torsion is given by

$$\tau_{L(p,q)} = \frac{16}{p} \sin^2\left(\frac{2\pi n}{p}\right) \quad \sin^2\left(\frac{2\pi q^* n}{p}\right) \tag{5.12}$$

Here one integer is enough to describe the flat (Abelian) connections.

A typical example where two different connections have the same Chern-Simons invariant is afforded by the Lens space L(12, 5). For L(12, 5) we have $q^* = 5$ and one can see for example that n = 1 and n = 5 give the same Chern-Simons invariant (mod 2π), namely

$$CS(n) = q^* n^2 / p = \frac{5}{12}$$
(5.13)

so we are not free to use the norm of the Chern-Simons theory in this case as a way to define the BF theory.

The discussion of this section shows that this definition of BF theory as a limit of the $G_{-k} \times G_k$ Chern-Simons theory needs to be handled with care and we might have to know more about the moduli space itself than we would care to. However, for the part of the moduli space where the connections are isolated and irreducible this appears to be an appropriate definition.

5.2 Definition 3: BF and the large k or s limit of $G_{\mathbb{C}}$ Chern-Simons theory

The final definition that we give involves the gauge group $G_{\mathbb{C}}$. Note that the classical groups TG and $G_{\mathbb{C}}$ are diffeomorphic (as spaces). The Inönü-Wigner contraction that establishes the Lie algebra relationship is given in appendix A. Furthermore, the $G_{\mathbb{C}}$ connection is

$$A_{\mathbb{C}} = A + iB \tag{5.14}$$

and as $G_{\mathbb{C}}$ contracts to the compact group G the connection A may be considered a G connection while B is then a Lie algebra \mathfrak{g} valued one-form. This is a much clearer decomposition than the mixing of objects that appeared in the $G_{-k} \times G_k$ theory of the previous section.

The $G_{\mathbb{C}}$ Chern-Simons action is

$$I(k,s) = \frac{k}{4\pi} \int_{M} \operatorname{Tr} \left(A \wedge dA + \frac{2}{3}A \wedge A \wedge A - B \wedge d_{A}B \right) - \frac{s}{2\pi} \int_{M} \operatorname{Tr} \left(B \wedge F_{A} - \frac{1}{3}B \wedge B \wedge B \right)$$
(5.15)

with $k \in \mathbb{Z}$ and we take $s \in \mathbb{R}$ [20]. One can see that the complex Chern-Simons action has a number of limits which formally lead to the BF action. The first, and most straightforward way, to arrive at the BF theory is to set k = 0 and consider the $s \to \infty$ limit where one sends $B \to B/s$ to formally arrive at

$$I(k,s) \to -\frac{1}{2\pi} \int_M \operatorname{Tr} \left(B \wedge F_A \right)$$
 (5.16)

Alternatively one can consider the large k limit, for finite s, and this time by scaling $B \to B/\sqrt{k}$. One does not get the BF action directly, but rather

$$I(k,s) \to \frac{k}{4\pi} \int_{M} \operatorname{Tr}\left(A \wedge dA + \frac{2}{3}A \wedge A \wedge A\right) - \frac{1}{4\pi} \int_{M} \operatorname{Tr}\left(B \wedge d_{A}B\right)$$
(5.17)

About an isolated flat connection this would be the same as the large k limit of Chern-Simons theory except that the integral over the field B has, formally the effect of replacing the square root of the Ray-Singer torsion of the flat connection with the Ray-Singer torsion itself and so would, in the limit, reproduce the partition function of BF theory with a Chern-Simons action contribution of the flat connection. Both limits (and indeed any other limits of k and s) need to be carefully taken as in principle one would most likely land on the moduli space of flat $G_{\mathbb{C}}$ connections and not just the flat G connections.

For finite k and with s = 0 Gukov and Pei [23] have evaluated the partition function on $M = \Sigma \times S^1$. Even though the Hilbert space is also infinite dimensional in this case, they 'filter' it by giving the B field a mass and so are able to extract meaningful results. Unlike the situation with the $G_{-k} \times G_k$, where one gets essentially the square of the compact case, here one sees that the results come from a non-compact group and formally diverges as $k \to \infty$ as we would expect for the BF theory.

Returning to the case at hand note that the corresponding finite dimensional integral to be performed in the case of complex Chern-Simons after Abelianisation [17] has as its action

$$I(k,s) = -\frac{k}{4\pi} \operatorname{Tr}(\phi^2 - \lambda^2) c_1(\mathcal{L}_M) + k \sum_{i=1}^N \frac{1}{a_i} \operatorname{Tr}(-i\phi \mathbf{n}_i + r_i \pi \mathbf{n}_i^2) + k \operatorname{Tr}(-i\phi \mathbf{n}_0) + \frac{s}{2\pi} \operatorname{Tr}\left(\lambda \left(\phi c_1(\mathcal{L}_M) + 2\pi i \sum_{i=1}^N \frac{1}{a_i} \mathbf{n}_i + 2\pi i \mathbf{n}_0\right)\right)$$
(5.18)

while the Ray-Singer Torsion goes over to the complex Ray-Singer Torsion

$$\tau_M(\phi) \to \sqrt{\tau_M(\phi + i\lambda)\tau_M(\phi - i\lambda)}$$
(5.19)

Formally the complex Ray-Singer Torsion under either scaling goes back to the 'real' Ray-Singer Torsion. The k = 0 and $s \to \infty$ limit lands us on the finite dimensional action that we have been using for BF theory. Here we would like to investigate the other limit.

The limit of interest then is large k and for $\lambda \to \lambda/\sqrt{k}$. As λ now only appears in the action quadratically one may integrate it out. In this limit the finite action goes over to

$$I(k,s) \rightarrow -\frac{k}{4\pi} \operatorname{Tr}(\phi^2) c_1\left(\mathcal{L}_M\right) + k \sum_{i=1}^N \frac{1}{a_i} \operatorname{Tr}(-i\phi \mathbf{n}_i + r_i \pi \mathbf{n}_i^2) + k \operatorname{Tr}(-i\phi \mathbf{n}_0)$$
(5.20)

which is just the standard Chern-Simons action for compact group G. The partition function, however, includes an extra factor of the square root of the Ray-Singer Torsion (arising from the integral over the field B in (5.17)) in the integral over t so this is not the partition function of compact Chern-Simons theory.

As the action agrees with that of G Chern-Simons theory then all the properties of the integrand (see appendix B for the symmetry properties) and hence the considerations that appear in [24] and [25] apply here too. This includes the choice of contour to take the large k limit, the only difference being that of considering a slightly different function that takes into account that we have the Ray-Singer Torsion not its square root in the integrand. The resulting large k asymptotics are, for Chern-Simons theory, just as predicted by Witten in [12], contributions around flat connections. Consequently, in the limit that we are considering in the $G_{\mathbb{C}}$ theory, this means that the asymptotic form will have contributions from flat connections in BF theory as anticipated in section 2.

Specifically in the case of SU(2) the large k limit is that of the flat connection contributions on page 302 of [24] but with F(y) there replaced with $F(y)^2$ in order to pass to the *BF* theory (and one should not include the framing as it ought not to arise in complex Chern-Simons theory) while for a general compact group one can refer to (4.17) in [25] from which one can deduce the large k behaviour. At this point this particular definition of *BF* theory is the most complete that we have.

If one would like to obtain more explicit formulae for the BF theory one could follow the explanation of [24] on how to arrive at Rozansky's formula for the large k expansion of SU(2) Chern-Simons theory [41] which reproduces Witten's expansion. Of course this also encodes some of the large k structure of the $G_{\mathbb{C}}$ theory itself so that by following different parts of the asymptotic expansion we would be able to obtain parts of the perturbation theory. In particular we also have in mind the contributions about the trivial connection. We leave these issues, and the question of how and whether complex connections contribute to the BF limit for the future.

Acknowledgments

M. Kakona thanks the External Activities Unit of the ICTP for a Ph.D. grant at EAIFR in the University of Rwanda and the HECAP group at the ICTP for supporting a visit to complete this work. He is also indebted to OFID and the STEP programme of the ICTP for funding and support in the initial stages of his studies. The group of M. Blau is supported by the NCCR SwissMAP (The Mathematics of Physics) of the Swiss Science Foundation.

A The tangent bundle group TG and its Lie algebra \mathfrak{tg}

A.1 Basic properties of TG and \mathfrak{tg}

let G be a compact Lie group and (to not unnecessarily complicate things) assume that G is connected, simply-connected and semi-simple. Via a right-invariant trivialisation of the tangent bundle, the group TG can be identified with the semi-direct product of G with its Lie algebra \mathfrak{g} ,

$$TG \simeq G \ltimes \mathfrak{g} \equiv G \times_{\mathsf{Ad}} \mathfrak{g}, \tag{A.1}$$

with multiplication

$$(g,v)(h,w) = (gh, v + \operatorname{Ad}_g w)\}, \qquad (A.2)$$

where the adjoint action of $g \in G$ on $w \in \mathfrak{g}$ is $\operatorname{Ad}_g w = gwg^{-1}$. Correspondingly, its Lie algebra has the form

$$\mathsf{Lie}(TG) \equiv \mathfrak{tg} \simeq \mathfrak{g} \oplus_{\mathsf{ad}} \mathfrak{g}_{Ab} \tag{A.3}$$

where \mathfrak{g}_{Ab} is the Abelian Lie algebra based on the underlying vector space \mathfrak{g} , i.e. the commutator is

$$[(x,v),(y,w)] = ([x,y],[x,w] + [v,y]) = ([x,y],\mathsf{ad}_x w - \mathsf{ad}_y v)$$
(A.4)

and the (inverse) adjoint action of TG on its Lie algebra is

$$(g,v)^{-1}(y,w)(g,v) = (\mathsf{Ad}_{g^{-1}}y, \mathsf{Ad}_{g^{-1}}(w+[y,v]))$$
(A.5)

Let ℓ_a be a (real) basis of the (real) Lie algebra \mathfrak{g} , with

$$[\ell_a, \ell_b] = f_{ab}^c \ell_c \,. \tag{A.6}$$

Then

$$j_a = (\ell_a, 0), \quad p_a = (0, \ell_a)$$
 (A.7)

are a basis of \mathfrak{tg} , and the Lie algebra commutation relations take the form

$$[j_a, j_b] = f_{ab}^c j_c, \quad [j_a, p_b] = [p_a, j_b] = f_{ab}^c p_c, \quad [p_a, p_b] = 0.$$
(A.8)

Here are two useful properties of \mathfrak{tg} :

- 1. The Lie algebra \mathfrak{tg} of TG can be obtained as a contraction of the Lie algebra of the group $G\times G$
 - Consider the $G \times G$ Lie algebra

$$\mathfrak{g}_+ \oplus \mathfrak{g}_- = \mathfrak{g} \oplus \mathfrak{g} \tag{A.9}$$

with generators

$$[j_a^{\pm}, j_b^{\pm}] = f_{ab}^c j_c^{\pm}, \quad [j_a^{+}, j_b^{-}] = 0.$$
 (A.10)

• Perform the redefinition

$$j_a^{\pm} = \frac{1}{2}(j_a \pm p_a/\epsilon) \quad \Leftrightarrow \quad j_a = j_a^{+} + j_a^{-}, \quad p_a = \epsilon(j_a^{+} - j_a^{-})$$
(A.11)

• Then the algebra $\mathfrak{g} \oplus \mathfrak{g}$ takes the form

$$[j_a, j_b] = f_{ab}^c j_c, \quad [j_a, p_b] = [p_a, j_b] = f_{ab}^c p_c, \quad [p_a, p_b] = \epsilon^2 f_{ab}^c j_c.$$
(A.12)

- In the limit $\epsilon \to 0$, this reduces to (A.8), the Lie algebra of TG.
- If one sets ε = iδ, the same reasoning shows that tg can be obtained as the contraction of the complexification g_C = g ⊕ ig of g.

2. Existence of Invariant Scalar Products on tg

Let us consider the case where the Lie algebra \mathfrak{g} is simple. In that case, \mathfrak{g} has a preferred (and up to a choice of scale unique) non-degenerate and ad-invariant metric scalar product, namely the Killing-Cartan form $\operatorname{Tr} \operatorname{ad}_x \operatorname{ad}_y$ (the trace in the adjoint representation). With respect to the basis ℓ_a of generators, this metric has the components

$$g_{ab} \equiv <\ell_a, \ell_b >= \operatorname{Tr} \operatorname{ad}_{\ell_a} \operatorname{ad}_{\ell_b} = f_{ac}^d f_{bd}^c \tag{A.13}$$

Turning to the Lie algebra

$$\mathfrak{tg} \simeq \mathfrak{g} \oplus_{\mathsf{ad}} \mathfrak{g}_{Ab} \,, \tag{A.14}$$

as it is not semi-simple, its Killing-Cartan form will be \mathfrak{tg} -invariant (by construction, i.e. by the Jacobi identity) but degenerate. Indeed, in terms of the generators (j_a, p_a) one has

$$\operatorname{Tr} \operatorname{ad}_{j_a} \operatorname{ad}_{j_b} = 2g_{ab} = 2f_{ac}^d f_{bd}^c \tag{A.15}$$

(from the adjoint action of $\mathsf{ad}_{j_a}\mathsf{ad}_{j_b}$ on j_c and on p_c), and

$$\operatorname{Tr} \operatorname{ad}_{j_a} \operatorname{ad}_{p_b} = \operatorname{Tr} \operatorname{ad}_{p_a} \operatorname{ad}_{p_b} = 0 \tag{A.16}$$

(because both $\operatorname{ad}_{j_a}\operatorname{ad}_{p_b}$ and $\operatorname{ad}_{p_a}\operatorname{ad}_{p_b}$ only act non-trivially on a j_c and take it to a linear combination of the p_c). This scalar product can therefore also be written as

$$\langle j_a, j_b \rangle = 2g_{ab}, \quad \langle j_a, p_b \rangle = \langle p_a, p_b \rangle = 0.$$
 (A.17)

In addition to the (degenerate) Killing-Cartan form, \mathfrak{tg} exceptionally also possesses a non-degenerate \mathfrak{ad} -invariant scalar product given by

$$\ll j_a, j_b \gg = \ll p_a, p_b \gg = 0, \quad \ll j_a, p_b \gg = g_{ab}.$$
 (A.18)

It is easily verified that this is indeed both invariant and non-degenerate.

The existence of this second invariant scalar product, or of the overall two-parameter family of invariant scalar products, can be understood in terms of the contraction of the Lie algebra $\mathfrak{g} \oplus \mathfrak{g}$ to \mathfrak{tg} mentioned above. Indeed, starting with the non-degenerate Killing-Cartan metric on the Lie algebra $\mathfrak{g} \oplus \mathfrak{g}$ (A.10) of $G \times G$, with coefficients c^{\pm} ,

$$\langle j_a^{\pm}, j_b^{\pm} \rangle = c^{\pm} g_{ab}, \quad \langle j_a^{+}, j_b^{-} \rangle = 0,$$
 (A.19)

and performing the redefinition (A.11), one finds

$$\langle j_a, j_b \rangle = (c^+ + c^-)g_{ab}, \quad \langle j_a, p_b \rangle = \epsilon(c^+ - c^-)g_{ab}, \quad \langle p_a, p_b \rangle = \epsilon^2(c^+ + c^-)g_{ab}$$
(A.20)

Taking the contraction $\epsilon \to 0$ with $c^+ = c^- = 1$, one finds (A.17), while taking $c^+ = -c^- = 1/2\epsilon$ one finds (A.18),

$$c^{+} = c^{-} = 1 \quad \Rightarrow \quad < ., . >$$

$$c^{+} = -c^{-} = 1/2\epsilon \quad \Rightarrow \quad \ll ., . \gg$$
(A.21)

A.2 Diagonalisation / Abelianisation in TG and \mathfrak{tg}

Let G be compact, T_G be a maximal torus, \mathfrak{t}_G its Lie algebra, a Cartan subalgebra. Then the following assertions are true:

1. For any $h \in G$ one can find a $g \in G$ such that

$$h \in G \quad \Rightarrow \quad \exists g \in G : \quad \operatorname{Ad}_g h = ghg^{-1} \in T_G.$$
 (A.22)

2. For any $y \in \mathfrak{g}$ one can find a $g \in G$ such that

$$y \in \mathfrak{g} \quad \Rightarrow \quad \exists g \in G : \quad \mathsf{Ad}_{g} y = g y g^{-1} \in \mathfrak{t}_{G}.$$
 (A.23)

As TG is not compact and not semi-simple, a priori the corresponding statements do not necessarily hold (and would usually be far from true for a generic such group). Nevertheless, it turns out that the above statements carry over literally to the case of the group TG, provided that we replace the maximal torus T_G of G and the Cartan subalgebra \mathfrak{t}_G of \mathfrak{g} by

$$T_{TG} = T(T_G) \simeq T_G \times_{\mathsf{Ad}} \mathfrak{t}_G = T_G \times \mathfrak{t}_G \tag{A.24}$$

$$\mathfrak{t}_{TG} = \mathsf{Lie}(T_G \times \mathfrak{t}_G) = \mathfrak{t}_G \oplus \mathfrak{t}_G \tag{A.25}$$

(since \mathfrak{t}_G is already an Abelian Lie algebra, here it is not necessary to write $(\mathfrak{t}_G)_{Ab}$ in the second factor / summand). Indeed, we can now prove the following two statements:

1. Diagonalisation / Abelianisation in TG

$$\forall (h,w) \in TG \simeq G \times_{\mathsf{Ad}} \mathfrak{g} \quad \exists (g,v) \in TG : \quad (g,v)^{-1}(h,w)(g,v) = (t,\tau_2) \in T_G \times \mathfrak{t}_G$$
(A.26)

2. Diagonalisation / Abelianisation in \mathfrak{tg}

$$\forall (y,w) \in \mathfrak{tg} \simeq \mathfrak{g} \oplus_{\mathsf{ad}} \mathfrak{g} \quad \exists (g,v) \in TG : \quad (g,v)^{-1}(y,w)(g,v) = (\tau_1,\tau_2) \in \mathfrak{t}_G \oplus \mathfrak{t}_G$$
(A.27)

Here are the proofs of these assertions:

- 1. Diagonalisation / Abelianisation in TG
 - The (inverse) Ad-action of TG on itself is

$$(g,v)^{-1}(h,w)(g,v) = (\mathsf{Ad}_{g^{-1}}h, \mathsf{Ad}_{g^{-1}}(w + \mathsf{Ad}_hv - v))$$
(A.28)

In particular, for the action of (g, 0) one finds

$$(g,0): \quad h \mapsto \operatorname{Ad}_{g^{-1}}h, \quad w \mapsto \operatorname{Ad}_{g^{-1}}w \tag{A.29}$$

and for that of (e, v) one has

$$(e,v): \quad h \mapsto h, \quad w \mapsto w + \mathsf{Ad}_h v - v.$$
 (A.30)

• First of all, we can and will choose $g \in G$ such that

$$h \mapsto g^{-1}hg = t \in T_G. \tag{A.31}$$

• Next we define S to be the stabiliser of t under the adjoint action of G,

$$S := \mathsf{Stab}_G(t) = \{ g \in G : g^{-1}tg = t \}.$$
(A.32)

One has

$$S \supseteq T_G$$
, (A.33)

with equality if t is generic (regular). In the non-generic case, S will be a product of simple factors and U(1)s. We use the convention that $T_S = T_G$, and likewise at the Lie algebra level,

$$T_S = T_G, \quad \mathfrak{t}_S = \mathfrak{t}_G. \tag{A.34}$$

At the Lie algebra level one has the (reductive, because \mathfrak{g} is compact) decomposition

$$\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{m} \quad \text{with} \quad [\mathfrak{s}, \mathfrak{m}] \subset \mathfrak{m} \,.$$
 (A.35)

• Now let us turn to the second entry in (A.28),

$$w \mapsto \mathsf{Ad}_{a^{-1}}(w + \mathsf{Ad}_h v - v) \,. \tag{A.36}$$

With $h = t \in T_G$ and thus $g = s \in S$, this becomes

$$w \mapsto \mathsf{Ad}_{s^{-1}}(w + \mathsf{Ad}_t v - v) \,. \tag{A.37}$$

Decomposing $v = v^{\mathfrak{s}} + v^{\mathfrak{m}}$ into its components in $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{m}$, we see that only the \mathfrak{m} -component of v contributes, since $\operatorname{Ad}_t v^{\mathfrak{s}} = v^{\mathfrak{s}}$,

$$\mathsf{Ad}_t v - v = (\mathsf{Ad}_t - 1)v^{\mathfrak{m}} \in \mathfrak{m}, \qquad (A.38)$$

and $v^{\mathfrak{m}}$ can be chosen to cancel the \mathfrak{m} -component $w^{\mathfrak{m}}$ of w (because by definition of \mathfrak{m} the operator $(\mathsf{Ad}_t - 1)$ is invertible on \mathfrak{m}).

• We are then left with

$$w \mapsto \mathsf{Ad}_{s^{-1}} w^{\mathfrak{s}}, \tag{A.39}$$

and we can now choose $s \in S$ such that

$$w \mapsto \mathsf{Ad}_{s^{-1}} w^{\mathfrak{s}} = \tau_2 \in \mathfrak{t}_S = \mathfrak{t}_G \,. \tag{A.40}$$

• Altogether, we have thus managed to conjugate

$$(h,w) \mapsto (t,\tau_2) \in T_G \times \mathfrak{t}_G, \qquad (A.41)$$

as announced.

- 2. Diagonalisation / Abelianisation in tg
 - The inverse adjoint action of TG on its Lie algebra is

$$(g,v)^{-1}(y,w)(g,v) = (\mathsf{Ad}_{g^{-1}}y, \mathsf{Ad}_{g^{-1}}(w+[y,v]))$$
(A.42)

In particular, for the action of (g, 0) one finds

$$(g,0): \quad y \mapsto \mathsf{Ad}_{g^{-1}}y \,, \quad w \mapsto \mathsf{Ad}_{g^{-1}}w \tag{A.43}$$

and for that of (e, v) one has

$$(e,v): \quad y \mapsto y, \quad w \mapsto w + [y,v]. \tag{A.44}$$

• First of all, we can and will choose $g \in G$ such that

$$y \mapsto g^{-1} y g = \tau_1 \in \mathfrak{t}_G. \tag{A.45}$$

• Next we define S to be the stabiliser of τ_1 under the adjoint action of G,

$$S := \mathsf{Stab}_G(\tau_1) = \{ g \in G : g^{-1}\tau_1 g = \tau_1 \}.$$
 (A.46)

One has

$$S \supseteq T_G$$
, (A.47)

with equality if τ_1 is generic (regular). As above, we set $T_S = T_G$ and $\mathfrak{t}_S = \mathfrak{t}_G$, with $\mathfrak{g} = \mathfrak{s} \oplus \mathfrak{m}$ and $[\mathfrak{s}, \mathfrak{m}] \subset \mathfrak{m}$.

• Now let us turn to the second entry in (A.42),

$$w \mapsto \mathsf{Ad}_{q^{-1}}(w + [y, v]) \tag{A.48}$$

With $y = \tau_1 \in \mathfrak{t}_G$, and thus $g = s \in S$, this becomes

$$w \mapsto \mathsf{Ad}_{s^{-1}}(w + [\tau_1, v]) \tag{A.49}$$

Decomposing $v = v^{\mathfrak{s}} + v^{\mathfrak{m}}$, only $v^{\mathfrak{m}}$ contributes to the commutator,

$$[\tau_1, v] = [\tau_1, v^{\mathfrak{m}}] \in \mathfrak{m}, \qquad (A.50)$$

and $v^{\mathfrak{m}}$ can be chosen to cancel $w^{\mathfrak{m}}$ (because by definition of \mathfrak{m} the operator ad_{τ_1} is invertible on \mathfrak{m}).

• We are then left with

$$w \mapsto \mathsf{Ad}_{s^{-1}} w^{\mathfrak{s}} \,, \tag{A.51}$$

and as in the proof above we can now choose $s \in S$ such that

$$w \mapsto \mathsf{Ad}_{s^{-1}} w^{\mathfrak{s}} = \tau_2 \in \mathfrak{t}_S = \mathfrak{t}_G \,. \tag{A.52}$$

• Altogether, we have thus managed to conjugate

$$(y,w) \mapsto (\tau_1, \tau_2) \in \mathfrak{t}_G \oplus \mathfrak{t}_G,$$
 (A.53)

as announced.

Remarks:

- 1. We see that for any $(h, w) \in TG$ or $(y, w) \in \mathfrak{tg}$
 - one needs $g \in G/S$ (meaning $g \in G$ modulo elements in S) to conjugate h or y into T_G
 - one needs $v^m \in \mathfrak{m} \subset \mathfrak{g}$ to map $w \mapsto w^{\mathfrak{s}} \in \mathfrak{s}$
 - one then needs $s \in S/T_G$ to conjugate $w^{\mathfrak{s}} \mapsto \tau_2 \in \mathfrak{t}_G$

Thus in total, for any g or y (regardless of regular or not) the parameters (g, v) lying in $TT_G \simeq T_G \times \mathfrak{t}_G \subset TG$ are not needed to accomplish the Abelianisation / diagonalisation.

2. In particular, in the generic regular case (g or y regular), one has $S = T_G$ and $\mathfrak{s} = \mathfrak{t}_G$. Thus the last step in the proof (consisting of conjugating $w^{\mathfrak{s}}$ into $\mathfrak{t}_S = \mathfrak{t}_G$) is empty. The entire ambiguity in the construction then lies in the first step, the conjugation of h into T_G (or y into \mathfrak{t}_G). The possible choices of t or τ_1 are related by the action of the Weyl Group W,

$$W = N_G(T_G)/T_G, \qquad (A.54)$$

where $N_G(T_G)$ is the normaliser of T_G in G.

3. In the opposite extreme case of g = e or y = 0, say, evidently the first step of the argument is empty, the stabiliser of g = e or y = 0 is of course all of G, S = G, and then in the last step there is then a W-fold degeneracy in the choice $\tau_2 \in \mathfrak{t}_G$.

B A symmetry of complex Chern-Simons theory on Seifert 3-manifolds

Here we establish that the symmetry that we need in section 3.4 to pass from having a background field and summing over integers at each orbifold point to having a background field with just one integer summation to perform exists at the level of Complex Chern-Simons theory. BF theory is then just a special limit.⁴ One, possibly surprising, outcome is that the two approaches do not agree at the level of holomorphic factorisation.

 $^{{}^{4}}$ It is straightforward to show that the symmetry is available directly in BF theory but we wanted to establish the more general result here.

We should also point out that these symmetries are a variation of those found by Lawrence and Rozansky [24] and by Mariño [25]. However, here the symmetries have a clear geometric meaning which we explain.

B.1 Discrete symmetries

The exponent in the finite dimensional integral that one arrives at in the case of complex Chern-Simons theory is

$$I(k,s) = -\frac{k}{4\pi} \operatorname{Tr}(\phi^2 - \lambda^2) c_1(\mathcal{L}_M) + k \sum_{i=1}^{N} \frac{1}{a_i} \operatorname{Tr}(-i\phi \mathbf{n}_i + r_i \pi \mathbf{n}_i^2) + k \operatorname{Tr}(-i\phi \mathbf{n}_0) + \frac{s}{2\pi} \operatorname{Tr}\left(\lambda \left(\phi c_1(\mathcal{L}_M) + 2\pi i \sum_{i=1}^{N} \frac{1}{a_i} \mathbf{n}_i + 2\pi i \mathbf{n}_0\right)\right)$$
(B.1)

(just use this instead of the exponent in (4.1)).

The exponential of this action as well as the complex Ray-Singer torsion enjoy a number of symmetries. We will present them here, but before doing that we note that the transformations below do not depend on the coupling constants k and s and so those parts of the action are independently invariant. Also one can see that the k-dependent part of the action (B.1), apart from the λ^2 term, is just what one gets from Chern-Simons theory on M with compact gauge group G. Furthermore, the field λ does not transform, so that the k-dependent part has the same invariance as for the G Chern-Simons theory.

First, we have the symmetry

$$\mathbf{n}_i \to \mathbf{n}_i + a_i \mathbf{m}_i, \ \mathbf{n}_0 \to \mathbf{n}_0 - \sum_{i=1}^N \mathbf{m}_i$$
 (B.2)

It is clear that (B.1) is not changed by these transformations (and it is consistent with our limit on the range of summation over the \mathbf{n}_i).

Remark. Note that geometrically this symmetry is the statement that, as the appropriate powers of the V-line bundle \mathcal{L}_i of degree $1/a_i$ at the *i*'th orbifold point is an honest line bundle

$$\mathcal{L}_i^{\otimes m_i a_i} \simeq \mathcal{L}_0^{\otimes m_i} \tag{B.3}$$

any orbifold line bundle \mathcal{L} satisfies

$$\mathcal{L} = \mathcal{L}_{0}^{\otimes n_{0}} \otimes \mathcal{L}_{1}^{\otimes n_{1}} \otimes \cdots \otimes \mathcal{L}_{N}^{\otimes n_{N}}$$
$$\simeq \mathcal{L}_{0}^{\otimes (n_{0} - \sum_{i} m_{i})} \otimes \mathcal{L}_{1}^{\otimes (n_{1} + a_{1} m_{1})} \otimes \cdots \otimes \mathcal{L}_{N}^{\otimes (n_{N} + a_{N} m_{N})}$$
(B.4)

so one has not changed the bundle but just expressed it in a different way.

Second, we also have the transformations

$$\mathbf{n}_i \to \mathbf{n}_i + b_i \mathbf{v}, \ \phi \to \phi - 2\pi i \mathbf{v}, \ \mathbf{n}_0 \to \mathbf{n}_0 + b_0 \mathbf{v}$$
 (B.5)

which form a symmetry of the exponential of the action.

Remark. Geometrically this says that the pair (\mathcal{L}, ϕ) of an orbifold bundle \mathcal{L} with curvature $\phi d\kappa$ satisfy

$$(\mathcal{L}, \phi) \simeq (\mathcal{L} \otimes \mathcal{L}_M^{\otimes v}, \phi - 2\pi i v)$$
 (B.6)

The first Chern class of $\mathcal{L}_M^{\otimes v}$ is

$$c_1(\mathcal{L}_M^{\otimes v}) = v.c_1(\mathcal{L}_M) \tag{B.7}$$

while

$$\frac{i}{2\pi} \int_{\Sigma_V} -2\pi i v \, d\kappa = v \int_{\Sigma_V} d\kappa = -v \cdot c_1(\mathcal{L}_M) \tag{B.8}$$

It is now possible to combine the symmetries and consider

$$\mathbf{n}_i \to \mathbf{n}_i + a_i \mathbf{u} + b_i \mathbf{v}, \ \phi \to \phi - 2\pi i \mathbf{v}, \ \mathbf{n}_0 \to \mathbf{n}_0 + b_0 \mathbf{v} - N \mathbf{u}$$
 (B.9)

One then has a constructive proof that this symmetry allows one to set all the \mathbf{n}_i for $i \neq 0$ to zero. Recall that if the gcd(a, b) = 1 then any integer n can be expressed as

$$n = au + bv \tag{B.10}$$

for some integers u. Furthermore, if gcd(a, b) = 1 and gcd(a, c) = 1 then gcd(a, bc) = gcd(a, b). gcd(a, c) = 1.

As a first step let $(\mathbf{u}, \mathbf{v}) = (\mathbf{u}_1, \mathbf{v}_1)$ so that

$$\mathbf{n}_1 + a_1 \mathbf{u}_1 + b_1 \mathbf{v}_1 = 0 \tag{B.11}$$

which is guaranteed to have a solution by (B.10). All the other \mathbf{n}_i are changed by this but are brought back into their appropriate ranges by the use of (B.2). Now we want to use (B.9) again to set $\mathbf{n}_2 = 0$ while keeping $\mathbf{n}_1 = 0$. If the transformation \mathbf{v} is proportional to a_1

$$\mathbf{n}_{1} \rightarrow 0 + a_{1}\mathbf{u}_{2} + b_{1}a_{1}\mathbf{v}_{2}$$

$$\mathbf{n}_{2} \rightarrow \mathbf{n}_{2} + a_{2}\mathbf{u}_{2} + b_{2}a_{1}\mathbf{v}_{2}$$

$$\mathbf{n}_{3} \rightarrow \mathbf{n}_{3} + a_{3}\mathbf{u}_{2} + b_{3}a_{1}\mathbf{v}_{2}$$

$$\dots \dots \dots$$

$$\mathbf{n}_{N} \rightarrow \mathbf{n}_{N} + a_{N}\mathbf{u}_{2} + b_{N}a_{1}\mathbf{v}_{2}$$
(B.12)

then the change in \mathbf{n}_1 is proportional to a_1 and by (B.2) is zero. We note that $gcd(a_2, b_2a_1) = 1$ so by (B.10) we can choose $(\mathbf{u}_2, \mathbf{v}_2)$ so the \mathbf{n}_2 maps to zero. Now we would like to set $\mathbf{n}_3 = 0$ without changing \mathbf{n}_1 and \mathbf{n}_2 . To not change \mathbf{n}_1 the vector \mathbf{v} must be proportional to a_1 , as we saw before, and not to change \mathbf{n}_2 it must also be proportional to a_2 so we perform a transformation

$$\mathbf{n}_{1} \rightarrow 0 + a_{1}\mathbf{u}_{3} + b_{1}a_{1}a_{2}\mathbf{v}_{3} \simeq 0$$

$$\mathbf{n}_{2} \rightarrow 0 + a_{2}\mathbf{u}_{3} + b_{2}a_{1}a_{2}\mathbf{v}_{3} \simeq 0$$

$$\mathbf{n}_{3} \rightarrow \mathbf{n}_{3} + a_{3}\mathbf{u}_{3} + b_{3}a_{1}a_{2}\mathbf{v}_{3}$$

$$\dots \dots \dots \dots$$

$$\mathbf{n}_{N} \rightarrow \mathbf{n}_{N} + a_{N}\mathbf{u}_{2} + b_{N}a_{1}a_{2}\mathbf{v}_{2}$$
(B.13)

Now $gcd(a_3, b_3a_1a_2) = 1$ so we can set $\mathbf{n}_3 = 0$ by a suitable choice of $(\mathbf{u}_3, \mathbf{v}_3)$.

Clearly the procedure can be repeated till we have set all the \mathbf{n}_i to zero and the parameters we use are

$$\mathbf{u} = \sum_{i=1}^{N} \mathbf{u}_i, \quad \mathbf{v} = \mathbf{v}_1 + a_1 \mathbf{v}_2 + \dots + a_1 \dots a_{N-1} \mathbf{v}_N \tag{B.14}$$

B.2 Holomorphic factorisation

The finite dimensional action for complex Chern-Simons theory (B.1) can be expressed in terms of complex fields $\phi + i\lambda$ and a complex coupling constant t = k + is as

$$I(t,\bar{t}) = -\frac{t}{8\pi}c_1(\mathcal{L}_M)\operatorname{Tr}\Phi^2 - i\frac{t}{2}\operatorname{Tr}(\Phi\widehat{\mathbf{q}}) + \frac{t\pi}{2}\sum_{i=1}^N\operatorname{Tr}(\mathbf{n}_i^2)$$

$$-\frac{\bar{t}}{8\pi}c_1(\mathcal{L}_M)\operatorname{Tr}\bar{\Phi}^2 + i\frac{\bar{t}}{2}\operatorname{Tr}(\bar{\Phi}\widehat{\mathbf{q}}) + \frac{\bar{t}\pi}{2}\sum_{i=1}^N\operatorname{Tr}(\mathbf{n}_i^2)$$
(B.15)

Now we note that the holomorphic part of the action

$$I(t) = -\frac{t}{8\pi}c_1(\mathcal{L}_M)\operatorname{Tr}\Phi^2 - i\frac{t}{2}\operatorname{Tr}(\Phi\widehat{\mathbf{q}}) + \frac{t\pi}{2}\sum_{i=1}^{N}\operatorname{Tr}(\mathbf{n}_i^2)$$
(B.16)

is not invariant under the first symmetry namely under

$$\mathbf{n}_i \to \mathbf{n}_i + a_i \mathbf{m}_i, \ \mathbf{n}_0 \to \mathbf{n}_0 - \sum_{i=1}^N \mathbf{m}_i$$
 (B.17)

even though both Φ and $\hat{\mathbf{q}}$ are invariant. The quadratic term \mathbf{n}_i^2 is not invariant. Furthermore, the exponential $\exp(iI(t))$ is also not invariant, unless s = 0, as the coupling constant is complex and so one does not just get a phase.

This implies that we can have two inequivalent holomorphic factorizations. The first is to take the partition function with all the \mathbf{n}_i switched on and then factorise with holomorphic action (B.16) while the second is to set all the $\mathbf{n}_i = 0$ for $i \neq 0$ and then get the factorisation in [17].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- M. Blau, M. Kakona and G. Thompson, Massive Ray-Singer torsion and path integrals, JHEP 08 (2022) 230 [arXiv:2206.12268] [INSPIRE].
- [2] A.S. Schwarz, The Partition Function of Degenerate Quadratic Functional and Ray-Singer Invariants, Lett. Math. Phys. 2 (1978) 247 [INSPIRE].
- [3] A.S. Schwarz, The Partition Function of a Degenerate Functional, Commun. Math. Phys. 67 (1979) 1 [INSPIRE].

- [4] A.S. Schwarz and Y.S. Tyupkin, Quantization of Antisymmetric Tensors and Ray-Singer Torsion, Nucl. Phys. B 242 (1984) 436 [INSPIRE].
- [5], R Torsion and the Laplacian on Riemannian manifolds, Adv. Math. 7 (1971) 145 [INSPIRE].
- [6] D. Ray and I. Singer, Analytic Torsion, Partial Differential Equations PSPM 23 (1971) 167.
- [7] M. Blau and G. Thompson, A New Class of Topological Field Theories and the Ray-singer Torsion, Phys. Lett. B 228 (1989) 64 [INSPIRE].
- [8] G. Horowitz, Exactly Soluble Diffeomorphism Invariant Theories Commun. Math. Phys. 125 (1989) 417.
- M. Blau and G. Thompson, Topological Gauge Theories of Antisymmetric Tensor Fields, Annals Phys. 205 (1991) 130 [INSPIRE].
- [10] M. Blau and G. Thompson, Do metric independent classical actions lead to topological field theories?, Phys. Lett. B 255 (1991) 535 [INSPIRE].
- [11] E. Witten, Topology Changing Amplitudes in (2+1)-Dimensional Gravity, Nucl. Phys. B 323 (1989) 113 [INSPIRE].
- [12] E. Witten, Quantum Field Theory and the Jones Polynomial, Commun. Math. Phys. 121 (1989) 351 [INSPIRE].
- [13] M. Blau and G. Thompson, Derivation of the Verlinde formula from Chern-Simons theory and the G/G model, Nucl. Phys. B 408 (1993) 345 [hep-th/9305010] [INSPIRE].
- [14] M. Blau and G. Thompson, Lectures on 2D gauge theories. Topological aspects and path integral techniques, hep-th/9310144 [DOI:10.48550/arXiv.hep-th/9310144].
- [15] M. Blau and G. Thompson, Chern-Simons theory on S1-bundles: Abelianisation and q-deformed Yang-Mills theory, JHEP 05 (2006) 003 [hep-th/0601068] [INSPIRE].
- [16] M. Blau and G. Thompson, Chern-Simons Theory on Seifert 3-Manifolds, JHEP 09 (2013)
 033 [arXiv:1306.3381] [INSPIRE].
- [17] M. Blau and G. Thompson, Chern-Simons Theory with Complex Gauge Group on Seifert Fibred 3-Manifolds, arXiv:1603.01149 [INSPIRE].
- [18] M. Blau, K.M. Keita, K.S. Narain and G. Thompson, Chern-Simons theory on a general Seifert 3-manifold, Adv. Theor. Math. Phys. 24 (2020) 279 [arXiv:1812.10966] [INSPIRE].
- [19] D.S. Freed and R.E. Gompf, Computer calculation of Witten's three manifold invariant, Commun. Math. Phys. 141 (1991) 79 [INSPIRE].
- [20] E. Witten, Quantization of Chern-Simons Gauge Theory With Complex Gauge Group, Commun. Math. Phys. 137 (1991) 29 [INSPIRE].
- [21] S. Gukov, Three-dimensional quantum gravity, Chern-Simons theory, and the A polynomial, Commun. Math. Phys. 255 (2005) 577 [hep-th/0306165] [INSPIRE].
- [22] T. Dimofte, S. Gukov, J. Lenells and D. Zagier, Exact Results for Perturbative Chern-Simons Theory with Complex Gauge Group, Commun. Num. Theor. Phys. 3 (2009) 363 [arXiv:0903.2472] [INSPIRE].
- [23] S. Gukov and D. Pei, Equivariant Verlinde formula from fivebranes and vortices, Commun. Math. Phys. 355 (2017) 1 [arXiv:1501.01310] [INSPIRE].
- [24] R. Lawrence and L. Rozansky, Witten-Reshetikhin-Turaev Invariants of Seifert Manifolds, Commun. Math. Phys. 205 (1999) 287.

- [25] M. Marino, Chern-Simons theory, matrix integrals, and perturbative three manifold invariants, Commun. Math. Phys. 253 (2004) 25 [hep-th/0207096] [INSPIRE].
- [26] E. Witten, Analytic Continuation Of Chern-Simons Theory, AMS/IP Stud. Adv. Math. 50 (2011) 347 [arXiv:1001.2933] [INSPIRE].
- [27] S. Gukov, M. Marino and P. Putrov, *Resurgence in complex Chern-Simons theory*, arXiv:1605.07615 [INSPIRE].
- [28] D. Bar-Natan and E. Witten, Perturbative expansion of Chern-Simons theory with noncompact gauge group, Commun. Math. Phys. 141 (1991) 423 [INSPIRE].
- [29] D. Fried, Lefschetz Formulas for Flows, in The Lefshetz Centennial Conference, Part III, Contemp. Math. 58 (1987) 19.
- [30] K. Furuta and B. Steer Seifert Fibred Homology 3-Spheres and the Yang-Mills Equations on Riemann Surfaces with Marked Points, Adv. Math. 96 (1992) 38.
- [31] M. Blau and G. Thompson, On diagonalization in map(M,G), Commun. Math. Phys. 171 (1995) 639 [hep-th/9402097] [INSPIRE].
- [32] D. Freed, Reidemeister Torsion, Spectral Sequences, and Brieskorn Spheres, J. Reine Angew. Math. 429 (1992) 75.
- [33] D. Ray, Reidemeister Torsion and the Laplacian on Lens Spaces, Adv. Math. 4 (1970) 109.
- [34] L.C. Jeffrey, Chern-Simons-Witten invariants of lens spaces and torus bundles, and the semiclassical approximation, Commun. Math. Phys. 147 (1992) 563 [INSPIRE].
- [35] P. Orlik, Seifert Manifolds, LNM 291, Springer-Verlag Berlin, Germany (1972).
- [36] P. Kirk and K. Klassen, Chern-Simons Invariants of 3-Manifolds and Representation Spaces of Knot Groups, Math. Ann. 287 (1990) 343.
- [37] D.R. Auckly, Topological methods to compute Chern-Simons invariants, Math. Proc. Cambridge Phil. Soc. 115 (1994) 229.
- [38] H. Nishi, SU(n)-Chern-Simons Invariants of Seifert Fibred 3-Manifolds, Int. J. Math. 9 (1998) 295.
- [39] R. Fintushel and R. Stern, Instanton Homology of Seifert Fibred Homology Three Spheres Proc. Lond. Math. Soc. 61 (1990) 107.
- [40] T. Kitano and Y. Yamaguchi, SL(2, R)-representations of a Brieskorn homology 3-sphere, arXiv:1602.07503.
- [41] L. Rozansky, A large k asymptotics of Witten's invariant of Seifert manifolds, Commun. Math. Phys. 171 (1995) 279 [hep-th/9303099] [INSPIRE].
- [42] E. Witten, On quantum gauge theories in two-dimensions, Commun. Math. Phys. 141 (1991) 153 [INSPIRE].