



Influence of Cooperation Work on Management Continuity of Ambulatory Cardiovascular Care: A Cross-Sectional Exploratory Study in Germany

RESEARCH AND
THEORY

ubiquity press

CHRISTINE ARNOLD 

PATRICK HENNRICH

FRANK PETERS-KLIMM

MICHEL WENSING

*Author affiliations can be found in the back matter of this article

ABSTRACT

Introduction: A wide range of factors influence coordination and continuity of care. The aim of this study was to explore how management continuity of cardiovascular-related ambulatory care is influenced by the following network characteristics: presence of a case coordinator, network reciprocity, network composition and team climate.

Methods: This cross-sectional observational study included three written surveys. The primary outcome management continuity of cardiovascular care was measured with the team/cross-boundary scale in the Nijmegen Continuity Questionnaire. The final analysis comprised a multivariate linear multilevel model with the predictors: presence of a case coordinator, network reciprocity, network composition and team climate.

Results: Eighteen general practices with 83 health workers and 340 patients participated. The linear multilevel regression analysis showed a positive influence of team climate on cross-boundary continuity of care (b-coefficient 0.44, 95% confidence interval 0.09–0.78, $p = 0.02$). No statistically significant influence was measured for the other predictors.

Discussion: To improve integrated care, therefore, emphasis should also be placed on promoting the team climate within individual practices. Regarding network characteristics, further research is needed, especially in larger practices.

Conclusion: This study showed that team climate had an independent, relevant and statistically significant association with cross-boundary continuity of cardiovascular ambulatory care.

CORRESPONDING AUTHOR:

Christine Arnold

Department of General
Practice and Health Services
Research, Heidelberg
University Hospital, Germany;
Division of Neonatology,
Department of Paediatrics,
Inselspital, Bern University
Hospital, University of Bern,
Switzerland

christine.arnold@insel.ch

KEYWORDS:

integrated care; continuity
of care; ambulatory
cardiovascular care; team
climate

TO CITE THIS ARTICLE:

Arnold C, Hennrich P, Peters-
Klimm F, Wensing M. Influence
of Cooperation Work on
Management Continuity of
Ambulatory Cardiovascular
Care: A Cross-Sectional
Exploratory Study in Germany.
*International Journal of
Integrated Care*, 2023; 23(4):
14, 1–8. DOI: [https://doi.
org/10.5334/ijic.7019](https://doi.org/10.5334/ijic.7019)

INTRODUCTION

Cardiovascular diseases continue to have a high prevalence and are one of the leading causes of death worldwide. With the increase in multimorbidity, the care of patients with cardiovascular diseases is becoming more complex and requires the integration of various health workers to achieve optimal patient care [1, 2]. Optimal patient care requires high continuity of care, which is a central characteristic of effective integrated care models. Integrated care is structured healthcare delivery to a specific population (e.g. coronary heart disease patients), involving various healthcare providers and informed by the best available evidence [3]. Higher continuity of care is associated with lower mortality rates [4, 5], lowered hospitalisations [6, 7] and increased feelings of security and confidence for the patients [8, 9].

Continuity of care has been defined by Haggerty et al. [10] as a three-dimensional concept: informational continuity (information sharing between health workers); relational continuity (trusting relationships with health workers); and management continuity (if multiple health workers are involved in care, their approach is consistent with one another's and congruent with patient needs) [10, 11].

The focus in this study is on management continuity in ambulatory cardiovascular care, which is challenging in healthcare systems that lack clinical guidance that is shared across healthcare providers. Many factors influence provider's views on clinical management, including vocational training and socialisation, the influence of peers, and other considerations. For example, a previous study with claims data regarding physicians-networks showed that the prescription of a new medication for heart failure is influenced by network structures. When physicians were connected to other physicians through common patients who were already prescribing the new medication, the likelihood of prescribing the new drug increased [12]. However, little is known about the views of healthcare providers are influenced by the teams and networks in which they are embedded. For example, an interview study of hospital admissions and discharges suggests that personal relationships among healthcare providers in different sectors promote alignment of practices [13].

In this study, we focused on the role of ambulatory care providers' interaction networks. Based on empirical and theoretical work, we developed a conceptual model to explain the impact of professional interaction networks of health workers on care delivery and healthcare outcomes. Health workers conduct various activities, including diagnosis, counselling, treatment and prevention, and these activities must be coordinated. We hypothesized that coordination of care is influenced by the following factors: presence of a case

coordinator (typically a generalist physician or nurse [14]), network reciprocity, team climate, and change of network composition. Together these factors influence cooperation work, which covers various processes such as convincement, competition, use of power, and selection of network members.

Over time, network reciprocity within interactions between network members increases mutual trust and reduces risk of defecting behaviours. Research in evolutionary biology has shown that, over time, network reciprocity is crucial for the emergence of altruistic cooperation and that it may even counterbalance short-term individual benefits of non-cooperation behaviours [15]. Under certain conditions, this also applies to anticipated rather than experienced reciprocity.

A moderate change of network composition as opposed to no change makes it possible to select co-operators and unselect non-co-operators, while maintaining the favourable effects of repeated interactions on cooperation [15]. This change is likely to influence the structure of referral and advice networks between primary care and other health professionals. It can also contribute to the homogeneity of network members.

In addition to these structural network factors, a good climate and team culture in teams can enhance management continuity as well as experience of healthcare providers [16]. According to Mundt et al. [17], a shared team vision prevails in small centralised networks, which are associated with higher quality of care and lower costs.

The aim of this study was to explore how management continuity of cardiovascular-related ambulatory care is influenced by the following network characteristics: presence of a case coordinator, network reciprocity, network composition and team climate.

METHODS

The study was conducted in accordance with the Declaration of Helsinki and received approval from the Ethics Committee of the Medical Faculty of Heidelberg (ID: S-726/2018) and from the respective State Medical Chambers. Due to the anonymity of the patient survey, the Ethics Committee of the Medical Faculty of Heidelberg approved a waiver for informed consent. Additionally, participants were informed about this waiver in writing and that returning the questionnaire was sufficient. We registered the study prospectively on 07/11/2019 at the German Clinical Trials Register under the ID: DRKS00019219. The STROBE reporting guideline [18] for observational studies was followed in the reporting of this study.

STUDY DESIGN AND STUDY POPULATION

This cross-sectional observational study explored the influence of coordination work on management continuity of cardiovascular care in German primary care and consisted of three written surveys. The three-year (2019–2022) ExKoCare project aimed to recruit a sample of 40 general practices in the German states of Baden-Wuerttemberg (approximately 11 million inhabitants; sampling took place in 10 of the 44 counties), Rhineland-Palatinate (approximately 4 million inhabitants; sampling in 13 of the 36 counties), and Saarland (approximately 1 million inhabitants; sampling in 2 of the 6 counties) [19]. The study included three samples: general practices; their physicians and non-physician health workers; and patients from these general practices with at least three recorded chronic diseases, one of which was coronary heart disease.

DATA COLLECTION

General practices

We aimed to recruit 40 general practices and anticipated a low participation rate of 5%. The general practices were recruited from a clustered, stratified sample from a total of 25 counties. The counties were chosen with regard to population density to ensure that rural and urban areas were equally represented. We aimed to contact each general practice via fax or email. The owners of each general practice were identified through the publicly available online physician's databases of the three states. This led to an initial sample of 1,617 practices (Baden-Wuerttemberg, 912 practices; Rhineland-Palatinate, 596 practices; Saarland, 109 practices) [20]. Some basic characteristics of the practices were documented in a practice questionnaire completed by the practice owner. These characteristics included the size of the practice, as measured by the number of cases per quarter, and the participation in case-management or disease-management programmes.

Health workers

After the practice owners consented to the ExKoCare study, all health workers over the age of 18 of the general practices were contacted personally in writing and invited ($n = 208$) to participate in the survey. Data were collected using a written pseudonymized questionnaire. All participants gave written informed consent for the study.

Patients

After the health worker survey was completed, all participating practices were contacted and invited to support patient recruitment. Initially, the aim was to recruit 15 patients from each general practice ($n = 600$). Based on previous research, we expected a response rate of 30%, and so we intended to invite 50 patients per practice. The research team assisted the practices

via phone in identifying potential study participants to ensure that the inclusion criteria (adult patients with three or more chronic conditions, including at least one atherosclerosis-related cardiovascular condition, only adult patients over the age of 18 are eligible to participate in this study, ability to consent to participate) were met. A list of potential participants was compiled from the physicians' billing systems. This system lists all patients who have been billed for services, regardless of whether they regularly visit this doctor. Then, up to 50 patients were selected by selecting every 3rd patient from a starting point specified by the researcher. If there were fewer than 50 patients on the list, all were selected. Before this selection, the physician was asked to check for the cognitive ability to complete a questionnaire and potential contra-indications. Patients were sent the questionnaire by post, and an anonymous return envelope addressed to the research department was included, so that the practices did not know which patients participated.

MEASURES

For each stage of the study, we used a newly developed questionnaire that included both validated and newly created parts derived from previous studies and guidelines. The three questionnaires used a pseudonym for each practice, so that the results could be assigned accordingly.

Outcome

The outcome of the study was management continuity of cardiovascular care, which was measured from the patient's perspective with the team/cross-boundary scale in the Nijmegen Continuity Questionnaire (NCQ) [21, 22]. The NCQ was developed in the Netherlands and has been validated in various countries and applied in different settings (e.g., in Norway in the field of rehabilitation) [23]. The 12-item NCQ includes the following subscales: personal continuity-1 ('care provider knows me', items 1–5), personal continuity-2 ('care provider shows commitment', items 6–8), and team/cross-boundary continuity (items 9–12). The items 9–12 include the following statements: "These care providers transfer information very well to each other.", "These care providers work together very well.", "The care of these care providers is very well connected." and "These care providers always know very well from each other what they do." In the ExKoCare project, the two subscales for personal continuity were used to assess the continuity regarding the family doctor and cardiologist, the team part of the team/cross-boundary subscale was used to assess the continuity in the general practice (continuity between physicians and practice assistants) and the cross-boundary part was used to assess the continuity between family doctor and cardiologist. Thus, the questionnaire contained six continuity of care scores and

a total of 24 items. Each question could be answered on a 5-point Likert-scale (1 = *strongly disagree*, 2 = *disagree*, 3 = *neutral*, 4 = *agree* to 5 = *strongly agree*) or with *I do not know*.

The English version of all 12 items in the NCQ was translated carefully and independently by CA and PH using a forward translation into German and a backwards translation into English. After the independent translations, consensus discussions were held and a common German version was produced. A test involving interviews with six patients did not identify any significant lack of clarity [24]. For this part of the project, we used only the scales for team and cross-boundary continuity, which served as the outcomes of the study. For the calculation of the two outcomes, one missing value was allowed in each case.

Predictors

Presence of care coordinator was measured in the practice questionnaire using a dichotomous item (yes/no) that indicated the presence of a case manager for cardiovascular care.

Reciprocity indicates the percentage of returned relationships in a directed network. The value range is between 0 and 1. The network was constructed based on the questionnaire in which the health workers were asked about the weekly exchange of information within their general practice. They were asked to mark the persons with whom they exchange information weekly [25].

Change of network composition was measured on a 5-point Likert scale, with 1 indicating little change in the last two years regarding the collaboration with cardiologists outside the health worker's own general practice and 5 indicating a high level of change. This predictor was also used to measure network changes within the general practice and among health workers outside the general practice.

Team climate within the general practice was measured with the short version of the Team Climate Inventory scale [26]. This questionnaire contains 14 items and four subscales: 1) clear and realistic objectives to which the team members are committed (called 'shared vision', 4 items); 2) interaction between team members (called 'psychological safety', 4 items); 3) commitment to high standards of performance and appraisal of weakness (called 'task orientation', 3 items); and 4) support for innovation (called 'innovation support', 3 items). The responses were given on a 5-point Likert scale from 1 = *disagree* to 5 = *agree*. We translated the questionnaire into German using the forward-backward method, which was completed in each part by two individuals. For each participant, the mean of the 14 items was calculated. One missing value was permitted. The mean value for each practice was then determined. Higher scores indicated a better team climate. Thus, a

mean with a standard deviation for each practice was available with a possible range of 1 to 5.

Potential confounders

We included *network size* that indicates the number of health workers in each general practice. In addition, the number of physicians and non-physician professionals was measured. From the patient questionnaire, we included the number of chronic diseases (from a list of ten).

STATISTICAL ANALYSES

First, we conducted a descriptive analysis of the study population. According to the measurement level, relative and absolute frequencies and mean values with SDs were calculated. Next, descriptive statistics were calculated for the predictors, and then the patient-reported outcome and team and cross-boundary continuity of care with means and SDs were calculated. The final analysis comprised univariate and multivariate linear multilevel regression analyses. Before this, the independent variables were tested for multicollinearity, which was assumed when the correlation coefficient was larger than 0.6. The two final models included the following predictors: presence of care coordinator (yes or no), reciprocity (range 0–1), change of network composition (range 1–5) and team climate (range 1–5), which should express the cooperation work. Additionally, the models were adjusted for network size and number of chronic diseases. In the first model, the dependent variable was the cross-boundary continuity of care between general practice and cardiologist practice; in the second model, it was team continuity of care within the general practice. Based on the nested data in GP practices, the interclass correlation coefficient was calculated to compute the proportion of variance explained by the GP practice in the total variance. All analyses were performed with the statistics software R (version 4.0.2) using RStudio (version 1.2.5033). The significance level was set at an alpha of 0.05.

RESULTS

Data were collected from November 2019 to December 2021 (health workers survey from November 2019 and January 2021 and patients from November 2020 to December 2021), that was predominantly during the COVID-19 pandemic. After exclusion due to unavailable fax numbers or incorrect deliveries, we contacted 1,511 family practices, from which 42 took part in the ExKoCare project (response rate 2.8%). Eighteen general practices had collected all data for this study and were included in the present analysis. Due to the increased workload during the COVID-19 pandemic, the data from the other practices ($n = 24$) were not complete, which led to their exclusion from

this analysis. From the 18 practices, a total of 93 health workers and 596 patients were invited to participate in the study. With response rates of 89.3% and 57.0%, respectively, 83 health workers and 340 patients were included in the study. Table 1 provides an overview of the characteristics of the participating practices and patients. Over 75% of the practices were single-handed practices with an average of 1.4 physicians and 3.8 non-physician health workers. The patients were on average 74.5 years old (range 45–94), and 247 (72.9%) participants were male.

The practices examined had an average network size of 5.2 (SD 1.2). In addition to the measured network characteristics (Table 2), the team climate score across all practices was on average 4.2 (SD 0.4). With a maximum value of 1, the average network reciprocity was 0.6 (SD 0.3). Patients (n = 231) reported cross-boundary continuity of care between family doctor and cardiologist with a mean of 3.8 (SD 0.8) and team continuity of care within the general practice (n = 270) with a mean of 4.0 (SD 0.7).

PRACTICE CHARACTERISTICS	N = 18
Type of practice, n (%)	
Not single-handed practice	4 (22.2)
Single-handed practice	14 (77.8)
Total number of health workers, mean (SD)	5.2 (1.2)
Physicians, mean (SD)	1.4 (0.5)
Non-physicians, mean (SD)	3.8 (1.0)
Cases per quarter, n (%)	
<500 cases	0 (0.0)
500–1,000 cases	7 (38.9)
1,001–1,500 cases	7 (38.9)
>1,500 cases	4 (22.2)
PATIENT CHARACTERISTICS	N = 340
Continuity of care	
Team, mean (SD)	4.0 (0.7)
Cross-boundary, mean (SD)	3.8 (0.8)
Age, mean (SD), n = 316	74.5 (9.1)
Sex, n (%), n = 339	
Female	92 (27.1)
Male	247 (72.9)
Number of chronic diseases, mean (SD),	1.6 (1.5)
Participation in DMP, n (%)	76 (24.8)

Table 1 Practice and patient characteristics.
SD = Standard deviation.
DMP = Disease management programme.

NETWORK CHARACTERISTICS	MEAN (SD)
	N = 18
Presence of case coordinator, n (%)	n = 17
No	12 (70.6)
Yes	5 (29.4)
Network reciprocity	0.6 (0.3)
Change of network composition	
Changes outside the general practice	1.7 (0.7)
Changes inside the general practice	2.2 (0.9)
Changes cardiology practice	1.6 (0.6)

Table 2 Descriptive data of network characteristics.

In the univariate regression analyses, only team climate and the number of chronic conditions showed a statistically significant impact. Therefore, we excluded the confounders age and sex from the final model. The linear multilevel regression analysis showed a positive influence of team climate on cross-boundary continuity of care (b-coefficient 0.44, 95% CI 0.09; 0.78, p = 0.02). No statistically significant influence was measured for the other predictors (see Table 3). The interclass correlation coefficient for clustering within practices was 0.02. In addition, the predictors in the model with the outcome team continuity of care within general practice were not statistically significant.

DISCUSSION

This exploratory study found a positive influence of the team climate within general practices on patient-reported cross-boundary continuity of cardiovascular ambulatory care. However, no effect of the hypothesized network mechanisms was found. In addition, team climate did not have a statistically significant impact on team continuity of care within the general practice. It may also be noted that German primary care practices are often small, so a practice typically has one team.

Previous studies have shown that team climate positively influences interprofessional collaboration, which is important for managing continuity of care [27]. While within the practice it may be possible to compensate for a lower team climate, it seems to be even more important that there is a good team climate within the practice in order to achieve cross-boundary continuity of care, which may explain the different results in the two measured team and cross-boundary continuity of care scores.

Outside of the practice, various arrangements must be made in which tasks may not be precisely assigned to practice members, making coordination more difficult,

	B-COEFFICIENT	SE	95% CI	P-VALUE
Presence of care coordinator (ref. no)	0.08	0.11	-0.16; 0.32	0.50
Reciprocity	-0.21	0.20	-0.65; 0.22	0.31
Change of network composition	-0.05	0.08	-0.23; 0.13	0.56
Team climate	0.44	0.16	0.09; 0.78	0.02*
Intercept	2.07	0.73	0.63; 3.51	<0.01

Table 3 Influence of cooperation work on cross-boundary continuity of cardiovascular care.

n = 231 patients and 17 general practices.

Model adjusted for network size and number of chronic diseases.

* Statistically significant at $\alpha = 0.05$.

ref. = reference group, CI = confidence interval, SE = standard error.

possibly leading to fewer agreements if the team climate is poorer and, ultimately, reducing the continuity of care.

Promoting the team climate in individual practices could therefore also contribute to improving integrated care. However, we are not aware of any studies that have examined the relationship between team climate and continuity of care in ambulatory practices.

The network mechanisms, which influence cooperation work, studied provided a first indication of influence on cross-boundary continuity of care, but the influence was not statistically significant.

LIMITATIONS

The study has limitations in terms of unknown measurement validity of several predictors and generalisability due to sampling procedures, so the descriptive figures should be carefully interpreted despite the use of a validated outcome measure. As the data collection took place during the corona pandemic, this could be of influence on the nature of the collected data and the low response rate. It follows that the findings should be interpreted as preliminary.

A weakness of the study is the relatively high non-response in the team/cross-boundary continuity of care items. For example, of the participating patients, only 231 (68%) completed the questions about cross-boundary continuity of cardiovascular care. A further issue is that only few confounders could be included in the analysis due to the sample size. It is therefore important to conduct further research with larger samples to demonstrate effects of network mechanisms on coordination work.

The findings suggest, that especially in smaller single-handed practices the network structures play a subordinate role compared to the team functioning, since all interact frequently and there is little explicit coordination. It is possible that we would have been more likely to find a statistically significant effect of network mechanisms in a sample with larger practices.

A specific issue is that single-handed practices constituted almost 80% of the sample of practices. Although Germany has a high proportion (60%) of individual practices [28], group practices are increasing due to rising costs for individual practices, and so increasing numbers of practice rooms and staff are shared. The number of practice members is expected to increase, and the structures of the practices will become more important.

Another limitation is that the team climate questionnaire was not explicitly validated for the setting due to low response rate.

CONCLUSION

This exploratory study showed that team climate had an independent, relevant and statistically significant association with cross-boundary continuity of cardiovascular ambulatory care. The hypotheses regarding network structures were not confirmed, but it can be supposed that the structures and division of tasks in practices are becoming increasingly important due to the expansion resulting from mergers of single-handed practices.

ACKNOWLEDGEMENTS

We would like to thank all patients for participating in our study, all GP practices for the support in the recruitment of the patients and our research assistant Pia Traulsen for her support of data entry and study administration.

REVIEWERS

Two anonymous reviewers.


FUNDING INFORMATION

This work was funded by the Deutsche Forschungsgemeinschaft (DFG) under grant number 416396249. The funder had no role in the design of the study, the collection, analysis and interpretation of the data, or the writing of the manuscript. The study protocol was independently reviewed by the funder.

COMPETING INTERESTS

The authors have no competing interests to declare.

AUTHOR AFFILIATIONS

Christine Arnold  orcid.org/0000-0003-4201-1800
Department of General Practice and Health Services Research, Heidelberg University Hospital, Germany; Division of Neonatology, Department of Paediatrics, Inselspital, Bern University Hospital, University of Bern, Switzerland

Patrick Henrich
Department of General Practice and Health Services Research, Heidelberg University Hospital, Germany

Frank Peters-Klimm
Department of General Practice and Health Services Research, Heidelberg University Hospital, Germany

Michel Wensing
Department of General Practice and Health Services Research, Heidelberg University Hospital, Germany

REFERENCES

- Townsend N, Wilson L, Bhatnagar P, Wickramasinghe K, Rayner M, Nichols M.** Cardiovascular disease in Europe: epidemiological update 2016. *Eur Heart J.* 2016; 37(42): 3232–45. DOI: <https://doi.org/10.1093/eurheartj/ehw334>
- Buddeke J, Bots ML, van Dis I, Visseren FL, Hollander M, Schellevis FG,** et al. Comorbidity in patients with cardiovascular disease in primary care: a cohort study with routine healthcare data. *Br J Gen Pract.* 2019; 69(683): e398–e406. DOI: <https://doi.org/10.3399/bjgp19X702725>
- World Health Organization.** Integrated care models: an overview Copenhagen; 2016. Available from: https://www.euro.who.int/__data/assets/pdf_file/0005/322475/Integrated-care-models-overview.pdf. Download (22.07.2022).
- Pereira Gray DJ, Sidaway-Lee K, White E, Thorne A, Evans PH.** Continuity of care with doctors- a matter of life and death? A systematic review of continuity of care and mortality. *BMJ Open.* 2018; 8: e021161. DOI: <https://doi.org/10.1136/bmjopen-2017-021161>
- Baker R, Freeman GK, Haggerty JL, Bankart MJ, Nockels KH.** Primary medical care continuity and patient mortality: a systematic review. *Br J Gen Pract.* 2020 Aug 27; 70(698): e600–e611. DOI: <https://doi.org/10.3399/bjgp20X712289>
- Barker I, Steventon A, Deeny SR.** Association between continuity of care in general practice and hospital admissions for ambulatory sensitive conditions: cross-sectional study of routinely collected, person level data. *BMJ.* 2017; 356: j84. DOI: <https://doi.org/10.1136/bmj.j84>
- Bazemore A, Petterson S, Peterson LE, Bruno R, Chung Y, Phillips RL.** Higher general practitioner continuity is associated with lower costs and hospitalizations. *Ann Fam Med.* 2018; 16: 492–7. DOI: <https://doi.org/10.1370/afm.2308>
- Haggerty JL, Roberge D, Freeman GK, Beaulieu C.** Experienced continuity of care when patients see multiple clinicians: a qualitative meta summary. *Ann Fam Med.* 2013; 11: 262–71. DOI: <https://doi.org/10.1370/afm.1499>
- Wilfling D, Warkentin N, Laag S, Goetz K.** “I have such a great care” – geriatric patients’ experiences with a new healthcare model: a qualitative study. *Patient Prefer Adherence.* 2021; 15: 309–15. DOI: <https://doi.org/10.2147/PPA.S296204>
- Haggerty JL, Reid RJ, Freeman GK, Starfield BH, Adair CE, McKendry R.** Continuity of care: a multidisciplinary review. *BMJ.* 2003; 327(7425): 1219–21. DOI: <https://doi.org/10.1136/bmj.327.7425.1219>
- Freeman GK, Olesen F, Hjortdahl P.** Continuity of care: an essential element of modern general practice? *Fam Pract.* 2003; 20(6): 623–7. DOI: <https://doi.org/10.1093/fampra/cm601>
- Arnold C, Koetsenruijter J, Forstner J, Peters-Klimm F, Wensing M.** Influence of physician networks on prescribing a new ingredient combination in heart failure: a longitudinal claim data-based study. *Implement Sci.* 2021 Aug 28; 16(1): 84. DOI: <https://doi.org/10.1186/s13012-021-01150-y>
- Forstner J, Bossert J, Weis A, Litke N, Strassner C, Szecsenyi J, Wensing M.** The role of personalised professional relations across care sectors in achieving high continuity of care. *BMC Fam Pract.* 2021 Apr 14; 22(1): 72. DOI: <https://doi.org/10.1186/s12875-021-01418-8>
- Uijen AA, Schers HJ, Schellevis FG, van den Bosch WJ.** How unique is continuity of care? A review of continuity and related concepts. *Fam Pract.* 2012; 29(3): 264–71. DOI: <https://doi.org/10.1093/fampra/cm104>
- Nowak MA.** Five rules for the evolution of cooperation. *Science.* 2006; 314(5805): 1560–3. DOI: <https://doi.org/10.1126/science.1133755>
- Espinoza P, Peduzzi M, Agreli HF, Sutherland MA.** Interprofessional team member’s satisfaction: a mixed methods study of a Chilean hospital. *Hum Resour Health.* 2018; 16(1): 30. DOI: <https://doi.org/10.1186/s12960-018-0290-z>
- Mundt MP, Gilchrist VJ, Fleming MF, Zakletskaia LI, Tuan W-J, Beasley JW.** Effects of primary care team social networks on quality of care and costs for patients with cardiovascular disease. *Ann Fam Med.* 2015; 13(2): 139–48. DOI: <https://doi.org/10.1370/afm.1754>

18. **Vandenbroucke JP, von Elm E, Altman DG, Gotzsche PC, Mulrow CD, Pocock SJ**, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Epidemiology*. 2007; 18(6): 805–35. DOI: <https://doi.org/10.1097/EDE.0b013e3181577511>
19. **Arnold C, Hennrich P, Koetsenruijter J, van Lieshout J, Peters-Klimm F, Wensing M**. Cooperation networks of ambulatory health care providers: exploration of mechanisms that influence coordination and uptake of recommended cardiovascular care (ExKoCare): a mixed-methods study protocol. *BMC Fam Pract*. 2020; 21(1): 168. DOI: <https://doi.org/10.1186/s12875-020-01229-3>
20. **Hennrich P, Arnold C, Wensing M**. Effects of personalized invitation letters on research participation among general practitioners: a randomized trial. *BMC Med Res Methodol*. 2021; 21(1): 247. DOI: <https://doi.org/10.1186/s12874-021-01447-y>
21. **Uijen AA, Schellevis FG, van den Bosch WJ, Mokkink HG, Van Weel C, Schers HJ**. Nijmegen Continuity Questionnaire: development and testing of a questionnaire that measures continuity of care. *J Clin Epidemiol*. 2011; 64(12): 1391–9. DOI: <https://doi.org/10.1016/j.jclinepi.2011.03.006>
22. **Uijen AA, Schers HJ, Schellevis FG, Mokkink HG, Van Weel C, van den Bosch WJ**. Measuring continuity of care: psychometric properties of the Nijmegen Continuity Questionnaire. *Br J Gen Pract*. 2012; 62(600): e949–e57. DOI: <https://doi.org/10.3399/bjgp12X652364>
23. **Hetlevik O, Hustoft M, Uijen A, Assmus J, Gjesdal S**. Patient perspectives on continuity of care: adaption and preliminary psychometric assessment of a Norwegian version of the Nijmegen Continuity Questionnaire (NCQ-N). *BMC Health Serv Res*. 2017; 17(1): 760. DOI: <https://doi.org/10.1186/s12913-017-2706-1>
24. **Hennrich P, Arnold C, Koetsenruijter J, Wensing M**. Measuring continuity of ambulatory cardiovascular care: a cross-sectional study on the applicability of the Nijmegen Continuity Questionnaire in Germany. *BMC Health Serv Res*. 2022 Oct 18; 22(1): 1258. DOI: <https://doi.org/10.1186/s12913-022-08612-z>
25. **Arnold C, Hennrich P, Wensing M**. Information exchange networks for chronic diseases in primary care practices in Germany: a cross-sectional study. *BMC Prim Care*. 2022; 23(1): 56. DOI: <https://doi.org/10.1186/s12875-022-01649-3>
26. **Kivimaki M, Elovainio M**. A short version of the Team Climate Inventory: Development and psychometric properties. *Journal of occupational and organizational psychology*. 1999; 72(2): 241–6. DOI: <https://doi.org/10.1348/0963179991666644>
27. **Ndibu Muntu Keba Kebe N, Chiochio F, Bamvita JM, Fleury MJ**. Variables associated with interprofessional collaboration: The case of professionals working in Quebec local mental health service networks. *J Interprof Care*. 2019; 33(1): 76–84. DOI: <https://doi.org/10.1080/13561820.2018.1515191>
28. **Busse R, Blümel M, Knieps F, Bärnighausen T**. Statutory health insurance in Germany: a health system shaped by 135 years of solidarity, self-governance, and competition. *The Lancet*. 2017; 390(10097): 882–97. DOI: [https://doi.org/10.1016/S0140-6736\(17\)31280-1](https://doi.org/10.1016/S0140-6736(17)31280-1)

TO CITE THIS ARTICLE:

Arnold C, Hennrich P, Peters-Klimm F, Wensing M. Influence of Cooperation Work on Management Continuity of Ambulatory Cardiovascular Care: A Cross-Sectional Exploratory Study in Germany. *International Journal of Integrated Care*, 2023; 23(4): 14 , 1–8. DOI: <https://doi.org/10.5334/ijic.7019>

Submitted: 01 September 2022 **Accepted:** 31 October 2023 **Published:** 06 December 2023

COPYRIGHT:

© 2023 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.

International Journal of Integrated Care is a peer-reviewed open access journal published by Ubiquity Press.