

# Acculturation and Disparities in Telemedicine Readiness: A National Study

The International Journal of Aging and Human Development  
2024, Vol. 99(1) 96–114  
© The Author(s) 2023



Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/00914150231219259  
journals.sagepub.com/home/ahd



Jorge M. Rodríguez-Fernández<sup>1</sup> ,  
Nicolas Hoertel<sup>2,3,4</sup>, Hugo Saner<sup>5</sup>,  
and Mukaila Raji<sup>6</sup>

## Abstract

Telemedicine provided older adults the ability to safely seek care during the coronavirus disease (COVID-19) pandemic. This study aimed to evaluate the potential impact of acculturation factors in telemedicine uptake between ethnic groups. As part of the National Health and Aging Trends Study 2018 survey, 303 participants ( $\geq 65$  years) were interviewed. We assessed the impact of acculturation on telemedicine readiness by race and ethnicity. Compared to the white non-Hispanic immigrant population, Hispanic and Asian/Pacific Islander (API) populations had significantly lower telemedicine readiness and uptake. Limited English proficiency or older age at the time of migration was associated with telemedicine unreadiness and uptake in the Hispanic and API populations. Our findings suggested that acculturation factors play a substantial role in telemedicine uptake among older adult immigrants in the United States. Therefore, acculturation factors should be considered when promoting and adopting telemedicine technologies in older adults.

<sup>1</sup>Department of Neurology, The University of Texas Medical Branch, Galveston, TX, USA

<sup>2</sup>AP-HP, Centre-Université de Paris, Hôpital Corentin-Celton, DMU Psychiatrie et Addictologie, Issy-les-Moulineaux, France

<sup>3</sup>INSERM, Institut de Psychiatrie et Neurosciences de Paris, Paris, France

<sup>4</sup>Faculté de Santé, UFR de Médecine, Université de Paris, Paris, France

<sup>5</sup>Institute of Social and Preventive Medicine (ISPM), University of Bern, Switzerland

<sup>6</sup>Division of Geriatrics, Internal Medicine, The University of Texas Medical Branch, Galveston, TX, USA

## Corresponding Author:

Jorge Mario Rodríguez-Fernández, The University of Texas Medical Branch, 301 University Blvd. 9.128 John Sealy Annex Galveston, TX 77555, USA.

Email: jormrodr@utmb.edu

**Keywords**

telemedicine, telemedicine readiness, acculturation, English proficiency, COVID-19

**Introduction**

Health disparities exist in the United States, particularly among minorities (Forrester et al., 2020; Thomson & Hoffman-Goetz, 2009). Health disparities in older adults, especially older minorities, reflect several factors, including limited access to care, poverty, unhealthy neighborhood, systemic racism, and acculturation (Arcaya et al., 2016; Lee et al., 2020a, 2020b; Ryvicker & Sridharan, 2018). Acculturation factors (defined as “the vehicle to assimilation by cultural adoption of the host society’s norms, values, behaviors and attitudes”)—unlike other factors in health disparity—have been less studied as a source of health disparity (Wallace et al., 2010). Proxy measures for acculturation include age at migration and language proficiency. The impact on this population of telemedicine remains unclear and poorly studied (Thomson & Hoffman-Goetz, 2009). It is currently estimated that about 5 million older adults across the United States possess limited English proficiency (NCLER, 2019), and despite current work addressing disparities among older adults (Robert & Ruel, 2006), language barriers limit the quality of care received (United States Census Bureau). Specifically, prior studies have shown an increased number of hospitalizations, emergency department visits, readmissions, longer lengths of stay, and a lack of health literacy (Baker et al., 2002). In addition, foreign-born (first generation) older adults perceive themselves as having a lower health quality compared to subsequent generations (Acevedo-Garcia et al., 2010). Therefore, adequate education and equitable deployment of telemedicine alternatives may offer opportunities to reduce these inequities (Fulmer et al., 2021; López et al., 2011).

The expansion of telemedicine during the COVID-19 pandemic could give a unique opportunity to close those gaps in care. Evidence shows telemedicine-related improvements in health outcomes (Chevance et al., 2020; Comín-Colet et al., 2016; Gandolfi et al., 2017; Hong et al., 2017; Karhula et al., 2015; Kim et al., 2014; Liu et al., 2011; Logan et al., 2012; Trief et al., 2013; Wakefield et al., 2011; Wu et al., 2015), quality of life (Fortney et al., 2007; Häggglund et al., 2015; Liu et al., 2011; Seto et al., 2012; van der Weegen et al., 2015), medication management (Menon et al., 2018; Moore et al., 2015; Patel et al., 2013; Trief et al., 2013), and health literacy (Liu et al., 2011; Patel et al., 2013) among patients using telehealth prior to the pandemic. Nevertheless, it remains unknown how acculturation factors play a role in telemedicine uptake among older adults, especially older minorities. Designing and implementing culture-appropriate interventions to increase technology uptake in older minorities requires a comprehensive study of the roles of acculturation factors in healthcare

technology readiness and uptake. Such a study has the potential to provide data to guide telemedicine approaches to narrow the digital divide in ethnic minorities. In this study, we aim to assess the association between acculturation and telemedicine uptake across race and ethnicity among older adults with limited English proficiency or who were foreign-born.

## **Methods**

### **Sample**

The present work is a cross-sectional study of community-dwelling participants from the 2018 National Health and Aging Trends Study (NHATS), which included 363 individuals (Freedman & Kasper, 2019). Of these, 12 (3.3%) participants were excluded because they were noncommunity residents (residential care or nursing home) and 48 (13.2%) because of cognitive decline, as detailed elsewhere (Rodríguez-Fernández et al., 2017; Rodríguez-Fernández et al., 2022). The remaining 303 individuals were included in the analysis. The NHATS protocol was approved by the the Johns Hopkins University Institutional Review Board, and all participants provided informed consent.

### **Race and Ethnicity**

National Health and Aging Trends Study participants were asked to mark one race and ethnicity with which they identified, including white, non-Hispanic; African American, Non-Hispanic; Hispanic/Latino; and non-Hispanic others, including Asians and Pacific Islanders (API). This categorization was used in the present study.

### **Acculturation Factors**

Per prior work (Kang et al., 2021), acculturation was measured in two different ways: First, English proficiency was measured by (a) speaking and (b) listening skills using a four-point Likert scale (1 = very well, 4 = not at all). Participants who spoke English well or very well and understood spoken English well or very well were categorized as highly English proficient. Second, age at migration was measured as a continuous variable ranging from 0 to 76.

### **Telemedicine Readiness and Unreadiness**

Telemedicine readiness was defined as one of the following: (1) contacting a medical provider online; (2) handling medical insurance matters online (coverage, comparing providers, bill status, or filing a claim); and (3) obtaining information about medical conditions online.

Telemedicine unreadiness was defined as any physical or technical factor that limited the communication between healthcare providers and the patient. Physical limitations included any of the following: difficulty hearing, difficulty watching television or reading a newspaper even with glasses, or problems speaking or making self-understood. Technical limitations included not owning a working telecommunication device (computer, cell phone, or telephone) or not knowing how to use them, or no recent use of email, texting, or Internet during the last month.

## Medical Conditions

Participants or their proxies were asked if they had been diagnosed with any of the following medical conditions: cancer, myocardial infarction, heart disease (separate from myocardial infarction), hypertension, stroke, diabetes, lung disease, osteoarthritis, and osteoporosis. Self-reports of chronic conditions have shown good diagnostic accuracy against claims and medical records (kappa ranging from 0.6 to 0.82) (Miller et al., 2008). Depression and anxiety symptoms were assessed using two validated screening questionnaires, the Patient Health Questionnaire-2 (PHQ-2) and the Generalized Anxiety Disorder-2 scale (GAD-2). We used a PHQ-2 score  $\geq 3$  to define substantial depression and a GAD-2 score  $\geq 3$  for anxiety symptoms; both questionnaires are validated clinical tools among older adults (Boyle et al., 2011; Wild et al., 2014).

## Statistical Analysis

Demographic characteristics and acculturation factors of community-dwelling participants were compared between each telemedicine group; those who met any of readiness criteria and those who did not, and between those with unreadiness (physical or technical) and those without this condition, using  $\chi^2$  tests. To account for the effects of age, we dichotomized age as younger than 80 versus older than 80 since the latter is more likely to lack technological exposure and have higher frailty levels limiting technological implementation and acquisition (Harvie et al., 2014; Harvie et al., 2016). Next, frequency rates of telemedicine readiness and unreadiness factors across acculturation factors were calculated. Adjusted logistic regression models were used to estimate telemedicine readiness and unreadiness due to physical or technical factors by acculturation factors. We adjusted for potentially confounding factors, including age, gender, education, marital status, and medical conditions (Rodríguez-Fernández et al., 2022). For all associations, we performed residual analyses to assess the fit of the data, checked assumptions, and examined the potential influence of outliers. Since our analyses were mainly exploratory, statistical significance was fixed a priori at a two-sided  $P$ -value  $< .05$ . Statistical analyses were conducted using Stata 12 SE version (Stata Corp, TX) and R version 4 (R Foundation for Statistical Computing, Vienna, Austria).

## Results

### Demographic Characteristics by Race and Ethnicity

Table 1 shows the demographics, clinical characteristics, and medical conditions of the study sample by race and ethnicity. No age or gender differences were noted across groups. White non-Hispanic and APIs were more likely to be married and have a college graduate degree compared to African American non-Hispanic and Hispanic. Regarding acculturation, the age of migration to the United States was significantly higher for African American non-Hispanic and API. Lack of English proficiency was more prevalent among APIs and Hispanics. Regarding medical conditions for African American non-Hispanic and Hispanics were more likely to report diabetes, depression, and anxiety; lung disease and osteoporosis were more prevalent among APIs and Hispanics.

### Telemedicine Factors by Race and Ethnicity

Table 2 shows the prevalence of each telemedicine category by race and ethnicity. In regard to telemedicine readiness, being able to look for medical information online and contact medical providers online were more prevalent in white non-Hispanics (47% and 43%, respectively). Regarding physical unreadiness, difficulty reading the newspaper even with glasses was noted to be more prevalent among APIs (13%) and Hispanics (11%) compared with white Non-Hispanic and African American Non-Hispanic. For technical unreadiness, no recent internet use in the last month was the most prevalent (52%), followed by no recent email or texting use in the last month (65%), not having a working computer (54%), among Hispanic older adults.

### Acculturation Factors Across Telemedicine Readiness and Unreadiness

The association between telemedicine readiness and race and ethnicity was evaluated while adjusting for age, sex, marital status, education level, and chronic medical conditions. As shown in Table 3, a higher age at migration to the United States was significantly and negatively associated among APIs (odds ratio [OR]=0.22, 95% confidence interval [CI]=0.08–0.64,  $P < .01$ ) and Hispanic older adults (OR = 0.22, 95%CI=0.09–0.58,  $P < .01$ ) with telemedicine readiness, while a positive association was observed for technical unreadiness in APIs (OR = 6.93, 95% CI = 2.50–19.23,  $P < .001$ ) and Hispanic (OR = 3.22, 95% CI = 1.38–7.51,  $P < .01$ ) older adults. Similarly, lack of English proficiency was significantly and negatively associated with APIs (OR = 0.14, 95% CI = 0.04–0.46,  $P < .01$ ) for telemedicine readiness. Concerning physical unreadiness, APIs had a strong association with an OR of 5.04 (95%CI = 1.50–16.98,  $P < .01$ ). Lastly, for technical unreadiness, a positive association was noted with Hispanic (OR = 2.70, 95%CI = 1.14–6.36,  $P < .05$ )

**Table 1.** Demographic, Acculturation Factors, and Clinical Characteristics by Race/Ethnicity.

Characteristics <sup>a</sup>	White, Non-Hispanic (N = 205)	African American, Non-Hispanic (N = 55)	Asian and Pacific Islanders (N = 52)	Hispanic (N = 143)	P value
Age					.57
<80	119 (58)	36 (65)	34 (65)	82 (57)	
>80	86 (42)	19 (35)	18 (35)	61 (43)	.567
Gender					
Male	102 (50)	27 (49)	29 (56)	64 (45)	
Female	103 (50)	28 (51)	23 (44)	79 (55)	.003
Marital status					
Unmarried	86 (42)	38 (69)	21 (40)	71 (50)	
Married	119 (58)	17 (31)	31 (60)	72 (50)	<.001
Education status					
High school graduate or less	44 (21)	24 (44)	14 (27)	105 (73)	
College graduate	161 (79)	31 (56)	38 (73)	38 (27)	
Acculturation factors					
Age of migration to the US, mean $\pm$ SD	25.526 $\pm$ 14.960	38.359 $\pm$ 15.484	37.178 $\pm$ 12.229	32.227 $\pm$ 15.538	<.001
Lack of English proficiency	3 (2.0)	0 (0)	6 (22)	49 (50)	<.001
Medical condition					
Hypertension	147 (72)	47 (85)	37 (71)	112 (78)	.129
Myocardial infarction	2 (1.0)	0 (0)	0 (0)	7 (4.9)	.061
Heart disease <sup>b</sup>	43 (21)	9 (16)	7 (13)	30 (21)	.566
Stroke	4 (2.0)	1 (1.8)	0 (0)	2 (1.4)	.946
Diabetes	50 (24)	33 (60)	18 (35)	73 (51)	<.001

(Continued)

Table 1. Continued

Characteristics <sup>a</sup>	White, Non-Hispanic (N = 205)	African American, Non-Hispanic (N = 55)	Asian and Pacific Islanders (N = 52)	Hispanic (N = 143)	P value
Cancer	9 (4.4)	3 (5.5)	0 (0)	3 (2.1)	.241
Lung disease	24 (12)	9 (16)	11 (21)	39 (27)	.003
Osteoarthritis	144 (70)	36 (65)	36 (69)	101 (71)	.905
Osteoporosis	73 (36)	11 (20)	25 (48)	56 (39)	.019
Depression	16 (7.9)	8 (15)	4 (7.7)	33 (23)	<.001
Anxiety	12 (5.9)	4 (7.4)	1 (2.0)	21 (15)	.011

<sup>a</sup>Data are given as number (percentage) for each group, unless noted.

<sup>b</sup>Heart disease is separate from myocardial infarction.

**Table 2.** Telemedicine Factors by Race/Ethnicity.

Telemedicine factors <sup>a</sup>	White, Non-Hispanic (N = 205)	African American, Non-Hispanic (N = 55)	Asian and Pacific Islanders (N = 52)	Hispanic (N = 143)	P value
<b>Readiness</b>					
Medical information online	71 (47)	11 (42)	7 (35)	10 (25)	.088
Contact medical provider online	65 (43)	9 (35)	6 (30)	8 (20)	.055
Medical insurance online	34 (22)	9 (35)	2 (10)	8 (20)	.246
<b>Physical unreadiness</b>					
Difficulty speaking or making self-understood	8 (3.9)	4 (7.3)	4 (7.7)	10 (7.0)	.409
Difficulty reading newspaper even with glasses	2 (1.0)	3 (5.5)	7 (13)	16 (11)	<.001
Unable to see TV across room with glasses	1 (9.1)	0 (0)	2 (33)	5 (33)	.341
Difficulty hearing phone	4 (2.0)	1 (1.8)	1 (1.9)	7 (4.9)	.436
<b>Technical unreadiness</b>					
No recent internet use in the last month	45 (23)	25 (49)	29 (59)	93 (70)	<.001
No recent email or texting use in the last month	39 (20)	17 (33)	15 (31)	87 (65)	<.001
Does not have a working computer	39 (19)	21 (39)	15 (31)	74 (54)	<.001
Does not have a working cellphone	15 (7.3)	5 (9.1)	4 (7.7)	15 (10)	.752
Has a computer, doesn't know how to use it	2 (1.2)	1 (2.9)	3 (8.1)	5 (7.2)	.026

<sup>a</sup>Data are given as number (percentage) for each group.



and API (OR = 6.68 95%CI = 2.35–18.93,  $P < .001$ ) older adults respectively, as compared with white older adults.

## Discussion

In this nationally representative sample of older adults, we found limited telemedicine uptake among APIs and Hispanics based on their age of migration or English proficiency. Our results confirmed and extended previous findings. For example, data from a large cohort of older adults from a large healthcare system in California indicated that Black, Latino, and Filipino elders and those aged 75 years and older were less likely than non-Hispanic whites and Chinese elders to engage in eHealth technologies or use the patient portal, in part due to not having or not knowing how to use the devices (computers and smartphones) (Gordon & Hornbrook, 2016). However, this study did not account for acculturation factors across race and ethnicity in older adults, which may play a role in telemedicine uptake in this community. Overall, the findings in our study of health inequalities in telemedicine access and use among API and Hispanic populations underscored the need to develop targeted healthcare access-improving interventions that account for acculturation factors unique to this population.

Interestingly, telemedicine uptake for both API and Hispanic immigrants was worse based on their age of migration to the United States and limited English proficiency. Several explanations could account for these results. We know that older migration age is independently associated with a decline in physical and mental health after migration (Kang et al., 2021), with higher rates of mental health symptoms (Mui et al., 2003), multiple chronic medical conditions (Seo et al., 2017), social isolation (Lee et al., 2020a, 2020b), discrimination (Li & Dong, 2017), and social disparities (Lee et al., 2020a, 2020b). These factors may present barriers to telemedicine access to Hispanic older adults compared to their peers during the COVID-19 pandemic (Ryskina et al., 2021). Further work has shown that older minorities experience difficulties in telemedicine, especially during video visits, with the difficulty being further accentuated when needing an interpreter (Schifeling et al., 2020). Also, a lack of English proficiency and limited health literacy were noted to result in important communication barriers among older, adult Chinese American immigrants (Tsoh et al., 2016). In addition, despite policy changes by the Centers for Medicare and Medicaid Services (CMS), including paying providers for telehealth visits, minorities with limited English proficiency still experience disparities in their care, especially during the COVID-19 pandemic (Wilkins et al., 2021). Moreover, up to a third of the US health systems do not offer language services despite known mandates (Chen et al., 2007; Schiaffino et al., 2014). These barriers not only limit telemedicine access but also accentuate care gaps and previously existing suboptimal healthcare access among older minorities.

A substantial proportion of US older adults are more likely to lack access to technology and electronic health records portals and to have low digital health

**Table 3. Adjusted Odds Ratios for Telemedicine Readiness and Unreadiness Across Race/Ethnicity and Acculturation Factors.**

Characteristic <sup>a</sup>	Readiness		Physical unreadiness		Technical unreadiness	
	Odds Ratio	95% CI	Odds Ratio	95% CI	Odds Ratio	95% CI
<b>Age came to the United States (ref = White, Non-Hispanic)</b>						
African American, Non-Hispanic	0.32	(0.10 - 1.03)	1.17	(0.25 - 5.60)	1.72	(0.58 - 5.16)
	252		281		277	
Asian and Pacific Islanders	0.22**	(0.08 - 0.64)	2.64	(0.69 - 10.08)	6.93***	(2.50 - 19.23)
	252		281		277	
Hispanic	0.22**	(0.09 - 0.58)	1.53	(0.48 - 4.88)	3.22**	(1.38 - 7.51)
	252		281		277	
<b>English Proficiency (ref = White, Non-Hispanic)</b>						
African American, Non-Hispanic	0.49	(0.18 - 1.37)	1.19	(0.22 - 6.61)	2.04	(0.69 - 6.05)
	286		288		288	
Asian and Pacific Islanders	0.14**	(0.04 - 0.46)	5.04**	(1.50 - 16.98)	6.68***	(2.35 - 18.93)
	286		288		288	
Hispanic	0.59	(0.25 - 1.37)	1.17	(0.36 - 3.82)	2.70*	(1.14 - 6.36)
	286		288		288	

<sup>a</sup>Odds ratios were estimated adjusting for age, gender, marital status, education level, and chronic medical conditions.

\*  $P < .05$ .

\*\*  $P < .01$ .

\*\*\*  $P < .001$ .

literacy (Smith, 2014). Currently, efforts by the National Institute on Aging include the promotion and support of research and other projects to improve the effectiveness of telemedicine for people with cognitive deficits, with the intention that such projects can be scaled and expanded to older adults without dementia (National Institute on Aging, 2020). In addition, the Federal Communications Commission (FCC) supported telehealth programs across the nation as part of the COVID-19 pandemic. Despite these efforts, current limitations include the limited infrastructure and providers supporting the offering of telemedicine services—especially in rural areas, a known limiting factor, as stated by the American Telemedicine Association (Humble et al., 2016). These prior efforts highlight the importance of developing interventions to enhance and optimize telemedicine uptake among older adults—especially older minorities. To achieve such optimization, telehealth experts offer two major recommendations: (1) optimizing telemedicine efforts outside health care (design and testing) and within healthcare (evaluating healthcare delivery, determinants of health, and increasing access) and (2) ensuring privacy and access as part of the 21st Century Cures Act (Humble et al., 2016).

Surpassing the existing limitations on telemedicine uptake includes the comprehensive evaluation of digital literacy, support for internet infrastructure (e.g., 5G wireless, fiber optic cable, and Wi-Fi, especially in rural and minority-dense areas), health literacy and language proficiency support, reduction of technical and physical barriers for all seniors, especially among API and Hispanic older adults. Our study, in particular, noted the strong effects of digital literacy and technical factors, consistent with prior work (Gordon & Hornbrook, 2016; Rodriguez et al., 2021; Wilcock et al., 2019). As per a recent systematic review (Kruse et al., 2020), the most common factor limiting telemedicine interventions included technical factors, motivation, and costs. These findings were replicated by our study even after we accounted for acculturation factors. However, technical factors do not account for the entire problem. Therefore, further research evaluating interventions across patient–provider communication and patient empowerment is needed, especially among older adult minorities experiencing communication barriers, as telemedicine continues to expand across all healthcare settings (clinics, nursing homes, postacute care, and hospitals).

Our results have several implications. From a clinical perspective, our work suggested that clinicians should be aware that immigrant older adults and those with limited English proficiency are more likely to experience difficulties with telemedicine, extending prior work on the barriers to telemedicine among older adults (Foster & Sethares, 2014; Kruse et al., 2020). With that awareness, clinicians may need to ask if a patient's family member or caregiver might be present to facilitate the visit. Alternatively, the visit might be recorded—thus allowing the patient to listen or watch the visit encounters by themselves with or without a more health-literate family member. Also, our findings highlighted the value of communication optimization for healthcare delivery, as English proficiency enhances healthcare access as per the Andersen Behavioral Model (Andersen, 1995). Interpreter availability should be increased to close the gap in healthcare access among older adult minorities. From a

public health perspective, our study highlighted the importance of the continued expansion of telemedicine services, especially in underserved communities, with the federal support of the FCC and payment approvals by Medicare and private healthcare insurance companies. Furthermore, reimbursement continuation and equality for telemedicine visits from commercial services and CMS need to be maintained as compared to the more complex earlier guidelines pre-COVID-19. Our results suggest that from an institutional level, providers should be aware of the high rate of limited health literacy and communication barriers in older adult minorities with low English proficiency or those who were foreign-born. They also should be aware of the need for the development of streamlined protocols for language interpreters to be available to facilitate patient-provider communication and decrease gaps in healthcare access. Finally, inflation of type I error might have occurred in analyses due to multiple testing, and the observed associations should not be interpreted as causal relationships (Le Strat & Hoertel, 2011).

The strengths of our study include a relatively large sample size and its national representativeness. Results should also be interpreted in light of limitations common to most large-scale surveys. Although the NHATS is a nationally representative sample, it does not include non-Medicare beneficiaries or undocumented immigrants. Moreover, since this survey was conducted in English and Spanish, it is possible that immigrants who were not familiar with either language may not have been included in this survey. In addition, reporting devices or technical issues can potentially lead to over- or underestimation of the true rates of readiness or unreadiness. Also, the severity and possible consequences of telemedicine unreadiness were not evaluated, leaving the impact on older adults' health overtime unclear. Longitudinal data are needed to examine the course of acculturation barriers and to evaluate older adults' responses to either readiness or unreadiness.

Despite these limitations, our study constituted a critical step in the importance of racial and ethnic disparities and acculturation factors in telemedicine readiness and unreadiness in the United States. In a large, nationally representative sample of older, foreign-born US adults, a lack of English proficiency was associated with a lack of telemedicine readiness for Hispanics and APIs. Given the ongoing efforts to provide and enhance care through telemedicine for older adults during the COVID-19 pandemic, these results support the need to consider acculturation factors by race and ethnicity to facilitate telemedicine usage and adoption in older adults.

### **IRB Review**

Since all the analyses were performed on de-identified NHATS data which is publicly available for download, IRB review was not required.


### **Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## ORCID iD

Jorge M. Rodríguez-Fernández  <https://orcid.org/0000-0003-4888-2041>

## References

- Acevedo-Garcia, D., Bates, L. M., Osypuk, T. L., & McArdle, N. (2010). The effect of immigrant generation and duration on self-rated health among US adults 2003–2007. *Social Science & Medicine*, *71*(6), 1161–1172. <https://doi.org/10.1016/j.socscimed.2010.05.034>
- Andersen, R. M. (1995). Revisiting the behavioral model and access to medical care: Does it matter? *Journal of Health and Social Behavior*, *36*(1), 1–10. <https://doi.org/10.2307/2137284>
- Arcaya, M., Tucker-Seeley, R., Kim, R., Schnake-Mahl, A., So, M., & Subramanian, S. (2016). Research on neighborhood effects on health in the United States: A systematic review of study characteristics. *Social Science & Medicine*, *168*, 16–29. <https://doi.org/10.1016/j.socscimed.2016.08.047>
- Baker, D. W., Gazmararian, J. A., Williams, M. V., Scott, T., Parker, R. M., Green, D., Ren, J., & Peel, J. (2002). Functional health literacy and the risk of hospital admission among Medicare managed care enrollees. *American Journal of Public Health*, *92*(8), 1278–1283. <https://doi.org/10.2105/AJPH.92.8.1278>
- Boyle, L. L., Richardson, T. M., He, H., Xia, Y., Tu, X., Boustani, M., & Conwell, Y. (2011). How do the phq-2, the phq-9 perform in aging services clients with cognitive impairment? *International Journal of Geriatric Psychiatry*, *26*(9), 952–960. <https://doi.org/10.1002/gps.2632>
- Chen, A. H., Youdelman, M. K., & Brooks, J. (2007). The legal framework for language access in healthcare settings: Title VI and beyond. *Journal of General Internal Medicine*, *22*(2), 362–367. <https://doi.org/10.1007/s11606-007-0366-2>
- Chevance, A., Gourion, D., Hoertel, N., Llorca, P.-M., Thomas, P., Bocher, R., Moro, M.-R., Benyamina, A., Fossati, P., Masson, M., Leaute, E., Leboyer, M., & Gaillard, R. (2020). Ensuring mental health care during the SARS-CoV-2 epidemic in France: A narrative review. *L'Encéphale*, *46*(3), 193–201. <https://doi.org/10.1016/j.encep.2020.04.005>
- Comín-Colet, J., Enjuanes, C., Verdú-Rotellar, J. M., Linas, A., Ruiz-Rodríguez, P., González-Robledo, G., Farré, N., Moliner-Borja, P., Ruiz-Bustilla, S., & Bruguera, J. (2016). Impact on clinical events and healthcare costs of adding telemedicine to multidisciplinary disease management programmes for heart failure: Results of a randomized controlled trial. *Journal of Telemedicine and Telecare*, *22*(5), 282–295. <https://doi.org/10.1177/1357633X15600583>

- Forrester, S. N., Taylor, J. L., Whitfield, K. E., & Thorpe, R. J. (2020). Advances in understanding the causes and consequences of health disparities in aging minorities. *Current Epidemiology Reports*, 7(2), 59–67. <https://doi.org/10.1007/s40471-020-00234-5>
- Fortney, J. C., Pyne, J. M., Edlund, M. J., Williams, D. K., Robinson, D. E., Mittal, D., & Henderson, K. L. (2007). A randomized trial of telemedicine-based collaborative care for depression. *The Journal of General Internal Medicine*, 22(8), 1086–1093. <https://doi.org/10.1007/s11606-007-0201-9>
- Foster, M. V., & Sethares, K. A. (2014). Facilitators and barriers to the adoption of telehealth in older adults: An integrative review. *Computers, Informatics, Nursing*, 32(11), 523–533. <https://doi.org/10.1097/CIN.0000000000000105>
- Freedman, V. A., & Kasper, J. D. (2019). Cohort profile: The National Health and Aging Trends Study (NHATS). *International Journal of Epidemiology*, 48(4), 1044. <https://doi.org/10.1093/ije/dyz109>
- Fulmer, T., Reuben, D. B., Auerbach, J., Fick, D. M., Galambos, C., & Johnson, K. S. (2021). Actualizing better health and health care for older adults: Commentary describes six vital directions to improve the care and quality of life for all older Americans. *Health Affairs*, 40(2), 219–225. <https://doi.org/10.1377/hlthaff.2020.01470>
- Gandolfi, M., Geroin, C., Dimitrova, E., Boldrini, P., Waldner, A., Bonadiman, S., Picelli, A., Regazzo, S., Stirbu, E., Primon, D., Bosello, C., Gravina, A. R., Peron, L., Trevisan, M., Garcia, A. C., Menel, A., Bloccari, L., Valè, N., Saltuari, L., & Smania, N. (2017). Virtual reality telerehabilitation for postural instability in Parkinson's disease: A multicenter, single-blind, randomized, controlled trial. *Biomed Research International*, 2017. <https://doi.org/10.1155/2017/7962826>
- Gordon, N. P., & Hornbrook, M. C. (2016). Differences in access to and preferences for using patient portals and other eHealth technologies based on race, ethnicity, and age: A database and survey study of seniors in a large health plan. *Journal of Medical Internet Research*, 18(3), e5105. <https://doi.org/10.2196/jmir.5105>
- Hägglund, E., Lyngå, P., Frie, F., Ullman, B., Persson, H., Melin, M., & Hagerman, I. (2015). Patient-centred home-based management of heart failure. *Scandinavian Cardiovascular Journal*, 49(4), 193–199. <https://doi.org/10.3109/14017431.2015.1035319>
- Harvie, G., Burmeister, O., & Eustace, K. (2014). Bringing the oldest-old into the digital age: 13th national conference of emerging researchers in ageing. 13th National Conference of Emerging Researchers in Ageing – Adelaide. 2014, 103–105.
- Harvie, G., Eustace, K., & Burmeister, O. K. (2016). Assistive technology devices for the oldest-old: Maintaining independence for the fourth age. In D. Kreps, G. Fletcher, & M. Griffiths (Eds.), *Technology and intimacy: Choice or coercion* (pp. 25–33). Springer International Publishing. [https://doi.org/10.1007/978-3-319-44805-3\\_3](https://doi.org/10.1007/978-3-319-44805-3_3)
- Hong, J., Kim, J., Kim, S. W., & Kong, H.-J. (2017). Effects of home-based tele-exercise on sarcopenia among community-dwelling elderly adults: Body composition and functional fitness. *Experimental Gerontology*, 87, 33–39. <https://doi.org/10.1016/j.exger.2016.11.002>
- Humble, J. R., Tolley, E. A., Krukowski, R. A., Womack, C. R., Motley, T. S., & Bailey, J. E. (2016). Use of and interest in mobile health for diabetes self-care in vulnerable populations. *Journal of Telemedicine and Telecare*, 22(1), 32–38. <https://doi.org/10.1177/1357633X15586641>

- Kang, G., Hajduk, A., Marottoli, R., & Nunez-Smith, M. (2021). Older immigrants perceived health after migration to the United States: Influence of age and level of acculturation. *Journal of the American Geriatrics Society*, 69(9), 2625–2637. <https://doi.org/10.1111/jgs.17363>
- Karhula, T., Vuorinen, A.-L., Rääpysjärvi, K., Pakanen, M., Itkonen, P., Tepponen, M., & Lähteenmäki, J. (2015). Telemonitoring and mobile phone-based health coaching among Finnish diabetic and heart disease patients: Randomized controlled trial. *The Journal of Medical Internet Research*, 17(6), e153. <https://doi.org/10.2196/jmir.4059>
- Kim, H.-S., Choi, W., Baek, E. K., Kim, Y. A., Yang, S. J., Choi, I. Y., & Cho, J.-H. (2014). Efficacy of the smartphone-based glucose management application stratified by user satisfaction. *Diabetes & Metabolism Journal*, 38(3), 204–210. <https://doi.org/10.4093/dmj.2014.38.3.204>
- Kruse, C., Fohn, J., Wilson, N., Patlan, E. N., Zipp, S., & Mileski, M. (2020). Utilization barriers and medical outcomes commensurate with the use of telehealth among older adults: Systematic review. *JMIR Medical Informatics*, 8(8), e20359. <https://doi.org/10.2196/20359>
- Lee, H., Kim, D., Lee, S., & Fawcett, J. (2020a). The concepts of health inequality, disparities and equity in the era of population health. *Applied Nursing Research*, 56, 151367. <https://doi.org/10.1016/j.apnr.2020.151367>
- Lee, J., Hong, J., Zhou, Y., & Robles, G. (2020b). The relationships between loneliness, social support, and resilience among Latinx immigrants in the United States. *Clinical Social Work Journal*, 48(1), 99–109. <https://doi.org/10.1007/s10615-019-00728-w>
- Le Strat, Y., & Hoertel, N. (2011). Correlation is no causation: Gymnasium proliferation and the risk of obesity. *Addiction*, 106(10), 1871–1872. <https://doi.org/10.1111/j.1360-0443.2011.03547.x>
- Li, L. W., & Dong, X. (2017). Self-reported discrimination and depressive symptoms among older Chinese adults in Chicago. *Journals of Gerontology: Series A*, 72(suppl\_1), S119–S124. <https://doi.org/10.1093/gerona/glw174>
- Liu, W.-T., Huang, C.-D., Wang, C.-H., Lee, K.-Y., Lin, S. M., & Kuo, H. P. (2011). A mobile telephone-based interactive self-care system improves asthma control. *European Respiratory Journal*, 37(2), 310–317. <https://doi.org/10.1183/09031936.00000810>
- Logan, A. G., Irvine, M. J., McIsaac, W. J., Tisler, A., Rossos, P. G., Easty, A., Feig, D. S., & Cafazzo, J. A. (2012). Effect of home blood pressure telemonitoring with self-care support on uncontrolled systolic hypertension in diabetics. *Hypertension*, 60(1), 51–57. <https://doi.org/10.1183/09031936.00000810>
- López, L., Green, A. R., Tan-McGrory, A., King, R. S., & Betancourt, J. R. (2011). Bridging the digital divide in health care: The role of health information technology in addressing racial and ethnic disparities. *The Joint Commission Journal on Quality and Patient Safety*, 37(10), 437–445. [https://doi.org/10.1016/s1553-7250\(11\)37055-9](https://doi.org/10.1016/s1553-7250(11)37055-9)
- Menon, V., Selvakumar, N., Kattimani, S., & Andrade, C. (2018). Therapeutic effects of mobile-based text message reminders for medication adherence in bipolar I disorder: Are they maintained after intervention cessation? *Journal of Psychiatric Research*, 104, 163–168. <https://doi.org/10.1016/j.jpsychires.2018.07.013>

- Miller, D. R., Rogers, W. H., Kazis, L. E., Spiro, I. I. I., Ren, A., Haffer, X. S., & C, S. (2008). Patients' self-report of diseases in the medicare health outcomes survey based on comparisons with linked survey and medical data from the veterans health administration. *The Journal of Ambulatory Care Management*, 31(2), 161–177. <https://doi.org/10.1097/01.JAC.0000314707.88160.9c>
- Moore, D. J., Poquette, A., Casaletto, K. B., Gouaux, B., Montoya, J. L., Posada, C., Rooney, A. S., Badice, J., Deutsch, R., Letendre, S. L., Depp, C. A., Grant, I., & Atkinson, J. H., & The HIV Neurobehavioral Research Program (HNRP) Group (2015). Individualized texting for adherence building (iTAB): Improving antiretroviral dose timing among HIV-infected persons with co-occurring bipolar disorder. *AIDS and Behavior*, 19(3), 459–471. <https://doi.org/10.1007/s10461-014-0971-0>
- Mui, A. C., Kang, S.-Y., Chen, L. M., & Domanski, M. D. (2003). Reliability of the geriatric depression scale for use among elderly Asian immigrants in the USA. *International Psychogeriatrics*, 15(3), 253–271. <https://doi.org/10.1017/S1041610203009517>
- National Institute on Aging (2020, September 23). *Telehealth: improving dementia care*. <https://www.nia.nih.gov/news/telehealth-improving-dementia-care>
- NCLER (2019). Language Access Rights: Tips for Advocates of Limited English Proficient Older Adults. Retrieved from [https://ncler.acl.gov/getattachment/Resources/Language-Access-Practice-Tip-\(1\).pdf.aspx?lang=en-US#:~:text=Over%20five%20million%20older%20adults,LEP%20older%20adults%20is%20growing](https://ncler.acl.gov/getattachment/Resources/Language-Access-Practice-Tip-(1).pdf.aspx?lang=en-US#:~:text=Over%20five%20million%20older%20adults,LEP%20older%20adults%20is%20growing)
- Patel, S., Jacobus-Kantor, L., Marshall, L., Ritchie, C., Kaplinski, M., Khurana, P. S., & Katz, R. J. (2013). Mobilizing your medications: An automated medication reminder application for mobile phones and hypertension medication adherence in a high-risk urban population. *Journal of Diabetes Science and Technology*, 7(3), 630–639. <https://doi.org/10.1177/193229681300700307>
- Robert, S. A., & Ruel, E. (2006). Racial segregation and health disparities between black and white older adults. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*, 61(4), S203–S211. <https://doi.org/10.1093/geronb/61.4.S203>
- Rodríguez-Fernández, J. M., Danies, E., Hoertel, N., Galanter, W., Saner, H., & Franco, O. H. (2022). Telemedicine readiness across medical conditions in a US national representative sample of older adults. *Journal of Applied Gerontology*, 41(4), 982–992. <https://doi.org/10.1177/07334648211056231>
- Rodríguez-Fernández, J. M., Danies, E., Martínez-Ortega, J., & Chen, W. C. (2017). Cognitive decline, body mass index, and waist circumference in community-dwelling elderly participants: Results from a nationally representative sample. *Journal of Geriatric Psychiatry and Neurology*, 30(2), 67–76. <https://doi.org/10.1177/0891988716686832>
- Rodriguez, J. A., Saadi, A., Schwamm, L. H., Bates, D. W., & Samal, L. (2021). Disparities in telehealth use among California patients with limited English proficiency: Study examines disparities in telehealth use among California patients with limited English proficiency. *Health Affairs*, 40(3), 487–495. <https://doi.org/10.1377/hlthaff.2020.00823>
- Ryskina, K. L., Shultz, K., Zhou, Y., Lautenbach, G., & Brown, R. T. (2021). Older adults' access to primary care: Gender, racial, and ethnic disparities in telemedicine. *Journal of the American Geriatrics Society*, 69(10), 2732–2740. <https://doi.org/10.1111/jgs.17354>



- Ryvicker, M., & Sridharan, S. (2018). Neighborhood environment and disparities in health care access among urban medicare beneficiaries with diabetes: A retrospective cohort study. *Inquiry: A Journal of Medical Care Organization, Provision and Financing*, 55, 0046958018771414. <https://doi.org/10.1177/0046958018771414>
- Schiaffino, M. K., Al-Amin, M., & Schumacher, J. R. (2014). Predictors of language service availability in US hospitals. *International Journal of Health Policy and Management*, 3(5), 259–268. <https://doi.org/10.1111/jgs.17354>
- Schifeling, C. H., Shanbhag, P., Johnson, A., Atwater, R. C., Koljack, C., Parnes, B. L., Vehar, M. M., Farrow, S. A., Phimphasone-Brady, P., & Lum, H. D. (2020). Disparities in video and telephone visits among older adults during the COVID-19 pandemic: Cross-sectional analysis. *JMIR Aging*, 3(2), e23176. <https://doi.org/10.2196/23176>
- Seo, J., Choi, B., Kim, S., Lee, H., & Oh, D. (2017). The relationship between multiple chronic diseases and depressive symptoms among middle-aged and elderly populations: Results of a 2009 Korean community health survey of 156,747 participants. *BMC Public Health*, 17(1), 1–10. <https://doi.org/10.1186/s12889-017-4798-2>
- Seto, E., Leonard, K. J., Cafazzo, J. A., Barnsley, J., Masino, C., & Ross, H. J. (2012). Mobile phone-based telemonitoring for heart failure management: A randomized controlled trial. *The Journal of Medical Internet Research*, 14(1), 844. <https://doi.org/10.1186/s12889-017-4798-2>
- Smith, A. (2014). *Older Adults and Technology Use*. <https://policycommons.net/artifacts/620165/older-adults-and-technology-use/1601368/>
- Thomson, M. D., & Hoffman-Goetz, L. (2009). Defining and measuring acculturation: A systematic review of public health studies with hispanic populations in the United States. *Social Science & Medicine*, 69(7), 983–991. <https://doi.org/10.1016/j.socscimed.2009.05.011>
- Trief, P. M., Izquierdo, R., Eimicke, J. P., Teresi, J. A., Goland, R., Palmas, W., Shea, S., & Weinstock, R. S. (2013). Adherence to diabetes self care for white, African-American and Hispanic American telemedicine participants: 5 year results from the IDEATel project. *Ethnicity & Health*, 18(1), 83–96. <https://doi.org/10.1080/13557858.2012.700915>
- Tsoh, J. Y., Sentell, T., Gildengorin, G., Le, G. M., Chan, E., Fung, L.-C., Pasick, R. J., Stewart, S., Wong, C., Woo, K., Burke, A., Wang, J., McPhee, S. J., & Nguyen, T. T. (2016). Healthcare communication barriers and self-rated health in older Chinese American immigrants. *Journal of Community Health*, 41(4), 741–752. <https://doi.org/10.1007/s10900-015-0148-4>
- United States Census Bureau. American Community Survey 5-year estimates 2013–2017: selected social characteristics in the United States. Retrieved from <https://data.census.gov/cedsci/table?q=dp02&tid=ACSDP5Y2019.DP02>
- van der Weegen, S., Verwey, R., Spreeuwenberg, M., Tange, H., van der Weijden, T., & de Witte, L. (2015). It's LiFe! Mobile and web-based monitoring and feedback tool embedded in primary care increases physical activity: A cluster randomized controlled trial. *The Journal of Medical Internet Research*, 17(7), e4579. <https://doi.org/10.2196/jmir.4579>
- Wakefield, B. J., Holman, J. E., Ray, A., Scherubel, M., Adams, M. R., Hillis, S. L., & Rosenthal, G. E. (2011). Effectiveness of home telehealth in comorbid diabetes and

- hypertension: A randomized, controlled trial. *TelemedicineJournal and e-Health*, 17(4), 254–261. <https://doi.org/10.2196/jmir.4579>
- Wallace, P. M., Pomery, E. A., Latimer, A. E., Martinez, J. L., & Salovey, P. (2010). A review of acculturation measures and their utility in studies promoting latino health. *Hispanic Journal of Behavioral Sciences*, 32(1), 38. <https://doi.org/10.1177/0739986309352341>
- Wilcock, A. D., Rose, S., Busch, A. B., Huskamp, H. A., Uscher-Pines, L., Landon, B., & Mehrotra, A. (2019). Association between broadband internet availability and telemedicine use. *JAMA Internal Medicine*, 179(11), 1580–1582. <https://doi.org/10.1001/jamainternmed.2019.2234>
- Wild, B., Eckl, A., Herzog, W., Niehoff, D., Lechner, S., Maatouk, I., Brenner, H., Müller, H., & Löwe, B. (2014). Assessing Generalized Anxiety Disorder in elderly people using the GAD-7 and GAD-2 scales: Results of a validation study. *The American Journal of Geriatric Psychiatry*, 22(10), 1029–1038. <https://doi.org/10.1016/j.jagp.2013.01.076>
- Wilkins, C. H., Friedman, E. C., Churchwell, A. L., Slayton, J. M., Jones, P., Pulley, J. M., & Kripalani, S. (2021). A systems approach to addressing COVID-19 health inequities. *NEJM Catalyst Innovations in Care Delivery*, 2(1). <https://doi.org/10.1056/CAT.20.0374>
- Wu, J.-M., Yu, H.-J., Ho, T.-W., Su, X.-Y., Lin, M.-T., & Lai, F. (2015). Tablet PC-enabled application intervention for patients with gastric cancer undergoing gastrectomy. *Computer Methods and Programs in Biomedicine*, 119(2), 101–109. <https://doi.org/10.1016/j.cmpb.2015.03.004>

## Author Biographies

**Jorge M. Rodríguez-Fernández**, MD, is an assistant professor in the Neurology Department at the University of Texas Medical Branch. He also serves as the associate chief medical information officer. His research is focused on aging, and predictive modelling. He is particularly interested in understanding what factors are linked to cognitive decline, and the impact of telemedicine during the COVID-19 pandemic.

**Nicolas Hoertel** is professor of Psychiatry at Paris Cité University, Corentin-Celton hospital at Issy-les-Moulineaux, and Institute of Psychiatry and Neuroscience of Paris (INSERM UMR\_S1266). His research focuses on epidemiology and pharmaco-epidemiology in the field of mental health, with an emphasis on old age psychiatry, psychiatric comorbidity, and suicide prevention.

**Hugo Saner** is a cardiologist and professor emeritus at the Medical Faculty University of Bern, former head of Cardiovascular Prevention, Rehabilitation and Sports Cardiology. He works as senior researcher at the ARTORG Center for Biomedical Engineering Research at the University of Bern and as an adjunct professor at the Institute for Social and Preventive Medicine at the University of Bern. His research focuses on eHealth and Telemedicine in the context of cardiovascular risk factors and diseases.

**Mukaila Raji** is Edgar Gnitzinger distinguished professor in Aging and Director of Geriatrics & Palliative Medicine at the University of Texas Medical Branch. He received his medical degree from the University of Ife in Nigeria and subsequently completed Internal Medicine Residency at Washington Hospital Center in Washington, DC, USA, a geriatric fellowship at Duke University in Durham, NC, USA, MSc in Pharmacology at the University of Alberta in Edmonton, Canada, and a Certificate in Medical & Health Care Management from Rice University in Houston, TX. His research focuses on Alzheimer's disease/related dementias, cognitive aging and functional outcomes, and geriatric pharmacology.