

Short-term association between air temperature and mortality in seven cities in Norway: A time series analysis.

Journal:	Scandinavian Journal of Public Health
Manuscript ID	SPUB2023-0092.R1
Manuscript Type:	Special issue: Climate change and health in the Nordic countries
Date Submitted by the Author:	26-Jan-2024
Complete List of Authors:	Vázquez Fernández, Liliana; Norwegian Institute of Public Health; University of Oslo, Department of Biostatistics, Institute of Basic Medical Sciences Diz-Lois Palomares, Alfonso; Norwegian Institute of Public Health; University of Oslo, Department of Mathematics Vicedo Cabrera, Ana María; University of Bern Institute of Social and Preventive Medicine; University of Bern, Oeschger Center of Climate Change Research Freiesleben de Blasio, Birgitte; Norwegian Institute of Public Health; University of Oslo, Department of Biostatistics, Institute of Basic Medical Sciences, Centre for Biostatistics and Epidemiology Di Ruscio, Francesco; Norwegian Institute of Public Health Wisløff, Torbjørn; Akershus University Hospital, Health Services Research Unit; University of Oslo, Institute of Clinical Medicine Rao, Shilpa; Norwegian Institute of Public Health
Problem Areas and Research:	Environmental health, Mortality/Morbidity
Discipline:	Epidemiology
Methodology:	Quantitative longitudinal

SCHOLARONE[™] Manuscripts

Title page

Title: Short-term association between air temperature and mortality in seven cities in Norway: A time series analysis.

Authors: Liliana Vázquez Fernández^{1,2}, Alfonso Diz-Lois Palomares^{1,3}, Ana Maria Vicedo Cabrera^{4,5}, Birgitte Freiesleben de Blasio^{1,6}, Francesco Di Ruscio¹, Torbjørn Wisløff^{7,8}, Shilpa Rao¹

Authors affiliations:

1 Norwegian Institute of Public Health, Oslo, Norway

2 Department of Biostatistics, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

3 Department of Mathematics, University of Oslo, Oslo, Norway

4 Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland

5 Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

6 Oslo Center for Biostatistics and Epidemiology, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

7 Health Services Research Unit, Akershus University Hospital, Lørenskog, Norway

8 Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Corresponding author: Liliana Vázquez Fernández, Postboks 222 Skøyen, Oslo, NO 0213.

Email address: lilianavazquezfernandez@gmail.com

Wordcount:

Abstract: 249

Manuscript:2999

Abstract

Background: The association between ambient air temperature and mortality has not been assessed in Norway. This study aimed to quantify for seven Norwegian cities (Oslo, Bergen, Stavanger, Drammen, Fredrikstad, Trondheim and Tromsø) the non-accidental, cardiovascular, and respiratory diseases mortality burden due to non-optimal ambient temperatures.

Methods: We used a historical daily dataset (1996-2018) to perform city-specific analyses with a distributed lag non-linear model with 14 days of lag, and pooled results in a multivariate metaregression. We calculated attributable deaths for heat and cold, defined as days with temperatures above and below the city -specific optimum temperature. We further divided temperatures into moderate and extreme using cut-offs at the 1st and 99th percentiles.

Results: We observed that 5.3% (95%CI 2.0-8.3) of the non-accidental related deaths, 11.8% (95%CI 6.4-16.4) of the cardiovascular and 5.9% (95%CI -4.0-14.3) of the respiratory were attributable to nonoptimal temperatures. Notable variations were found between cities and subgroups stratified by sex and age. The mortality burden related to cold dominated in all three health outcomes (5.1%, 2.0-8.1, 11.4%, 6.0-15.4, and 5.1%, -5.5-13.8 respectively). Heat had a more pronounced effect on the burden of respiratory deaths (0.9%, 0.2-1.0). Extreme cold accounted for 0.2% of non-accidental deaths, and 0.3% of cardiovascular and respiratory deaths, while extreme heat contributed to 0.2% of nonaccidental to 0.3% of respiratory deaths.

Conclusions: Most of the burden could be attributed to the contribution of moderate cold. This evidence has significant implications for enhancing public-health policies to better address health consequences in the Norwegian setting.

Keywords: Climate

Norway

Cold Temperature / adverse effects* Hot Temperature / adverse effects* Cardiovascular Diseases / mortality Respiratory Tract Diseases / mortality

Humans

Mortality*

Risk Assessment / methods

Introduction

Norway is experiencing increasing temperatures and evident effects of extreme weather events (1, 2). Particularly, the Artic region has seen a steep increase, warming three times faster than the global average (3). The country has distinct differences between urban and non-urban areas and a wide range of climatic zones, with its population steadily moving to the cities and gradually ageing (4). This suggests a heightened vulnerability to climate change as older populations are particularly susceptible to thermal stress. Consequently, there is a pressing need to comprehend the effects of temperature on mortality.

The leading cause of death in Norway is ischaemic heart disease, with 104.53 deaths per 100 000 population in 2018, followed by dementia and chronic obstructive pulmonary disease (COPD) (5). One in five Norwegians live with a confirmed cardiovascular disease (CVD) or are at a high risk of developing one (6), and around 6% of the population over 40 years of age has been diagnosed with COPD (7).

CVD and respiratory diseases are particularly affected by non-optimal temperature. Exposure to cold air temperature compromises the pulmonary mechanics, triggers vasoconstriction and may lead to an increase in cardiovascular and respiratory mortality (8, 9). Hot temperatures trigger a widening of blood vessels (vasodilation), the rate at which blood pumps increases, and sweating, which may originate in dehydration, and a series of pathological events that can result in a cardiovascular impairment (10) and induce airways inflammation or bronchoconstriction (11).

Specific research on the effects of extreme temperatures on mortality in Nordic cities is still limited. Some examples of recent studies in Nordic include Åström et al. (12), Orru et al. (13) and Fonseca et al. (14). Few studies on heat effects in Sweden and Finland have found significant excess morbidity and mortality, especially among older people, and mainly attributable to cardiovascular and respiratory diseases (15-17). According to (16), resilience to heat may have decreased over the last two decades, along with the increase in temperature and ageing population.

In this study, we aim to assess the temperature related mortality burden in seven urban municipalities in Norway for different causes of death in the 1996-2018 period. We focus on non-accidental and cause-specific mortality due to cardiovascular and respiratory diseases and stratify by sex and groups of age (18). To our knowledge, this is the first study of its kind in Norway.

Data

We chose the seven largest cities in Norway by their population in 2018. These are in decreasing order: Oslo (673 469 inhabitants), Bergen, Trondheim, Stavanger, Drammen, Fredrikstad and Tromsø (75 638). The cities represent three different climates in the Köppen-Geiger climate classification: Humid continental, oceanic and subarctic (19). See Table S1 for more details.

For the temperature exposures at the city level, we used a high-resolution observational dataset of daily temperature for Norway, which is available at a 1 km resolution and is continuously updated (20). Making use of the geographical borders of each city, we averaged the mean daily temperatures for each city from 1st January 1996 to 31st December 2018.

We also used mean relative humidity (24-hour) data provided by the Norwegian Centre for Climate Services (NCCS) (21) and air pollution data, in particular measurements of ozone (O_3), particular matter with a diameter of 10 µm and 2.5 µm or less (PM_{10} and $PM_{2.5}$) and nitrogen dioxide (NO_2) at a 1 km resolution. Pollution measurements for all cities come from the Nordic DEHM-UBM (Danish Eulerian Hemispheric Model- Urban Background Model) setup. All pollutants were computed as 24-hour averages but O_3 , calculated as the daily maximum 8-hour running average from hourly measurements.

Concentrations of pollutants were weighted according to the population grid GHS 2019A (GHS-POP R2019A: GHS population grid multitemporal (1975-1990-2000-2015) https://ghsl.jrc.ec.europa.eu/ghs_pop2019.php), utilizing the year 2015 and averaging the daily values within the municipal borders as for the temperature exposure.

The health data comprised daily mortality counts for non-accidental, cardiovascular and pulmonary mortality (International Classification of Diseases, ICD-10: A00-R99, I00-I99 and J00-J99 respectively) over the study period of 1996 to 2018, provided by the Norwegian Death Registry (22). The data was also disaggregated by age groups (0-74, 75+) and sex.

Methods

We used a two-stage methodology. We first analysed the relationship between the mortality outcome and the average mean daily temperature using a quasi-Poisson distributed lag non-linear model for each city to derive estimates of city-specific temperature-mortality associations, reported as relative risk (RR). The first-stage regression included a natural cubic B-spline of time with 8 degrees of freedom per year to control for seasonal and long-term trends, and indicators for the day of the week and bank holidays. The temperature was modelled with a quadratic B-spline with 3 internal knots placed at 10th, 75th, and 90th percentiles of location-specific distributions and the lag-response curve with a cubic Bspline with an intercept and 3 knots placed at equally spaced values in the log scale. We opted to extend the lag period to 14 days (14) instead of 21 (23) in order to preserve statistical power and account for any harvesting effect, while also capturing the delayed impact of cold temperatures. This parametrization is commonly used (23) and yielded the final model:

No. daily deaths ~ intercept + cb(Avg.Temp) + ns(time) + holidays + weekday

In the second stage, we pooled the city-specific relative risks using a multivariate meta-regression model with the city-specific range of temperature as a covariate and fixed effects to obtain the overall pooled estimate. Then we calculated the best linear unbiased predictors (BLUPs) (24), which could provide more precise estimations, particularly for the cities with smaller numbers of deaths. We next derived the minimum mortality temperature (MMT) in each city from the BLUP of the overall cumulative association between temperature and each mortality outcome. We referred to the minimum mortality temperature as the optimum temperature and used it as the reference for calculating the attributable risk (23). The quadratic B-spline of temperature in the first stage analysis was then re-centred according to each city-specific MMT to obtain accurate risk estimates at a given temperature. We explored heterogeneity with the I² statistic and the Cochrane's Q test.

We conducted sensitivity analyses to explore the impact of different modelling choices on nonaccidental mortality, while accounting for air pollution and area-level vulnerability factors, such as the proportion of the population aged 65 years and above. In the first stage models, pollutants were included as daily covariates, whereas the vulnerability factors were incorporated in the second stage as 2018 annual values. Additional results and further details can be found in the Supplementary Material (Table S2 and Methods S1).

The overall effect of temperature on a specific day on the RR of death was defined as the accumulation of the risk during the lag period (14 days). We calculated the attributable number of deaths and fractions of total deaths using the overall cumulative RR corresponding to each day's temperature in the next 14 days, and the respective empirical confidence intervals using Monte Carlo simulations assuming a normal distribution of the BLUPs of the reduced coefficients (25), using a previously described and applied method (23, 26). The sum of the contributions gives the total number of

attributable deaths due to non-optimal temperatures and the ratio with the total number of deaths gives the total attributable fraction (AF).

We aggregated the contributions from the days below and above the point of MMT to calculate the AF due to cold and heat. Additionally, we classified AFs into moderate and extreme contributions by defining extreme temperatures as those falling below the 1st and exceeding the 99th location-specific percentiles.

We report pooled effects, defined as the cold and heat effects, represented by the RR at the 1st and 99th percentiles of the exposure-response association relative to the MMT, accompanied by 95% confidence intervals, for each outcome and subgroup. The model was separately fitted for sex and the two age groups: 0-74 years, and 75 years or older, based on cause of death.

All analyses were performed in software R version 4.1.0 using the packages *data.table, splines, dlnm* for the first stage and *mixmeta* for the second stage meta-analysis.

Results

The analysis included 230 979 non-accidental deaths, of which 83 051 were from cardiovascular causes and 24 217 were attributed to respiratory diseases. The average daily deaths from non-accidental mortality spanned from 12 in Oslo to 1 in Tromsø. The populations of the different cities are subject to different ranges of temperatures characterised by the specific climates (Table 1).

The overall exposure-response functions (ERF) curves (best linear unbiased predictions) presented in Figure 1 for non-accidental, cardiovascular, and respiratory mortality demonstrate risk increases for temperatures below and above the MMT, except for cardiovascular-related mortality above the MMT and the oceanic cities below the MMT for cardiovascular and respiratory disease mortality. Remarkably, oceanic cities exhibit a stronger effect with decreasing temperatures for non-accidental mortality. Additionally, at higher temperatures, the respiratory ERFs yield higher risks than the non-accidental ERFs in all cities.

We report the pooled predicted overall cumulative association for the seven cities in Figure 2, denoted as the RR at the 1st and 99th percentiles. These percentiles represent the extreme temperature thresholds that define the excess mortality attributed to non-optimal temperatures, specifically cold and heat effects. Regarding non-accidental mortality, statistically significant effects were observed in the entire population, as well as in women and the age group 75+. Both the female group and the oldest age group, 75+, exhibited a similar pattern with statistically significant associations between non-optimal weather conditions and mortality. This suggests that they are the most vulnerable among the groups analysed, including the effects of cold on CVD mortality and the effects of heat on pulmonary mortality.

In contrast, the 0-74 age group exhibited only one significant pooled effect concerning non-accidental mortality attributed to low temperatures. Remarkably, significant associations were observed solely for cold effects concerning CVD mortality, while heat effects were linked to respiratory disease mortality. Conversely, no significant pooled effects were found for males.

By city, statistically significant heat effects were identified in all cities except Bergen and Stavanger for non-accidental and pulmonary mortality in the 75+, women, and overall groups. Concerning cardiovascular mortality, we observed only one significant heat effect in Oslo for women. A similar pattern emerged for cold effects in the non-accidental and cardiovascular groups. Notably, all cities except Oslo yielded a significant cold effect in the non-accidental women's group. Additionally, the younger group exhibited significant cold effects related to pulmonary mortality in Oslo, Trondheim, and Drammen.

The largest significant heat effect was found in Trondheim, with a RR of 1.6 (95%CI 1.2-2.2) for the females, whereas for cold effects, Oslo had the highest impact with a RR of 1.9 (95%CI 1.2-3.0) in the younger group, both associated with respiratory mortality. In contrast, there were no statistically significant effects in any of the male groups and the younger cardiovascular group. All outcomes are available in Table S4 of the Supplementary Material.

The overall non-accidental mortality AF due to non-optimal temperatures was 5.3% (95%CI 2.0-8.3), which translated into 12 286 deaths in 23 years (1996-2018). Of these deaths 11 729 were attributed to cold temperatures and 557 to heat. Examining causes of death, overall AF for CVD mortality was 11.8% (95%CI 6.4-16.4) and 5.9% (95%CI -4.0-14.3) for respiratory mortality. Cold was responsible for most of the mortality burden (AF for non-accidental 4.9% (95%CI 1.7-7.8), cardiovascular 11.4% (95%CI 6.0-15.4) and respiratory deaths 5.1% (95%CI -5.5-13.8)).

The city-specific attributable fractions, further itemised into extreme and moderate heat and cold components, are presented in Figure 3. Additionally, results categorised by sex and age groups can be found in the Supplementary Material (Tables S5-S8 and Figures S7-S12). Extreme temperatures were associated with relatively smaller risks portions compared to moderate temperatures, aligning with the ERFs depicted in Figure 1 and the city-specific temperature distributions.

The MMTs shifted to higher temperatures in the CVD groups compared to non-accidental and respiratory in most of the cities. These corresponded to an average minimum mortality percentile (MMP) per cause, rising from 70 to 94 for non-accidental, 80 to 99 for cardiovascular diseases, and 49 to 91 for respiratory diseases mortality. Consequently, this explains that the absence of heat effects in cardiovascular diseases. MMTs are detailed per cause, group, and city in Tables S6, S7 and S8 of the Supplementary Material.

We also plotted the joint distributed lag graphs using a maximum lag of 14 days for the RRs comparing the effects of extreme cold and heat to the MMT on both non-accidental and cause-specific mortality. Cold effects peaked in general on the first day (lag 1) and lasted longer, while the heat effects were immediate. Cold effects were significant for non-accidental and respiratory mortality, and heat effects were statistically significant for all three causes of mortality in females (Figures S13-S17 of the Supplementary Material).

Our sensitivity analyses suggested that our results were consistent and independent of modelling assumptions, also when controlled for PM_{25} , PM_{10} , NO_2 , ozone, humidity, and vulnerability factors. These results are shown in Tables S2 and S3 of the Supplementary Material.

Discussion

We examined the associations between exposure to air temperature and mortality due to nonaccidental, cardiovascular, and respiratory diseases in Norwegian urban areas from 1996-2018. Additionally, we analysed these associations in different subgroups, stratifying by age and sex. to ensure the accuracy and integrity of our results, we also examined climatic and geographical characteristics as potential confounders and selectively included only those variables that acted as confounders in our analyses, prioritising their impact. Our findings indicate that non-optimal temperatures were associated with an increased mortality risk for all the causes and subgroups analysed. Particularly moderate cold temperatures were responsible for a higher proportion of the mortality burden in the seven Norwegian cities. We estimated higher relative risks for colder

temperatures in contrast to warmer temperatures, except within the pulmonary groups: overall, those aged 75 and above, and men, for which the relative risks for hotter temperatures were slightly higher. In an Estonian study by Orru et al. (13), greater effects of heat were noted compared to cold for mortality due to external causes, whereas Aström DO et al. (12) in the Baltics found no association between respiratory disease mortality and cold effects but identified an adverse effect with cold temperatures on CVD mortality. In contrast, our findings reveal more pronounced cold and heat effects for respiratory mortality compared to CVD mortality, with the exception of males experiencing cold effects. Additionally, only the heat effect for non-accidental mortality in the younger group was larger than the effect on respiratory related mortality. Changes linked to age have been associated with physiological differences in thermoregulation, which could account for differences in our results (9). Gender differences are evident in our analysis, with women displaying greater susceptibility to temperature-related mortality. However, it is worth considering that the observed gender disparities could be influenced by various additional factors, such as the social context (15). These results in which the estimated pooled overall heat effects are larger for respiratory than CVD mortality, are consistent with other studies (27). In our study, we did not identify a significant increase in cardiovascular mortality due to heat, similarly to this multi-city study (28); however, another found a clear high effect in southern Swedish locations and not in the northern ones (29).

Consisting with other studies (27), we found that cold effects lasted longer than heat effects. On average, the effect of cold temperatures peaked at lag one day and lasted almost ten days, while heat effects appeared immediately and lasted two or three days, followed by a mortality displacement.

We found that temperature is responsible for a substantial fraction of 5.3% of the non-accidental mortality burden. This percentage translates to 469 deaths in 2018 alone for the seven cities. Other studies have reported similar mortality burden due to cold temperatures, and in similar latitudes to the overall we describe. Åström et al. (30) recount that temperature is responsible for 4.5% in Stockholm, Sweden for all-cause mortality in the period 1990-2002 and Gasparrini et al. (23) 3.7% for the same period. Another study conducted in Tallin and Riga reported AF values of 7.4% and 8.3% respectively, although only November to March months were analysed from the years 1997-2015 and 2009-2015 (12). The former finding aligns more closely with our results for non-accidental mortality in Oslo, Trondheim and Drammen (7%, 6.8% and 6.2% respectively). Achebak et al. (18) documented an overall AF of 13.1% for CVD mortality which is comparable to our own findings of 11.8% (6.4-16.4).

Our analysis had several strengths. As the first study to focus exclusively on Norwegian cities, it captures their distinctive idiosyncrasy stemming from subpolar latitudes and comparatively milder temperatures than Russia or Canada. Our results provide novel and robust evidence on mortality risk and burden from both non-accidental causes and cardiopulmonary diseases. Another considerable strength of our data is the completeness of Norwegian registries, with close to 100% registrations of data such as residence addresses and time of death. Additionally, the reliability of the Norwegian death certification system bolsters our findings' credibility, supported by careful diagnosis and cause attribution procedures.

We acknowledge some limitations inherent to our study. Our scope centred on urban populations. Similar to numerous ecological epidemiological studies, assessing individual exposures remained unfeasible. The small number of daily deaths in some locations, particularly for respiratory mortality, produced wide confidence intervals and thus low precision estimates. Therefore, the reported effects should be interpreted cautiously, particularly the cold effects as we have not considered the possible confounding during the cold season for influenza activity. Lastly, an underlying assumption was made that children's deaths attributable to temperature are exceedingly scarce, thereby exerting minimal influence on our findings.

Conclusions

This study confirmed that non-optimal temperatures are associated with increased mortality for nonaccidental and cardiopulmonary-related causes in Norway. We further identified that moderate temperatures are responsible for most deaths, and women and the elderly are the most vulnerable groups. Further research is needed to delimit the excess mortality that could be preventable. These results could contribute to improving public health preparedness and response measures for both cold and hot seasons.

caus . deatns, . to delimit the .ing public health .

Acknowledgements: All authors acknowledge funding from European Union's Horizon 2020 research and innovation programme under grant agreement No 820655 (EXHAUSTION). We extend our acknowledgment to Camilla Geels and Lise Marie Frohn of the University of Aarhus for providing the DEHM air pollution data pertaining to Norway. We are also deeply grateful to the reviewers whose insightful comments greatly contributed to the enhancement of this paper.

Mari. uorway. W .ed to the enh.

References

1. Ipcc. Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C, Berger S, et al., editors. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2021.

2. Romanello M, McGushin A, Di Napoli C, Drummond P, Hughes N, Jamart L, et al. The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future. Lancet (London, England). 2021;398(10311):1619-62.

3. Vikhamar-Schuler D, Isaksen K, Haugen JE, Tømmervik H, Luks B, Schuler TV, et al. Changes in Winter Warming Events in the Nordic Arctic Region. Journal of Climate. 2016;29(17):6223-44.

4. Health Nutrition and Population Statistics: Population estimates and projections. The World Data Bank. [Internet]. [cited 10/11/2022]. Available from: https://databank.worldbank.org/home.aspx.

5. Global Health Estimates 2020: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2019. Geneva, World Health Organization; 2020 [Internet]. [cited 10/11/2022]. Available from: https://www.who.int/data/gho/data/themes/mortality-and-global-health-estimates/ghe-leading-causes-of-death.

6. Ariansen IKH, Kvåle, R., Olsen, K., & Selmer, R. M. . Hjerte- og karsykdommer i Norge. Norwegian Institute of Public Health; 2021 26.11.2021.

7. Health NIOP. Chronic Obstructive Pulmonary Disease (COPD) in Norway. 2018.

8. Ryti NR, Guo Y, Jaakkola JJ. Global Association of Cold Spells and Adverse Health Effects: A Systematic Review and Meta-Analysis. Environmental health perspectives. 2016;124(1):12-22.

9. Schneider A, Rückerl R, Breitner S, Wolf K, Peters A. Thermal Control, Weather, and Aging. Current Environmental Health Reports. 2017;4(1):21-9.

10. Liu J, Varghese BM, Hansen A, Zhang Y, Driscoll T, Morgan G, et al. Heat exposure and cardiovascular health outcomes: a systematic review and meta-analysis. The Lancet Planetary Health. 2022;6(6):e484-e95.

11. Miao J, Feng S, Wang M, Jiang N, Yu P, Wu Y, et al. Life-time summer heat exposure and lung function in young adults: A retrospective cohort study in Shandong China. Environment international. 2022;160:107058.

12. Åström DO, Veber T, Martinsone Ž, Kaļužnaja D, Indermitte E, Oudin A, et al. Mortality Related to Cold Temperatures in Two Capitals of the Baltics: Tallinn and Riga. Medicina (Kaunas). 2019;55(8).

13. Orru H, Åström DO. Increases in external cause mortality due to high and low temperatures: evidence from northeastern Europe. International journal of biometeorology. 2017;61(5):963-6.

14. Fonseca-Rodríguez O, Lundevaller EH, Sheridan SC, Schumann B. Association between Weather Types based on the Spatial Synoptic Classification and All-Cause Mortality in Sweden, 1991⁻2014. International journal of environmental research and public health. 2019;16(10).

15. Astrom DO, Forsberg B, Rocklov J. Heat wave impact on morbidity and mortality in the elderly population: a review of recent studies. Maturitas. 2011;69(2):99-105.

16. Oudin Åström D, Schifano P, Asta F, Lallo A, Michelozzi P, Rocklöv J, et al. The effect of heat waves on mortality in susceptible groups: a cohort study of a mediterranean and a northern European City. Environmental Health. 2015;14(1):30.

17. Åström DO, Tornevi A, Ebi KL, Rocklöv J, Forsberg B. Evolution of Minimum Mortality Temperature in Stockholm, Sweden, 1901-2009. Environmental health perspectives. 2016;124(6):740-4.

18. Achebak H, Devolder D, Ballester J. Trends in temperature-related age-specific and sexspecific mortality from cardiovascular diseases in Spain: a national time-series analysis. The Lancet Planetary health. 2019;3(7):e297-e306.

19. Beck HE, Zimmermann NE, McVicar TR, Vergopolan N, Berg A, Wood EF. Publisher Correction: Present and future Köppen-Geiger climate classification maps at 1-km resolution. Scientific Data. 2020;7(1):274.

20. Lussana C, Tveito OE, Dobler A, Tunheim K. seNorge_2018, daily precipitation, and temperature datasets over Norway. Earth Syst Sci Data. 2019;11(4):1531-51.

21. Norwegian Centre for Climate Services. [Internet]. [cited 10/10/2022]. Available from: https://seklima.met.no/.

22. Resgistry NCoD. Norwegian Institution of Public Healfh. Norwegian Cause of Death Registry. https://wwwfhino/en/hn/health-registries/cause-of-death-registry/.

23. Gasparrini A, Guo Y, Hashizume M, Lavigne E, Zanobetti A, Schwartz J, et al. Mortality risk attributable to high and low ambient temperature: a multicountry observational study. Lancet (London, England). 2015;386(9991):369-75.

24. Sera F, Armstrong B, Blangiardo M, Gasparrini A. An extended mixed-effects framework for meta-analysis. Stat Med. 2019;38(29):5429-44.

25. Tobías A, Armstrong B, Gasparrini A. Brief Report: Investigating Uncertainty in the Minimum Mortality Temperature: Methods and Application to 52 Spanish Cities. Epidemiology (Cambridge, Mass). 2017;28(1):72-6.

26. Gasparrini A, Leone M. Attributable risk from distributed lag models. BMC Med Res Methodol. 2014;14:55.

27. Iñiguez C, Royé D, Tobías A. Contrasting patterns of temperature related mortality and hospitalization by cardiovascular and respiratory diseases in 52 Spanish cities. Environmental research. 2021;192:110191.

28. Baccini M, Biggeri A, Accetta G, Kosatsky T, Katsouyanni K, Analitis A, et al. Heat Effects on Mortality in 15 European Cities. Epidemiology (Cambridge, Mass). 2008;19(5):711-9.

29. Fonseca-Rodríguez O, Sheridan SC, Lundevaller EH, Schumann B. Hot and cold weather based on the spatial synoptic classification and cause-specific mortality in Sweden: a time-stratified case-crossover study. International journal of biometeorology. 2020;64(9):1435-49.

30. Oudin Åström D, Ebi KL, Vicedo-Cabrera AM, Gasparrini A. Investigating changes in mortality attributable to heat and cold in Stockholm, Sweden. International journal of biometeorology. 2018;62(9):1777-80.

Table 1. Descriptive statistics by city.

City	Total Non- accidental deathsTotal CVD deathsDaily Avg.(range)Total CVD deaths		Total Resp. deaths Daily Avg. (range)	Temperatures Daily Avg. (1 st – 99 th percentiles)
Oclo	99 123	35 516	10 581	5.27 (-13.42 -
USIO	12 (1 - 29)	4 (0 - 16)	1 (0 - 8)	20.35)
Porgon	43 526	15 424	4 380	6.48 (-7.02 -
Bergen	5 (0 - 16)	2 (0 - 9)	1 (0 - 6)	19.00)
Trandhaim	27 394	9 895	2 801	4.68 (-13.89 -
Ironaneim	3 (0 - 11)	1 (0 - 8)	0 (0 - 6)	19.35)
Chavanaar	19 401	6 602	2 054	7.94 (-5.07 -
Stavanger	2 (0 - 10)	1 (0 - 6)	0 (0 - 4)	44977)
Drammon	17 437	6 514	1 863	5.39 (-13.18 -
Drammen	2 (0 - 10)	1 (0 - 5)	0 (0 - 3)	20.17)
Fue duileete d	15 915	6 210	1 682	7.44 (-10.42 -
Fredrikstad	2 (0 - 8)	1 (0 - 6)	0 (0 - 3)	21.13)
Troport	8 183	2 890	856	1.22 (-15.49 -
romsø	1 (0 - 6)	0 (0 - 4)	0 (0 - 2)	16.15)





Figure 1. Overall cumulative exposure–response associations for non-accidental and cardiopulmonary mortality in seven cities and the respective pooled association. Exposure–response associations are showed as best linear unbiased predictions. RR=Relative risk.



Figure 2. Pooled effects for non-accidental, cardiovascular, and respiratory mortality are reported at the 1st and 99th percentiles as cold and heat effects, respectively, with 95% confidence intervals. These effects are stratified by sex and age group.



Figure 3. Proportion of non-accidental and cardiopulmonary mortality attributed to moderate and extreme hot and cold temperatures across different cities. Extreme and moderate high and low temperatures were determined using the minimum mortality temperature at the 1st and 99th percentiles of temperature as thresholds.







Supplementary Material

For Peer Review Only

3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
20	
27	
20	
30	
21	
22	
2∠ 22	
22 24	
54 25	
22	
30	
3/	
38	
39	
40	
41	
42	
43	
44	
45	
46	
47	
48	
49	
50	
51	
52	
53	
54	
55	
56	
57	
58	
59	
60	

Short-term association between air temperature and mortality in seven cities in Norway: A time series analysis.

Liliana Vázquez Fernández^{1,2}, Alfonso Diz-Lois Palomares^{1,3}, Ana Maria Vicedo Cabrera^{4,5}, Birgitte Freiesleben de Blasio^{1,6}, Francesco Di Ruscio¹, Torbjørn Wisløff^{7,8}, Shilpa Rao¹

- 1 Norwegian Institute of Public Health, Oslo, Norway
- 2 Department of Biostatistics, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway
- 3 Department of Mathematics, University of Oslo, Oslo, Norway
- 4 Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland
- 5 Oeschger Center for Climate Change Research, University of Bern, Bern, Switzerland

6 Oslo Center for Biostatistics and Epidemiology, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway

Perez on

7 Health Services Research Unit, Akershus University Hospital, Lørenskog, Norway

8 Institute of Clinical Medicine, University of Oslo, Oslo, Norway

Table of content

Methods

Methods S1: First stage sensitivity analysis

Tables

Table S1: Extended descriptive statistics

Table S2. First stage sensitivity analysis.

 Table S3. Second-stage meta-regression models.

Table S4. The relative risks of cold temperature and hot temperature on deaths at lags of 0–14 days in the seven cities.

Table S5. Attributable fractions on non-accidental and cardiopulmonary deaths in the seven cities for cold and heat.

Table S6. Attributable fractions on non-accidental, cardiovascular and respiratory deaths in the seven cities for cold and heat.

Table S7. Attributable fractions on non-accidental, cardiovascular and respiratory deaths by sex in the seven cities for moderate cold and heat.

Table S8. Attributable fractions on non-accidental, cardiovascular and respiratory deaths by age group in the seven cities for moderate cold and heat.

Table S9. Attributable number of deaths (AN) for non-accidental, cardiovascular, and respiratory disease mortality by sex and age group in each city.

Table S10. Pooled attributable fractions and number of deaths (AN) for non-accidental, cardiovascular, and respiratory disease mortality.

Figures

Figure S1. Overall cumulative exposure–response associations for non-accidental mortality in seven cities by sex.

Figure S2. Overall cumulative exposure-response associations for CVD mortality in seven cities by sex.

Figure S3. Overall cumulative exposure–response associations for respiratory disease mortality in seven cities by sex.

Figure S4. Overall cumulative exposure–response associations for non-accidental mortality in seven cities by age group.

Figure S5. Overall cumulative exposure–response associations for CVD mortality in seven cities by age group.

Figure S6. Overall cumulative exposure–response associations for respiratory disease mortality in seven cities by age group.

Figure S7. Fractions of non-accidental mortality attributable to moderate and extreme hot and cold temperature by city and age group.

Figure S8. Fractions of cardiovascular mortality attributable to moderate and extreme hot and cold temperature by city and age group.

Figure S9. Fractions of respiratory disease mortality attributable to moderate and extreme hot and cold temperature by city and age group.

Figure S10. Fractions of non-accidental mortality attributable to moderate and extreme hot and cold temperature by city and sex.

Figure S11. Fractions of cardiovascular mortality attributable to moderate and extreme hot and cold temperature by city and sex.

Figure S12. Fractions of respiratory disease mortality attributable to moderate and extreme hot and cold temperature by city and sex.

Figure S13. Overall lag structure in effects of extreme cold temperature on daily cause specific mortality in seven Norwegian cities, 1996-2018, by cause of death.

Figure S14. Overall lag structure in effects of extreme hot temperature on daily cause specific mortality in seven Norwegian cities, 1996-2018, by cause of death.

Figure S15. Overall lag structure in effects of extreme cold and hot temperature on daily non-accidental mortality in seven Norwegian cities, 1996-2018, by sex.

Figure S16. Overall lag structure in effects of extreme cold and hot temperature on daily cardiovascular mortality in seven Norwegian cities, 1996-2018, by sex.

Figure S17. Overall lag structure in effects of extreme cold and hot temperature on daily respiratory disease mortality in seven Norwegian cities, 1996-2018, by sex.

Figure S18. Directed Acyclic Graph (DAG)

References



Methods S1: First stage sensitivity analysis

An initial sensitivity analysis was performed on the non-accidental mortality models, where we checked the city-specific average and range temperature as meta-predictors. We further considered two modelling frameworks: fixed-effects and random-effects (restricted maximum likelihood) metaanalysis and sorted them according to their Akaike Information Criteria (AIC). The model with the lowest AIC was invariably the one with no covariates, followed by the temperature range singlepredictor model. We also varied the choice of lag from 1 to 28 days and internal knots of the temperature distribution from 2 to 4. We chose a lag period of 14 days and three internal knots to model the exposure-response curves based on previous studies (1). Table S2 contains results from the tests of these modelling choices for this sensitivity analysis.

With regards to the cut-offs for defining the extreme temperatures, we did not choose the 2.5th and 97.5th as the minimum mortality temperature in colder weathers is normally very high, in the range of the 90ths' percentiles (averaged 93rd in non-accidental mortality). In temperate regions the temperature percentile of minimum mortality varies from about the 80th to the 90th percentile (1).

Ren Review Only

http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Table S1. Extended descriptive statistics

City	Köppen climate classification	Latitude	Population (2018)	Total Non- accidental	Total CVD No.	Total Resp. No.	Min. Temp	Avg. Temp.	Max. Temp.	1 st perc. Temp	99 th perc. Temp
				NO.							
Oslo	Humid continental	59.98	673 469	99 123	35 516	10 581	-22.41	5.27	26.02	-13.42	20.35
Bergen	Oceanic	60.36	279 792	43 526	15 424	4 380	-13.95	6.48	24.06	-7.02	19.00
Trondheim	Humid continental	63.37	199 595	27 394	9 895	2 801	-23.07	4.68	24.4	-13.89	19.35
Stavanger	Oceanic	59.14	141 186	19 401	6 602	2 054	-10.52	7.94	26.19	-5.07	20.02
Drammen	Humid continental	59.71	100 302	17 437	6 514	1 863	-21.69	5.39	24.67	-13.18	20.17
Fredrikstad	Humid continental	59.19	80 977	15 915	6 210	1 682	-17.43	7.44	26.13	-10.42	21.13
Tromsø	Subarctic	69.74	75 638	8 183	2 890	856	-20.78	1.22	22.73	-15.49	16.15
		Average:	221 566	32 997	11 864	3 460	-18.55	5.49	24.89	-11.21	19.45
		Source:	Statistics Bureau of Norway	Cause of	Death Re	gistry	Norwegi	an Meteor	ological Ins	titute (MET	Norway)

Table S2. Sensitivity analysis.

Computed on the fraction (%) attributable to temperature (total, heat, and cold components), by varying modelling choices, fitting the models to non-accidental mortality, and controlling for PM25, PM10, NO2, ozone, humidity, and vulnerability factors. Brackets indicate 95% empirical confidence intervals.

	Total (%)	Cold (%)	Heat (%)
Main model	5.32 (1.99-8.25)	5.08 (2.04-8.10)	0.24 (0.13-0.33)
Knots for exposure-response: 10th and 90th	5.21 (2.26-7.95)	4.97 (2.09-7.79)	0.24 (0.12-0.35)
Knots for exposure-response: 10th, 50th, and 90th	5.32 (2.05-8.29)	4.98 (1.84-7.68)	0.34 (-0.08-0.73)
Knots for exposure-response: 10th, 50th, 75th and 90th	5.48 (2.54-8.34)	5.12 (2.13-8.15)	0.36 (-0.12-0.77)
Lag period: 7 days	4.76 (2.48-6.89)	4.45 (2.23-6.67)	0.32 (0.14-0.49)
Lag period: 14 days	5.32 (1.99-8.25)	5.08 (2.04-8.10)	0.24 (0.13-0.33)
Lag period: 21 days	7.82 (3.11-11.84)	7.64 (3.44-11.57)	0.18 (0.04-0.28)
Lag period: 28 days	13.22 (6.59-18.54)	13.18 (7.33-18.34)	0.04 (-0.07-0.12)
Df/year for seasonal control: 6	6.45 (3.23-9.16)	6.20 (3.08-8.89)	0.25 (0.16-0.33)
Df/year for seasonal control: 10	6.05 (2.98-8.87)	5.12 (1.97-7.85)	0.93 (0.05-1.67)
Controlling for:			
NO2	6.34 (2.58-9.59)	5.99 (2.49-9.51)	0.35 (0.18-0.51)
03	6.65 (3.01-10.03)	6.28 (2.56-9.73)	0.37 (0.11-0.59)
PM10	5.32 (1.95-8.27)	5.07 (2.10-7.97)	0.24 (0.13-0.33)
PM2.5	5.57 (1.90-8.77)	5.29 (1.91-8.59)	0.28 (0.13-0.42)
Humidity + NO2	5.99 (2.02-9.46)	5.65 (2.14-9.16)	0.35 (0.17-0.51)
Humidity + O3	5.21 (1.76-8.10)	4.98 (1.68-8.14)	0.23 (0.12-0.32)
Humidity + PM10	5.49 (1.64-8.74)	5.21 (1.76-8.61)	0.28 (0.07-0.44)
Humidity + PM2.5	5.25 (1.54-8.60)	4.98 (1.59-8.37)	0.27 (0.11-0.39)
% Densely populated	5.62 (2.05-8.77)	5.29 (1.98-8.51)	0.34 (-0.04-0.66)
proportion 65+	6.05 (2.36-9.38)	5.73 (2.29-9.08)	0.32 (0.15-0.47)
% Daily smoker (16-74 years)	5.58 (2.18-8.49)	5.26 (2.18-8.37)	0.32 (-0.04-0.64)

Table S3. Second-stage meta-regression models.

AIC, BIC, multivariate Cochran Q test for heterogeneity (p-value), and I² statistic (%) in different metaregression models.

Model	Predictor	AIC	BIC	Q test	l ²
Intercept-only	-	3.33	11.11	0.1910	17.9%
Single predictor	Average temperature	11.62	27.17	0.0914	28.2%
	Temperature range	9.30	24.85	0.1439	23.1%

Table S4. The relative risks of cold temperature and hot temperature on deaths at lags of 0–14 days in the seven cities. Estimates were calculated from the non-linear model. Cold effects (1st VS minimum-mortality temperature (MMT)) and heat effects (99th VS MMT) correspond to Figure 2 where the pooled ones are shown.

8 9	City	Outcome	Cold effects	Heat effects
10	Oslo	Non-accidental	1.14 (1.02-1.28)	1.11 (1.03-1.19)
11	Bergen		1.13 (0.97-1.30)	1.05 (0.95-1.15)
12	Trondheim		1.14 (1.03-1.27)	1.12 (1.04-1.21)
13	Stavanger		1.15 (0.96-1.37)	1.03 (0.95-1.11)
14 15	Drammen		1 14 (1 04-1 26)	1 12 (1 04-1 20)
16	Eredrikstad		1.17(1.071.20) 1 15 (1 05-1 27)	1 07 (1 02-1 13)
17	Tromsø		1.15(1.05 1.27) 1.16(1.05 1.20)	$1.07 (1.02 \ 1.13)$
18		Non assidental mon	$1.10(1.05 \cdot 1.25)$	1.00 (0.08 1.22)
19	DSIO	Non-accidental men	1.19 (0.99-1.43)	1.09 (0.98-1.22)
20	Bergen		1.04 (0.90-1.20)	1.02 (0.86-1.21)
21	Irondheim		1.16 (0.97-1.39)	1.07 (0.98-1.15)
23	Stavanger		1.02 (0.83-1.25)	1.03 (0.85-1.24)
24	Drammen		1.16 (0.99-1.35)	1.09 (0.98-1.21)
25	Fredrikstad		1.06 (0.88-1.27)	1.03 (0.94-1.13)
26	Tromsø		1.01 (0.76-1.34)	1.01 (0.95-1.09)
27	Oslo	Non-accidental women	1.12 (0.96-1.31)	1.13 (1.03-1.25)
29	Bergen		1.26 (1.04-1.52)	1.09 (0.97-1.23)
30	Trondheim		1.16 (1.00-1.35)	1.11 (1.03-1.20)
31	Stavanger		1.37 (1.08-1.75)	1.07 (0.95-1.21)
32	Drammen		1.15 (1.01-1.31)	1.14 (1.04-1.24)
33	Fredrikstad		1.30 (1.11-1.53)	1.11 (1.02-1.21)
35	Tromsø		1.51 (1.17-1.95)	1.08 (1.00-1.17)
36	Oslo	Non-accidental 0-74	1 10 (0 89-1 36)	1 10 (0 97-1 25)
37	Bergen		1.10(0.051.00) 1.32(1.00-1.74)	1 03 (0 90-1 18)
38	Trondheim		1.32(1.001.74) 1 12(0 92-1 37)	1 12 (0 97-1 30)
39	Stavanger		$1.12(0.52 \cdot 1.57)$	1.12 (0.02 1 10)
40 41	Dramman		1.38(0.99-1.92)	1.01 (0.93-1.10)
42	Drammen Friedrikete d		1.14 (0.95-1.37)	1.11 (0.97-1.27)
43	Fredrikstad		1.21 (1.01-1.45)	1.06 (0.97-1.15)
44	Iromsø		1.22 (1.01-1.48)	1.05 (0.97-1.12)
45	Oslo	Non-accidental 75+	1.16 (1.01-1.32)	1.11 (1.02-1.2)
46 47	Bergen		1.11 (0.99-1.24)	1.11 (0.96-1.28)
48	Trondheim		1.15 (1.02-1.30)	1.13 (1.03-1.23)
49	Stavanger		1.13 (0.98-1.31)	1.11 (0.95-1.30)
50	Drammen		1.14 (1.02-1.28)	1.12 (1.03-1.22)
51	Fredrikstad		1.13 (1.02-1.27)	1.09 (1.02-1.16)
52	Tromsø		1.14 (1.02-1.28)	1.08 (1.01-1.14)
53 54	Oslo	CVD	1.27 (1.06-1.52)	1.13 (0.99-1.27)
55	Bergen		1.34 (1.02-1.75)	1.00 (1.00-1.00)
56	Trondheim		1.26 (1.06-1.48)	1.09 (0.96-1.24)
57	Stavanger		1.31 (0.96-1.79)	1.00 (1.00-1.00)
58	Drammen		1.24 (1.07-1.44)	1.06 (0.94-1.19)
59 60	Fredrikstad		1.22 (1.02-1.45)	1.00 (1.00-1.00)
58 59 60	Drammen Fredrikstad		1.24 (1.07-1.44) 1.22 (1.02-1.45)	1.06 (0.94-1.19) 1.00 (1.00-1.00)

3	Tromsø		1.21 (1.01-1.45)	1.00 (1.00-1.00)
4 5	Oslo	CVD men	1.24 (0.95-1.63)	1.03 (0.86-1.23)
5	Bergen		1.43 (0.95-2.17)	1.00 (1.00-1.00)
7	Trondheim		1.23 (0.91-1.67)	1.00 (0.99-1.00)
8	Stavanger		1 37 (0 85-2 20)	1 00 (1 00-1 00)
9	Drammon		1.37 (0.05 2.20) 1.25 (0.05 1.66)	1.00 (1.00 1.00)
10	Diaminen		1.23 (0.93-1.00)	1.00 (1.00-1.00)
11	Fredrikstad		1.27 (0.97-1.64)	1.00 (1.00-1.00)
12	Iromsø	-	1.24 (0.95-1.63)	1.00 (1.00-1.00)
14	Oslo	CVD women	1.27 (1.00-1.62)	1.20 (1.01-1.42)
15	Bergen		1.28 (0.88-1.84)	1.00 (1.00-1.00)
16	Trondheim		1.27 (1.02-1.58)	1.17 (0.99-1.39)
17	Stavanger		1.27 (0.83-1.94)	1.00 (1.00-1.00)
18	Drammen		1.27 (1.04-1.55)	1.14 (0.97-1.34)
19 20	Fredrikstad		1.28 (1.04-1.58)	1.09 (0.96-1.25)
20	Tromsø		1.29 (1.04-1.60)	1.10 (0.97-1.26)
22	Oslo	CVD 0-74	1.24 (0.84-1.84)	1.10 (0.83-1.45)
23	Bergen		1.13 (0.64-1.98)	1.04 (0.79-1.37)
24	Trondheim		1 21 (0 84-1 73)	1 03 (0 78-1 36)
25	Stavanger		1 10 (0 55-2 18)	1.03 (0.86-1.23)
20 27	Drammon		1.10 (0.55-2.18)	1.00 (0.00-1.20)
28	Diaminen		1.17 (0.64-1.02)	1.00 (0.77-1.30)
29	Fredrikstad		1.10 (0.75-1.62)	1.00 (1.00-1.01)
30	Iromsø		1.08 (0.73-1.61)	1.00 (0.99-1.00)
31	Oslo	CVD 75+	1.29 (1.05-1.58)	1.13 (0.98-1.29)
32	Bergen		1.41 (1.04-1.92)	1.00 (1.00-1.00)
33 34	Trondheim		1.28 (1.06-1.54)	1.10 (0.96-1.27)
35	Stavanger		1.38 (0.97-1.96)	1.00 (1.00-1.00)
36	Drammen		1.27 (1.07-1.50)	1.06 (0.93-1.21)
37	Fredrikstad		1.26 (1.03-1.53)	1.00 (1.00-1.00)
38	Tromsø		1.25 (1.02-1.53)	1.00 (1.00-1.00)
39	Oslo	Respiratory	1.24 (0.88-1.76)	1.33 (1.07-1.67)
40 41	Bergen		1.24 (0.81-1.89)	1.26 (0.90-1.77)
42	Trondheim		1.25 (0.90-1.74)	1.39 (1.08-1.79)
43	Stavanger		1 21 (0 71-2 04)	1 23 (0 86-1 75)
44	Drammen		1 23 (0 93-1 65)	1 38 (1 10-1 74)
45	Erodrikstad		$1.23(0.35^{-}1.03)$	1.38 (1.10-1.74)
46 47	Freurikstau		1.25 (0.95-1.04)	1.26 (1.00-1.54)
47	Tromsø	D	1.24 (0.92-1.67)	1.24 (1.04-1.49)
49	Oslo	Resp. men	1.36 (1.00-1.85)	1.17 (0.71-1.92)
50	Bergen		1.04 (0.56-1.95)	1.33 (0.79-2.23)
51	Trondheim		1.32 (0.98-1.79)	1.20 (0.74-1.95)
52	Stavanger		1.00 (0.99-1.00)	1.55 (0.64-3.74)
53 54	Drammen		1.23 (0.94-1.61)	1.18 (0.76-1.82)
55	Fredrikstad		1.08 (0.73-1.61)	1.16 (0.87-1.53)
56	Tromsø		1.07 (0.70-1.64)	1.16 (0.88-1.54)
57	Oslo	Resp. women	1.26 (0.82-1.95)	1.57 (1.14-2.16)
58	Bergen		1.62 (0.87-3.01)	1.32 (0.87-2.00)
59	Trondheim		1.28 (0.85-1.91)	1.61 (1.17-2.20)
60			- (, _ ()

2				
3	Stavanger		1.69 (0.80-3.55)	1.20 (0.85-1.70)
4	Drammen		1.30 (0.90-1.88)	1.58 (1.18-2.11)
5	Fredrikstad		1.37 (0.92-2.04)	1.35 (1.09-1.68)
7	Tromsø		1 38 (0.91-2.10)	1 30 (1 07-1 58)
8	Oslo	Resp. 0.74	1.30(0.312.10) 1.03(1.24-3.02)	1.87 (0.84-4.16)
9	Borgon	Nesp. 0-74	1.55(1.24-5.02)	1.07(0.04-4.10)
10			1.00 (0.47-5.90)	1.00 (1.00-1.00)
11	Tronaneim		1.88 (1.20-2.92)	1.68 (0.79-3.58)
12	Stavanger		1.27 (0.31-5.17)	1.01 (1.00-1.02)
14	Drammen		1.65 (1.11-2.44)	1.43 (0.70-2.92)
15	Fredrikstad		1.33 (0.82-2.17)	1.01 (0.53-1.93)
16	Tromsø		1.39 (0.79-2.45)	1.07 (0.56-2.01)
17	Oslo	Resp. 75+	1.25 (0.85-1.85)	1.32 (1.03-1.70)
18	Bergen		1.37 (0.85-2.20)	1.42 (0.98-2.06)
19	Trondheim		1.25 (0.87-1.80)	1.39 (1.06-1.84)
20	Stavanger		1.37 (0.76-2.48)	1.35 (0.91-1.99)
21	Drammen		1 27 (0 92-1 75)	1 42 (1 10-1 83)
23	Eredrikstad		1.27(0.92 1.73) 1.28(0.92 1.77)	$1.42(1.10\ 1.00)$
24	Tromed		$1.20(0.95 \cdot 1.77)$	1.33(1.09-1.02)
25	Tromsø		1.28 (0.91-1.80)	1.29 (1.07-1.56)
26				
27				
20				
30				
31				
32				
33				
34				
35				
37				
38				
39				
40				
41				
42				
43 44				
45				
46				
47				

Table S5. Attributable fractions (%) on non-accidental and cardiopulmonary deaths in the seven cities for cold and heat defined as temperatures below and above the minimum mortality temperature (MMT) and extremes defined as cut-offs at the 1st and 99th percentiles. Brackets indicate 95% empirical confidence intervals. AFs for CVD are presented in italics.

City	Causes	Extreme Cold	Moderate Cold	Moderate Heat	Extreme Heat	Overall
Oslo	Non-accidental	0.15 (0-0.28)	6.59 (0.80-11.7)	0.14 (0.01-0.26)	0.14 (0.04-0.23)	7.01 (1.54-12.52)
	CVD	0.29 (0.06-0.47)	10.01 (1.90-16.89)	0.70 (-0.24-1.59)	0.09 (-0.07-0.21)	11.08 (3.08-18.33)
	Respiratory	0.35 (-0.16-0.69)	2.26 (-19.94-19.87)	0.40 (-0.1-0.84)	0.33 (0.05-0.51)	3.33 (-16.34-20.01)
Bergen		0.18 (-0.01-0.35)	1.82 (-7.40-10.44)	0.06 (-0.07-0.18)	0.09 (-0.13-0.25)	2.15 (-7.21-10.10)
		0.28 (-0.0 <mark>6</mark> -0.52)	16.47 (-1.87-29.70)	0 (0-0)	-0.12 (-0.39-0.07)	16.63 (-1.63-30.58)
		0.16 (-0.68-0.62)	9.63 (-12.43-25.14)	0.96 (-1.82-3.18)	0.25 (-0.32-0.54)	10.99 (-11.13-27.84)
Trondheim		0.16 (0-0.29)	6.28 (0.34-11.62)	0.19 (0.06-0.31)	0.16 (0.05-0.26)	6.78 (0.89-12.16)
		0.27 (0.06-0.44)	8.98 (2.16-15.31)	0.63 (-0.26-1.43)	0.06 (-0.15-0.21)	9.93 (2.35-16.58)
		0.37 (-0.18-0.70)	2.29 (-16.93-18.29)	0.51 (0.09-0.88)	0.34 (0.08-0.51)	3.50 (-17.22-20.45)
Stavanger		0.22 (0-0.40)	1.21 (-9.26-10.56)	0.02 (-0.07-0.10)	0.05 (-0.08-0.16)	1.50 (-8.48-10.77)
		0.23 (-0.26-0.53)	15.8 (-2.5-30.59)	0 (0-0)	-0.07 (-0.20-0.04)	15.96 (-2.31-31.37)
		0.08 (-1.57-0.71)	11.41 (-15.81-30. <mark>5</mark> 2)	0.77 (-1.71-2.53)	0.16 (-0.16-0.35)	12.42 (-17.4-30.79)
Drammen		0.16 (0.03-0.26)	5.71 (0.56-10.29)	0.17 (0.05-0.28)	0.15 (0.05-0.24)	6.18 (0.82-10.88)
		0.25 (0.06-0.41)	8.36 (1.65-13.97)	0.52 (-0.30-1.30)	0.02 (-0.15-0.16)	9.14 (2.40-15.34)
		0.32 (-0.13-0.63)	3.22 (-13.71-17.00)	0.56 (0.03-1.04)	0.34 (0.08-0.5)	4.43 (-12.21-17.91)
Fredrikstad		0.19 (0.05-0.31)	4.42 (-0.41-9.11)	0.10 (0.01-0.18)	0.12 (0.03-0.19)	4.83 (-0.18-9.36)
		0.21 (-0.03-0.40)	7.62 (-2.90-16.30)	0 (0-0)	-0.04 (-0.14-0.03)	7.78 (-2.15-16.36)
		0.31 (-0.33-0.71)	5.63 (-8.90-15.94)	0.47 (-0.91-1.57)	0.29 (0.09-0.43)	6.69 (-6.89-18.78)
Tromsø		0.18 (0.05-0.30)	4.57 (0.01-9.38)	0.08 (0.02-0.14)	0.10 (0.03-0.17)	4.93 (-0.30-9.64)
		0.24 (-0.03-0.45)	7.04 (-3.00-15.73)	0 (0-0)	-0.05 (-0.15-0.03)	7.23 (-3.87-16.03)
		0.20 (-0.19-0.48)	5.67 (-7.37-16.68)	0.53 (-0.88-1.94)	0.26 (0.09-0.39)	6.66 (-6.92-17.14)

Table S6. Attributable fractions (%) on non-accidental, cardiovascular and respiratory deaths in the seven cities for moderate cold and heat defined as temperatures below and above the minimum mortality temperature (MMT) and extremes defined as cut-offs at the 1st and 99th percentiles. Brackets indicate 95% empirical confidence intervals.MMP is minimum mortality percentile.

City	Outcome	MMP	MMT	Extreme cold	Moderate cold	Moderate heat	Extreme heat	Overall
Oslo	Non-accidental	92	16	0.15 (0.00-0.28)	6.59 (0.80-11.7)	0.14 (0.01-0.26)	0.14 (0.04-0.23)	7.01 (1.54-12.52)
Bergen		94	16	0.18 (-0.01-0.35)	1.82 (-7.40-10.44)	0.06 (-0.07-0.18)	0.09 (-0.13-0.25)	2.15 (-7.21-10.10)
Trondheim		92	15	0.16 (0.00-0.29)	6.28 (0.34-11.62)	0.19 (0.06-0.31)	0.16 (0.05-0.26)	6.78 (0.89-12.16)
Stavanger		95	17	0.22 (0.00-0.40)	1.21 (-9.26-10.56)	0.02 (-0.07-0.10)	0.05 (-0.08-0.16)	1.50 (-8.48-10.77)
Drammen		92	16	0.16 (0.03-0.26)	5.71 (0.56-10.29)	0.17 (0.05-0.28)	0.15 (0.05-0.24)	6.18 (0.82-10.88)
Fredrikstad		93	18	0.19 (0.05-0.31)	4.42 (-0.41-9.11)	0.10 (0.01-0.18)	0.12 (0.03-0.19)	4.83 (-0.18-9.36)
Tromsø		93	12	0.18 (0.05-0.30)	4.57 (0.01-9.38)	0.08 (0.02-0.14)	0.10 (0.03-0.17)	4.93 (-0.30-9.64)
Oslo	CVD	81	14	0.29 (0.06-0.47)	10.01 (1.90-16.89)	0.7 (-0.24-1.59)	0.09 (-0.07-0.21)	11.08 (3.08-18.33)
Bergen		99	19	0.28 (-0.06-0.52)	16.47 (-1.87-29.70)	0 (0-0)	-0.12 (-0.39-0.07)	16.63 (-1.63-30.58)
Trondheim		82	12	0.27 (0.06-0.44) 🗸	8.98 (2.16-15.31)	0.63 (-0.26-1.43)	0.06 (-0.15-0.21)	9.93 (2.35-16.58)
Stavanger		99	20	0.23 (-0.26-0.53)	15.80 (-2.50-30.59)	0 (0-0)	-0.07 (-0.20-0.04)	15.96 (-2.31-31.37)
Drammen		80	13	0.25 (0.06-0.41)	8.36 (1.65-13.97)	0.52 (-0.30-1.30)	0.02 (-0.15-0.16)	9.14 (2.40-15.34)
Fredrikstad		99	21	0.21 (-0.03-0.40)	7.62 (-2.90-16.30)	0 (0-0)	-0.04 (-0.14-0.03)	7.78 (-2.15-16.36)
Tromsø		99	16	0.24 (-0.03-0.45)	7.04 (-3.00-15.73)	0 (0-0)	-0.05 (-0.15-0.03)	7.23 (-3.87-16.03)
Oslo	Resp.	91	16	0.35 (-0.16-0.69)	2.26 (-19.94-19.87)	0.40 (-0.10-0.84)	0.33 (0.05-0.51)	3.33 (-16.34-20.01) 10.99 (-11.13-
Bergen		80	12	0.16 (-0.68-0.62)	9.63 (-12.43-25.14)	0.96 (-1.82-3.18)	0.25 (-0.32-0.54)	27.84)
Trondheim		91	15	0.37 (-0.18-0.70)	2.29 (-16.93-18.29)	0.51 (0.09-0.88)	0.34 (0.08-0.51)	3.50 (-17.22-20.45) 12.42 (-17.40-
Stavanger		81	14	0.08 (-1.57-0.71)	11.41 (-15.81-30.52)	0.77 (-1.71-2.53)	0.16 (-0.16-0.35)	30.79)
Drammen		90	16	0.32 (-0.13-0.63)	3.22 (-13.71-17.00)	0.56 (0.03-1.04)	0.34 (0.08-0.50)	4.43 (-12.21-17.91)
Fredrikstad		79	15	0.31 (-0.33-0.71)	5.63 (-8.90-15.94)	0.47 (-0.91-1.57)	0.29 (0.09-0.43)	6.69 (-6.89-18.78)
Tromsø		80	9	0.20 (-0.19-0.48)	5.67 (-7.37-16.68)	0.53 (-0.88-1.94)	0.26 (0.09-0.39)	6.66 (-6.92-17.14)

Table S7. Attributable fractions (%) on non-accidental, cardiovascular and respiratory deaths by sex in the seven cities for moderate cold and heat defined as temperatures below and above the minimum mortality temperature and extremes defined as cut-offs at the 1st and 99th percentiles. Brackets indicate 95% empirical confidence intervals.

City	Outcome	MMP	MMT	Extreme cold	Moderate cold	Moderate heat	Extreme heat	Overall
Oslo	Non-accidental men	94	17	0.18 (-0.05-0.37)	7.98 (-1.61-16.03)	0.08 (-0.05-0.2)	0.14 (-0.02-0.28)	8.37 (-1.72-17.38)
Bergen		42	5	-0.01 (-0.26-0.21)	0.89 (-1.20-2.93)	1.40 (-3.32-5.72)	0.04 (-0.32-0.29)	2.32 (-2.21-6.64)
Trondheim		94	16	0.17 (-0.06-0.35)	7.51 (-1.57-15.74)	0.12 (-0.05-0.28)	0.17 (-0.04-0.31)	7.96 (-1.55-16.79)
Stavanger		41	6	-0.05 (-0.46-0.24)	0.90 (-1.27-2.93)	1.80 (-3.54-6.72)	0.03 (-0.23-0.23)	2.68 (-2.30-7.29)
Drammen		94	17	0.15 (-0.06-0.32)	6.63 (-2.06-14.3)	0.10 (-0.04-0.23)	0.15 (-0.02-0.29)	7.02 (-1.09-14.43)
Fredrikstad		95	19	0.07 (-0.18-0.25)	4.36 (-3.51-11.7)	0.04 (-0.05-0.13)	0.08 (-0.07-0.19)	4.55 (-3.34-11.90)
Tromsø		96	14	0.07 (-0.17-0.26)	4.52 (-3.50-11.83)	0.03 (-0.03-0.08)	0.07 (-0.04-0.17)	4.69 (-3.57-12.25)
Oslo	Non-accidental women	84	14	0.14 (-0.06-0.31)	7.23 (0.10-13.48)	0.35 (-0.29-0.94)	0.15 (0.03-0.25)	7.87 (1.26-14.15)
Bergen		91	15	0.33 (0.13-0.51)	4.02 (-6.90-12.89)	0.15 (-0.18-0.46)	0.16 (-0.09-0.34)	4.66 (-6.2-13.97)
Trondheim		85	13	0.16 (-0.03-0.31)	6.90 (1.14-12.3)	0.37 (-0.18-0.89)	0.17 (0.05-0.28)	7.60 (1.19-13.50)
Stavanger		92	16	0.43 (0.14-0.64)	3.72 (-8.38-14.44)	0.09 (-0.19-0.33)	0.09 (-0.07-0.23)	4.33 (-7.90-14.60)
Drammen		84	12	0.18 (0.03-0.31)	6.47 (0.94-11.33)	0.35 (-0.17-0.86)	0.17 (0.05-0.27)	7.16 (1.51-12.35)
Fredrikstad		86	16	0.29 (0.11-0.43)	5.61 (0.31-10.18)	0.22 (-0.22-0.61)	0.16 (0.05-0.26)	6.27 (0.28-11.10)
Tromsø		87	10	0.27 (0.12-0.40)	5.66 (0.33-10.39)	0.19 (-0.18-0.58)	0.13 (0.05-0.21)	6.25 (0.09-11.41)
Oslo	CVD men	82	14	0.24 (-0.14-0.52)	9.51 (-3.27-19.79)	0.35 (-1.03-1.62)	0.00 (-0.30-0.20)	10.10 (-2.60-20.85)
Bergen		99	19	0.24 (-0.33-0.60)	26.20 (0.86-42.98)	0 (0-0)	-0.03 (-0.38-0.18)	26.41 (2.13-44.64)
Trondheim		99	19	0.22 (-0.22-0.50)	9.85 (-9.21-26.13)	0 (0-0)	-0.04 (-0.25-0.11)	10.03 (-9.83-26.92)
Stavanger		99	20	0.23 (-0.17-0.48)	12.40 (-4.91-28.04)	0 (0-0)	-0.03 (-0.21-0.10)	12.59 (-6.12-26.57)
Drammen		99	20	0.15 (-0.92-0.63)	27.03 (1.44-45.51)	0 (0-0)	-0.03 (-0.22-0.11)	27.15 (1.28-46.38)
Fredrikstad		99	21	0.18 (-0.23-0.46)	16.57 (2.07-27.88)	0 (0-0)	-0.03 (-0.18-0.08)	16.72 (2.99-27.86)
Tromsø		99	16	0.21 (-0.29-0.57)	16.25 (2.27-27.34)	0 (0-0)	-0.03 (-0.16-0.08)	16.43 (2.65-27.83)
Oslo	CVD women	80	13	0.29 (-0.03-0.53)	10.05 (-0.93-18.90)	1.00 (-0.30-2.16)	0.13 (-0.07-0.27)	11.47 (0.68-20.98)
Bergen		99	19	0.31 (-0.17-0.63)	9.29 (-18.46-28.00)	0 (0-0)	-0.23 (-0.72-0.06)	9.37 (-18.69-29.14)
Trondheim		81	12	0.29 (-0.02-0.50)	9.36 (0.37-17.31)	1.03 (-0.17-2.04)	0.13 (-0.11-0.31)	10.80 (0.50-19.50)
Stavanger		99	20	0.28 (-0.36-0.61)	6.81 (-21.55-27.74)	0 (0-0)	-0.12 (-0.34-0.03)	6.95 (-21.15-29.07)

Page	29	of	54
------	----	----	----

2									
3	Drammen		79	13	0.29 (0.03-0.48)	9.19 (0.37-16.51)	1.05 (-0.05-2.07)	0.07 (-0.16-0.23)	10.59 (1.75-18.5)
4 5	Fredrikstad		78	15	0.30 (-0.01-0.51)	8.03 (-0.26-14.72)	1.01 (-0.17-1.99)	0.02 (-0.18-0.18)	9.35 (0.08-16.80)
6	Tromsø		79	8	0.31 (0.02-0.54)	7.83 (-0.45-14.78)	0.95 (-0.05-1.98)	0.03 (-0.19-0.19)	9.11 (0.53-15.99)
7	Oslo	Resp. Men	31	1	0.49 (-0.05-0.83)	1.55 (-2.34-4.68)	3.72 (-9.13-13.67)	0.25 (-0.45-0.57)	6.00 (-7.88-15.42)
8									16.24 (-18.17-
9 10	Bergen		78	12	-0.26 (-3.17-0.58)	13.37 (-20.29-33.34)	3.38 (-0.25-5.79)	-0.25 (-3.70-0.44)	38.55)
10	Trondheim		30	1	0.52 (-0.09-0.90)	1.22 (-1.92-3.79)	3.10 (-9.17-12.12)	0.22 (-0.43-0.50)	5.05 (-7.04-14.42)
12									22.67 (-35.02-
13	Stavanger		1	-5	-0.45 (-2.71-0.15)	0 (0-0)	23.00 (-33.52-55.01)	0.13 (-1.12-0.58)	51.72)
14	Drammen		38	2	0.35 (-0.24-0.73)	1.58 (-2.57-5.50)	2.20 (-7.79-10.42)	0.16 (-0.35-0.41)	4.28 (-6.24-11.91)
15 16	Fredrikstad		72	13	0.06 (-0.93-0.57)	2.74 (-16.22-15.32)	1.01 (-1.16-2.72)	0.08 (-0.36-0.33)	3.89 (-13.3-17.58)
10	Tromsø		74	7	0.04 (-0.84-0.42)	2.90 (-15.66-16.63)	1.29 (-1.19-3.45)	0.11 (-0.28-0.35)	4.34 (-14.47-18.76)
18									10.29 (-13.29-
19	Oslo	Resp. Women	78	13	0.31 (-0.43-0.74)	8.30 (-15.92-24.76)	1.28 (-1.25-3.26)	0.40 (0.11-0.58)	28.03)
20									19.80 (-19.63-
21	Bergen		94	16	0.49 (-0.47-0.90)	18.53 (-20.04-41.39)	0.33 (-0.25-0.81)	0.46 (-0.36-0.78)	42.69)
22	Trondheim		80	12	0.30 (-0.37-0.67)	7.76 (-11.42-22.70)	1.21 (-1.00-2.96)	0.43 (0.15-0.59)	9.69 (-12.03-25.78)
23									22.24 (-22.62-
24	Stavanger		95	17	0.57 (-1.15-1.10)	21.37 (-20.47-47.53)	0.11 (-0.17-0.37)	0.20 (-0.21-0.4)	48.31)
26	Drammen		80	13	0.33 (-0.22-0.68)	8.39 (-10.81-21.84)	1.08 (-1.04-2.88)	0.47 (0.17-0.66)	10.25 (-9.25-25.13)
27	Fredrikstad		91	17	0.53 (-0.52-1.08)	10.23 (-9.33-24.54)	0.34 (-0.06-0.71)	0.41 (0.08-0.60)	11.51 (-7.55-27.63)
28									11.42 (-10.56-
29	Tromsø		91	12	0.31 (-0.24-0.63)	10.49 (-11.03-26.24)	0.31 (-0.07-0.69)	0.32 (0.09-0.47)	27.79)
50									

Table S8. Attributable fractions (%) on non-accidental, cardiovascular and respiratory deaths by age group in the seven cities for moderate cold and heat defined as temperatures below and above the minimum mortality temperature and extremes defined at the 1st and 99th percentiles. Brackets indicate 95% empirical confidence intervals.

City	Outcome	MMP	MMT	Extreme cold	Moderate cold	Moderate heat	Extreme heat	Overall
Oslo	Non-accidental 0-74	93	17	0.09 (-0.22-0.32)	7.67 (-3.18-17.96)	0.12 (-0.06-0.3)	0.15 (-0.06-0.30)	8.02 (-2.64-17.84)
Bergen		97	17	0.31 (0.01-0.54)	14.59 (-1.73-26.84)	0.02 (-0.08-0.12)	0.11 (-0.27-0.37)	15.04 (-0.52-27.08)
Trondheim		93	15	0.11 (-0.17-0.30)	8.61 (-1.61-17.87)	0.15 (-0.07-0.34)	0.17 (-0.05-0.34)	9.03 (-1.62-18.93)
Stavanger		98	19	0.36 (-0.08-0.61)	15.36 (-1.09-29.31)	0.00 (-0.03-0.04)	0.05 (-0.17-0.22)	15.76 (-0.69-30.02)
Drammen		94	17	0.14 (-0.11-0.32)	8.88 (-0.94-17.97)	0.12 (-0.05-0.28)	0.17 (-0.05-0.34)	9.31 (-0.47-17.72)
Fredrikstad		95	19	0.20 (-0.03-0.37)	10.47 (1.62-17.64)	0.05 (-0.05-0.15)	0.11 (-0.05-0.24)	10.83 (2.24-18.68)
Tromsø		96	14	0.24 (-0.05-0.45)	10.88 (1.36-19.31)	0.03 (-0.03-0.10)	0.08 (-0.03-0.17)	11.23 (1.11-19.65)
Oslo	Non-accidental 75+	91	16	0.17 (-0.01-0.33)	6.36 (-0.88-12.82)	0.16 (-0.03-0.34)	0.14 (-0.03-0.24)	6.83 (0.25-13.01)
Bergen		45	5	0.16 (-0.01-0.31)	0.68 (-1.22-2.40)	0.98 (-2.49-4.14)	0.12 (-2.49-0.30)	1.94 (-1.28-4.77)
Trondheim		90	14	0.17 (0.00-0.33)	5.58 (-0.54-11.25)	0.22 (0.00-0.42)	0.16 (0.00-0.27)	6.13 (-0.09-12.12)
Stavanger		41	6	0.20 (-0.06-0.40)	0.72 (-1.24-2.51)	1.30 (-2.75-5.30)	0.10 (-2.75-0.24)	2.32 (-1.67-6.22)
Drammen		90	15	0.16 (0.02-0.29)	4.75 (-0.97-9.93)	0.20 (0.01-0.40)	0.15 (0.01-0.25)	5.26 (-0.25-10.46)
Fredrikstad		83	16	0.18 (0.01-0.32)	2.74 (-1.96-6.90)	0.18 (-0.32-0.64)	0.13 (-0.32-0.21)	3.22 (-2.15-7.85)
Tromsø		85	10	0.16 (0.02-0.29)	2.73 (-2.11-7.18)	0.16 (-0.27-0.61)	0.12 (-0.27-0.2)	3.16 (-1.97-7.69)
Oslo	CVD 0-74	79	13	0.30 (-0.29-0.65)	11.54 (-7.09-24.53) 🦉	1.86 (-0.37-3.73)	-0.07 (-0.66-0.23)	13.62 (-4.31-27.97)
Bergen		97	17	0.05 (-1.06-0.58)	7.69 (-34.27-32.92)	0.02 (-0.19-0.21)	0.13 (-0.74-0.51)	7.89 (-34.24-32.91)
Trondheim		79	11	0.22 (-0.27-0.50)	9.64 (-4.84-21.5)	1.46 (-0.58-3.10)	-0.13 (-0.72-0.2)	11.19 (-5.38-23.88)
Stavanger		78	13	0.20 (-0.31-0.52)	7.63 (-7.71-19.53)	1.18 (-0.76-2.84)	-0.18 (-0.88-0.23)	8.84 (-6.50-21.37)
Drammen		98	19	-0.04 (-2.02-0.57)	8.80 (-34.12-37.24)	0.01 (-0.06-0.07)	0.09 (-0.38-0.38)	8.86 (-33.69-37.86)
Fredrikstad		99	21	0.07 (-0.46-0.38)	4.77 (-20.44-23.34)	0 (0-0)	-0.05 (-0.30-0.10)	4.78 (-19.09-22.63)
Tromsø		99	16	0.10 (-0.77-0.67)	3.81 (-22.11-22.33)	0 (0-0)	-0.04 (-0.20-0.08)	3.87 (-23.34-23.15)
Oslo	CVD 75+	83	14	0.29 (0.03-0.50)	9.96 (0.56-18.00)	0.42 (-0.52-1.31)	0.12 (-0.52-0.25)	10.79 (1.67-18.96)
Bergen		99	19	0.34 (-0.03-0.59)	20.16 (0.24-34.19)	0 (0-0)	-0.19 (0.00-0.04)	20.30 (0.44-35.04)
Trondheim		84	12	0.28 (0.02-0.47)	9.15 (1.24-16.35)	0.41 (-0.47-1.17)	0.10 (-0.47-0.26)	9.94 (1.27-17.45)
Stavanger		99	20	0.29 (-0.26-0.61)	18.8 (-1.17-34.62)	0 (0-0)	-0.11 (0.00-0.02)	18.98 (-1.12-35.48)

http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Drammen		81	14	0.26 (0.04-0.44)	8.78 (1.09-14.98)	0.35 (-0.55-1.19)	0.05 (-0.55-0.19)	9.43 (1.63-16.4)
Fredrikstad		99	21	0.25 (-0.02-0.47)	9.10 (-2.46-19.02)	0 (0-0)	-0.04 (0-0.05)	9.31 (-2.15-19.17)
Tromsø		99	16	0.25 (-0.01-0.47)	8.49 (-3.10-18.02)	0 (0-0)	-0.05 (0-0.06)	8.68 (-3.47-18.43)
Oslo	Resp. 0-74	9	-6	0.99 (0.45-1.24)	1.43 (0.34-2.37)	16.47 (-11.61-34.59)	0.52 (-0.35-0.78)	19.4 (-11.87-37.12)
Bergen		99	19	0.38 (-3.87-1.09)	55.18 (-20.02-79.58)	0 (0-0)	-2.83 (-142.82-0.05)	52.73 (-114.46- 71.93)
Trondheim		8	-6	0.89 (0.40-1.12)	0.85 (0.09-1.47)	15.56 (-13.48-33.69)	0.47 (-0.71-0.79)	17.76 (-11.68- 36.43)
Stavanger		8	-6	0.57 (0.16-0.78)	0.84 (0.14-1.41)	14.45 (-11.36-33.63)	0.36 (-1.08-0.86)	16.21 (-11.66- 35.23)
Drammen		99	20	-0.05 (-9.88-0.77)	49.88 (-32.32-78.48)	0 (0-0)	-5.81 (-1162.43-0.02)	44.01 (-1568.18- 70.48)
								11.82 (-21.14-
Fredrikstad		7	-4	0.40 (-0.44-0.72)	0.35 (-0.36-0.89)	11.78 (-18.52-32.68)	-0.70 (-9.29-0.63)	32.84) 12 87 (-14 17-
Tromsø		8	-10	0.41 (-0.45-0.76)	0.50 (-0.56-1.35)	11.99 (-13.28-31.37)	-0.04 (-0.53-0.16)	31.66)
Oslo	Resp. 75+	91	16	0.25 (-0.42-0.66)	7.29 (-16.22-25.51)	0.40 (-0.18-0.90)	0.32 (-0.18-0.52)	8.25 (-12.62-25.53)
Bergen		81	12	0.29 (-0.61-0.71)	9.47 (-16.09-26.77)	0.83 (-2.31-3.15)	0.45 (-2.31-0.70)	30.98)
Trondheim		90	14	0.28 (-0.42-0.69)	6.78 (-13.11-22.55)	0.49 (-0.07-0.99)	0.33 (-0.07-0.51)	7.86 (-13.48-24.74)
Stavanger		82	14	0.31 (-1.38-0.89)	10.98 (-20.88-32.67)	0.58 (-2.02-2.4)	0.27 (-2.02-0.44)	33.25)
Drammen		90	15	0.29 (-0.32-0.67)	7.70 (-10.59-22.35)	0.57 (0.01-1.10)	0.32 (0.01-0.49)	8.87 (-8.74-22.29)
Fredrikstad		84	16	0.32 (-0.52-0.81)	8.54 (-8.18-20.26)	0.35 (-0.66-1.22)	0.30 (-0.66-0.42)	9.49 (-5.93-22.96)
Tromsø		85	10	0.18 (-0.32-0.49)	8.26 (-6.82-20.72)	0.44 (-0.72-1.56)	0.37 (-0.72-0.52)	9.24 (-6.34-21.07)

Table S9. Attributable number of deaths (AN) for non-accidental, cardiovascular, and respirator
mortality by sex and age group in each city. Brackets indicate 95% empirical confidence intervals.

City	Outcome	Total	Cold	Heat
Oslo	Non-accidental	6949 (1526-12410)	6671 (952-11796)	278 (79-446)
Bergen		936 (-3138-4396)	875 (-3143-4313)	65 (-74-183)
Trondheim		1857 (244-3331)	1761 (110-3180)	96 (33-153)
Stavanger		291 (-1645-2089)	277 (-1647-2051)	14 (-29-50)
Drammen		1078 (143-1897)	1022 (71-1869)	56 (17-85)
Fredrikstad		769 (-29-1490)	734 (-86-1450)	35 (8-59)
Tromsø		403 (-25-789)	389 (-20-781)	15 (4-25)
Oslo	Non-accidental men	3619 (-744-7515)	3528 (-670-7074)	95 (-22-195)
Bergen		463 (-441-1325)	176 (-253-583)	287 (-607-1137)
Trondheim		1013 (-197-2137)	978 (-281-2014)	36 (-9-73)
Stavanger		240 (-206-653)	76 (-145-263)	164 (-354-656)
Drammen		560 (-87-1152)	541 (-136-1143)	19 (-5-42)
Fredrikstad		340 (-249-888)	331 (-239-865)	9 (-7-24)
Tromsø		190 (-144-496)	186 (-123-478)	4 (-2-11)
Oslo	Non-accidental women	4398 (704-7908)	4119 (246-7489)	274 (-84-592)
Bergen		1098 (-1461-3293)	1025 (-1544-3203)	73 (-38-170)
Trondheim		1115 (175-1980)	1034 (158-1890)	79 (-6-160)
Stavanger		452 (-825-1524)	433 (-818-1477)	19 (-20-51)
Drammen		677 (143-1167)	628 (79-1094)	48 (-5-91)
Fredrikstad		530 (24-938)	498 (41-929)	32 (-4-67)
Tromsø		259 (4-472)	245 (21-460)	13 (-3-30)
Oslo	Non-accidental 0-74	2338 (-769-5200)	2259 (-950-5048)	76 (-29-160)
Bergen		1854 (-64-3339)	1838 (-192-3410)	16 (-43-59)
Trondheim		760 (-136-1592)	733 (-209-1502)	27 (-11-56)
Stavanger		914 (-40-1741)	911 (-106-1744)	3 (-12-15)
Drammen		509 (-26-969)	493 (-40-952)	16 (-5-34)
Fredrikstad		508 (105-876)	500 (61-877)	8 (-4-19)
Tromsø		328 (32-574)	325 (56-576)	3 (-2-7)
Oslo	Non-accidental 75+	4779 (175-9104)	4569 (-217-8852)	210 (35-357)
Bergen		605 (-399-1488)	262 (-346-830)	346 (-849-1348)
Trondheim		1164 (-17-2301)	1091 (-125-2177)	72 (15-118)
Stavanger		316 (-227-846)	125 (-159-390)	190 (-398-714)
Drammen		629 (-30-1252)	588 (-142-1221)	42 (7-67)
Fredrikstad		361 (-241-881)	328 (-184-855)	35 (-20-91)
Tromsø		166 (-104-405)	152 (-101-391)	15 (-9-39)
Oslo	CVD all	3935 (1094-6510)	3655 (767-6240)	281 (-57-568)
Bergen		2565 (-251-4717)	2584 (-231-4706)	-19 (-60-12)
Trondheim		983 (233-1641)	914 (198-1531)	68 (-23-148)
Stavanger		1054 (-153-2071)	1058 (-187-2050)	-5 (-14-2)
Drammen		595 (156-999)	560 (107-933)	35 (-18-82)
Fredrikstad		483 (-134-1016)	486 (-181-1056)	-2 (-9-2)
Tromsø		209 (-112-463)	210 (-69-463)	-1 (-4-1)
Oslo	CVD men	1527 (-393-3152)	1474 (- <u>467-</u> 3077)	- (· +) 53 (-157-234)
Bergen		1803 (145-3047)	1805 (51-2012)	-2 (-31-13)
Trondheim		<u>464 (-455-1745)</u>	465 (-458-1729)	-2 (-12-6)
nonuneiiii		·J+ (+JJ 12+J)	105 (1 50 ±255)	- (0)

1	_				
2	Stavanger		380 (-185-802)	381 (-151-816)	-1 (-6-3)
3 A	Drammen		826 (39-1412)	827 (11-1397)	-1 (-7-3)
5	Fredrikstad		476 (85-793)	477 (54-817)	-1 (-5-2)
6	Tromsø		236 (38-399)	236 (40-399)	0 (-3-1)
7	Oslo	CVD women	2340 (139-4280)	2109 (-155-4051)	228 (-33-457)
8	Bergen		806 (-1607-2505)	825 (-1546-2501)	-20 (-63-5)
9	Trondheim		569 (26-1028)	508 (-6-967)	61 (-3-117)
10	Stavanger		249 (-758-1042)	254 (-790-1031)	-4 (-13-1)
12	Drammen		367 (61-642)	329 (9-583)	39 (0-71)
13	Fredrikstad		314 (3-565)	280 (-2-536)	34 (-3-70)
14	Tromsø		133 (8-233)	118 (3-232)	14 (-1-29)
15	Oslo	CVD 0-74	971 (-307-1993)	844 (-462-1847)	127 (-34-249)
17	Bergen		253 (-1096-1053)	248 (-1068-1106)	5 (-31-22)
18	Trondheim		253 (-121-539)	223 (-144-505)	30 (-18-67)
19	Stavanger		117 (-86-283)	104 (-110-256)	13 (-13-33)
20	Drammen		138 (-524-589)	136 (-585-595)	2 (-7-7)
21	Fredrikstad		61 (-244-290)	62 (-270-313)	-1 (-4-1)
23	Tromsø		31 (-188-187)	32 (-154-184)	0(-2-1)
24	Oslo	CVD 75+	3063 (171-5383)	2010 (273-5232)	153 (-116-395)
25	Dorgon		2491 (54 4292)	2510(275-3232)	22 (67 6)
26	Trandhaim		2401 (34-4203)	2304 (00-4293)	-25(-07-0)
27	Ctouronger		759 (97-1333)	720 (30-1302) 1009 (97 1959)	39 (-32-100)
29	Stavanger		1002 (-59-1873)	1008 (-87-1858)	-0 (-15-1)
30	Drammen		468 (81-813)	448 (53-763)	19 (-26-58)
31	Fredrikstad		459 (-106-945)	461 (-111-974)	-2 (-7-2)
32	Tromsø	- "	181 (-72-384)	182 (-37-383)	-1 (-4-1)
33 34	Oslo	Resp. all	352 (-1729-2117)	275 (-1985-2016)	77 (10-128)
35	Bergen		481 (-487-1219)	428 (-569-1141)	53 (-71-152)
36	Trondheim		98 (-482-573)	74 (-523-523)	24 (6-37)
37	Stavanger		255 (-357-632)	236 (-341-601)	19 (-35-59)
38	Drammen		83 (-227-334)	66 (-269-327)	17 (3-26)
39 40	Fredrikstad		113 (-116-316)	100 (-122-297)	13 (-9-32)
41	Tromsø		57 (-59-147)	50 (-60-144)	7 (-6-19)
42	Oslo	Resp. men	264 (-347-680)	89 (-83-227)	175 (-433-619)
43	Bergen		317 (-355-753)	256 (-390-670)	61 (-36-107)
44	Trondheim		65 (-90-185)	22 (-17-57)	43 (-127-166)
45 46	Stavanger		213 (-329-486)	-4 (-24-2)	217 (-345-513)
47	Drammen		36 (-52-99)	16 (-23-46)	20 (-72-88)
48	Fredrikstad		32 (-111-147)	23 (-124-143)	9 (-8-23)
49	Tromsø		18 (-59-77)	12 (-65-70)	6 (-5-15)
50	Oslo	Resp. women	635 (-821-1731)	532 (-958-1636)	104 (-48-223)
51	Bergen		480 (-476-1036)	461 (-481-1055)	19 (-18-39)
53	Trondheim		147 (-182-391)	122 (-197-368)	25 (-10-52)
54	Stavanger		248 (-252-538)	244 (-232-540)	3 (-1-8)
55	Drammon		270 (-232-330) 105 (-05-358)	274 (-232-340) 89 (-115-221)	3 (- 4 -0) 16 (_7_22)
56	Erodriketad		103 (-33-230) 08 (-61-221)	01 (-01 220)	10 (-7-33) 6 (2-10)
57 58		Resp. 0.74	50 (-04-254)	10 (10 1 10)	0 (2-10)
59	USIU	Nesp. 0-74	JI (-4/-IZ4)	40 (-40-122)	5 (U-5)
60	Bergen		384 (-235-734)	48 (19-69)	530 (-239-704)
	irondheim		429 (-931-585)	452 (-191-663)	-23 (-1236-1)

2 3 4	Stavanger Drammen Fredrikstad		99 (-65-204) 65 (-47-141) 103 (6884 200)	10 (4-14) 6 (1-9) 210 (158 247)	90 (-72-195) 59 (-47-135)
5	Tramad		195 (-0004-509)	219 (-156-547)	-20 (-8112-0)
6	Tromsø	D	41 (-74-115)	3 (-3-6)	39 (-88-113)
7	Oslo	Resp. 75+	24 (-26-58)	2 (-2-4)	22 (-28-55)
8	Bergen		710 (-1086-2197)	649 (-1266-2092)	61 (1-108)
9 10	Trondheim		394 (-561-1105)	348 (-552-1000)	46 (-61-132)
11	Stavanger		176 (-302-554)	158 (-326-523)	18 (2-30)
12	Drammen		200 (-378-550)	186 (-365-517)	14 (-33-48)
13	Fredrikstad		126 (-124-317)	114 (-162-317)	13 (2-21)
14	Tromsø		126 (-79-306)	118 (-98-284)	9 (-5-21)
15			(````````	(,	- ()
16					
17					
19					
20					
21					
22					
23					
24					
25 26					
27					
28					
29					
30					
31					
32					
33 34					
35					
36					
37					
38					
39					
40					
41 47					
43					
44					
45					
46					
47					

http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Table S10. Pooled attributable fractions (%) and number of deaths (AN) for non-accidental, cardiovascular, and respiratory mortality. Brackets indicate 95% empirical confidence intervals.

		AF	AN
Non-accidental	Extreme cold	0.17 (0.09-0.23)	384 (204-534)
mortality	Moderate cold	4.91 (1.73-7.75)	11350 (4007-17899)
	Moderate heat	0.12 (0.05-0.18)	274 (125-407)
	Extreme heat	0.12 (0.06-0.17)	284 (138-396)
CVD mortality	Extreme cold	0.27 (0.14-0.37)	223 (118-310)
	Moderate cold	11.14 (5.92-15.41)	9248 (4920-12794)
	Moderate heat	0.42 (0.00-0.80)	346 (-2-667)
	Extreme heat	0.01 (-0.07-0.08)	11 (-60-68)
Respiratory mortality	Extreme cold	0.28 (-0.03-0.46)	69 (-6-112)
	Moderate cold	4.80 (-7.13-13.83)	1162 (-1727-3350)
	Moderate heat	0.57 (0.01-1.03)	137 (2-249)
	Extreme heat	0.30 (0.13-0.39)	73 (32-95)

http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Figure S1

Overall cumulative exposure–response associations for non-accidental mortality in seven cities by sex. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.



Figure S2

Overall cumulative exposure–response associations for CVD mortality in seven cities by sex. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.





Figure S3

Overall cumulative exposure–response associations for respiratory disease mortality in seven cities by sex. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.



Figure S4

Overall cumulative exposure–response associations for non-accidental mortality in seven cities by age group. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.



Figure S5

Overall cumulative exposure–response associations for CVD mortality in seven cities by age group. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.



Figure S6

Overall cumulative exposure–response associations for respiratory mortality in seven cities by age group. Exposure–response associations as best linear unbiased prediction (with 95% empirical CI, shaded) in the seven cities. RR=Relative risk.



Figure S7

Fractions of non-accidental mortality attributable to moderate and extreme hot and cold temperature by city and age group. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.





http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Figure S8

Fractions of cardiovascular mortality attributable to moderate and extreme hot and cold temperature by city and age group. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.

CVD deaths: 0-74 years >75 years Trondheim Tromsø Stavanger Oslo Fredrikstad Drammen Bergen Percentage (%) Extreme cold Moderate cold Moderate heat Extreme heat



Figure S9

 Fractions of respiratory mortality attributable to moderate and extreme hot and cold temperature by city and age group. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.

Respiratory deaths: 0-74 years >75 years Trondheim Tromsø Stavanger Oslo Fredrikstad Drammen Bergen Percentage (%) Extreme cold Moderate cold Moderate heat Extreme heat



Figure S10

Fractions of non-accidental mortality attributable to moderate and extreme hot and cold temperature by city and sex. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.

Non-accidental deaths: men women Trondheim Tromsø Stavanger Oslo Fredrikstad Drammen Bergen Percentage (%) Extreme cold Moderate cold Moderate heat Extreme heat



Figure S11

Fractions of cardiovascular mortality attributable to moderate and extreme hot and cold temperature by city and sex. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.





Figure S12

Fractions of respiratory mortality attributable to moderate and extreme hot and cold temperature by city and sex. Extreme and moderate high and low temperatures were defined with the minimum mortality temperature and the 1st and 99th percentiles of temperature as cut-offs.





http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Figure S13

 Overall lag structure in the effects of extreme cold temperature on daily cause-specific mortality in seven Norwegian cities, from 1996 to 2018, classified by cause of death. The effects were determined by comparing the risks at -11.2°C (that is, the mean of the 1st centile of temperature distributions) to the mean minimum temperature (MMT). The solid lines represent the mean estimates, while the shaded areas represent the 95% confidence intervals. Natural refers to all non-accidental mortality.



Figure S14

Overall lag structure in the effects of extreme hot temperature on daily cause-specific mortality in seven Norwegian cities, from 1996 to 2018, classified by cause of death. The effects were determined by comparing the risks at 19.5°C (that is, the mean of the 99th centile of temperature distributions) to the estimated minimum mortality temperature (MMT). The solid lines represent the mean estimates, whilst the shaded areas represent the 95% confidence intervals. Natural refers to all non-accidental mortality.



http://mc.manuscriptcentral.com/spub E-mail: sjpheditorial@sagepub.com

Figure S15

Overall lag structure in the effects of extreme cold and hot temperatures on daily non-accidental mortality in seven Norwegian cities, from 1996 to 2018, distinguished by sex. The effects were determined by comparing the risks at -11.2°C and 19.5°C (that is, the mean of the 1st and 99th centiles of temperature distributions) to the estimated minimum mortality temperature. The solid lines represent the mean estimates, while the shaded areas represent the 95% confidence intervals. Natural refers to all non-accidental mortality.



Figure S16

Overall lag structure in the effects of extreme cold and hot temperatures on daily cardiovascular mortality in seven Norwegian cities, from 1996 to 2018, distinguished by sex. The effects were determined by comparing the risks at -11.2°C and 19.5°C (that is, the mean of the 1st and 99th centiles of temperature distributions) to the estimated minimum mortality temperature. The solid lines represent the mean estimates, while the shaded areas represent the 95% confidence intervals.



Figure S17

Overall lag structure in the effects of extreme cold and hot temperatures on daily respiratory disease mortality in seven Norwegian cities, from 1996 to 2018, distinguished by sex. The effects were determined by comparing the risks at -11.2°C and 19.5°C (that is, the mean of the 1st and 99th centiles of temperature distributions) to the estimated minimum mortality temperature. The solid lines represent the mean estimates, while the shaded areas represent the 95% confidence intervals.







References

Gasparrini A, Guo Y, Hashizume M, Lavigne E, Zanobetti A, Schwartz J, et al. Mortality risk 1. attributable to high and low ambient temperature: a multicountry observational study. Lancet (London, England). 2015;386(9991):369-75.

.rature: a π. .s.