

3-BRAID KNOTS WITH MAXIMAL 4-GENUS

SEBASTIAN BAADER, LUKAS LEWARK, FILIP MISEV, AND PAULA TRUÖL

ABSTRACT. We classify 3-braid knots whose topological 4-genus coincides with their Seifert genus, using McCoy’s twisting method and the Xu normal form. In addition, we give upper bounds for the topological 4-genus of positive and strongly quasipositive 3-braid knots.

1. INTRODUCTION

Four decades after Freedman’s celebrated work on 4-manifolds [Fre82], the topological 4-genus $g_4^{\text{top}}(K)$ of knots K remains difficult to determine. The first challenge is posed by the figure-eight knot 4_1 , which satisfies $g_4^{\text{top}}(4_1) = 1$, although its second power $4_1 \# 4_1$ bounds an embedded disc in the 4-ball. This causes all its additive lower bounds on the 4-genus to be trivial. In particular, the signature invariant $\sigma(4_1)$ is zero. Due to this example, there is little reason to believe that the inequality

$$|\sigma(K)| \leq 2g_4^{\text{top}}(K)$$

has much to tell us about the (topological) 4-genus of knots in general. In this note, we show that closures of 3-braids are exceptional in this respect. More precisely, we will see that the figure-eight knot is exceptional among closed 3-braids in that it is the only 3-braid knot K that satisfies $|\sigma(K)| < 2g(K)$, yet $g_4^{\text{top}}(K) = g(K)$, where $g(K)$ denotes the ordinary Seifert genus.

Theorem 1.1. *Let K be a 3-braid knot other than the figure-eight knot. Then*

$$|\sigma(K)| = 2g(K) \iff g_4^{\text{top}}(K) = g(K).$$

These equalities hold precisely if K or its mirror is one of the following knots:

- $T(2, 2m + 1) \# T(2, 2n + 1)$, with $m, n \geq 0$,
- $P(2p, 2q + 1, 2r + 1, 1)$, with $p \geq 1, q, r \geq 0$,
- $T(3, 4)$, or $T(3, 5)$.

The question arises whether the equivalence of $|\sigma(K)| = 2g(K)$ and $g_4^{\text{top}}(K) = g(K)$ holds for other families of knots K . Indeed, it is also true for braid positive knots K [Lie16]. Moreover, we do not know if there exists a knot K of braid index 4 with $g_4^{\text{top}}(K) = g(K)$, but $|\sigma(K)| < 2g(K)$. One may check that such a knot would have to be prime and have crossing number at least 13. For braid index 5 however, there are several knots K in the table with $\sigma(K) = 0 < 2g(K) = 2$ and $g_4^{\text{top}}(K) = g(K) = 1$, such as $K = 8_1$ [LM].

Received by the editors April 21, 2023, and, in revised form, November 22, 2023.

2020 *Mathematics Subject Classification.* Primary 57K10.

The second author was supported by the Emmy Noether Programme of the DFG, Project number 412851057. The fourth author gratefully acknowledges support from the Swiss National Science Foundation Grant 181199.

The proof of Theorem 1.1 is based on a technique called twisting, used by McCoy for estimating the topological 4-genus from above [McC19]. We will also make use of a special presentation for 3-braids called Xu normal form, which we will describe in the next section. The third section contains the proof of Theorem 1.1, as well as a complementary result (Theorem 3.2): a sharp lower bound on the difference $g(K) - g_4^{\text{top}}(K)$ of so-called strongly quasipositive 3-braid knots, in terms of two characteristic quantities associated with their Xu normal form. In the last section, we determine the topological 4-genus of various families of braid positive 3-braid knots (almost) exactly, and display examples where our technique comes to a limit.

2. THE XU NORMAL FORM OF 3-BRAIDS

Our tool to handle 3-braids is what we call their *Xu normal form*. It was developed by Xu [Xu92] (who called it *representative symbol*), as a variation of the Garside normal form [Gar69]. Using the Xu normal form, one may decide whether two given 3-braids are conjugate [Xu92], and whether their closures are equivalent links [BM93, BM08]. Later, the Xu normal form was generalized to braids on arbitrarily many strands by Birman–Ko–Lee [BKL98]. The Garside, Xu and BKL normal forms are all examples of Garside structures on (braid) groups [DDG⁺15]. In this section, we will introduce the Xu normal form and show how it determines the signature invariant σ of its closure (Proposition 2.9), as well as strong quasipositivity and braid positivity (Proposition 2.11).

A *3-braid* β is an element of the braid group $B_3 = \langle a, b \mid aba = bab \rangle$. By closing off the three strands of β and interpreting a and b as positive crossings between the first two and last two strands, respectively, β gives rise to a link called *closure of β* , denoted $\text{cl}(\beta)$. Note that throughout the text, links are oriented. Let us write $x := a^{-1}ba \in B_3$ and $\delta := ba = ax = xb \in B_3$. By a *Xu word* or simply *word*,




FIGURE 1. The generators a and b of the braid group B_3 and the elements $x = a^{-1}ba$, $\delta = ba$ and $\Delta = aba$. The latter is used for the Garside normal form (further down in the text).

we mean a word with letters a, b, x, δ (see Figure 1) and their inverses. We reserve the equality sign $=$ for equality of braids, and use a dotted equality sign \doteq for equality of words. Moreover, we write $\beta \sim \gamma$ if the two braids β, γ are conjugate. For efficiency, let us also introduce the following notation: for any $i \in \mathbb{Z}$, set $\tau_i \doteq a$ if $i \equiv 1 \pmod{3}$, $\tau_i \doteq b$ if $i \equiv 2 \pmod{3}$ and $\tau_i \doteq x$ if $i \equiv 0 \pmod{3}$.

Definition 2.1. Let w be a word of the form

$$(1) \quad w \doteq \delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t} \quad \text{for } n \in \mathbb{Z}, t \geq 0, u_i \geq 1.$$

We say that w is in *Xu normal form* if the tuple $(-n, t, u_1, \dots, u_t)$ is lexicographically minimal among all words of the form (1) representing the same conjugacy class of 3-braids.

The condition of lexicographic minimality means, in particular, that the Xu normal form maximizes n , and afterwards minimizes t . The term ‘normal form’ is justified by the following.

Theorem 2.2. *Every 3-braid is conjugate to a unique word in Xu normal form. \square*

Lemma 2.3 gives a criterion to easily decide whether a word is in Xu normal form.

Lemma 2.3. *A word $w \doteq \delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ for some $n \in \mathbb{Z}$, $t \geq 0$, $u_i \geq 1$ is in Xu normal form if and only if one of the following conditions holds:*

- (a) $t = 0$. In this case $w \doteq \delta^n$.
- (b) $t = 1$, and if $n \equiv 1 \pmod{3}$ then $u_1 = 1$. In this case, $w \doteq \delta^{3k} a^{u_1}$, or $w \doteq \delta^{3k+1} a$, or $w \doteq \delta^{3k+2} a^{u_1}$.
- (c) $t \geq 2$, $n + t \equiv 0 \pmod{3}$ and the tuple (u_1, \dots, u_t) is lexicographically minimal among its cyclic permutations.

The proofs of Theorem 2.2 and Lemma 2.3 are essentially contained in Xu’s paper [Xu92, Section 4], albeit with slightly different conventions. In our setup, Theorem 2.2 is not actually hard to prove, and makes a good exercise to get acquainted with the calculus of Xu words. The same is true for the ‘only if’ direction of Lemma 2.3. Let us provide two hints. Firstly, the easily verifiable rules

$$(2) \quad \delta = \tau_i \tau_{i-1}, \quad \tau_i \delta^n = \delta^n \tau_{i+n}, \quad \tau_i^{-1} \delta^n = \delta^{n-1} \tau_{i+n+1}$$

allow to find Xu words for every 3-braid without the letters a^{-1}, b^{-1}, x^{-1} , and to ‘pull all $\delta^{\pm 1}$ to the left’ in a Xu word. In this way, one can find a Xu word of the form $\delta^n \tau_{m+1}^{u_1} \dots \tau_{m+t}^{u_t}$ with $m \in \mathbb{Z}$ and $u_i \geq 1$ for any 3-braid. Secondly, note that

$$\begin{aligned} \delta^n \tau_1^{u_1} \dots \tau_t^{u_t} &= \tau_{1-n}^{u_1} \delta^n \tau_2^{u_2} \dots \tau_t^{u_t} \sim \delta^n \tau_2^{u_2} \dots \tau_t^{u_t} \tau_{1-n}^{u_1} \\ &\sim \delta^{n+1} \tau_2^{u_2} \dots \tau_t^{u_t} \tau_{1-n}^{u_1} \delta^{-1} = \delta^n \tau_1^{u_2} \dots \tau_{t-1}^{u_t} \tau_{-n}^{u_1}. \end{aligned}$$

So cyclically permuting the tuple (u_1, \dots, u_t) results in a conjugate braid if $-n \equiv t \pmod{3}$.

Proof of the ‘if’ direction of Lemma 2.3. Xu proves that condition (c) is sufficient for w to be in Xu normal form (see the definition of the representative symbol and Theorem 5 in [Xu92]), but omits a discussion of conditions (a) and (b). So suppose w satisfies (a) or (b), $w' = \delta^{n'} \tau_1^{u'_1} \dots \tau_{t'}^{u'_{t'}}$ is in Xu normal form, and $w' \sim w$. We need to show that $w \doteq w'$. Let us distinguish the following cases.

- If w satisfies (a), then $w \doteq \delta^n$. The words w and w' must have the same writhe, that is, the same image under the abelianisation homomorphism $B_3 \rightarrow \mathbb{Z}$, which maps both a and b to 1. Therefore, $2n = 2n' + u'_1 + \dots + u'_{t'}$. But since w' is in Xu normal form, $n' \geq n$. Because $u'_i \geq 0$, it follows that $n' = n$ and $t' = 0$, thus $w \doteq w'$ as desired.

- If w satisfies (b) and $w \doteq \delta^n a$, a similar argument applies: the equality of the writhes of w and w' now reads as $2n + 1 = 2n' + u'_1 + \dots + u'_{t'}$. Since the left hand side is odd, we must have $t' \geq 1$ and $u'_1 \geq 1$, so $2n + 1 \geq 2n' + 1$. On the other hand $n' \geq n$, again because w' is in Xu normal form, so $2n + 1 \leq 2n' + 1$. It follows that $n' = n$, $t' = 1$, $u'_1 = 1$ and $w \doteq w'$ as desired.

- The remaining case is that $w \doteq \delta^n a^{u_1}$ satisfies (b), $u_1 \geq 2$, and $n \not\equiv 1 \pmod{3}$. Now the so-called r -index of w defined below Lemma 5 in [Xu92] is 0,

and so [Xu92, Theorem 4] implies that n is maximal. It follows that $n' = n$, and hence the minimality of t' implies $t' \leq t = 1$. Finally, from the equality of the writhes of w and w' , it follows that $t' = 1$ and $u_1 = u'_1$, and thus $w \doteq w'$. \square

To get a link invariant from the Xu normal form, we need to understand the relationship between conjugacy classes of 3-braids and link equivalence classes of their closures. Birman–Menasco have shown that with a few well-understood exceptions, this relationship is one-to-one:

Theorem 2.4 ([BM93, BM08]). *Two 3-braids are conjugate if their closures are equivalent links, except in the following cases:*

- (1) *The non-conjugate braids $ab, ab^{-1}, a^{-1}b^{-1}$ have the unknot as closure.*
- (2) *For $n \in \mathbb{Z} \setminus \{\pm 1\}$, the non-conjugate braids $a^n b, a^n b^{-1}$ have the $T(2, n)$ torus link as closure.*
- (3) *For pairwise distinct integers $p, q, r \in \mathbb{Z} \setminus \{0, -1, -2\}$, the two non-conjugate braids $\beta = a^p b^q x^r$ and $\gamma = a^p b^r x^q$ have the $P(p, q, r, 1)$ pretzel link as closure (see Figure 2); and the two non-conjugate braids β^{-1} and γ^{-1} have the $P(-p, -q, -r, -1)$ pretzel link as closure.*

Corollary 2.5 allows us to sidestep the exceptional cases (1), (2), (3) in the above theorem by focusing on links of braid index 3 instead of 3-braid links (the latter class of links includes links with braid index 1 and 2, i.e. 2-stranded torus links). Let the *reverse* of a braid $\beta \in B_3$, denoted by $\text{rev}(\beta)$, be the braid given by reading β backwards and switching a with b , and a^{-1} with b^{-1} . Note that $\text{cl}(\text{rev}(\beta))$ is obtained from $\text{cl}(\beta)$ by reversing the link’s orientation.

Corollary 2.5. *Let L be a link of braid index 3.*

- (1) *Either there is a unique conjugacy class of 3-braids with closure L , or there are two of them, such that one consists of the reverses of the braids contained in the other.*
- (2) *The numbers n, t , and $U = u_1 + \dots + u_t$ of the Xu normal form of a braid with closure L do not depend on the choice of braid. Thus n, t and U are link invariants of links with braid index 3.*




FIGURE 2. Isotopy (denoted \approx) from the closure of the braid $a^{u_1} b^{u_2} x^{u_3}$ to the pretzel knot $P(u_1, 1, u_2, u_3) \approx P(u_1, u_2, u_3, 1)$; here $(u_1, u_2, u_3) = (4, 3, 5)$.

Proof. Claim (1) follows quickly from Theorem 2.4, since the unknot and the two-stranded torus links have braid index less than 3, and $\text{rev}(a^p b^q x^r) = x^r a^p b^q$ which is conjugate to $a^p b^r x^q$. For (2), note that $\text{rev}(x) = x$ and so $\text{rev}(\tau_i) = \tau_{-i}$. Also $\text{rev}(\delta) = \delta$. Thus the reverse of $\delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ is $\tau_{-t}^{u_t} \dots \tau_{-1}^{u_1} \delta^n$, which has Xu normal form $\delta^n \tau_1^{u_t} \tau_2^{u_{t-1}} \dots \tau_t^{u_1}$ (up to cyclically permuting the exponents u_t, \dots, u_1). So the numbers n, t and U do not change under braid reversal. Together with (1), this implies (2). \square

Xu calls n and t the *power* and *syllable length*, while Birman–Ko–Lee use the terms *infimum* and *canonical length*, respectively.

Since the Xu normal form determines the link type, all link invariants may be read off it. Let us first prove a formula for the signature invariant. We will need the Garside normal form [Gar69], which we introduce now. The reader will note many parallels between the Garside and Xu normal forms. Let $\Delta := aba$, as in Figure 1 on the right. A *Garside word* is a word with letters a, b, Δ and their inverses. Again, we use \doteq for equality of words. We also use the following notation: for any $i \in \mathbb{Z}$, set $\sigma_i \doteq a$ if $i \equiv 1 \pmod{2}$ and $\sigma_i \doteq b$ if $i \equiv 0 \pmod{2}$.

Proposition 2.6 ([Tru23, Proposition 3.2]). *Every 3-braid contains in its conjugacy class a unique Garside word v in Garside normal form, i.e. a word*

$$v \doteq \Delta^\ell \sigma_1^{p_1} \sigma_2^{p_2} \dots \sigma_r^{p_r},$$

with $\ell \in \mathbb{Z}, r \geq 0, p_i \geq 1$, satisfying one of the following conditions:

- (A) ℓ is even and $r \in \{0, 1\}$, i.e. $v \doteq \Delta^{2k} a^{\geq 0}$,
- (B) ℓ is even, $r = 2, p_1 \in \{1, 2, 3\}$ and $p_2 = 1$, i.e. $v \doteq \Delta^{2k} a^{\{1,2,3\}} b$,
- (C)/(D) $r \geq 1, p_i \geq 2, \ell \equiv r \pmod{2}$, and the tuple (p_1, \dots, p_r) is lexicographically minimal among its cyclic permutations.

We refer by (C) and (D) to the case that ℓ is even and odd, respectively.

Lemma 2.7 tells us how to convert between the Xu and the Garside normal forms.

Lemma 2.7. *Let a word $w \doteq \delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ in Xu normal form be given. Then the unique word v in Garside normal form representing the same conjugacy class of 3-braids as w is given by the following table.*

Case in Lemma 2.3	Xu normal form w	Garside normal form v	Case in Proposition 2.6
(a)	δ^{3k}	Δ^{2k}	(A)
(a)	δ^{3k+1}	$\Delta^{2k} ab$	(B)
(a)	δ^{3k+2}	$\Delta^{2k} a^3 b$	(B)
(b)	$\delta^{3k} a^{u_1}$	$\Delta^{2k} a^{u_1}$	(A)
(b)	$\delta^{3k+1} a$	$\Delta^{2k} a^2 b$	(B)
(b)	$\delta^{3k+2} a^{u_1}$	$\Delta^{2k+1} a^{1+u_1}$	(D)
(c)	$\delta^n \tau_1^{u_1} \dots \tau_t^{u_t}$	$\Delta^{(2n-t)/3} \sigma_1^{1+u_1} \dots \sigma_t^{1+u_t}$	(C)/(D)

Proof. All rows in the table except for the last one may be checked quickly, using $\delta^3 = \Delta^2$. Let us now prove $w \sim v$ for the words w and v in the last row. Let $n = 3k + m$ and $t = 3s + 3 - m$ for $k, s \in \mathbb{Z}, s \geq 0, m \in \{1, 2, 3\}$. In the Xu word w , replace δ^n by $(ba)^m \Delta^{2k}$. Moreover, replace every x^u by $\Delta^{-1} ab^{1+u} a$. These replacements yield a Garside word v_1 with $v_1 = w$ and

$$v_1 \doteq (ba)^m \Delta^{2k} a^{u_1} b^{u_2} (\Delta^{-1} ab^{1+u_3} a) a^{u_4} \dots (\Delta^{-1} ab^{1+u_{3s}} a) a^{u_{3s+1}} b^{u_{3s+2}},$$

where we set $u_i = 0$ if $i > t$. Now proceed by ‘pulling all the Δ^{-1} to the right’, i.e. replacing $\Delta^{-1} a$ by $b\Delta^{-1}$ and $\Delta^{-1} b$ by $a\Delta^{-1}$ as long as possible. These replacements produce a word v_2 with $v_2 = v_1$, where v_2 starts with $(ba)^m \Delta^{2k} a^{u_1} b^{u_2} (ba^{1+u_3} b) b^{u_4} \dots$. Using the σ_i -notation and noting that there are precisely s occurrences of Δ^{-1} in v_1 , we have

$$\begin{aligned} v_2 &\doteq (ba)^m \Delta^{2k} \sigma_1^{u_1} \sigma_2^{1+u_2} \sigma_3^{1+u_3} \sigma_4^{1+u_4} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1}^{1+u_{3s+1}} \sigma_{3s+2}^{u_{3s+2}} \Delta^{-s} \\ &\sim v_3 \doteq \Delta^{-s} (ba)^m \Delta^{2k} \sigma_1^{u_1} \sigma_2^{1+u_2} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1}^{1+u_{3s+1}} \sigma_{3s+2}^{u_{3s+2}}. \end{aligned}$$

Let us now consider the three possibilities for m case by case.

– If $m = 3$, then $u_{3s+2} = u_{3s+1} = 0$ and

$$\begin{aligned} v_3 &\doteq \Delta^{-s} (ba)^3 \Delta^{2k} \sigma_1^{u_1} \sigma_2^{1+u_2} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1} \\ &= \Delta^{2k+2-s} \sigma_1^{u_1} \sigma_2^{u_2+1} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1} \\ &\sim v_4 = \Delta^{2k+2-s} \sigma_1^{1+u_1} \sigma_2^{u_2+1} \dots \sigma_{3s}^{1+u_{3s}}. \end{aligned}$$

We have $v_4 \doteq v$ as desired, since $2k + (m - 1) - s = (2n - t)/3$.

– If $m = 2$, then $u_{3s+2} = 0$ and

$$\begin{aligned} v_3 &\doteq \Delta^{-s} (ba)^2 \Delta^{2k} \sigma_1^{u_1} \sigma_2^{1+u_2} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1}^{1+u_{3s+1}} \\ &= \Delta^{2k+1-s} \sigma_1^{1+u_1} \sigma_2^{1+u_2} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1}^{1+u_{3s+1}} \doteq v. \end{aligned}$$

– If $m = 1$, then

$$\begin{aligned} v_3 &\doteq \Delta^{-s} ba \Delta^{2k} \sigma_1^{u_1} \sigma_2^{1+u_2} \dots \sigma_{3s}^{1+u_{3s}} \sigma_{3s+1}^{1+u_{3s+1}} \sigma_{3s+2}^{u_{3s+2}} \\ &= \sigma_s \Delta^{2k-s} \sigma_1^{1+u_1} \sigma_2^{1+u_2} \dots \sigma_{3s+1}^{1+u_{3s+1}} \sigma_{3s+2}^{u_{3s+2}} \sim v. \end{aligned}$$

□

We rely on the signature formula for 3-braids in Garside normal form deduced by the fourth author [Tru23, Remark 1.6, Proposition 4.2, Remark 4.3] from a result by Erle [Erl99, Theorem 2.6].

Proposition 2.8 ([Tru23]). *Let K be a knot that is the closure of a Garside normal form $\Delta^\ell \sigma_1^{p_1} \dots \sigma_r^{p_r}$ in case (C)/(D) of Proposition 2.6. Then*

$$\sigma(K) = -2\ell + r - \sum_{i=1}^r p_i.$$

We are now ready to state and prove our signature formula for 3-braids in Xu normal form.

Proposition 2.9. *Let $\delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ be the Xu normal form of a 3-braid whose closure is a knot K of braid index 3. Set $U = u_1 + \dots + u_t$. If $t > 0$ (equivalently, if K is not a torus knot), then*

$$\sigma(K) = -U - \frac{4}{3}n + \frac{2}{3}t.$$

In the case $t = 0$, i.e. $U = 0$ and $K = T(3, n)$, the value $\sigma(K) = -\frac{4}{3}n$ given by the above formula is only approximately true, with an error of at most $\frac{4}{3}$. In fact, in that case we have

$$\sigma(K) = -\frac{4}{3}n + \left(2 + 4 \left\lfloor \frac{1}{6}n \right\rfloor - \frac{2}{3}n\right) = 2 - 2n + 4 \left\lfloor \frac{1}{6}n \right\rfloor.$$

Proof. Our signature formula for torus knots may be seen to agree with the formula given e.g. in [Mur74, Proposition 9.1]. So we are left with the case $t \geq 1$, i.e. cases (b) and (c) in Lemma 2.3. Denote the Xu normal form in question by w . Let us first consider case (c). Then w has Garside normal form $\Delta^{(2n-t)/3} \sigma_1^{1+u_1} \dots \sigma_t^{1+u_t}$, see Lemma 2.7. By Proposition 2.8, we have $\sigma(K) = -4n/3 + 2t/3 + t - (U + t)$, which is equal to the claimed formula. In case (b), since the closure of w is a knot, the only possibility is $w \doteq \delta^{3k+2} a^{u_1}$. Then the Garside normal form of w is $\Delta^{2k+1} a^{1+u_1}$, which has the desired signature, again by Proposition 2.8. \square

Next, let us give complete criteria to decide braid positivity and strong quasipositivity for links of braid index 3. A *braid positive* link is the closure of some *positive word*, i.e. a word in positive powers of the standard generators a_1, \dots, a_{k-1} of the braid group B_k on some number k of strands. Similarly, a link is called *strongly quasipositive* [Rud83, Rud90] if it is the closure of a *strongly quasipositive word* in some B_k , i.e. a word in positive powers of

$$a_{ij} = a_i^{-1} a_{i+1}^{-1} \dots a_{j-1}^{-1} a_j a_{j-1} \dots a_i$$

with $1 \leq i \leq j \leq k-1$. Note that for $k = 3$, positive words are words in $a = a_1$, $b = a_2$ and strongly quasipositive words are words in $a = a_{11}$, $b = a_{22}$ and $x = a_{12}$. It is well-known and straightforward to show that a k -braid can be written as a (strongly quasi-)positive braid word if and only if the power of Δ (δ) in its Garside (Birman–Ko–Lee) normal form is non-negative, respectively. This makes (strong quasi-)positivity decidable for braids. For links however, the problem is harder because a priori, a braid positive (strongly quasipositive) link with braid index k need not be the closure of a (strongly quasi-)positive word on k strands. For $k = 3$, however, this is the case:

Theorem 2.10 ([Sto17, Theorem 1.1 and 1.3]). *The following hold.*

- (1) *If a strongly quasipositive link is the closure of some 3-braid, then it is the closure of a strongly quasipositive 3-braid.*
- (2) *If a braid positive link is the closure of some 3-braid, then it is the closure of a positive 3-braid.*

We are now ready to prove our positivity characterizations. For braid positivity, we will once again need the Garside normal form introduced above.

Proposition 2.11. *Let $\delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ be in Xu normal form, with closure a link L of braid index 3. Then the following hold.*

- (1) *L is a strongly quasipositive link if and only if $n \geq 0$.*
- (2) *L is a braid positive link if and only if $n \geq t/2$ or $n = 0, t = 1$.*

Proof. Part (1). If $n \geq 0$, then the Xu normal form yields a strongly quasipositive word for L , so L is strongly quasipositive. For the other direction, assume L is strongly quasipositive and a Xu normal form w with closure L is given. By Theorem 2.10, there is a strongly quasipositive word in B_3 representing L . It

may be transformed to its Xu normal form w' just by replacing $\tau_{i+1}\tau_i \rightarrow \delta$ and $\tau_i\delta \leftrightarrow \delta\tau_{i+1}$, and by passing from $y\tau_i$ to $\tau_i y$ for y a word in positive powers of a, b, x, δ . None of these transformations create negative powers of δ , and so we find that the Xu normal form w' has $n \geq 0$. By Corollary 2.5(2), all Xu normal forms with closure L have the same n , so w has $n \geq 0$ as well, which was to be proven.

Part (2). If $n \geq t/2$, then one may check using Lemma 2.7 that the Garside normal form of $\delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t}$ starts with a non-negative power of Δ , and thus yields a positive word with closure L . If $n = 0$ and $t = 1$, then the Xu normal form is a^{u_1} , which is already a positive word with closure L . In both cases, it follows that L is braid positive. For the other direction, assume L is braid positive and let a Xu normal form w with closure L be given. By Theorem 2.10, there is a positive word in B_3 representing L . Similarly as in the proof of part (1), one sees that the Garside normal form of this positive word starts with Δ^ℓ with $\ell \geq 0$. By going through the rows of the table in Lemma 2.7, one sees that this implies that $n \geq t/2$, with the sole exception in the fourth row if $k = 0$ and $u_1 \neq 0$: then, the Xu normal form is a^{u_1} , so $n = 0$ and $t = 1$. Again using Corollary 2.5(2), it follows that w also satisfies $n \geq t/2$, or $n = 0$ and $t = 1$. \square

If a knot K is the closure of a strongly quasipositive 3-braid in Xu normal form $\delta^n \tau_1^{u_1} \dots \tau_t^{u_t}$, there is also a simple formula for the Seifert genus of K :

$$(3) \quad g(K) = \frac{U}{2} + n - 1,$$

where $U = u_1 + \dots + u_t$. This follows from the Bennequin inequality [Ben83].

3. PROOFS OF MAIN THEOREMS

Before beginning with the proof of Theorem 1.1, we describe a technique that we use to detect topological 4-genus defect in a given knot K , that is, to show $g_4^{\text{top}}(K) < g(K)$. The main ingredient is the so-called *generalized crossing change*, also known as *null-homologous twist*, or simply *twist*. A null-homologous twist consists in performing a ± 1 Dehn surgery on the boundary circle of an embedded disc $D \subset S^3$, such that D intersects K transversely in a finite number of interior points, with total algebraic intersection count 0. While ± 1 Dehn surgery on an unknot in S^3 gives back S^3 , the effect on K is that a (left- or right-handed) full twist is introduced into the strands of K that cross D (cf. Figure 3). The *untwisting number* $\text{tu}(K)$ of K , introduced by Ince [Inc16], is defined as the minimal number of null-homologous twists needed to turn K into the unknot. Clearly, $\text{tu}(K) \leq u(K)$, since crossing changes are special cases of null-homologous twists. McCoy [McC19]




FIGURE 3. Two examples of twists, at the locations marked by \star . Here, the boundary of the respective disc D is drawn in blue; in subsequent figures, it will be omitted.




FIGURE 4. Left: Saddle move. Middle: How to use isotopy and a saddle move to add or remove a braid crossing. Right: Example of a cobordism between $cl(\alpha)$ and $cl(\alpha \cdot abx)$ that consists of three saddle moves, for some 3-braid α .

showed that $g_4^{\text{top}}(K) \leq \text{tu}(K)$, that is, the untwisting number of K is an upper bound on the topological 4-genus of K . His result is based on Freedman’s theorem, which states that knots of Alexander polynomial 1 are topologically slice [Fre82, FQ90]. This bound can now be used to find topological 4-genus defect: If we find a way to turn a knot K into the unknot with strictly less than $g(K)$ null-homologous twists, this will show $g_4^{\text{top}}(K) < g(K)$. This method was already applied by Baader–Banfield–Lewark [BBL20] to 3-stranded torus knots. For the proof of Theorem 1.1 below, we use a slightly refined version of the method, as follows.

Assume we find a cobordism $C \subset S^3 \times [0, 1]$ from a given knot K to some knot K' such that $g(C) = g(K) - g(K')$. If such a C exists, we will write $K \rightsquigarrow K'$. Assume that further $\text{tu}(K') < g(K')$. Then, by McCoy’s result, $g_4^{\text{top}}(K') \leq \text{tu}(K') < g(K')$. Composing the cobordism C with a topological slice surface for K' , we obtain

$$g_4^{\text{top}}(K) \leq g_4^{\text{top}}(K') + g(C) < g(K') + g(C) = g(K).$$

In particular, for the topological 4-genus defect we find

$$(4) \quad g(K) - g_4^{\text{top}}(K) \geq g(K') - g_4^{\text{top}}(K') \geq g(K') - \text{tu}(K').$$

One way to construct cobordisms is to apply saddle moves to knot diagrams, as in Figure 4. Note that such cobordisms will always be smooth. Suppose the knot K is the closure of a strongly quasipositive 3-braid of the form $\beta = \delta^n \tau_1^{u_1} \dots \tau_t^{u_t}$ with $n \geq 0$ and $u_1, \dots, u_t \geq 1$. Out of saddle moves, one may build a cobordism C that lowers the exponents u_i and n (one saddle move per letter τ_i , two saddle moves per letter δ), or transforms δ into τ_i (one saddle move). Suppose the exponents remain non-negative, and C is a cobordism from K to another knot K' . Then K' is also strongly quasipositive, and it follows from the Bennequin inequality, see (3), that $g(C) = g(K) - g(K')$, i.e. $K \rightsquigarrow K'$.

Proof of Theorem 1.1. We organize the proof into two parts, which consist in verifying the following statements.

- (1) $|\sigma(K)| = 2g(K)$ for all K in the list of Theorem 1.1 and their mirrors,
- (2) $g_4^{\text{top}}(K) < g(K)$ for all other 3-braid knots except the figure-eight.

In light of Kauffman and Taylor’s signature bound $|\sigma(J)| \leq 2g_4^{\text{top}}(J)$, valid for all knots J , see [KT76, Pow17], these two statements together imply the theorem.

Part (1). The genera and signatures of torus knots are well understood [Lit79]; in particular, we know that the torus knots K with $|\sigma(K)| = 2g(K)$ are precisely the knots $T(3, 4), T(3, 5), T(2, 2n + 1), n \geq 0$, and their mirrors. In fact, the signature of $T(2, 2n + 1), n \geq 0$, is known to be $-2n$. Both Seifert genus and signature are additive under connected sum of knots; hence $|\sigma(K)| = 2g(K)$ for all knots K of the first type listed. If K is one of the listed pretzel knots $K = P(2p, 2q + 1, 2r + 1, 1)$ with $p \geq 1, q, r \geq 0$, then K has a positive and alternating, hence special alternating diagram; see Figure 2. Murasugi shows in this case that $|\sigma(K)| = 2g(K)$; see [Mur65, Corollary 10.3].

Part (2). Let K be a 3-braid knot other than the figure-eight knot such that neither K nor its mirror appears in the list of Theorem 1.1. The goal is to show that $g_4^{\text{top}}(K) < g(K)$. We distinguish several cases.

- If K is the closure of a positive 3-braid, this is a special case of the analogous statement about all positive braids, on any number of strands, by Liechi [Lie16, Theorem 1, Corollary 2].

- Next, we consider the case in which K is strongly quasipositive without being braid positive; this is the main part of the proof. By Lemma 2.3 and Proposition 2.11, K is the closure of a 3-braid β in Xu normal form $\beta = \delta^n \tau_1^{u_1} \tau_2^{u_2} \cdots \tau_t^{u_t}$ with $u_1, \dots, u_t \geq 1, n \geq 0, t \geq 2, n + t \equiv 0 \pmod 3$, and $2n < t$ (note that the case $n = 0, t = 1$ is excluded since we assume that $\text{cl}(\beta)$ is a knot). These conditions on n and t leave the following possibilities: $(n, t) = (0, 3)$, or $(n, t) = (1, 5)$, or $t \geq 6$. First, if $(n, t) = (0, 3)$, then $K = P(u_1, u_2, u_3, 1)$; see Figure 2. If more than one of u_1, u_2, u_3 is even, K is a link of more than one component; the same is true if all three parameters are odd. We may therefore assume that $u_1 = 2p$ is even and $u_2 = 2q + 1, u_3 = 2r + 1$ are odd, with $p \geq 1, q, r \geq 0$, in which case K is a pretzel knot from the list, which we excluded.

Secondly, if $(n, t) = (1, 5)$, we have $\beta = \delta a^{u_1} b^{u_2} x^{u_3} a^{u_4} b^{u_5}$. If the exponents u_i are all odd, β closes to a two component link, a contradiction to K being a knot. Therefore at least one of the u_i is even. We may assume that u_1 is even, because the exponents u_1, u_2, \dots, u_t may be cyclically permuted without changing the braid closure, as explained in Section 2 after Lemma 2.3. In particular, we may assume that $u_1 \geq 2$. Then, since $\delta = xb$,

$$\beta \sim a^{u_1} b^{u_2} x^{u_3} a^{u_4} b^{u_5} x b \rightsquigarrow a^2 b x a b x b = a(abx)^2 b \rightsquigarrow ab,$$

where ‘ \sim ’ denotes conjugation, ‘ \rightsquigarrow ’ denotes the existence of a cobordism whose genus equals the difference of the Seifert genera of the knots it connects and ‘ \rightsquigarrow ’ is shown in Figure 5. Here, the cobordism ‘ \rightsquigarrow ’ is built from saddles decreasing the exponents u_i . Since the closure of $a(abx)^2 b$ has Seifert genus 3 by (3), while only 2 twists are used in ‘ \rightsquigarrow ’, we obtain $g_4^{\text{top}}(K) \leq g(K) - 1 < g(K)$.

Finally, if $t \geq 6$, we proceed similarly as in the previous case. Again, the parity of the exponents u_1, u_2, \dots, u_t determines whether β closes to a knot or a multi-component link. Indeed, a brief case study shows that the closure K of β is a link of two or three components whenever all the u_i are odd. We may therefore assume via conjugation that u_1 is even. If $t > 6$ or $n > 0$, then

$$\beta \rightsquigarrow a^{u_1} b^{u_2} x^{u_3} a^{u_4} b^{u_5} x^{u_6} b \rightsquigarrow a(abx)^2 b \rightsquigarrow ab.$$

If $t = 6$ and $n = 0$, then $\beta = a^{u_1} b^{u_2} x^{u_3} a^{u_4} b^{u_5} x^{u_6}$. In this case, in order for K to be a knot, at least two of the exponents u_1, \dots, u_6 , say u_i and u_j , need to be even. We may assume that $(i, j) = (1, 2)$ or $(i, j) = (1, 4)$. To see this, use cyclic permutation (as in the case $(n, t) = (1, 5)$) and the fact that u_1, u_3 cannot be the only even exponents, again because β would not close to a knot if they were. Therefore, smooth cobordisms bring us to $a^2 b^2 x a b x$ or to $a^2 b x a^2 b x$, whose closures have Seifert genus 3 by (3). In the first case,

$$a^2 b^2 x a b x = a^2 b \delta^{-1} \delta b x a b x = a^2 b \delta^{-1} b (a b x)^2 \rightsquigarrow a^2 b \delta^{-1} b = a^2 b a^{-1} \sim ab.$$

For the second case, Figure 6 shows how to turn $a(abx a^2 b x)$ into $a\gamma$ using two twists. Here, γ is the tangle shown in the top right corner of the figure. Note that $a\gamma$ describes the unknot when closed like a braid. Since the closure of $a^2 b x a^2 b x$ has Seifert genus 3 (see (3)), we obtain $g_4^{\text{top}}(K) \leq g(K) - 1 < g(K)$. This concludes the case that K is strongly quasipositive.

– If the mirror of K is strongly quasipositive, we apply the above argument to the mirror of K ; since both g_4^{top} and g are invariant under taking mirror images, we obtain $g_4^{\text{top}}(K) < g(K)$ again.

– If K or its mirror is the knot $T(2, 2m + 1) \# T(2, -2n - 1)$, with $m \geq n \geq 1$, it has a Seifert surface S of genus $m + n = g(K)$ that contains a copy of the ribbon knot $R := T(2, 2n + 1) \# T(2, -2n - 1)$, bounding a subsurface of S of genus $g(R) = 2n$. A surgery that cuts this subsurface off S and replaces it with a slice disk for R gives rise to a smooth surface of genus $m + n - 2n = m - n$ embedded in the four-ball, with boundary K . This shows that $g_4(K) \leq m - n \leq m + n - 2 < g(K)$, because $n \geq 1$ by assumption. Here, we use the standard notation $g_4(K)$ for the smooth 4-genus. In particular, we obtain $g_4^{\text{top}}(K) < g(K)$.




FIGURE 5. $abx a b x \rightsquigarrow \emptyset$ using one twist on four strands, followed by another twist on two strands, at the locations marked \star (cf. Figure 3). The first step is an isotopy, moving the blue strand. The last step is a crossing change near \star , followed by an isotopy fixing the endpoints of the braid strands.

– What remains are the 3-braid knots K such that neither K nor its mirror is among the following: strongly quasipositive, a connected sum of the form $T(2, 2m + 1) \# T(2, -2n - 1)$, $m \geq n \geq 1$, or the figure-eight knot. For such K , Lee–Lee [LL13] prove the following bound on the unknotting number: $u(K) < g(K)$. Since $g_4^{\text{top}}(J) \leq g_4(J) \leq u(J)$ holds for all knots J , this implies $g_4^{\text{top}}(K) < g(K)$ and completes the proof. \square

For comparison, we note the following analog of Theorem 1.1, in which the smooth 4-genus $g_4(K)$ replaces $g_4^{\text{top}}(K)$, and Rasmussen’s invariant $s(K)$ from Khovanov homology [Ras10] (or any other slice-torus invariant [Liv04], e.g. the Heegaard–Floer τ -invariant) plays the role of the signature $\sigma(K)$.

Proposition 3.1. *Let K be a 3-braid knot other than the figure-eight knot. Then*

$$|s(K)| = 2g(K) \iff g_4(K) = g(K).$$

These equalities hold precisely when K or its mirror is strongly quasipositive.

Proof. If K or its mirror is strongly quasipositive, then $|s(K)| = 2g(K)$ by [Shu07, Proposition 1.7]. Moreover, the implication $|s(K)| = 2g(K) \Rightarrow g_4(K) = g(K)$ holds for all knots K , because of the inequalities $|s(K)| \leq 2g_4(K) \leq 2g(K)$. It remains to prove that if K is a 3-braid knot other than the figure-eight knot with $g_4(K) = g(K)$, then K or its mirror is strongly quasipositive. This follows from Lee–Lee’s results [LL13]. More precisely, for a 3-braid knot K with $g_4(K) = g(K)$ Theorem 1.1 in [LL13] implies $u(K) = g(K)$. By Theorem 1.3 of the same paper, K or its mirror is either strongly quasipositive or a connected sum of two-strand torus knots $T(2, 2m + 1) \# T(2, -2n - 1)$ with $m \geq n \geq 1$. The latter can be excluded as




FIGURE 6. How to turn the braid $abxa^2bx$ (top left) into the tangle γ (top right) using one twist on four strands, followed by another twist on two strands, at the locations marked \star (cf. Figure 3).

in the proof of our Theorem 1.1 by showing $g_4(K) \leq m - n \leq m + n - 2 < g(K)$, a contradiction to $g_4(K) = g(K)$. \square

Theorem 3.2. *Let K be a strongly quasipositive 3-braid knot, written as the closure of a 3-braid β in Xu normal form $\beta = \delta^n \tau_1^{u_1} \tau_2^{u_2} \cdots \tau_t^{u_t}$, with $u_1, \dots, u_t \geq 1$ and $n \geq 0$. Then the topological 4-genus defect of K is bounded as follows:*

$$\frac{n}{3} + \frac{t}{3} - 1 \geq g(K) - g_4^{\text{top}}(K) \geq \frac{n}{3} + \frac{t}{6} - 3.$$

The constants $\frac{1}{3}$ and $\frac{1}{6}$ in the second inequality are optimal in the following sense: Whenever $C > \frac{1}{3}$ or $D > \frac{1}{6}$, and $E \in \mathbb{R}$, there exists a 3-braid in Xu normal form as above with $n \geq 0$ such that its braid closure K satisfies

$$g(K) - g_4^{\text{top}}(K) < Cn + Dt - E.$$

The case $t = 0$ in Theorem 3.2, in which $K = T(3, n)$, is covered by [BBL20, Theorem 1]. The equality

$$g(K) - g_4^{\text{top}}(K) = n - 1 - \left\lfloor \frac{2n}{3} \right\rfloor$$

directly implies the claimed upper and lower bounds in this case.

Proof. We only consider the case $t > 0$ and begin with the upper bound $\frac{n}{3} + \frac{t}{3} - 1 \geq g(K) - g_4^{\text{top}}(K)$. We first use Proposition 2.9 to compute the absolute value of the signature $|\sigma(K)| = -\sigma(K) = U + \frac{4}{3}n - \frac{2}{3}t$, where $U = u_1 + \cdots + u_t$, and recall that $g(K) = \frac{U}{2} + n - 1$ from (3), Section 2. The bound then follows directly from Kauffman and Taylor’s classical bound $|\sigma(K)| \leq 2g_4^{\text{top}}(K)$.

To establish the lower bound, we first apply a smooth cobordism from K to a knot K' , by suitably lowering the exponents n, u_1, u_2, \dots, u_t in β , as explained in Figure 4. First, assume $n > 0$. We set $K' := \text{cl}(\delta^m(abx)^{\frac{s}{3}})$, where

$$s = 6 \left\lfloor \frac{t}{6} \right\rfloor \quad \text{and} \quad m = \begin{cases} n - 2 & \text{if } n \equiv 0 \pmod{3} \\ n - 1 & \text{if } n \equiv 2 \pmod{3} \\ n & \text{if } n \equiv 1 \pmod{3}. \end{cases}$$

We have $K \rightsquigarrow K'$, and so this is a smooth cobordism which does not increase the topological 4-genus defect. In other words,

$$g(K) - g_4^{\text{top}}(K) \geq g(K') - g_4^{\text{top}}(K'),$$

as in (4). Next, we apply the untwisting move $(abx)^2 \rightarrow \emptyset$ from Figure 5 exactly $\frac{s}{6}$ times to the knot K' , resulting in $\text{cl}(\delta^m) = T(3, m)$. Since $m \equiv 1 \pmod{3}$, this is a knot again. For $m \geq 4$, [BBL20, Lemma 5 (1)] yields $\text{tu}(T(3, m)) \leq \frac{2}{3}m + \frac{1}{3}$. This inequality still holds for $m = 1$ and adds up to

$$g_4^{\text{top}}(K') \leq \text{tu}(K') \leq 2 \cdot \frac{s}{6} + \frac{2}{3}m + \frac{1}{3}.$$

Since $g(K') = \frac{s}{2} + m - 1$ by (3), and since $\frac{s}{6} \geq \frac{t}{6} - \frac{5}{6}$ and $m \geq n - 2$, we obtain

$$g(K) - g_4^{\text{top}}(K) \geq \frac{s}{6} + \frac{m}{3} - \frac{4}{3} \geq \frac{t}{6} + \frac{n}{3} - 3.$$

In the case $n = 0$, the above procedure fails because m , as defined above, is negative and the smooth cobordism to K' might therefore increase the 4-genus defect. However, a simple cosmetic modification allows for a cobordism that only increases

the 4-genus defect by at most one. Specifically, we set $m = 1$ (instead of $m = -2$) and $K' = \text{cl}(\delta(abx)^{\frac{5}{3}})$. The cobordism from K to K' is then given by lowering the exponents u_1, u_2, \dots, u_t in β as above while increasing the exponent of δ from 0 to 1. This gives

$$g(K) - g_4^{\text{top}}(K) \geq g(K') - g_4^{\text{top}}(K') - 1.$$

Now we apply the $\frac{5}{6}$ untwisting moves $(abx)^2 \rightsquigarrow \emptyset$ as above. The result is the unknot $\text{cl}(\delta)$. Since $n = 0$, we again obtain the claimed bound

$$g(K) - g_4^{\text{top}}(K) \geq \frac{s}{6} - 1 \geq \frac{t}{6} - \frac{5}{6} - 1 \geq \frac{t}{6} + \frac{n}{3} - 3.$$

In order to demonstrate optimality of the constants, we consider the two special families of 3-braids δ^n and $(abx)^n$, which we slightly modify to δ^{3k+1} and $(abx)^{2k}abx^2abx^2$, to make sure that their braid closures are connected.

– For $K = T(3, 3k + 1)$ with $k \geq 1$, the closure of the braid δ^{3k+1} in Xu normal form with $n = 3k + 1$ and $t = 0$, we have $g(K) = 3k$ and $g_4^{\text{top}}(K) = 2k + 1$ by [BBL20], hence $g(K) - g_4^{\text{top}}(K) = k - 1$. Whenever $C > \frac{1}{3}$ and D, E are arbitrary constants, we will therefore have

$$g(K) - g_4^{\text{top}}(K) = k - 1 < C \cdot (3k + 1) + D \cdot 0 - E$$

for sufficiently large k .

– If K is the closure of $(abx)^{2k}abx^2abx^2$, which is in Xu normal form with $n = 0$ and $t = 6k + 6$, then $g(K) = 3k + 3$. We will now invoke the Levine–Tristram signature function [Lev69, Tri69], which associates to each knot J a function $[0, 1] \rightarrow \mathbb{Z}$, $t \mapsto \sigma_{e^{2\pi it}}(J)$ with $\sigma_1(J) = 0$ for $t = 0$. By Gambaudo–Ghys [GG05, Corollary 4.4], for any 3-braid β with closure J , the Levine–Tristram signature function of J grows linearly on $(0, \frac{1}{3})$ with slope -2 times the writhe of β , up to a pointwise error of at most 2 (see e.g. Figure 7 at the end of the paper). For strongly quasipositive β with J a knot, that slope is $-4(g(J) + 1)$, see (3), and hence

$$(5) \quad \widehat{\sigma}(J) := \max_{\omega \in S^1 \setminus \Delta_J^{-1}(0)} |\sigma_\omega(J)| \geq \frac{4}{3}(g(J) + 1) - 2.$$

Since $\frac{1}{2}|\sigma_\omega(J)| \leq g_4^{\text{top}}(J)$ if $\omega \in S^1$ is not a root of the Alexander polynomial of J (see [KT76, Pow17]), we obtain $\frac{2}{3}(g(K) + 1) - 1 = 2k + \frac{5}{3} \leq g_4^{\text{top}}(K)$. Hence, if C, E are arbitrary constants and $D > \frac{1}{6}$, we have

$$g(K) - g_4^{\text{top}}(K) \leq 3k + 3 - 2k - \frac{5}{3} = k + \frac{4}{3} < C \cdot 0 + D \cdot (6k + 6) - E,$$

for sufficiently large k . In this case, it does not suffice to consider the (classical) signature bound on $g_4^{\text{top}}(K)$. Indeed, $|\sigma(K)| = 2k + 4$ (see Proposition 2.9) is roughly half the Levine–Tristram signature $\sigma_{e^{2\pi i/3}}(K)$. Substituting $\sigma(K)$ for $\sigma_{e^{2\pi i/3}}(K)$ in the above argument would therefore not work. \square

Remark 3.3. The proof of Theorem 3.2 shows that for K the closure of $(abx)^{2k}abx^2abx^2$, $k \geq 0$, we have

$$2k + \frac{5}{3} \leq \frac{1}{2}|\sigma_{e^{2\pi i/3}}(K)| \leq \frac{1}{2}\widehat{\sigma}(K) \leq g_4^{\text{top}}(K).$$

On the other hand, from K we obtain the closure of $abx^2abx^2 \sim a^2bxa^2bx$ using $2k$ twists (using k times our trick $abxabx \rightsquigarrow \emptyset$ from Figure 5), which can be untwisted

with two twists (compare the case $t = 6, n = 0$ in the proof of Theorem 1.1). Thus $g_4^{\text{top}}(K) \leq \text{tu}(K) \leq 2k + 2$ and we obtain

$$\frac{1}{2}|\sigma_{e^{2\pi i/3}}(K)| = \frac{1}{2}\widehat{\sigma}(K) = g_4^{\text{top}}(K) = \text{tu}(K) = 2k + 2.$$

4. THE 4-GENUS OF CLOSURES OF POSITIVE 3-BRAIDS

Our results so far provide infinitely many strongly quasipositive 3-braid knots K satisfying $\frac{1}{2}\widehat{\sigma}(K) = g_4^{\text{top}}(K)$, see Theorem 1.1 and Remark 3.3. This leads us to the following.

Question 4.1. Does the equality $\frac{1}{2}\widehat{\sigma}(K) = g_4^{\text{top}}(K)$ hold for all strongly quasipositive 3-braid knots? If not, does it at least hold for all braid positive 3-braid knots?

In this section, we exhibit large families of positive 3-braid knots for which $\frac{1}{2}\widehat{\sigma}(K) = g_4^{\text{top}}(K)$ holds up to an error of 1, see Proposition 4.4. In certain cases, we can even determine g_4^{top} exactly (see Examples 4.2 and 4.3 and Remark 4.5). For the knots covered by Proposition 4.4, $\widehat{\sigma}$ equals $|\sigma|$ or $|\sigma| + 2$, whereas in the example discussed in Remark 3.3, $\widehat{\sigma}$ equals $|\sigma_{e^{2\pi i/3}}|$. However, as we will see in Remark 4.7, the differences $\widehat{\sigma} - |\sigma|$ and $\widehat{\sigma} - |\sigma_{e^{2\pi i/3}}|$ can be arbitrarily large for positive 3-braid knots. This fact contributes to the difficulty of Question 4.1.

Throughout, we will use the Xu normal form of 3-braids from Section 2. Recall that we write $\delta = ba = ax = xb$ such that $a\delta = \delta b, b\delta = \delta x$ and $x\delta = \delta a$ (see (2) in Section 2).

Example 4.2. Let K be a knot that is the closure of a 3-braid in Xu normal form $\beta = \delta^{3\ell+2}a^{u_1}$ for $\ell \geq 0, u_1 \geq 1$. Note that u_1 must be even for K to be a knot. We claim that

$$(6) \quad g_4^{\text{top}}(K) = \text{tu}(K) = \frac{|\sigma(K)|}{2} = \frac{u_1}{2} + 2\ell + 1.$$

The last equality follows directly from Proposition 2.9. To prove Equation (6), using $\frac{1}{2}|\sigma(K)| \leq g_4^{\text{top}}(K) \leq \text{tu}(K)$ (as explained in the beginning of Section 3) it is enough to show that $\text{tu}(K) \leq \frac{1}{2}|\sigma(K)|$. By $\frac{u_1-2}{2}$ crossing changes from positive crossings of β to negative crossings we obtain the braid $\delta^{3\ell+2}a^2$. We will prove by induction that this braid can be untwisted with $2\ell + 2$ twists, which implies $\text{tu}(K) \leq \frac{u_1}{2} + 2\ell + 1$. For $\ell = 0$, we have $\delta^2a^2 = baba^3 = aba^4$ which becomes ab (with closure the unknot) by two crossing changes. For $\ell = 1$, we have

$$\begin{aligned} \delta^5a^2 &= \delta^4axa^2 = \delta^3x^2bxa^2 = \delta^2b^3abxa^2 = \delta a^4xabxa^2 = x^5bxabxa^2 \\ &\sim b^5abxabxx \rightsquigarrow b^5x, \end{aligned}$$

which turns into $bx \sim \delta$ using two crossing changes. Recall that the two twists needed to untwist $abxabx \rightsquigarrow \emptyset$ are shown in Figure 5 of Section 3. Now, for $\ell \geq 2$, we have

$$\begin{aligned} \delta^{3\ell+2}a^2 &= \delta^{3\ell-3}x^5bxabxa^2 \sim \delta^{3\ell-3}b^5abxabxx \rightsquigarrow \delta^{3\ell-3}b^5x \\ &\sim \delta^{3\ell-3}\delta b^4 = \delta^{3\ell-3}a\delta b^3 \sim \delta^{3\ell-1}b^2 \sim \delta^{3(\ell-1)+2}a^2, \end{aligned}$$

where we again used two twists for \rightsquigarrow . Inductively this shows that $\delta^{3\ell+2}a^2$ can be untwisted with $2\ell + 2$ twists as claimed.

Example 4.2 combined with the results from [BBL20] for 3-strand torus knots shows that the equalities $g_4^{\text{top}} = \text{tu} = \frac{\widehat{\sigma}}{2}$ hold for all strongly quasipositive 3-braid knots in Xu normal form (a) or (b) from Lemma 2.3, where $\widehat{\sigma} = |\sigma|$ except for certain torus knots of braid index 3; see also Remark 4.7. We next consider a sub-case of case (c) from Lemma 2.3.

Example 4.3. Let K be a knot that is the closure of a 3-braid in Xu normal form $\beta = \delta^{3\ell+1}a^{u_1}b^{u_2}$ for $\ell \geq 0, u_1, u_2 \geq 1$. Note that u_1 and u_2 must both be even for K to be a knot. We claim that

$$g_4^{\text{top}}(K) = \text{tu}(K) = \frac{|\sigma(K)|}{2} = \frac{u_1 + u_2}{2} + 2\ell.$$

The proof works as in Example 4.2. After $\frac{u_1+u_2-4}{2}$ positive to negative crossing changes in β we obtain the braid $\delta^{3\ell+1}a^2b^2$, which we can untwist with $2\ell+2$ twists as follows. For $\ell = 0$, the braid δa^2b^2 turns into δ by two crossing changes. For $\ell = 1$, we have

$$\begin{aligned} \delta^4 a^2 b^2 &= \delta^3 x a x a b^2 = \delta^2 b x^2 b x a b^2 = \delta a b^3 a b x a b^2 = x a^4 x a b x a b^2 \\ &\sim a b^4 a b x a b x x \rightsquigarrow a b^4 x, \end{aligned}$$

which can be untwisted using two crossing changes. For $\ell \geq 2$, we have

$$\begin{aligned} \delta^{3\ell+1} a^2 b^2 &= \delta^{3\ell-3} x a^4 x a b x a b^2 \sim \delta^{3\ell-3} a b^4 a b x a b x x \rightsquigarrow \delta^{3\ell-3} a b^4 x \\ &= \delta^{3\ell-4} x a^3 b a b x = \delta^{3\ell-5} b x^3 a^2 x a b x = \delta^{3\ell-6} a b^3 x^3 b x a b x \\ &\sim \delta^{3\ell-6} a^3 b^3 a b x a b x \rightsquigarrow \delta^{3\ell-6} a^3 b^3 \sim \delta^{3\ell-5} a^2 b^2 = \delta^{3(\ell-2)+1} a^2 b^2, \end{aligned}$$

which we can untwist inductively using the two base cases above.

Proposition 4.4 improves the statement from Theorem 3.2 for braid positive 3-braid knots under the additional assumption $u_i \geq 2$ for the exponents in the Xu normal form of their braid representatives. In fact, we can determine $g_4^{\text{top}}(K)$ in this case up to an error of 1, using $\frac{1}{2}|\sigma(K)|$ as a lower bound.

Proposition 4.4. *Let K be a knot that is the closure of a 3-braid in Xu normal form*

$$\delta^n \tau_1^{u_1} \tau_2^{u_2} \dots \tau_t^{u_t} \quad \text{for } t \geq 1, n \geq \frac{t}{2}, u_1, \dots, u_t \geq 2.$$

Then K is a braid positive knot and

$$(7) \quad \frac{n+t}{3} - 1 = g(K) - \frac{|\sigma(K)|}{2} \geq g(K) - g_4^{\text{top}}(K) \geq \frac{n+t}{3} - 2.$$

Proof. Set $U = u_1 + \dots + u_t$. Proposition 2.11 implies that K is braid positive. Moreover, we have $\frac{|\sigma(K)|}{2} = \frac{U}{2} + \frac{2n}{3} - \frac{t}{3}$ by Proposition 2.9 and $g(K) = \frac{U}{2} + n - 1$ by (3). Using $\frac{1}{2}|\sigma(K)| \leq g_4^{\text{top}}(K)$, it remains to show that $g_4^{\text{top}}(K) \leq \frac{|\sigma(K)|}{2} + 1$.

We distinguish two cases depending on the parity of t . First, let $t = 2r$ be even for $r \geq 1$. The conditions $2n \geq t$ and $n+t \equiv 0 \pmod{3}$ imply that we can write $n = 3\ell + r$ for $\ell \geq 0$, and K is the closure of

$$\beta = \delta^{3\ell+r} \tau_1^{u_1} \tau_2^{u_2} \dots \tau_{2r}^{u_{2r}}.$$

The case $r = 1$ ($t = 2$) is covered by Example 4.3, so we can further assume that $r \geq 2$. There is a smooth cobordism of Euler characteristic $4r - U - 4$ from K to the knot that is the closure of

$$\beta' = \tau_{1-r} \delta^{3\ell+r-1} \prod_{i=1}^{r-2} \tau_i^2 \tau_{r-1} \tau_r \tau_{r+1} \prod_{i=r+2}^{2r} \tau_i^2.$$

Indeed, we can use $U - 4r + 3$ saddle moves to replace all but three of the exponents u_i by 2 and the other three by 1. We use a last saddle move to replace δ by τ_{1-r} . We will prove by induction on r that β' turns into $\delta^{3\ell+1}$ by $2r - 2$ twists. Since the closure of $\delta^{3\ell+1}$ is the torus knot $T(3, 3\ell + 1)$, this will imply

$$(8) \quad \text{tu}(\text{cl}(\beta')) \leq \text{tu}(T(3, 3\ell + 1)) + 2r - 2 = \begin{cases} 2\ell + 2r - 1 & \text{if } \ell \geq 1, \\ 2r - 2 & \text{if } \ell = 0, \end{cases}$$

where the equality follows from [BBL20, Lemma 5 and Theorem 1]. Recall that $\delta = \tau_{i+1} \tau_i$, $\tau_i \delta = \delta \tau_{i+1}$ (see (2)) and $\tau_i = \tau_{i+3m}$ for all $m \in \mathbb{Z}, i \in \mathbb{Z}$. For $r = 2$, we thus have $\tau_{1-r} = \tau_2$ and

$$\begin{aligned} \beta' &= \tau_2 \delta^{3\ell+1} \tau_1 \tau_2 \tau_3 \tau_4^2 = \delta^{3\ell} \tau_2 \tau_1 \tau_0 \tau_1 \tau_2 \tau_3 \tau_4^2 = \delta^{3\ell} \tau_0 \tau_{-1} \tau_0 \tau_1 \tau_2 \tau_3 \tau_4^2 \\ &\sim \delta^{3\ell} \tau_2 \tau_1 \tau_2 \tau_3 \tau_4 \tau_5 \tau_6^2 \rightsquigarrow \delta^{3\ell} \tau_2 \tau_6 \sim \delta^{3\ell+1}, \end{aligned}$$

so β' indeed turns into $\delta^{3\ell+1}$ using $2r - 2 = 2$ twists in this case. Now, for $r \geq 3$, consider

$$\begin{aligned} \beta' &= \tau_{1-r} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_i^2 \delta \tau_{r-2}^2 \tau_{r-1} \tau_r \tau_{r+1} \tau_{r+2}^2 \prod_{i=r+3}^{2r} \tau_i^2 \\ &= \tau_{1-r} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_i^2 \tau_{r-3} \tau_{r-2} \tau_{r-3} \tau_{r-2} \tau_{r-1} \tau_r \tau_{r+1} \tau_{r+2}^2 \prod_{i=r+3}^{2r} \tau_i^2 \\ &\sim \tau_{(1-r)-(r-4)} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_{i-1-(r-4)}^2 \tau_1 \tau_2 \tau_1 \tau_2 \tau_3 \tau_4 \tau_5 \tau_6^2 \prod_{i=r+3}^{2r} \tau_{i-(r-4)}^2 \\ &\rightsquigarrow \tau_{(1-r)-r+1} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_{i-r}^2 \tau_1 \tau_2 \tau_3 \prod_{i=r+3}^{2r} \tau_{i-r+1}^2 \\ &\sim \tau_{(1-r)+1} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_i^2 \tau_{r-2} \tau_{r-1} \tau_r \prod_{i=r+1}^{2(r-1)} \tau_i^2. \end{aligned}$$

The braid β' hence turns into $\tau_{(1-r)+1} \delta^{3\ell+r-2} \prod_{i=1}^{r-3} \tau_i^2 \tau_{r-2} \tau_{r-1} \tau_r \prod_{i=r+1}^{2(r-1)} \tau_i^2$ by two twists and inductively we get that β' turns into

$$\tau_{(1-r)+r-2} \delta^{3\ell+1} \tau_1 \tau_2 \tau_3 \tau_4^2$$

by $2(r - 2)$ twists. Since $\tau_{(1-r)+r-2} = \tau_2$, this braid is the same as the one from the base case $r = 2$ and therefore can be untwisted with two twists. We obtain that β'

becomes $\delta^{3\ell+1}$ by $2r - 2$ twists. Equation (8) follows and we get

$$\begin{aligned} g_4^{\text{top}}(K) &\leq g_4^{\text{top}}(\text{cl}(\beta')) + \frac{U - 4r + 4}{2} \leq \text{tu}(\text{cl}(\beta')) + \frac{U}{2} - 2r + 2 \\ &\leq \begin{cases} \frac{U}{2} + 2\ell + 1 = \frac{|\sigma(K)|}{2} + 1 & \text{if } \ell \geq 1, \\ \frac{U}{2} = \frac{|\sigma(K)|}{2} & \text{if } \ell = 0. \end{cases} \end{aligned}$$

Next, let $t = 2r + 1$ be odd for $r \geq 0$. The conditions $2n \geq t$ and $n + t \equiv 0 \pmod{3}$ imply that we can write $n = 3\ell + r + 2$ for $\ell \geq 0$, and K is the closure of

$$\beta = \delta^{3\ell+r+2} \tau_1^{u_1} \tau_2^{u_2} \dots \tau_{2r+1}^{u_{2r+1}}.$$

The case $t = 1$ is covered by Example 4.2, so we can further assume that $r \geq 1$. There is a smooth cobordism of Euler characteristic $4r - U - 2$ from K to the knot that is the closure of

$$\beta' = \tau_{1-r} \delta^{3\ell+r+1} \prod_{i=1}^{r-1} \tau_i^2 \tau_r \tau_{r+1} \tau_{r+2} \prod_{i=r+3}^{2r+1} \tau_i^2,$$

similar to the cobordism considered in the above case. We prove by induction on r that β' turns into $\delta^{3(\ell+1)+1}$ by $2r - 2$ twists, hence

$$\text{tu}(\text{cl}(\beta')) \leq \text{tu}(T(3, 3(\ell + 1) + 1)) + 2r - 2 = 2\ell + 2r + 1.$$

For $r = 1$, we have

$$(9) \quad \beta' = \tau_0 \delta^{3\ell+2} \tau_1 \tau_2 \tau_3 = \delta^{3\ell+3} \tau_2 \tau_3 \sim \delta^{3(\ell+1)+1}.$$

For $r = 2$, we have

$$\begin{aligned} \beta' &= \tau_2 \delta^{3\ell+3} \tau_1^2 \tau_2 \tau_3 \tau_4 \tau_5^2 = \tau_2 \delta^{3\ell+2} \tau_0 \tau_1 \tau_0 \tau_1 \tau_2 \tau_3 \tau_4 \tau_5^2 \\ &\sim \tau_0 \delta^{3\ell+2} \tau_1 \tau_2 \tau_1 \tau_2 \tau_3 \tau_4 \tau_5 \tau_6^2 \rightsquigarrow \tau_0 \delta^{3\ell+2} \tau_1 \tau_2 \tau_6 \sim \delta^{3(\ell+1)+1} \end{aligned}$$

using Equation (9) in the last step. Now, for $r \geq 3$, consider

$$\begin{aligned} \beta' &= \tau_{1-r} \delta^{3\ell+r} \prod_{i=1}^{r-2} \tau_i^2 \delta \tau_{r-1}^2 \tau_r \tau_{r+1} \tau_{r+2} \tau_{r+3}^2 \prod_{i=r+4}^{2r+1} \tau_i^2 \\ &= \tau_{1-r} \delta^{3\ell+r} \prod_{i=1}^{r-2} \tau_i^2 \tau_{r-2} \tau_{r-1} \tau_{r-2} \tau_{r-1} \tau_r \tau_{r+1} \tau_{r+2} \tau_{r+3}^2 \prod_{i=r+4}^{2r+1} \tau_i^2 \\ &\sim \tau_{(1-r)-(r-3)} \delta^{3\ell+r} \prod_{i=1}^{r-2} \tau_{i-1-(r-3)}^2 \tau_1 \tau_2 \tau_1 \tau_2 \tau_3 \tau_4 \tau_5 \tau_6^2 \prod_{i=r+4}^{2r+1} \tau_{i-(r-3)}^2 \\ &\rightsquigarrow \tau_{(1-r)-r} \delta^{3\ell+r} \prod_{i=1}^{r-2} \tau_{i-r-1}^2 \tau_1 \tau_2 \tau_3 \prod_{i=r+4}^{2r+1} \tau_{i-r}^2 \\ &\sim \tau_{(1-r)+1} \delta^{3\ell+r} \prod_{i=1}^{r-2} \tau_i^2 \tau_{r-1} \tau_r \tau_{r+1} \prod_{i=r+2}^{2(r-1)+1} \tau_i^2. \end{aligned}$$

Inductively we get that β' turns into

$$\tau_{(1-r)+r-2} \delta^{3\ell+3} \tau_1^2 \tau_2 \tau_3 \tau_4 \tau_5^2 = \tau_2 \delta^{3\ell+3} \tau_1^2 \tau_2 \tau_3 \tau_4 \tau_5^2$$

by $2(r-2)$ twists, so into $\delta^{3(\ell+1)+1}$ by $2r-2$ twists. We obtain

$$\begin{aligned} g_4^{\text{top}}(K) &\leq g_4^{\text{top}}(\text{cl}(\beta')) + \frac{U-4r+2}{2} \leq \text{tu}(\text{cl}(\beta')) + \frac{U}{2} - 2r + 1 \\ &\leq \frac{U}{2} + 2\ell + 2 = \frac{|\sigma(K)|}{2} + 1. \end{aligned}$$

□

Remark 4.5. The proof of Proposition 4.4 (more precisely, the first case with $\ell = 0$) shows that the first inequality in (7) is an equality when $2n = t$.

Example 4.6. For knots K of low Seifert genus arising as closure of positive 3-braids, we can often apply the untwisting moves shown in Section 3 and show $\frac{1}{2}\widehat{\sigma}(K) = \text{tu}(K)$, thus determining $g_4^{\text{top}}(K)$. Here is a selection of positive 3-braids that close off to knots of Seifert genus 6 and 7, for which that strategy did not succeed:

$$\begin{aligned} \delta^3 a^2 b^2 x a b x &\sim a^3 b^3 a^2 b^2 a^2 b^2 \\ \delta^4 a^2 b x a b &\sim \Delta a^3 b^2 a^2 b^2 a^2 \\ \delta^4 a^4 b x a b &\sim \Delta a^5 b^2 a^2 b^2 a^2 \\ \delta^4 a^2 b^2 x a^2 b &\sim \Delta a^3 b^3 a^2 b^3 a^2 \\ \delta^6 a^2 b x &\sim \Delta^3 a^3 b^2 a^2. \end{aligned}$$

First, we note that for all of these knots K , we have the lower bound $\frac{1}{2}\widehat{\sigma}(K) = \frac{1}{2}|\sigma(K)| = g(K) - 2 \leq g_4^{\text{top}}(K)$. Second, in the search for upper bounds, we are able to find a knot J such that $K \rightsquigarrow J$ and $\text{tu}(J) = g(J) - 1$, thus proving $g_4^{\text{top}}(K) \leq g(K) - 1$, for each of these knots K . However, we are unable to find J with $K \rightsquigarrow J$ and $\text{tu}(J) = g(J) - 2$. Nevertheless, we can prove $g_4^{\text{top}}(K) = g(K) - 2$ for all of these knots K in a different way, which is practical for individual knots with low Seifert genus. Namely, a computer search [Lew23] reveals that the *algebraic genus* $g_{\text{alg}}(K)$, which is defined in terms of Seifert matrices of K and provides an upper bound $g_4^{\text{top}}(K) \leq g_{\text{alg}}(K)$ [FL18], satisfies $g_{\text{alg}}(K) \leq g(K) - 2$ for all our knots K . Yet it remains an open question whether $\text{tu}(K) = g(K) - 1$ or $\text{tu}(K) = g(K) - 2$ for those knots K .

Remark 4.7. The maximal Levine–Tristram signature, see (5) in Section 3, provides a good computable lower bound for g_4^{top} :

$$\widehat{\sigma}(K) = \max_{\omega \in S^1 \setminus \Delta_K^{-1}(0)} |\sigma_\omega(K)| \leq 2g_4^{\text{top}}(K).$$

The function $S^1 \rightarrow \mathbb{Z}$, $\omega \mapsto \sigma_\omega(K)$, is piecewise constant and jumps only at zeroes of the Alexander polynomial Δ_K . A priori, its maximum absolute value may be assumed anywhere on $S^1 \setminus \{1\}$.

In the above examples and Proposition 4.4 we have seen that for certain families of 3-braid knots, $\widehat{\sigma} = |\sigma| = 2g_4^{\text{top}}$, where $\sigma = \sigma_{-1} = \sigma_{e^{\pi i}}$ is the classical knot signature. This also holds for the $T(3, 3k+m)$ torus knots with $m \in \{1, 2\}$ and odd $k \geq 1$ [BBL20]. For even k on the other hand, e.g. for $T(3, 7)$, one finds $\widehat{\sigma} = |\sigma_\omega| = |\sigma| + 2$ for ω chosen only one jump-point of the Levine–Tristram signature away from $e^{\pi i}$, i.e. $\omega = e^{2\pi i t}$ for

$$t \in \left(\frac{1}{2} - \frac{5}{18k+6m}, \frac{1}{2} - \frac{1}{18k+6m} \right).$$




FIGURE 7. In blue, the graph of the Levine–Tristram signature $\sigma_{e^{2\pi i t}}(K)$ for $t \in [0, \frac{1}{2}]$ and K the closure of the 3-braid $(a^2 b^2)^8 (a^5 b^5)^4 \sim \delta^{12} (a^4 b^4 x^4)^2 a^4 b^4 (xab)^5 x$. In black, the linear approximation by [GG05, Corollary 4.4] for $t \in [0, \frac{1}{3}]$. The maximum absolute value $\hat{\sigma}(K)$ of $\sigma_{e^{2\pi i t}}(K)$, which equals $|\sigma(K)| + 4 = |\sigma_{e^{2\pi i/3}}(K)| + 4$, is assumed between 0.3599 and 0.3826 (rounded). This graph was drawn with sage [Sag23], by computing the signatures of the Hermitian matrices $(1 - \omega)A + (1 - \bar{\omega})A^\top$, for A a Seifert matrix of K , and ω between the roots of Δ_K on S^1 .

This observation relies on the fact that the jumps of the Levine–Tristram signatures of torus knots are well understood [Lit79, Ban22]. We have also seen examples where $\hat{\sigma} = |\sigma_{e^{2\pi i/3}}|$, namely the closures of $(abx)^{2k} abx^2 abx^2$ for $k \geq 0$; see Remark 3.3. Overall, whenever we could precisely determine the topological 4-genus of a 3-braid knot K , then the maximum absolute value of $\sigma_{e^{2\pi i t}}$ was either assumed at $t = \frac{1}{3}$, or at $t = \frac{1}{2}$, or close to $t = \frac{1}{3}$.

However, $\hat{\sigma}(K) - \max(|\sigma_{e^{2\pi i/3}}(K)|, |\sigma(K)|)$ is unbounded for K ranging over positive 3-braid knots. Indeed, let K_n be the closure of $(a^2 b^2)^{2n} (a^5 b^5)^n$, which is a knot if n is not a multiple of 3. The Levine–Tristram signatures of K_4 are shown in Figure 7. Using Proposition 2.8 and [GG05, Corollary 4.4] (as in (5)), we find

$$(10) \quad \sigma(K_n) = -12n, \quad \sigma_{e^{2\pi i/3}}(K_n) \approx -12n,$$

with an error of at most 2. Let us now estimate $\hat{\sigma}(K_n)$. We use the fact that K_n can be transformed by $n + 10$ saddle moves into the link

$$J_n = T(2, 10)^{\#(n-1)} \# T(3, 6n) \# T(2, -4n).$$

The latter satisfies, up to a constant error (i.e. an error that is independent of n),

$$\sigma_{e^{6\pi i/7}}(J_n) \approx -9n - 8n + \frac{6}{7} \cdot 4n = -(13 + \frac{4}{7})n,$$

which implies that (again with constant error)

$$\hat{\sigma}(K_n) \geq |\sigma_{e^{6\pi i/7}}(K_n)| \geq |\sigma_{e^{6\pi i/7}}(J_n)| - n - 10 \approx (12 + \frac{4}{7})n.$$

This estimate, combined with (10), shows that

$$\lim_{n \rightarrow \infty} \widehat{\sigma}(K_n) - \max(|\sigma_{e^{2\pi i/3}}(K_n)|, |\sigma(K_n)|) = +\infty.$$

This last example shows that determining $\widehat{\sigma}$ and g_4^{top} for all 3-braid knots, or even just closures of positive 3-braids, could be a hard problem.

ACKNOWLEDGMENT

The fourth author thanks Peter Feller for his constant encouragement.

REFERENCES

- [Ban22] I. M. Banfield, *The maximum Levine–Tristram signature of torus knots*, 2022, Preprint available at [arXiv:2212.09604](https://arxiv.org/abs/2212.09604).
- [BBL20] S. Baader, I. Banfield, and L. Lewark, *Untwisting 3-strand torus knots*, Bull. Lond. Math. Soc. **52** (2020), no. 3, 429–436, DOI 10.1112/blms.12335. MR4171377
- [Ben83] D. Bennequin, *Entrelacements et équations de Pfaff* (French), Third Schnepfenried geometry conference, Vol. 1 (Schnepfenried, 1982), Astérisque, vol. 107, Soc. Math. France, Paris, 1983, pp. 87–161. MR753131
- [BKL98] J. Birman, K. H. Ko, and S. J. Lee, *A new approach to the word and conjugacy problems in the braid groups*, Adv. Math. **139** (1998), no. 2, 322–353, DOI 10.1006/aima.1998.1761. MR1654165
- [BM93] J. S. Birman and W. W. Menasco, *Studying links via closed braids. III. Classifying links which are closed 3-braids*, Pacific J. Math. **161** (1993), no. 1, 25–113. MR1237139
- [BM08] J. S. Birman and G. Tian, *Preface [In memory of Xiao-Song Lin]*, Commun. Contemp. Math. **10** (2008), no. suppl. 1, v–vii, DOI 10.1142/S0219199708003150. MR2468363
- [DDG⁺15] P. Dehornoy, F. Digne, E. Godelle, D. Krammer, and J. Michel, *Foundations of Garside theory*, EMS Tracts in Mathematics, vol. 22, European Mathematical Society (EMS), Zürich, 2015. Author name on title page: Daan Kramer, DOI 10.4171/139. MR3362691
- [Erl99] D. Erle, *Calculation of the signature of a 3-braid link*, Kobe J. Math. **16** (1999), no. 2, 161–175. MR1745024
- [FL18] P. Feller and L. Lewark, *On classical upper bounds for slice genera*, Selecta Math. (N.S.) **24** (2018), no. 5, 4885–4916, DOI 10.1007/s00029-018-0435-x. MR3874707
- [Fre82] M. H. Freedman, *The topology of four-dimensional manifolds*, J. Differential Geometry **17** (1982), no. 3, 357–453. MR679066
- [FQ90] M. H. Freedman and F. Quinn, *Topology of 4-manifolds*, Princeton Mathematical Series, vol. 39, Princeton University Press, Princeton, NJ, 1990. MR1201584
- [Gar69] F. A. Garside, *The braid group and other groups*, Quart. J. Math. Oxford Ser. (2) **20** (1969), 235–254, DOI 10.1093/qmath/20.1.235. MR248801
- [GG05] J.-M. Gambaudo and É. Ghys, *Braids and signatures* (English, with English and French summaries), Bull. Soc. Math. France **133** (2005), no. 4, 541–579, DOI 10.24033/bsmf.2496. MR2233695
- [Ine16] K. Ince, *The untwisting number of a knot*, Pacific J. Math. **283** (2016), no. 1, 139–156, DOI 10.2140/pjm.2016.283.139. MR3513844
- [KT76] L. H. Kauffman and L. R. Taylor, *Signature of links*, Trans. Amer. Math. Soc. **216** (1976), 351–365, DOI 10.2307/1997704. MR388373
- [Lev69] J. Levine, *Knot cobordism groups in codimension two*, Comment. Math. Helv. **44** (1969), 229–244, DOI 10.1007/BF02564525. MR246314
- [Lew23] L. Lewark, *galg-computations*, (2023), computer program available at <https://github.com/LLewark/galg-computations>.
- [Lie16] L. Liechti, *Positive braid knots of maximal topological 4-genus*, Math. Proc. Cambridge Philos. Soc. **161** (2016), no. 3, 559–568, DOI 10.1017/S0305004116000670. MR3569161
- [Lit79] R. A. Litherland, *Signatures of iterated torus knots*, Topology of low-dimensional manifolds (Proc. Second Sussex Conf., Chelwood Gate, 1977), Lecture Notes in Math., vol. 722, Springer, Berlin, 1979, pp. 71–84. MR547456

- [Liv04] C. Livingston, *Computations of the Ozsváth-Szabó knot concordance invariant*, *Geom. Topol.* **8** (2004), 735–742, DOI 10.2140/gt.2004.8.735. MR2057779
- [LL13] E.-K. Lee and S.-J. Lee, *Unknotting number and genus of 3-braid knots*, *J. Knot Theory Ramifications* **22** (2013), no. 9, 1350047, 18, DOI 10.1142/S0218216513500478. MR3105306
- [LM] C. Livingston and A. H. Moore, *KnotInfo: Table of Knot Invariants*, <https://knotinfo.math.indiana.edu>, retrieved April 17, 2023.
- [McC19] D. McCoy, *Null-homologous twisting and the algebraic genus*, 2019–20 *MATRIX annals*, *MATRIX Book Ser.*, vol. 4, Springer, Cham, [2021] ©2021, pp. 147–165. MR4294766
- [Mur65] K. Murasugi, *On a certain numerical invariant of link types*, *Trans. Amer. Math. Soc.* **117** (1965), 387–422, DOI 10.2307/1994215. MR171275
- [Mur74] K. Murasugi, *On closed 3-braids*, *Memoirs of the American Mathematical Society*, vol. No. 151, American Mathematical Society, Providence, RI, 1974. MR0356023
- [Pow17] M. Powell, *The four-genus of a link, Levine-Tristram signatures and satellites*, *J. Knot Theory Ramifications* **26** (2017), no. 2, 1740008, 28, DOI 10.1142/S0218216517400089. MR3604490
- [Ras10] J. Rasmussen, *Khovanov homology and the slice genus*, *Invent. Math.* **182** (2010), no. 2, 419–447, DOI 10.1007/s00222-010-0275-6. MR2729272
- [Rud83] L. Rudolph, *Braided surfaces and Seifert ribbons for closed braids*, *Comment. Math. Helv.* **58** (1983), no. 1, 1–37, DOI 10.1007/BF02564622. MR699004
- [Rud90] L. Rudolph, *A congruence between link polynomials*, *Math. Proc. Cambridge Philos. Soc.* **107** (1990), no. 2, 319–327, DOI 10.1017/S0305004100068584. MR1027784
- [Sag23] The Sage Developers, *Sagemath, the Sage Mathematics Software System (Version 10.1)*, (2023), <http://www.sagemath.org>.
- [Shu07] A. N. Shumakovitch, *Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots*, *J. Knot Theory Ramifications* **16** (2007), no. 10, 1403–1412, DOI 10.1142/S0218216507005889. MR2384833
- [Sto17] A. Stoimenow, *Properties of closed 3-braids and braid representations of links*, *SpringerBriefs in Mathematics*, Springer, Cham, 2017. MR3729301
- [Tri69] A. G. Tristram, *Some cobordism invariants for links*, *Proc. Cambridge Philos. Soc.* **66** (1969), 251–264, DOI 10.1017/s0305004100044947. MR248854
- [Tru23] P. Truöl, *The epsilon invariant at 1 of 3-braid knots*, *Algebr. Geom. Topol.* **23** (2023), no. 8, 3763–3804, DOI 10.2140/agt.2023.23.3763. MR4665283
- [Xu92] P. Xu, *The genus of closed 3-braids*, *J. Knot Theory Ramifications* **1** (1992), no. 3, 303–326, DOI 10.1142/S0218216592000185. MR1180404

MATHEMATISCHES INSTITUT, SIDLERSTRASSE 5, 3012 BERN, SWITZERLAND
Email address: sebastian.baader@unibe.ch

UNIVERSITÄT REGENSBURG, 93053 REGENSBURG, GERMANY
Email address: lukas@lewark.de
URL: <http://www.lewark.de/lukas>

UNIVERSITÄT REGENSBURG, 93053 REGENSBURG, GERMANY
Email address: filip.misev@mathematik.uni-regensburg.de

ETH ZÜRICH, RÄMISTRASSE 101, 8092 ZÜRICH, SWITZERLAND
Email address: paulagtruoel@gmail.com
URL: <https://people.math.ethz.ch/~ptruoel/>