

REVIEW ARTICLE

Differential diagnosis of contact dermatitis: A practical-approach review by the EADV Task Force on contact dermatitis

David Pesqué¹  | Olivier Aerts² | Mojca Bizjak³  | Margarida Gonçalo^{4,5}  |
Aleksandra Dugonik⁶ | Dagmar Simon⁷ | Suzana Ljubojević-Hadzavdić⁸  |
Laura Malinauskienė⁹  | Mark Wilkinson¹⁰ | Magdalena Czarnecka-Operacz¹¹ |
Beata Krecisz¹² | Swen M. John¹³ | Anna Balato¹⁴ | Fabio Ayala¹⁵ |
Thomas Rustemeyer¹⁶ | Ana M. Giménez-Arnau¹⁷ 

Correspondence

Ana M Giménez Arnau, Department of Dermatology, Hospital del Mar, Barcelona, Spain.
Email: anamariagimenezarnau@gmail.com

Abstract

The diagnosis of eczema ('dermatitis') is mostly clinical and depends on the clinical history and exploratory objective findings (primary lesions, patterns). Contact dermatitis remains as an important condition in the group of eczematous disorders, with important socioeconomic and occupational relevance. Although irritant and allergic contact dermatitis have a different pathogenesis, both are characterized by a rather typical morphology, are triggered by external factors and tend to occur primarily in the area of contact with the exogenous agent. In addition, allergic and irritant dermatitis may also co-exist. The importance of diagnosing contact dermatitis, especially when allergic in nature, is both due to the possibility of avoiding the trigger, and due to its role in aggravating other skin conditions. Nevertheless, the heterogeneity of clinical presentations in daily practice may pose an important challenge for the suspicion and correct diagnosis of contact dermatitis. Furthermore, other conditions, with different pathogenesis and treatment, may clinically simulate contact dermatitis. The Task Force aims to conduct a review of the unifying clinical features of contact dermatitis and characterize its main clinical phenotypes, and its simulators, in order to contribute to an early suspicion or recognition of contact dermatitis and enable a correct differential diagnosis.

BACKGROUND

Contact dermatitis (CD) is a type of skin inflammatory reaction that occurs secondary to contact with a specific substance. CD is mostly eczematous in nature. The definition of eczema relies on morphologic traits.¹ 'Eczema' may also be referred to as 'eczematous dermatitis' – or simply 'dermatitis', with equivalent denotation in clinical practice.² Eczematous

dermatitis presents with a remarkable diversity of primary lesions, such as erythema, macules, papules, vesicles and bullae, as well as secondary lesions like scaling, erosions, crusts, lichenification, hyperkeratosis and hypo- and/or hyperpigmentation. Erythema, vesicles, oedema and exudative changes are more typical in acute phases, whereas more papular, fissured, scaly and even hyperkeratotic states are seen in subacute and chronic phases (Figure 1a).³ Furthermore,

David Pesqué, Olivier Aerts and Mojca Bizjak have contributed equally as first authors.

Thomas Rustemeyer and Ana M. Giménez-Arnau are co-senior authors.

For affiliations refer to page 14.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Authors. *Journal of the European Academy of Dermatology and Venereology* published by John Wiley & Sons Ltd on behalf of European Academy of Dermatology and Venereology.



FIGURE 1 The multi-faceted presentations of contact dermatitis. (a) Pulpitis due to allergic contact dermatitis from diallyl disulfide present in garlic. (b) Airborne contact dermatitis due to potassium dichromate in a welder with retroauricular, submental, nasolabial and mild frontal involvement. (c) Photoallergic contact dermatitis from ketoprofen presenting as sharply demarcated eczematous lesions on the leg. (d) Violaceous erythema and areas of hyperpigmentation after healing along with areas of vesicles and erosions on the back of a patient as a manifestation of phototoxic contact dermatitis induced by common rue (*Ruta graveolens*).

overlap between acute and chronic lesions may also occur, for example during flares in chronic dermatitis.³ It is essential to highlight that history taking and physical examination, supplemented with additional diagnostic tests, and, if applicable, along with studying the effect of avoidance and re-exposition to the suspected culprit(s), are essential for establishing an accurate diagnosis of contact dermatitis (CD) and providing optimal care for the patient.⁴

The aim of this practical review is to provide an up-to-date guidance for dermatologists on how to adequately suspect and diagnose CD.

Classification of eczematous disorders

Eczematous disorders include CD and photo-induced CD, atopic dermatitis, dyshidrotic eczema, asteatotic eczema, seborrheic dermatitis, nummular eczema, stasis dermatitis, eczematous cutaneous adverse drug reactions and protein contact dermatitis (PCD).² Identifying the precise disorder(s) may be challenging because of overlapping clinical patterns and the difficulty in histopathological differentiation of eczematous conditions. Thus, diagnosis is mostly based on clinical findings and, if possible, complementary tests.

Clinical forms of contact dermatitis

Contact dermatitis may present with many different and subtle manifestations.⁵ Irritant contact dermatitis (ICD), allergic contact dermatitis (ACD), photo-induced CD (including

photoallergic CD and phototoxic CD) and PCD emerge as different categories within the spectrum of CD. The differentiation of these categories is facilitated by the presence of differences in terms of triggers and pathogenic mechanisms. Clinical manifestations of CD may vary by factors such as the triggering culprit, route and duration of exposure, skin type and duration of inflammation.³ Clinical examination *per se* is generally regarded as having limited reliability for the differentiation of these CD subtypes, although some clinical clues may occasionally provide guidance (see below).³

Irritant contact dermatitis is believed to be the most common subgroup of CD, accounting for even 80% of all cases.⁶ However, taking into account the occurrence of potential pitfalls in the diagnosis of ACD, this percentage could include misdiagnoses of other subgroups of CD. It may be occupational, especially related to wet-work conditions, and several factors may contribute to the susceptibility of ICD (age, sex, atopy, body region). Its clinical features are very polymorphous.⁷ The heterogeneity of ICD is related to the complex genetic and environmental mechanisms, but it is always characterized by the direct or indirect skin injury induced by external stimuli, activating the innate immune system.⁶ From a clinical perspective, lesions are considered to be well-demarcated and confined to the area of contact with the irritant, even if in selected cases some spreading may occur (e.g., from the hands to the forearms).⁸ In ICD of the hands, the interdigital spaces and dorsal side of the hands and fingers are often involved, yet this is not a specific sign.³ Lesions of ICD heal rather quickly if the triggering agent(s) are removed (*'decrescendo'* phenomenon).^{9,10} Chronic ICD may present as dry, erythematous, scaly lesions

with lichenification,⁹ a form often entailing a poor prognosis.¹¹ The predominating subjective symptom of ICD is not pruritus, but burning sensations, pain and skin soreness in acute ICD,¹¹ while pruritus and eczema may also be present in more chronic ICD.

Allergic contact dermatitis is also very common in daily practice, and an increase in its prevalence has been evidenced in the last years.¹² Skin involvement in ACD is the result of a specific, immune-mediated delayed reaction against haptens that come in contact with the skin. In its acute form, it has a tendency to develop erythema, papules, vesicles and, occasionally, bullae, whereas more evolved forms usually present with the features of chronic dermatitis.⁹ Pruritus is the cardinal symptom of ACD. The most frequently involved body areas are the face, hands and forearms, with legs and trunk often being other frequent sites of involvement.² ACD may typically present with extension to areas outside of the area that is in direct contact with the allergen ('spreading reaction; id-reaction'),³ possibly due to allergen persistence, extensive exposure to the allergen, areas with resident-memory T cells that had previous contact with the allergen, or even due to haematogenous spread. In contrast with ICD, a 'crescendo' phenomenon may occur in ACD, even after allergen removal.⁹ However, these diagnostic clues do not fully reflect the clinical diversity and complexity of ACD. In fact, several publications have already previously attempted to elucidate the clinical patterns/forms of presentation of ACD.^{1,5,12}

One consensus classification of ACD includes 13 main categories⁵: direct exposure ACD, exacerbating pre-existing ACD, multifactorial dermatitis including ACD, ACD by proxy, ACD mimicking angioedema, airborne ACD, photo-induced ACD, systemic ACD, PCD, allergic contact stomatitis, erythroderma/exfoliative dermatitis, non-eczematous ACD (e.g., lymphomatoid ACD) and ACD with additional respiratory/mucosal symptoms. Additional minor forms of ACD presentation include, for example, hair loss, increased hair growth, tattoo reactions, granulomatous reactions, among many others. The aim of this article is not to revisit each of these categories, but to outline the clinical features that may be of utmost importance for clinicians.

ACD by proxy or connubial/consort dermatitis is a form of ACD with a particular mechanism of allergen exposure. In this case, the allergen source is another individual and the exposure is by close contact.¹³ An illustrative example would be cheek ACD after transfer of a product from the partner's lips. This ectopic mechanism should be considered when no clear rationale for personal contact with a demonstrated allergen is found. The causative agents are commonly topical medications.¹³

Airborne ACD is a diagnostic challenge for many dermatologists. The fact that allergens contact with the skin is under the form of small, air-distributed particles, vapours and gases, may easily lead to confusion with photo-induced CD. Airborne ACD typically involves the face, neck and/or the neckline without any spared areas (Figure 1b), in contrast to phototoxic and/or photoallergic CD.⁹ Clinical clues to look for are involvement behind the ears, under the chin,

under the nose and on the upper eyelids, which are the so-called 'shadow zones', typically unaffected in photo-induced CD.⁹ Nevertheless, both airborne and photo-induced CD may co-exist, as has been reported for photo-aggravated, airborne ACD from isothiazolinone-containing paints and detergents.¹⁴ Moreover, as the eczematous lesions in airborne ACD are often quite ill-defined, the diagnosis may be challenging.⁹

Photo-induced CD includes two separate conditions. **Photoallergic CD** results from skin inflammation caused by a photosensitive allergen combined with sunlight or artificial light.¹⁵ It is characterized by relatively ill-defined, eczematous lesions in light-exposed areas (Figure 1c), such as face, ears, V-area of the neckline and below the cuffs of the two upper limbs, whereas 'shadow zones' (see above) remain spared.⁹ **Phototoxic reactions** involve rather non-specific immune mechanisms and present mostly as an exaggerated sharp-delimited sunburn that tends to precede subsequent hyperpigmentation (Figure 1d). Overlap features of both conditions may present in a patient.¹⁶

Systemic CD includes a spectrum of clinical presentations that occur after systemic exposure to a contact allergen. Several different allergens may lead to such pictures of ACD, such as drugs, plant products and metals.¹⁷ Clinical pictures are characterized by widespread/generalized eczema, often involving body folds, or by acute vesicular palmoplantar dermatitis (resembling dyshidrotic eczema or 'pompholyx'). Systemic ACD may have some additional, relatively specific skin manifestations, such as the recurrence of dermatitis at the site of previous contact after systemic exposure to the allergen, also observed in experimental oral challenge trials.¹⁸ Systemic ACD to drugs is a type of delayed IVa (T-helper 1 inflammation) or IVc (cytotoxic T lymphocytes) hypersensitivity reaction, caused by topical or systemic drugs.^{17,19} It presents with widespread eczematous eruptions, worsening/reactivation of eczema, with symmetrical drug-related intertriginous and flexural exanthema (SDRIFE), or with erythroderma. SDRIFE is characterized by a sharply defined symmetrical erythema (red to purple-red) of the buttocks and inner thighs, in the latter with V-shaped distribution, with involvement of at least one other flexural location and without systemic symptoms and signs (Figure 2a,b).^{20,21}

Protein contact dermatitis belongs to the group of immediate hypersensitivity skin reactions and follows repeated episodes of contact urticaria, although the latter is not always evident for the patient.²² Pruritus, erythema, wheals or angioedema occurring in a matter of minutes after skin contact with a culprit agent are characteristic in the acute phase, while the subacute phase is followed by vesicles and other eczematous features.²³ After the acute phase, chronic hand dermatitis (erythema, lichenification, fissures and sometimes residual scales) may be seen.²³ PCD shows a predilection for the hands (especially the fingertips), sometimes extends to the wrists and arms and is typically, yet not exclusively, occupational in origin. Rarely, the face and other locations may be involved.²⁴




FIGURE 2 Symmetrical Drug-related Intertriginous and Flexural Erythema (SDRIFE) secondary to azithromycin. (a) Symmetrical erythema of the axilla, here only showing the left axilla. (b) Symmetrical V-shaped erythema in the inguinal area.

Non-eczematous and other minor clinical forms of presentation of ACD are addressed in the section of 'Uncommon clinical findings of CD'. Even though these different clinical forms of CD share many common findings, their varied clinical presentations often lead to diagnostic delay and/or unnecessary complementary tests.

DIFFERENTIAL DIAGNOSES OF CONTACT DERMATITIS

To correctly diagnose CD, it is essential to understand its clinical presentations and to ensure an adequate diagnostic approach, which includes the differential diagnosis of a wide range of conditions. In accordance to guidelines, this process requires a comprehensive assessment of past medical history, questioning on the rash development, occupational and private exposures and, if applicable, the evaluation of the interrelationship.²⁵ Further anamnesis may be required depending on the clinical suspicion. This section presents the main differential diagnoses of CD and highlights clinical features to take into account for a complete anamnesis and exploration during the differential diagnosis process for eczematous and non-eczematous disorders.

Inflammatory conditions

Several entities may resemble the clinical presentation of CD due to the eczematous lesions and/or the pattern(s) of distribution. In addition, clinical situations such as the lack of response to conventional therapies or unexplained flares could raise the possibility of additional (superimposed) CD.

Other eczematous disorders

The diagnosis of **atopic dermatitis** is based on a series of clinical criteria.²⁶ The diagnosis is made in the setting of

eczematous, relapsing lesions with a typical morphology that correspond to the atopic dermatitis age-specific pattern (cheeks, forehead and extensor involvement in infantile forms, and later flexural, head and neck and hand lesions in adolescents and adults), together with the hallmark symptom of pruritus. Other important features are an early age of onset and personal/family history of atopy. In addition, increased serum IgE, the presence of minor atopy-associated cutaneous features (e.g., keratosis pilaris, hyperlinear palms and generalized xerosis) and susceptibility to cutaneous infections (e.g., *Staphylococcus aureus* and Herpes simplex) may also guide our clinical suspicion. With respect to atopic dermatitis, ACD should be considered as either an alternative diagnosis and/or an exacerbator.²⁶ ACD in patients with atopic dermatitis should be taken into consideration in cases when history and/or physical examination are suggestive of ACD, such as flares, or an inappropriate response to topical treatment (e.g., topical corticosteroids) or emollients, or when specific patterns are observed that reflect exposure to a causative agent. In this regard, ACD could be suggested by marked facial or eyelid involvement, increased severity in neck folds and vesicles on dorsal hands and fingertips.²⁷ Moreover, systemic ACD may occasionally result in a phenotype mimicking atopic dermatitis.²⁸ Patch testing is recommended in these situations, or in the setting of known atopic dermatitis in the absence of improvement to standard treatments.²⁷

Dyshidrotic eczema is defined as chronic, intermittent, highly pruritic episodes of vesicles with faint erythema that involve hands (palms, lateral aspects of fingers) and feet (soles). Less frequently, bullae may be seen.²⁹ Scaling occurs when vesicles dry, but an erythematous background of lesions occurs less frequently. It usually runs a seasonal course and is occasionally related to different triggers such as hyperhidrosis, contact irritants and even metal hypersensitivity.³⁰ The possible relationship with metal contact allergy and the common clinical patterns of dyshidrosis and acute ACD may hamper the correct diagnosis.³⁰ Moreover, acute palmoplantar vesicular eruptions may also be a clinical presentation

of systemic ACD that could mimic dyshidrotic eczema (see above).

Asteatotic eczema, or **craquelé dermatitis**, is the result of extremely dry skin after excessive epidermal water loss. Dry, cracked skin that can evolve into fissures and erythematous patches with the typical, almost diagnostic, polygonal or curvilinear pattern in fissured skin is seen.³¹

The diagnosis of **seborrheic dermatitis** remains a clinical one, presenting with the characteristic symmetrical erythema and yellowish-to-white 'greasy' scaling that may occur in nasolabial and retroauricular folds, eyebrows, scalp, and in interscapular and pre-sternal areas. Concomitance of ACD and seborrheic dermatitis may occur.³² Seborrheic dermatitis refractory to therapy, or worsening upon topical products, should be regarded as clues for secondary ACD. ACD may also mimic seborrheic dermatitis (e.g., from cosmetics and nail lacquers).³³

Nummular eczema is defined by the sharply defined, coin-shaped, eczematous plaques, usually symmetrically distributed, with an anatomical preference for the lower and upper limbs.³⁴ Contact allergy may be associated with nummular eczema in up to 25% of patients, particularly from nickel and preservatives.^{35,36} ACD with a nummular pattern has also been described in relation to persulfates, and contact allergens present in antiseptics.^{37,38}

Stasis dermatitis is the consequence of chronic venous insufficiency. Erythematous and eczematous patches and plaques restricted to the lower legs, without clear-cut borders, and classically involving the medial malleoli, are the most common findings.³⁹ Pruritus, scaling and lichenification may occur. Brown hyperpigmentation is a consequence of the dermal deposition of hemosiderin. Patients with stasis dermatitis, especially those with leg ulcers, may readily develop ACD to the personal care products, topical medications and wound dressings.^{40,41}

Delayed-type hypersensitivity drug reactions include a maculopapular eruptions, fixed drug eruption, acute generalized (localized) exanthematous pustulosis (AGEP/ALEP), drug reaction with eosinophilia and systemic symptoms (DRESS), systemic CD, erythema multiforme and Stevens–Johnson/toxic epidermis necrolysis (SJS/TEN) syndrome.^{42,43} It is important, when evaluating potential drug eruptions, to take into account that CD may occasionally simulate such drug eruptions (e.g., erythema multiforme-like CD^{44–47} or ACD mimicking AGEP^{48,49}), and that, occasionally, these drug reactions may also mimic CD.^{50,51} Notably, eczematous cutaneous adverse drug reactions, or eczematous drug eruptions, are general terms used to define rashes with eczematous features that occur following the administration of systemic medications.⁵² These reactions may develop in the context of systemic CD (see above).⁵² The diagnosis may be difficult as the exact clinical presentation may vary for each drug, but most develop within weeks to months after the onset of the treatment.⁵²

Table 1 outlines the main eczematous disorders included in the differential diagnosis of CD with anatomical and clinical clues.

Non-eczematous disorders

The differentiation of **psoriasis** (and its variants) from CD may pose several challenges in clinical practice, with previous literature citing the intricately clinical and pathological differences and similarities.^{53,54} It is important to stress the difficulty in distinguishing chronic plantar/palmar dermatitis from plantar/ palmar psoriasis. Furthermore, the term 'eczema in psoriatico' is frequently used to define the co-occurrence of psoriasis and eczematous disorders, such as CD.⁵⁵ Although it has been suggested that patients with psoriasis could be less prone to ACD due to differences in the inflammatory milieu,⁵⁶ clinicians need to be aware that ACD can certainly complicate psoriasis, with ACD sometimes presenting psoriatic features, namely through the Köebner phenomenon.⁵⁷

Lichen planus and its variants share common clinical and sometimes histopathological lichenoid features. Dermoscopy, with the rather typical Wickham striae, may contribute to the distinction.^{58,59} However, CD may also mimic lichen planus ('lichenoid CD').⁶⁰ In regard to cutaneous lesions, the eruption initially involves contact sites, followed by a characteristic spreading reaction, yet with less characteristic polygonal lilac papules, and often without mucosal lesions.⁶¹ Moreover, the onset of lichenoid CD is rather acute with sometimes frank eczematous lesions being noticeable in the contact site.⁶¹ Nevertheless, the clinical, histopathologic and dermatoscopic resemblance may be high, and distinction can remain difficult.⁶² In addition, lichenoid contact allergy has been particularly described in the oral mucosa, mostly due to dental amalgam.⁶³ Interestingly, positive patch test reactions may initially be eczematous and later on become lichenoid.⁶¹

Lichen simplex chronicus (LSC) is a form of localized itch with secondary thick lichenified plaques in the areas of scratching.⁶⁴ It is symmetrically found in easily accessible body regions, such as the ankles, shins, dorsal hands, upper back, neck and anogenital regions. Occasionally, it has been suggested that ACD from hair dye may precede in some cases the development of LSC, and it can be anticipated that long-lasting ACD may, in some cases, easily transform into LSC.⁶⁵

Regarding **acne vulgaris**, **perioral dermatitis** and **rosacea-like dermatitis**, one should remember that almost all the subgroups of CD may have perioral and face involvement, and that some cases may effectively mimic flares of the aforementioned conditions.^{66,67} Recalcitrant rosacea, including its oedematous complication, referred to as 'Morbihan disease', or perioral dermatitis should raise the possibility of superimposed CD and should lead to the evaluation of face cosmetic and topical products. If cheilitis and/or perioral PCD are present, the use of musical instruments, toothpastes and even food allergens should also be evaluated. Moreover, some reports have highlighted that ACD from fragrances and formaldehyde releasers may resemble or aggravate rosacea, whereas others have stressed that acne vulgaris/rosacea with superimposed ACD can mainly present as a flare-up of the original dermatosis.⁶⁸

TABLE 1 Diagnostic clues of eczematous conditions included in the differential diagnosis of contact dermatitis.

Condition	Predilection sites	Clinical (anamnesis and exploratory) clues
Allergic CD	It may affect any area (face and neck, scalp, hands/feet, trunk, arms/legs, nails, body folds, genital and/or be generalized)	Pruritic dermatitis, with partial or no response to corticosteroids, with recurrent flares. Ill-defined lesions, except for the volar palmar and wrists borders, ²⁵ with spread beyond areas of direct exposure. ⁹ 'In crescendo' phenomenon ⁹
Irritant CD	Mostly hands, but occasionally in arms, folds, genitals	Well-demarcated, rash in the area of contact with irritants. ⁶ Worsening in cold season ⁶ and 'decrecendo' phenomenon. ⁹ Burning or stinging sensations may be present
Photo-induced CD	Photo-exposed areas, mostly face and neck, scalp or generalized in exposed trunk and extremities	Identification of light as a trigger and evaluation of the use of topical or systemic drugs with phototoxic and/or photosensitizing properties. Photoallergic CD is ill-defined in contrast to phototoxic CD and may spread to covered body parts ²⁵
Systemic CD	Folds, hands/feet, generalized	Investigation of the use of topical and/or systemic drugs as well as contact with other products such as metals, plants and drugs. ¹⁷ When affecting the palmoplantar regions, a vesicular presentation may predominate the clinical picture
Protein CD	Mostly in hands	Atopic skin diathesis. Local immediate symptoms upon contact (mostly itch and erythema or contact urticaria) in initial stages and vesicles later ²³
Atopic dermatitis	Face and neck and/or hands/feet and/or arms/legs (flexures) and/or be generalized	Personal and familial history on atopy with early age of onset (though it can develop at any age). Specific age distribution patterns associating dry skin and intense itch
Dyshidrotic eczema	Hands and feet	Existence of hyperhidrosis, seasonal flares or contact with irritants
Asteatotic eczema	Arms and legs or generalized	Investigation of deficient use of moisturizers. Polygonal or curvilinear pattern in fissured skin
Seborrheic dermatitis	Face and neck, scalp, genital or be generalized (trunk)	Investigation of triggers (e.g., stress, alcohol and neurological diseases). Seborrheic areas with less pruritus. Stinging/burning may be associated
Nummular eczema	Arms and legs or be generalized	Coin-shaped, eczematous, symmetrical plaques
Stasis dermatitis	Legs	Well-demarcated macular patches and plaques in a background of chronic venous insufficiency of the legs and local symptoms of venous insufficiency
Delayed-type hypersensitivity drug eruption	Face and neck (DRESS, AGEP, SJS/TEN, FDE), hands and feet (erythema multiforme), genital (FDE), generalized (most types, including maculopapular eruptions)	Detailed history and timing of systemic medication Clinically very heterogeneous

Abbreviations: AGEP, acute generalized exanthematous pustulosis; CD, contact dermatitis; DRESS, drug reaction with eosinophilia and systemic symptoms; FDE, fixed drug eruption; SJS/TEN, Stevens–Johnson syndrome/Toxic epidermal necrolysis.

Polymorphic light eruption is characterized by a days' duration rash of usually smooth-topped erythematous papules that tend to form plaques, even if the rash may be heterogeneous.⁶⁹ The diagnosis is clinical, and many clues differentiate it from CD, such as the sun-related trigger, and especially the shorter duration and typically relapsing nature of the rash. Nevertheless, photo-induced CD—alone or superimposed (e.g., from sunscreens), remains an important consideration, especially when the eruption is acute, without any previous history, and/or when relapses do not have the same characteristic morphology.

Within the spectrum of auto-immune diseases, **lupus erythematosus** and **dermatomyositis** have to be considered in the differential diagnosis of CD. In regard to lupus, acute malar rash and some forms of subacute and discoid lupus erythematosus are part of the differential diagnosis of CD. CD has very occasionally complicated cutaneous lupus erythematosus.⁷⁰ Moreover, as in psoriasis and rosacea, a secondary ACD may also trigger a relapse of an underlying cutaneous lupus erythematosus, and, similar to lichen planus, patch tests may then also transform from an eczematous patch into a lesion with histopathological

characteristics of lupus ('contact lupus').⁷¹ Lymphomatoid ACD may also completely mimic cutaneous lupus, or related disorders (e.g., Jessner lymphocytic infiltration of the skin/Jessner-Kanof's disease, which, similar to CD, may histopathologically show the presence of eosinophils). Since cutaneous lesions of dermatomyositis affect the scalp, face, eyelids and hands (mechanic's hands), and associate pruritus, the differential diagnosis between CD and dermatomyositis based on clinical signs and symptoms is often challenging. Furthermore, ACD can also be superimposed and even delay the diagnosis of this particular auto-immune disease.⁷² In addition, both lupus erythematosus and dermatomyositis, when cutaneous lesions are photo-aggravated, need to be considered as differential diagnoses and even mimickers of photo-induced CD.⁷³

Infectious diseases

Skin infections may also resemble CD. However, an accurate examination and complementary tests will normally lead to the elucidation of the diagnosis. In contrast to inflammatory conditions, overlap or association with CD is uncommon.

Tinea infections, (Figure 3a,b) including **Tinea incognita**, have been described as mimickers of ACD, particularly in children.⁷⁴ To rule out the possibility of tinea, KOH test and/or culture of scrapings need to be performed.

Scabies can present with very unspecific signs (e.g., eczematous dermatitis, urticarial eruptions, intense pruritus and excoriations), yet may also have some specific dermatoscopic hallmarks (e.g., white lines and delta wing sign), and suggestive clinical signs (e.g., linear skin tunnels in the palmar creases and wrists and scabietic nodules in the genital and umbilical region). Importantly, superimposed CD to some topical treatments of scabies has been described.⁷⁵ Therefore, lack of improvement despite correct treatment will require to rule out not only the persistence/re-infection but also irritancy/contact allergy to the ingredients of topical treatments.

Non-bullous and bullous **impetigo** may resemble, or complicate CD,⁷⁶ causing impetiginized dermatitis. Bacterial cultures can be used for diagnostic confirmation.

In addition, severe cases of ACD, with severely exudating skin lesions, may mimic impetigo.⁶⁶

Erythrasma is a bacterial infection characterized by dark erythematous-brownish patches and fine scaling that primarily affects interdigital spaces and intertriginous areas. It may be confused with (A)CD (e.g., from fragrances and textiles). Wood's lamp examination may show a characteristic 'coral red' fluorescence.

Other entities such as **viral exanthems**, including **pityriasis rosea**, **herpetic lesions** and **candidiasis** are regularly included in the differential diagnosis of CD, especially in early stages. Cutaneous candidiasis affects mostly body folds (inguinal, inframammary, anogenital) with one characteristic finding, the presence of peripheral papules and pustules, accompanying erythema and maceration. However, differentiation from 'pustular' ACD (e.g., from fragrances) may need to be considered.⁷⁷ Herpes simplex is common in clinical practice and could be a misdiagnosis with intense, vesicular ACD, and vice versa, although the associated pain should direct the clinician to the right diagnosis.⁷⁸

Other diseases

This section includes a myriad of conditions that may occasionally be part of the differential diagnosis of CD.

Genetic conditions, such as **Hailey–Hailey disease** (autosomal dominant due to mutations in ATP2C1 mutations) and **Darier's disease** (autosomal dominant disease due to ATP2A2 mutations), may clinically resemble CD and even co-exist.⁷⁹

Grover's disease, although not considered a genodermatosis, is another dermatosis that might mimic CD. All three conditions are histopathologically characterized by acantholysis, which is useful to differentiate those from spongiotic conditions. Other genetic syndromes such as **peeling skin syndrome type B**, **Severe dermatitis—multiple Allergies—Metabolic wasting (SAM) syndrome**, **Netherton syndrome** and **Omenn syndrome** present with recalcitrant atopic dermatitis-like manifestations, and with additional cutaneous and non-cutaneous features suggesting their diagnosis.⁸⁰ The underlying skin barrier alteration could predispose these patients to secondary CD, even if evidence is lacking to date.




FIGURE 3 Tinea faciei in a child and Tinea incognita. (a) Erythematous annular and macular patch with a scaly, spreading border that also contains papules in a child with Tinea faciei. (b) Photo-distributed rash with eczematous appearance in a woman who had already been treated with topical corticosteroids. A detailed anamnesis and a closer look, altogether with microbiological studies, led to the diagnosis of Tinea incognita.

Premalignant and malignant disorders may also mimic CD. Importantly, any 'atypical' eczema which fails to respond to topical corticosteroids may represent a premalignant or malignant disorder, namely **cutaneous T-cell lymphoma, parapsoriasis en plaque, mammary and extramammary Paget's disease, and in situ squamous cell carcinoma**, including **Bowen's disease** and **erythroplasia of Queyrat**.¹ Therefore, in such cases, a skin biopsy is mandatory to establish the exact nature of the lesion.⁸¹ Mycosis fungoides, with its multiple clinical variants, and pathogenesis linked to 'persistent allergen stimulation', has been associated with an increased prevalence of delayed hypersensitivity.⁸² In addition, the clinical/histopathological differentiation between mycosis fungoides and lymphomatoid ACD may pose a significant challenge,⁸³ since both may be indistinguishable and lymphomatoid ACD is mainly T cell-based, and only rarely B-cell based.⁸⁴

Other conditions that have seldom been included in the differential diagnosis of CD are **bullous pemphigoid, porphyria cutanea tarda** and psychodermatologic diseases such as **delusions and dermatitis artefacta**.

Table 2 outlines the main non-eczematous disorders included in the differential diagnosis of CD with anatomical and clinical clues.

Differential diagnosis for special locations (face and neck, hands and feet and intertriginous areas) with diagnostic clues

Some anatomic areas entail difficulty during the differential diagnosis process due to specific clinical features of conditions in these areas. Table 3 highlights some clinical clues to consider in clinical practice. Figures 4 and 5 show different conditions on the face and hands/feet, respectively.

UNCOMMON CLINICAL FINDINGS OF CONTACT DERMATITIS

It is difficult to summarize rather uncommon forms of CD, since our knowledge mainly stems from case reports or small case series. Uncommon clinical forms do not correspond to primary clear-cut lesions of eczema, yet a diagnosis is achieved through complementary tests (patch testing, in particular, but also others, such as skin biopsy) and correlation with exposure. A myriad of skin manifestations has been reported including: **(de)pigmented dermatitis**,^{98–102} **purpuric dermatitis**,^{103–105} **skin and oral lichenoid dermatitis**,^{61,62} **erythema multiforme-like dermatitis**,^{44–47} **lymphomatoid dermatitis**,^{106–109} **granulomatous dermatitis and cheilitis**,^{110–112} **pustular dermatitis**,^{113–117} **neutrophilic and eosinophilic dermatitis**,¹¹⁸ **bullous dermatitis**^{119–121} and **sclerodermoid dermatitis**.¹²² In addition, in rare cases, ACD may not present as dermatitis, but rather with nail or hair alterations (hair loss

or increased hair growth).^{123,124} Furthermore, delayed contact allergy may also present with extracutaneous manifestations with or without associated skin findings, as in the case of allergy to prosthetic or external materials¹²⁵ (e.g. pseudotumors¹²⁶ and implant failure^{127,128}). A special clinical manifestation of CD includes 'angioedema-like' contact dermatitis, which is sometimes referred to as oedematous CD, or 'bullous' CD. Although this is often misinterpreted as classic angioedema, the observed oedema (often facial) is itchy, lasts longer than 3 days and often resolves leaving a residually scaly dermatitis or dry skin, all three being features not consistent with a classic angioedema. Different allergens have been associated with this particular clinical presentation, including paraphenylenediamine (PPD) and methylisothiazolinone.^{129,130}

CONTACT DERMATITIS: THE GREAT MIMICKER AND ITS SIMULATORS

This section includes a schematic presentation of (i) CD presenting with a morphology and/or distribution suggesting another skin condition, and (ii) CD simulators, that is, conditions with CD-like features. The first situation is the result of the variable clinical manifestations of CD and stresses the diagnostic difficulty in clinical practice. In contrast, CD simulators are conditions that may exhibit eczematous features and/or present a pattern or distribution that suggests CD.¹ For further examples, see Table 4.

COMPLEMENTARY TESTS

The diagnostic approach will also entail complementary tests, which will mostly be, for CD, patch and photopatch testing and, if needed, histopathological exam.

Patch and photopatch testing

Patch and photopatch testing are essential procedures in daily routine to investigate allergic and photoallergic CD. Patch testing is considered the gold standard to diagnose contact allergy resulting from type IV hypersensitivity.¹⁷² It should be performed in all patients with clinical manifestations of contact allergy, including its unusual forms.¹⁷² In addition, some clinical situations in which patch testing is proposed are previously well-controlled chronic dermatoses that lose response to treatment, unexplained flares, dermatitis that manifests in an atypical distribution and new-onset dermatitis that responds to therapy but returns rapidly after treatment discontinuation.¹⁷³ Counselling patients regarding the allergen(s) involved, after interpretation of the clinical relevance of positive results, is essential to allow allergen avoidance. Furthermore, follow-up consultation can help to ensure the understanding and

TABLE 2 Diagnostic clues of non-eczematous conditions included in the differential diagnosis of contact dermatitis.

Condition	Predilection sites	Clinical (anamnesis and exploratory) clues
Lichen planus	Hands and wrists, feet and ankles, scalp, nails, generalized, mucosal (oral and genital)	Investigate past or associated disease (e.g., hepatitis C), systemic drugs (e.g., ACEIs, NSAIDs and thiazides) and use of dental amalgam. In nails: nail plate thinning, trachyonychia, dorsal nail pterygium. In mucosa: erosions, Wickham striae
LSC	Easily accessible locations	Investigate scratching habit and associated lesions
Psoriasis	Any area, including scalp and nails, may be generalized	Investigate family history, arthritis, obesity, triggers (infections, stress, tobacco, alcohol, medications). At exploration: pinpoint bleeding (Auspitz sign), presence of Koebner phenomenon, characteristic dermatoscopic features (regular dotted vessels in a red background with silvery scales) ⁸⁵
Lupus erythematosus	Face (acute or discoid), scalp (discoid), trunk (subacute), arms/legs (profundus) or generalized	Evaluate photosensitivity, associated systemic symptoms (mostly in acute and subacute forms) and baseline diseases. If associated systemic lupus erythematosus, consider systemic clinical hallmarks. ⁸⁶ Wide spectrum of skin lesions depending on lupus subtype (acute, subacute or chronic) ⁸⁷
Dermatomyositis	Face, scalp, arms and legs, nails	Photosensitivity and presence of systemic symptoms or blood test hallmarks, even though amyopathic forms exist. ⁸⁸ Skin findings: Gottron papules, Gottron sign, Holster sign, mechanic hands, heliotrope rash, hallux patch
PLE	Photo-exposed areas	Photosensitivity; usually spares the face and presents with recurrent episodes
Acne—Rosacea	Face	Use of occlusive topical products in the face and triggers (e.g., UV-exposure, alcohol, hot beverages and spicy foods)
Tinea	Face, scalp, hands/feet, arms/legs, generalized	Assess contact with affected individuals and/or animals; check the feet Dermoscopy may be helpful in the differential diagnosis (e.g., tinea capitis). ⁸⁹
Scabies	Hands and feet or generalized	Assess contact with affected individuals. Intense nocturnal itch. It spares the face (except children, immunocompromised). Typical hallmarks (e.g., delta wing sign)
Impetigo	Mostly face	Investigation of contact with affected individuals. Non-bullous: vesicular phase leading to honey-coloured crusts. Bullous: flaccid bullae with subsequent erosions
Erythrasma	Body folds	Evaluate obesity and hyperhidrosis. Clinical diagnosis favoured by Wood's lamp
Candidiasis	Body folds and genitals	Evaluate obesity, diabetes and/or hyperhidrosis. Erythema, maceration, peripheral 'milky-white' pustules and papules
Viral exanthems	Mostly generalized	Heterogenous clinical presentation with systemic prodromes
Herpes simplex	Localized lesions	Painful recurrent vesicular lesions in the same region
Hailey-Hailey disease	Body folds	Presence of hyperhidrosis or friction in the area
PCT	Mostly in hands and forearms	Evaluate alcohol use, oestrogen use, viral infections, iron overload and smoking Skin blisters, vesicles, bullae and increased fragility of the skin on sun-exposed body areas with secondary hyper- and hypo pigmented scars and milia. ⁹⁰
Bullous pemphigoid	Hands, forearms or generalized	Evaluate medication and neurological baseline diseases. Initially urticarial and papular lesions with pruritus, which are followed by vesicles and bullae on normal or erythematous skin. Usually spares the head region.
CTCL	Localized or generalized	Heterogenous presentation ⁹¹
Mammary and extramammary Paget's disease	Body folds, nipple and areola, and/or genital	Evaluate baseline diseases and presence of local symptoms that could guide towards an underlying malignancy.
Bowen disease	Mostly extremities	Evaluate sun-exposure habits. Red, scaly patches/plaques with variable associated features (erosion, fissures, pigmented, nodular) ⁹²
Queyrat erythroplasia	Genital	Evaluate the presence of local symptoms Well-demarcated, velvety, shiny, bright red, plaque-like appearance ⁹²

Abbreviations: ACEI, angiotensin-converting enzyme inhibitor; CTCL, cutaneous T-cell lymphoma; LSC, lichen simplex chronicus; NSAID, nonsteroidal anti-inflammatory drug; PCT, porphyria cutanea tarda; PLE, polymorphic light eruption.

proper avoidance of identified allergens and causes of CD, which can result in reinforcing self-management skills of the patients. Patch testing improves the quality of life in patients with ACD.¹⁷⁴ Photopatch testing is indicated

in case a photoallergic CD is suspected, since additional ultraviolet exposure is needed to induce the reaction. Its value as diagnostic tool has been recognized in the study of any dermatitis in photo-exposed areas, or for evaluating

TABLE 3 Clinical clues in the differential diagnosis of body regions with specific features (face and neck, hand and feet and body folds).

Condition	Location	Specific clues
Allergic CD	Face and neck	Rinse-off pattern (preauricular and mandibular areas) ¹² Geographic pattern (area of application of a topical product) ¹² Bilateral patchy pattern on the face (due to cosmetics) ¹² Airborne pattern (upper eyelids, nasolabial fold, retroauricular and submental areas. The bridge and tip of the nose may be unaffected) ¹² Drip pattern (lower eyelids and cheeks, due to topical eyedrops) ¹² Contour pattern (bilateral periocular, mostly due to cosmetics) ¹²
	Hands and/or feet	Pincer grasp pattern (due to allergic CD from acrylates, diallyl disulfide and food or irritant CD due to friction). See Figure 1a . Periungual pattern (allergic CD from nail cosmetics) ⁹³ Shoe pattern (dorsum of feet sparing interdigital webs due to (photo)allergic CD to topical medications) ¹² Plantar pattern (the arches and interdigital of skin may be spared) ¹²
Irritant CD	Hands and/or feet	Apron pattern (interdigital spaces and extending to palmar and dorsal surfaces) Ring pattern (under the ring area). ¹² Rule out metal allergy if unfavourable course Glove pattern (dorsum of the hand, wrists and fingers eczema without interdigital involvement). ¹² It can be seen in allergic CD
	Body folds	Examine contact with body fluids and use of diapers/daily pads
Atopic dermatitis	Face and neck	Cheek dermatitis in children Retroauricular dermatitis with fissuring ⁹⁴ (also seen in irritant CD)
	Hands and/or feet	Dorsal hand with involvement of fingers and wrists. Less palmar involvement. ⁹⁵
Seborrheic dermatitis	Body folds	Ill-defined salmon-coloured thin patches
Photoallergic CD	Face and neck	On the bridge of the nose, forehead and cheeks. Upper eyelids, retroauricular and submental areas are spared. ¹²
Psoriasis	Hands and/or feet	Palmar grasp pattern . Concomitant involvement of the dorsum of the hands and wrists may point to allergic CD. ¹²
	Body folds	Inverse psoriasis : symmetrical well-defined shiny red patches Fissures/erosions in folds may be indicative of psoriasis, ⁹⁶ especially in the intergluteal crease.
Lichen planus	Hands and/or feet	Typical clinical features, particularly on wrists and ankles, but also with a hyperkeratotic papular pattern, erythematous scaly pattern or even verrucous lichen planus. ^{1,97}
Lupus erythematosus	Face and neck	Butterfly shaped malar rash (acute), annular/psoriasiform lesions on the neck (subacute) and/or discoid atrophic lesions in head and neck and scalp (chronic). ⁸⁷
Dermatomyositis	Face and neck	Scalp seborrheic-like rash, heliotrope rash, facial erythema, shawl sign
Dyshidrotic eczema	Hands and/or feet	Vesicles on lateral and dorsal sides on fingers. Palmar grasp pattern may be associated. ¹²
Erythrasma	Body folds	Brown or dark red, asymptomatic, persistent patches
Hailey-Hailey disease	Body folds	Erythematous plaque containing painful vesicles that rupture and form erosions, typically in the border of the lesions
Mammary and extramammary Paget's disease	Body folds, nipple and areola	Well-demarcated plaques with erosive, ulcerated, scaly or eczematous pattern

Abbreviation: CD, contact dermatitis.

photosensitivity due to systemic drugs.¹⁷⁵ Although elaborate information about these techniques is beyond the scope of this article, detailed recommendations are found in previous reference works.^{172,175}

Histopathological findings of contact dermatitis and its differential diagnoses

This section details the main histopathological hallmarks of CD and the most representative conditions included in its differential diagnosis.

The histopathology term 'spongiotic dermatoses' encompasses eczematous disorders, highlighting the predominant epidermal changes in the form of spongiosis, which is the intercellular oedema between keratinocytes.^{176,177} Spongiosis may associate with dermal and epidermal T-cell infiltration.^{176,177} Spongiosis, however, does not occur only in eczematous disorders, but can be present in many other inflammatory and infectious disorders (e.g., viral exanthems, psoriasis and lichenoid reactions).¹⁷⁷

The differentiation of eczematous disorders by means of other histopathological characteristics is very difficult. In addition, depending on the features of the reaction, the



FIGURE 4 Dermatoses involving the face. (a) Contour pattern in allergic contact dermatitis from cosmetics (b) Dermatomyositis with subtle cheek erythema and heliotrope periorcular oedema and erythema (c) Facial eczema herpeticum (d) Severe facial eczema in an atopic patient, which should raise suspicion about associated allergic contact dermatitis.



FIGURE 5 Dermatoses involving hands or feet. (a) Erosive phase of a bullous fixed drug eruption. The differential diagnosis of this lesion could include shoe allergic contact dermatitis; however the evaluation with anamnesis of drug intake and previous episodes led to the right diagnosis. (b) Palmar psoriasis with hyperkeratosis, desquamation and induration. (c) Acquired palmoplantar keratoderma which was diagnosed as mycosis fungoides after histopathological exam. (d) Lichen planus involvement of the proximal palmar aspect of the hand with the rather typical lilac papules on the wrist.

histopathologic hallmarks are variable. Acute dermatitis is characterized by severe spongiosis with intraepidermal vesicles,¹⁷⁸ and even eosinophilic spongiosis, particularly

in ACD.¹⁷⁷ In subacute dermatitis, the spongiosis becomes less intense, even very mild in chronic dermatitis. Furthermore, chronic stages will show more acanthosis,

TABLE 4 Contact dermatitis: the great mimicker and its simulators.

	CD as a mimicker	CD simulators
Inflammatory	<ul style="list-style-type: none"> • Other eczematous disorders (atopic dermatitis, seborrheic dermatitis)^{28,131,132} • Drug reactions (e.g., acute generalized exanthematous pustulosis and anti-IL-17 reactions)^{48,49,133,134} • Psoriasis^{135–137} • Lichen planus¹³⁸ • Lichen nitidus¹³⁹ • Prurigo nodularis¹⁴⁰ • Folliculitis decalvans¹⁴¹ • Pemphigus vulgaris¹⁴² • Aphthous stomatitis¹⁴³ • Angioedema and contact urticaria^{144–147} • Lupus erythematosus^{*73} • Dermatomyositis^{*73} • Polymorphic light eruption^{**140} • Actinic prurigo^{**148} 	<ul style="list-style-type: none"> • Other eczematous disorders (atopic dermatitis, dyshidrotic eczema, asteatotic eczema, seborrheic dermatitis, nummular eczema)^{1,2,7} • Drug reactions (e.g., erythema multiforme and fixed drug eruption)^{50,51} • Psoriasis¹ • Lichen planus¹⁶² • Hailey-Hailey disease¹ • Bullous pemphigoid^{163,164} • Rosacea¹⁶⁵ • Sweet syndrome¹⁶⁶ • Prurigo pigmentosa¹⁶⁷ • Lupus erythematosus^{*73} • Dermatomyositis^{*73,168} • Polymorphic light eruption^{**73} • Solar urticaria^{**73} • Actinic prurigo^{**73} • Hydroa vacciniforme^{**73} • Cutaneous porphyrias^{**73,169} • Dermatitis herpetiformis
Neoplastic	<ul style="list-style-type: none"> • Mammary and extramammary Paget's disease¹⁴⁹ • Cutaneous T-cell lymphoma (mycosis fungoides)^{150–153} • Basal cell carcinoma¹⁵⁴ 	<ul style="list-style-type: none"> • Cutaneous T-cell lymphoma (mycosis fungoides) • Parapsoriasis en <i>plaque</i> • Bowen's disease • Mammary and extramammary Paget's disease • Erythroplasia of Queyrat
Infectious	<ul style="list-style-type: none"> • Herpes simplex^{155,156} • Herpes zoster¹⁵⁷ • Impetigo¹⁵⁸ 	<ul style="list-style-type: none"> • Herpes zoster • Impetigo • Scabies¹⁷⁰ • Tinea⁷⁴ • Erythrasma • Cellulitis¹⁷¹
Others	<ul style="list-style-type: none"> • Child abuse¹⁵⁹ • Burn¹⁶⁰ • Ischaemia¹⁶¹ 	<ul style="list-style-type: none"> • Dermatitis artefacta

Note: (*): contact dermatitis and/or photocontact dermatitis; (**): photocontact dermatitis.

Abbreviation: CD, contact dermatitis.

parakeratosis and hyperkeratosis.¹⁷⁹ Spongiotic dermatoses may encompass combined histopathological changes of acute, subacute and chronic dermatitis. Other non-specific findings are perivascular lymphocytes and eosinophils and neutrophils in the stratum corneum (e.g., if secondarily impetiginized).¹⁷⁹ Some particular features may be more characteristic of some types of eczema, which are outlined in Table 5.

Examples of frequently used additional diagnostic tools

The repeated-open-application-test (ROAT) aims to clarify the relevance of a selected doubtful or positive patch test to a product—or its ingredients. This technique can occasionally be more sensitive than patch testing, especially when it concerns ACD from low-concentrated ingredients in the tested product (patch test negative, ROAT positive). The test involves applying the product, or ingredient, which is

suspected as a cause of ACD, two times a day on the volar side of the forearm for at least 10 days. Cosmetics and topical drugs are typically tested with this method.^{172,188}

The semi-open test has been suggested as a method to study products that may contain irritants (e.g., shampoos, detergents and paints). Normally, it is used with patient-supplied products. Before testing an aqueous product, its pH needs to be verified. Following full evaporation of the product (and observation of any immediate reactions), the completely dried site is covered with tape and readings are performed as with regular patch tests.¹⁸⁹

Open tests have been suggested as the first step for testing poorly defined products, such as patients' own materials (if they effectively come in contact with the skin), and for products suspected to cause immediate contact reactions, before proceeding to prick testing.

Prick tests, including prick-by-prick tests (e.g., food), are useful diagnostic tools for the diagnosis of contact urticaria and PCD, as well as identifying specific IgE against suspected allergens.

TABLE 5 Histopathological clues of contact dermatitis and its main differentials.

Conditions	Histopathology
Eczematous disorders	
Allergic CD	<ul style="list-style-type: none"> Spongiosis is more prominent in the lower epidermis^{176,177} Spongiotic vesicles present at different epidermal levels (distinctive from nummular dermatitis)¹⁷⁷ Intraepidermal Langerhans cells persist longer and are more distributed¹⁷⁶
Irritant CD	<ul style="list-style-type: none"> Changes vary depending on the causative agent and concentration of the product.¹⁸⁰ Ballooning of keratinocytes in the upper and mid epidermis¹⁷⁷ Variable epidermal necrosis and associated neutrophils¹⁷⁷ Psoriasiform hyperplasia in chronic phases¹⁷⁷
Atopic dermatitis	<ul style="list-style-type: none"> Dermal eosinophils frequently seen¹⁷⁷ Intraepidermal collections of Langerhans cells are rare¹⁸¹
Seborrheic dermatitis	<ul style="list-style-type: none"> In acute stages, overlying scale crust containing neutrophils¹⁷⁷ In subacute stages, less spongiosis and more psoriasiform hyperplasia and acanthosis
Photo-induced CD	<ul style="list-style-type: none"> Photoallergic CD is similar to allergic CD Phototoxic dermatitis: abundant apoptotic keratinocytes, analogous to a sunburn¹⁸²
Dyshidrotic eczema	<ul style="list-style-type: none"> Intraepidermal vesicle formation Vesicles displace acrosyringia at the periphery of the bleb¹⁷⁷
Stasis dermatitis	<ul style="list-style-type: none"> Variable epidermal features: spongiosis, acanthosis, parakeratosis Hallmark dermal changes: extravasated erythrocytes, hemosiderin-laden macrophages and proliferation of dilated small blood vessels in the papillary dermis¹⁸³
Psoriasis ¹⁸⁴	<ul style="list-style-type: none"> Regular acanthosis, with elongated rete ridges Thinning of the suprapapillary layer of the epidermis Areas of parakeratosis with collections of neutrophils (Munro microabscesses) Collections of neutrophils in the spinosum plate (spongiform pustules of Kogoj) Alternating hypo/hypergranulosis
Lichen planus ¹⁸⁴	<ul style="list-style-type: none"> Compact hyperkeratosis without parakeratosis Hypergranulosis and irregular acanthosis Basal layer liquefaction Lichenoid lymphocytes in the papillary dermis Direct immunofluorescence: immunoglobulins, complement and fibrin stain some keratinocytes in the deeper epidermis and outermost dermis
Drug reactions	<ul style="list-style-type: none"> Very variable features Dermal eosinophils and eosinophilic exocytosis may be noted¹⁸⁴ Spongiotic features. In morbilliform eruptions, spongiosis and exocytosis may be limited to basal portions of the epidermis.¹⁸⁵ Papillary dermal oedema Apoptotic keratinocytes
Lupus erythematosus ¹⁸⁴	<ul style="list-style-type: none"> Variable epidermal findings (from atrophy to hyperplasia), depending on the cutaneous form of lupus. Basal layer liquefaction (interface dermatitis); thickening of the basal membrane Perivascular and peri-adnexal lymphocytes, sometimes acquiring a lichenoid pattern Increased mucin deposition Usually no neutrophils or eosinophils (except in neutrophilic or bullous lupus; eosinophils may be present in lupus tumidus)
Dermatomyositis	<ul style="list-style-type: none"> Similar to cutaneous lupus erythematosus, but without vascular ectasia, fibrin and C5b-9 deposition in dermal vasculature and dermo-epidermal junction.^{186,187}
Mycosis fungoides	<ul style="list-style-type: none"> Epidermotropism with Pautrier microabscesses Most common phenotype: CD2+, CD3+, βF1+, CD4+, CD8-, CD45RO with loss of mature T-lymphocytes markers (CD7, CD2 o CD5)⁹¹

Abbreviation: CD, contact dermatitis, except in the immunophenotype of mycosis fungoides.

CONCLUSION

Contact dermatitis is a multi-faceted and very frequently encountered disease in our clinical practice. Because CD mimics other skin conditions, and as other skin diseases may also simulate CD, a good understanding of diagnostic clues pointing towards CD, and its differential diagnoses, is crucial. To this end, we have here presented a practical

review, including visual materials, to facilitate its diagnosis. As discussed, the evaluation of several clinical parameters, as well as ruling out other skin conditions, and performing complementary tests are required. Pragmatically, including CD in the differential diagnosis of many inflammatory and non-inflammatory skin conditions may lead to earlier diagnosis, especially of atypical cases, and might even reduce the need to perform further, sometimes invasive,

complementary tests. Similarly, if an external factor is found, and a relevant (photo-)contact allergy is present, counselling and eviction of the demonstrated (photo)allergens can be pursued, with a change in the natural course of the disease and even the avoidance of systemic treatments to control the condition.

AFFILIATIONS

¹Dermatology Department, Hospital del Mar Research Institute, Department of Medicine, Universitat Autònoma de Barcelona (UAB), Barcelona, Spain

²Department of Dermatology, University Hospital Antwerp (UZA) and Research Group Immunology, University of Antwerp, Antwerp, Belgium

³Division of Allergy, University Clinic of Respiratory and Allergic Diseases Golnik, Golnik, Slovenia

⁴Department of Dermatology, Coimbra University Hospital, Coimbra, Portugal

⁵Faculty of Medicine, University of Coimbra, Coimbra, Portugal

⁶Department of Dermatology, University Medical Centre Maribor, Maribor, Slovenia

⁷Department of Dermatology, Inselspital University Hospital of Bern, Bern, Switzerland

⁸Department of Dermatology and Venereology, University Hospital Center Zagreb, School of Medicine University of Zagreb, Zagreb, Croatia

⁹Faculty of Medicine, Institute of Clinical Medicine, Clinic of Chest Diseases, Immunology and Allergology, Vilnius University, Vilnius, Lithuania

¹⁰Leeds Centre for Dermatology, Leeds Teaching Hospitals NHS Trust, Chapel Allerton Hospital, Leeds, UK

¹¹Department of Dermatology, Medical University of Poznan, Poznan, Poland

¹²Collegium Medicum, Jan Kochanowski University Kielce, Kielce, Poland

¹³Department of Dermatology, Environmental Medicine, Osnabrueck University, Osnabrueck, Germany

¹⁴Dermatology Unit, University of Campania "Luigi Vanvitelli", Naples, Italy

¹⁵Department of Clinical Medicine and Surgery, University of Naples Federico II, Naples, Italy

¹⁶Dermato-Allergology and Occupational Dermatology, Amsterdam University Medical Centers, Amsterdam, Netherlands

¹⁷Dermatology Department, Hospital del Mar Research Institute, Universitat Pompeu Fabra (UPF), Barcelona, Spain

ACKNOWLEDGEMENTS

All patients or legal tutors have given informed consent to publish the clinical pictures herein presented.

FUNDING INFORMATION

None to declare.

CONFLICT OF INTEREST STATEMENT

David Pesqué has benefited from a research grant of Leo-Pharma, outside the submitted work. Olivier Aerts was a speaker and/or advisor for Abbvie, Sanofi, L'Oréal/La Roche Posay and Bioderma/NAOS and has participated in research from Leo Pharma, outside the submitted work. Mojca Bizjak was a speaker and/or advisor for Novartis, outside the submitted work. Margarida Gonçalo was a speaker or advisor for Abbvie, Astrazeneca, Eli Lilly, Leo Pharma, Novartis, Pfizer, Sanofi and Takeda, outside the submitted work. Aleksandra Dugonik was a speaker and/or advisor for Abbvie, L'Oréal and Pfizer, outside the submitted work. Dagmar Simon was a speaker and/or advisor for Abbvie, Amgen, Astrazeneca, Galderma, Incyte, Leo Pharma, Eli Lilly, Novartis, Pfizer and Sanofi Genzyme, outside the submitted work. Suzana Ljubojević-Hadzavdić was a speaker and/or advisor for Eli Lilly, Abbvie, Sanofi, Novartis, Pliva, Bayer, Berlin-Chemie and Dr. Wolf, and has received grants or contracts from Abbvie, Amgen and Novartis, outside the

submitted work. Laura Malinauskiene was a speaker and/or advisor for Abbvie, Eli Lilly, Pfizer and Sanofi, outside the submitted work. Magdalena Czarnecka-Operacz was a speaker and/or advisor for Abbvie, Novartis, Eli Lilly, Leo Pharma, Astrazeneca and Sanofi-Aventis, outside the submitted work. Beata Krecisz was a speaker and/or advisor for Abbvie, Novartis, Eli Lilly, Leo Pharma, Astrazeneca and Sanofi, outside the submitted work. Swen M. John was a speaker and/or advisor for Leo Pharma and Sanofi, outside the submitted work. Anna Balato was a speaker and/or advisor for Abbvie, Almirall, Amgen, Boehringer Ingelheim, Eli Lilly, Novartis and UCB and has received grants or contracts from Almirall and BMS, outside the submitted work. Thomas Rustemeyer was a speaker and/or advisor for Atos, Novartis, Eli Lilly, Leo Pharma, Sanofi, Vifor and Ypsomed, outside the submitted work. Ana M. Giménez-Arnau is or recently was a speaker and/or advisor for and has received research funding or participated in research from Almirall, Amgen, AstraZeneca, Avene, Celldex, Escient Pharmaceuticals, Genentech, GSK, Instituto Carlos III-FEDER, Leo Pharma, Menarini, Mitsubishi Tanabe Pharma, Novartis, Sanofi-Regeneron, Servier, Thermo Fisher Scientific and Uriach Pharma/Neucor, outside the submitted work. All other authors declare no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICS STATEMENT

The patients or the patient's parents/guardians in this manuscript have given written informed consent to the publication of their case details.

ORCID

David Pesqué  <https://orcid.org/0000-0002-5821-9780>

Mojca Bizjak  <https://orcid.org/0000-0003-2595-468X>

Margarida Gonçalo  <https://orcid.org/0000-0001-6842-1360>

Suzana Ljubojević-Hadzavdić  <https://orcid.org/0000-0002-3640-9567>

Laura Malinauskiene  <https://orcid.org/0000-0002-3255-7669>

Ana M. Giménez-Arnau  <https://orcid.org/0000-0001-5434-7753>

REFERENCES

1. Kränke B, Schuster C. Contact dermatitis: relevant differential diagnoses, simulators, and variants. *J Dtsch Dermatol Ges.* 2015;13(11):1073–88.
2. Chan CX, Zug KA. Diagnosis and Management of Dermatitis, including atopic, contact, and hand eczemas. *Med Clin North Am.* 2021;105(4):611–26.
3. Scheinman PL, Vocanson M, Thyssen JP, Johansen JD, Nixon RL, Dear K, et al. Contact dermatitis. *Contact Dermatitis Nat Rev Dis Primers.* 2021;7(1):38.
4. Bailiff OA, Mowad CM. Mimics of dermatitis. *Immunol Allergy Clin North Am.* 2021;41(3):493–515.

5. Pongpairoj K, Ale I, Andersen KE, Bruze M, Diepgen TL, Elsner PU, et al. Proposed ICDRG classification of the clinical presentation of contact allergy. *Dermatitis*. 2016;27(5):248–58.
6. Bains SN, Nash P, Fonacier L. Irritant contact dermatitis. *Clin Rev Allergy Immunol*. 2019;56(1):99–109.
7. Li Y, Li L. Contact dermatitis: classifications and management. *Clin Rev Allergy Immunol*. 2021;61(3):245–81.
8. Williams J, Cahill J, Nixon R. Occupational autoeczematization or atopic eczema precipitated by occupational contact dermatitis? *Contact Dermatitis*. 2007;56(1):21–6.
9. Ale SI, Maibach HI. Irritant contact dermatitis versus allergic contact dermatitis. In: Chew AL, Maibach HI, editors. *Irritant dermatitis*. Volume 2008. Berlin, Heidelberg: Springer; 2008. p. 11–8.
10. Lachapelle JM. Allergic contact dermatitis: clinical aspects. *Rev Environ Health*. 2014;29(3):185–94.
11. Seyfarth F, Schliemann S, Antonov D, Elsner P. Dry skin, barrier function, and irritant contact dermatitis in the elderly. *Clin Dermatol*. 2011;29(1):31–6.
12. Rozas-Muñoz E, Gamé D, Serra-Baldrich E. Allergic contact dermatitis by anatomical regions: diagnostic clues. *Actas Dermosifiliogr (Engl Ed)*. 2018;109(6):485–507.
13. Lee J, Guo S, Dinalo J, DeLeo VA, Adler BL. Consort allergic contact dermatitis: a systematic review. *Dermatitis*. 2022;33(3):181–6.
14. Aerts O, Goossens A, Marguery MC, Castelain M, Boursault L, Giordano-Labadie F, et al. Photoaggravated allergic contact dermatitis and transient photosensitivity caused by methylisothiazolinone. *Contact Dermatitis*. 2018;78(4):241–5.
15. Kerr A, Ferguson J. Photoallergic contact dermatitis. *Photodermatol Photoimmunol Photomed*. 2010;26(2):56–65.
16. Pesqué D, Pérez-Manich J, García-Díez I, Segura S, Ferran M, González-Farré M, et al. Drug-induced ultraviolet B photosensitivity due to hydroxychloroquine: the unexpected side effect. *Contact Dermatitis*. 2023;88(2):168–70.
17. de Groot AC. Systemic allergic dermatitis (systemic contact dermatitis) from pharmaceutical drugs: a review. *Contact Dermatitis*. 2022;86(3):145–64.
18. Thyssen JP, Maibach HI. Drug-elicited systemic allergic (contact) dermatitis-update and possible pathomechanisms. *Contact Dermatitis*. 2008;59(4):195–202.
19. Woodruff CM, Botto N. The role of patch testing in evaluating delayed hypersensitivity reactions to medications. *Clin Rev Allergy Immunol*. 2022;62(3):548–61.
20. Häusermann P, Harr T, Bircher AJ. Baboon syndrome resulting from systemic drugs: is there strife between SDRIFE and allergic contact dermatitis syndrome? *Contact Dermatitis*. 2004;51(5–6):297–310.
21. de Risi-Pugliese T, Barailler H, Hamelin A, Amsler E, Gaouar H, Kurihara F, et al. Symmetrical drug-related intertriginous and flexural exanthema: a little-known drug allergy. *J Allergy Clin Immunol Pract*. 2020;8(9):3185–3189.e4.
22. Barbaud A, Poreaux C, Penven E, Waton J. Occupational protein contact dermatitis. *Eur J Dermatol*. 2015;25(6):527–34.
23. Giménez-Arnau AM, Pesqué D, Maibach HI. Contact Urticaria syndrome: a comprehensive review. *Curr Dermatol Rep*. 2022;11(4):194–201.
24. Pesqué D, Canal-García E, Rozas-Muñoz E, Pujol RM, Giménez-Arnau AM. Non-occupational protein contact dermatitis induced by mango fruit. *Contact Dermatitis*. 2021;84(6):458–60.
25. Dickel H, Bauer A, Brehler R, Mahler V, Merk HF, Neustädter I, et al. German S1 guideline: contact dermatitis. *J Dtsch Dermatol Ges*. 2022;20(5):712–34.
26. Eichenfield LF, Tom WL, Chamlin SL, Feldman SR, Hanifin JM, Simpson EL, et al. Guidelines of care for the management of atopic dermatitis: section 1. Diagnosis and assessment of atopic dermatitis. *J Am Acad Dermatol*. 2014;70(2):338–51.
27. Sidbury R, Tom WL, Bergman JN, Cooper KD, Silverman RA, Berger TG, et al. Guidelines of care for the management of atopic dermatitis: section 4. Prevention of disease flares and use of adjunctive therapies and approaches. *J Am Acad Dermatol*. 2014;71(6):1218–33.
28. Aerts O, Cattaert N, Lambert J, Goossens A. Airborne and systemic dermatitis, mimicking atopic dermatitis, caused by methylisothiazolinone in a young child. *Contact Dermatitis*. 2013;68(4):250–1.
29. Lofgren SM, Warshaw EM. Dyshidrosis: epidemiology, clinical characteristics, and therapy. *Dermatitis*. 2006;17(4):165–81.
30. Stuckert J, Nedorost S. Low-cobalt diet for dyshidrotic eczema patients. *Contact Dermatitis*. 2008;59(6):361–5.
31. Specht S, Persaud Y. *Asteatotic Eczema*. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023.
32. Silverberg JI, Hou A, Warshaw EM, Maibach HI, Belsito DV, DeKoven JG, et al. Allergens in patients with a diagnosis of seborrheic dermatitis, north American contact dermatitis group data, 2001–2016. *J Am Acad Dermatol*. 2022;86(2):460–3.
33. Kuznetsov AV, Erlenkeuser-Uebelhoe I, Thomas P. Contact allergy to propylene glycol and dodecyl gallate mimicking seborrheic dermatitis. *Contact Dermatitis*. 2006;55(5):307–8.
34. Leung AKC, Lam JM, Leong KF, Leung AAM, Wong AHC, Hon KL. Nummular eczema: an updated review. *Recent Pat Inflamm Allergy Drug Discov*. 2020;14(2):146–55.
35. Bonamonte D, Foti C, Vestita M, Ranieri LD, Angelini G. Nummular eczema and contact allergy: a retrospective study. *Dermatitis*. 2012;23(4):153–7.
36. Silverberg JI, Hou A, Warshaw EM, Maibach HI, Belsito DV, DeKoven JG, et al. Prevalence and trend of allergen sensitization in patients with nummular (discoid) eczema referred for patch testing: north American contact dermatitis group data, 2001–2016. *Contact Dermatitis*. 2021;85(1):46–57.
37. Foti C, Caliano G, Mazzotta S, Guarneri F, Romita P. Allergic contact dermatitis to clostridiopeptidase a with a nummular eczema-like spread. *Contact Dermatitis*. 2022;86(2):122–3.
38. Tissinier-Vivent M, Houle MC, Du-Thanh A, Raison-Peyron N. Persulphates allergic contact dermatitis from spa: a new aetiology of nummular contact dermatitis. *Contact Dermatitis*. 2023;89(2):128–30.
39. Sundaresan S, Migden MR, Silapunt S. Stasis dermatitis: pathophysiology, evaluation, and management. *Am J Clin Dermatol*. 2017;18(3):383–90.
40. Nedorost ST. Allergic contact sensitization in healthy skin differs from sensitization in chronic dermatitis: atopic, occupational wet work, and stasis dermatitis. *Dermatol Clin*. 2020;38(3):301–30.
41. Silverberg JI, Hou A, Warshaw EM, Maibach HI, Belsito DV, DeKoven JG, et al. Prevalence and trend of allergen sensitization in patients with a diagnosis of stasis dermatitis referred for patch testing, north American contact dermatitis group data, 2001–2016. *Arch Dermatol Res*. 2022;314(9):857–67.
42. Barbaud A, Castagna J, Soria A. Skin tests in the work-up of cutaneous adverse drug reactions: a review and update. *Contact Dermatitis*. 2022;86(5):344–56.
43. Joshi SR, Khan DA. Non-IgE-mediated drug hypersensitivity reactions. *Curr Allergy Asthma Rep*. 2021;21(7):41.
44. Park S, Cha NR, Kim SC, Oh SH, Kim J. Erythema multiforme-like allergic contact dermatitis to rosewood. *Contact Dermatitis*. 2021;85:242–4. <https://doi.org/10.1111/cod.13819>
45. Lin N, Tang X, Zhou H, Han J. Erythema multiforme-like contact dermatitis caused by herbal medicine. *Contact Dermatitis*. 2019;80(3):189–90.
46. Abdulmula A, Mufti A, Ho JSS, Kashetsky N, Yeung J, Maibach HI. Erythema Multiforme-like contact dermatitis: a systematic review of characteristics and treatment outcomes. *Dermatitis*. 2021;32(6):e161–e165.
47. Martínez-Fernández A, Sánchez-Herreros C, Cabana-Navia R, Camino-Salvador JM, Palomo-Cousido S, Arribas-Granados P, et al. Erythema multiforme-like allergic contact dermatitis from bay leaves in a home-made natural remedy. *Contact Dermatitis*. 2023;89(2):118–20.
48. Traineau H, Benassaia E, Diaz E, Ingen-Housz-Oro S, Assier H. Allergic contact dermatitis from benzyl alcohol mimicking acute generalized exanthematous pustulosis. *Contact Dermatitis*. 2022;87(1):100–2.

49. Broesby-Olsen S, Clemmensen O, Andersen KE. Allergic contact dermatitis from a topical corticosteroid mimicking acute generalized exanthematous pustulosis. *Acta Derm Venereol*. 2005;85(5):444–5.
50. Abdelmalek M, Mahindrakar S, Wisner E. Lamotrigine-induced erythema multiforme mimicking contact dermatitis. *J Drugs Dermatol*. 2006;5(1):76–9.
51. Deno R, Nakagawa Y, Itoi-Ochi S, Kotobuki Y, Kiyohara E, Wataya-Kaneda M, et al. Fixed drug eruption caused by allylisopropylacetylurea mimicking contact dermatitis of the face. *Contact Dermatitis*. 2020;82(1):56–7.
52. Blum AE, Burgin S. Eczematous Drug Eruptions. *Am J Clin Dermatol*. 2021 May;22(3):349–66.
53. Quaranta M, Eyerich S, Knapp B, Nasorri F, Scarponi C, Mattii M, et al. Allergic contact dermatitis in psoriasis patients: typical, delayed, and non-interacting. *PLoS One*. 2014;9(7):e101814.
54. Lugović-Mihić L, Novak-Bilić G, Vučić M, Japundžić I, Bukvić I. CD44 expression in human skin: high expression in irritant and allergic contact dermatitis and moderate expression in psoriasis lesions in comparison with healthy controls. *Contact Dermatitis*. 2020;82(5):297–306.
55. Kolesnik M, Franke I, Lux A, Quist SR, Gollnick HP. Eczema in Psoriatico: an important differential diagnosis between chronic allergic contact dermatitis and psoriasis in palmoplantar localization. *Acta Derm Venereol*. 2018;98(1):50–8.
56. Bangsgaard N, Engkilde K, Thyssen JP, Linneberg A, Nielsen NH, Menné T, et al. Inverse relationship between contact allergy and psoriasis: results from a patient- and a population-based study. *Br J Dermatol*. 2009;161(5):1119–23.
57. Silverberg JI, Hou A, DeKoven JG, Warshaw EM, Maibach HI, Atwater AR, et al. Prevalence and trend of allergen sensitization in patients referred for patch testing with a final diagnosis of psoriasis: north American contact dermatitis group data, 2001–2016. *Contact Dermatitis*. 2021;85(4):435–45.
58. Tziotzios C, Lee JYW, Brier T, Saito R, Hsu CK, Bhargava K, et al. Lichen planus and lichenoid dermatoses: clinical overview and molecular basis. *J Am Acad Dermatol*. 2018;79(5):789–804.
59. Lim SS, Chamberlain A, Hur K, Ohn J, Han B, Mar V, et al. Dermoscopic evaluation of inflammatory nail disorders and their mimics. *Acta Derm Venereol*. 2021;101(9):adv00548. doi:10.2340/00015555-3917
60. Kuroda K, Hisanaga Y. The diagnosis of lichen-planus-like contact dermatitis to chlorpheniramine maleate. *Dermatology*. 2002;205(3):281–4.
61. Bonamonte D, Foti C, Vestita M, Angelini G. Noneczematous contact dermatitis. *ISRN Allergy*. 2013;2013:361746.
62. Navarro-Triviño FJ, Jadcak P, Llamas-Molina JM, Ruiz-Villaverde R. Lichenoid contact dermatitis caused by *Morus nigra* L. *Contact Dermatitis*. 2021;84(6):484–6.
63. Rahat S, Kashetsky N, Bagit A, Sachdeva M, Lytvyn Y, Mufti A, et al. Can we separate Oral lichen planus from allergic contact dermatitis and should we patch test? A systematic review of chronic Oral lichenoid lesions. *Dermatitis*. 2021;32(3):144–50.
64. Ju T, Vander Does A, Mohsin N, Yosipovitch G. Lichen simplex Chronicus itch: an update. *Acta Derm Venereol*. 2022;102:adv00796. <https://doi.org/10.2340/actadv.v102.4367>
65. Chey WY, Kim KL, Yoo TY, Lee AY. Allergic contact dermatitis from hair dye and development of lichen simplex chronicus. *Contact Dermatitis*. 2004;51(1):5–8.
66. Linauskiene K, Isaksson M. Allergic contact dermatitis from formaldehyde mimicking impetigo and initiating rosacea. *Contact Dermatitis*. 2018;78(5):359–61.
67. D'Erme AM, Gola M. Allergic contact dermatitis induced by topical hydrocortisone-17-butyrate mimicking papular rosacea. *Dermatitis*. 2012;23(2):95–6.
68. Goossens A, Aerts O. Contact allergy to and allergic contact dermatitis from formaldehyde and formaldehyde releasers: a clinical review and update. *Contact Dermatitis*. 2022;87(1):20–7.
69. Oakley AM, Ramsey ML. Polymorphic light eruption. *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2023. p. 2023.
70. Kosboth M, Chin-Loy A, Lyons R, Wesson SK, Reeves WH. Malar rash caused by metal allergy in a patient with systemic lupus erythematosus. *Nat Clin Pract Rheumatol*. 2007;3(4):240–5.
71. Van Aerde E, Kerre S, Goossens A. Discoid lupus triggered by allergic contact dermatitis caused by a hair dye. *Contact Dermatitis*. 2016;74(1):61–4.
72. Darnall A, Hamann CR, Hamann D. Eyelid allergic contact dermatitis masking dermatomyositis. *Contact Dermatitis*. 2023;88(2):164–6.
73. Snyder M, Turrentine JE, Cruz PD Jr. Photocontact dermatitis and its clinical mimics: an overview for the allergist. *Clin Rev Allergy Immunol*. 2019;56(1):32–40.
74. Cunningham EP, Carter NF. Tinea incognito "mask"erasing as allergic contact dermatitis due to COVID-19 facial covering in children. *Pediatr Dermatol*. 2022;39(2):326–7.
75. Kaminska R, Mörtenhumer M. Nummular allergic contact dermatitis after scabies treatment. *Contact Dermatitis*. 2003;48(6):337.
76. Zhong LS, Huang BQ, Tang WT, Zhuo ZQ. Bullous impetigo-like irritant contact dermatitis caused by perfume. *Pediatr Dermatol*. 2022;39(2):320–1.
77. Verma A, Tancharoen C, Tam MM, Nixon R. Pustular allergic contact dermatitis caused by fragrances. *Contact Dermatitis*. 2015;72(4):245–8.
78. Martina E, Diotallevi F, Radi G, Campanati A, Offidani A. A tricky case of contact dermatitis: simultaneous allergic reaction to topical and oral acyclovir, and scar gel. *Contact Dermatitis*. 2021;84(3):203–5.
79. Mori N, Mizawa M, Hara H, Norisugi O, Makino T, Nakano H, et al. Hailey-Hailey disease diagnosed based on an exacerbation of contact dermatitis with topical crotamiton. *Eur J Dermatol*. 2014;24(2):263–4.
80. Napolitano M, Fabbrocini G, Martora F, Genco L, Noto M, Patruno C. Children atopic dermatitis: diagnosis, mimics, overlaps, and therapeutic implication. *Dermatol Ther*. 2022;35(12):e15901.
81. Menendez M, Menendez C, Anzalone G. Paget's disease of the breast mimicking a chronic wound. *Wounds*. 2008;20(12):321–4.
82. O'Donnell M, Shalabi D, Gaspari AA, Nikbakht N. Determining the role of allergic contact dermatitis in mycosis fungoides. *Int J Dermatol*. 2020;59(6):e221–e223.
83. Park JW, Jung Y, Han JH, Kim M. Folliculotropic mycosis fungoides clinically resembling allergic contact dermatitis. *Contact Dermatitis*. 2021;85:236–9. <https://doi.org/10.1111/cod.13816>
84. Smets K, Busschots A, Hauben E, Goossens A. B-cell lymphomatoid contact dermatitis caused by methylisothiazolinone and methylchlorisothiazolinone. *Eur J Dermatol*. 2018;28(1):91–3.
85. Golińska J, Sar-Pomian M, Rudnicka L. Dermoscopic features of psoriasis of the skin, scalp and nails - a systematic review. *J Eur Acad Dermatol Venereol*. 2019;33(4):648–60.
86. Aringer M, Leuchten N, Johnson SR. New criteria for lupus. *Curr Rheumatol Rep*. 2020;22(6):18. <https://doi.org/10.1007/s11926-020-00896-6>
87. Sánchez-Schmidt JM, Pujol-Vallverdú RM. Diagnóstico diferencial de las lesiones cutáneas en el lupus eritematoso. *Semin Fund Esp Reumatol*. 2006;7:12–26.
88. Batista M, Gonçalves M. The rash that presents as a red swollen face. *Clin Dermatol*. 2020;38(1):63–78. <https://doi.org/10.1016/j.clindermatol.2019.10.017>
89. Pirmez R. The dermatoscope in the hair clinic: Trichoscopy of scarring and nonscarring alopecia. *J Am Acad Dermatol*. 2023;89(2S):S9–S15.
90. Muschalek W, Hermasch MA, Poblete-Gutiérrez P, Frank J. The Porphyrias. *J Dtsch Dermatol Ges*. 2022;20(3):316–31.
91. Pujol RM, Gallardo F. Cutaneous lymphomas - part I: mycosis fungoides, Sézary syndrome, and CD30+ cutaneous lymphoproliferative disorders. *Actas Dermosifiliogr (Engl Ed)*. 2021;112(1):14–23.

92. Palaniappan V, Karthikeyan K. Bowen's Disease. *Indian Dermatol Online J.* 2022;13(2):177–89.
93. Pesqué D, Curto-Barredo L, Pastor-Nieto MA, Pujol RM, Giménez-Arnau AM. Allergic contact dermatitis due to over-the-counter antifungal nail brush: is Resacetophenone an under-recognized allergen? *Dermatitis.* 2023;35:190–1. <https://doi.org/10.1089/derm.2023.0103>
94. Shi M, Zhang H, Chen X, Guo Y, Tao J, Qi H, et al. Clinical features of atopic dermatitis in a hospital-based setting in China. *J Eur Acad Dermatol Venereol.* 2011;25(10):1206–12.
95. Simpson EL, Thompson MM, Hanifin JM. Prevalence and morphology of hand eczema in patients with atopic dermatitis. *Dermatitis.* 2006;17(3):123–7.
96. Micali G, Verzi AE, Giuffrida G, Panebianco E, Musumeci ML, Lacarrubba F. Inverse psoriasis: from diagnosis to current treatment options. *Clin Cosmet Investig Dermatol.* 2019;12:953–9.
97. Rieder E, Hale CS, Meehan SA, Leger M. Palmoplantar lichen planus. *Dermatol Online J.* 2014;20(12). <http://dx.doi.org/10.5070/D32012025062>
98. Hassan S, Chen JK. A case of pigmented contact dermatitis caused by Ethylhexylglycerin. *Dermatitis.* 2022;33(3):e23–e24.
99. Kwong HL, Lim SPR. Pigmented contact dermatitis in the axillae caused by hydroperoxides of limonene. *JAAD Case Rep.* 2020;6(6):476–8.
100. Alagappan U, Tay YK, Lim SP. Pigmented contact dermatitis secondary to benzyl salicylate. *Acta Derm Venereol.* 2013;93(5):590.
101. Hayakawa R, Matsunaga K, Arima Y. Depigmented contact dermatitis due to incense. *Contact Dermatitis.* 1987;16(5):272–4.
102. Kim MS, Chung BY, Chang SE, Oh SH, Ryu HJ, Kim DH, et al. Pigmented contact dermatitis and hair dyes: a retrospective case-control multicenter study in Korea. *J Eur Acad Dermatol Venereol.* 2023;37(12):2543–9.
103. Marcant P, Sobaszek A, Nisse C, Staumont-Sallé D. Occupational purpuric contact dermatitis to methylchloroisothiazolinone / methylisothiazolinone. *Contact Dermatitis.* 2020;83(1):69–71.
104. Navarro-Triviño FJ, Ruiz-Villaverde R. Purpuric contact dermatitis caused by N-isopropyl-N'-phenyl-p-phenylenediamine from a Samsung fit strap. *Contact Dermatitis.* 2020;83(1):68–9.
105. Carlet C, Castelain F, Pelletier F, Aubin F, Raison-Peyron N. Postoperative purpuric contact dermatitis: think about antiseptics. *Contact Dermatitis.* 2019;81(3):215–6.
106. Knackstedt TJ, Zug KA. T cell lymphomatoid contact dermatitis: a challenging case and review of the literature. *Contact Dermatitis.* 2015;72(2):65–74.
107. Viedma-Martinez M, Gallo-Pineda G, Jimenez-Anton A, Garcia-Dorado J, Alonso-Fraile P, Santos-Briz A. Lymphomatoid contact dermatitis caused by *Anemonia viridis* sting. *Contact Dermatitis.* 2023;88(5):410–2.
108. Fernández-Canga P, Sánchez-Sambucety P, Valladares-Narganes LM, Varas-Meis E, Rodríguez-Prieto MA. Lymphomatoid contact dermatitis induced by acrylates mimicking Lymphomatoid papulosis. *Dermatitis.* 2018;29(3):167–8.
109. Coleman E, Bhawan J. Baby wet wipes: an unusual culprit of Lymphomatoid contact dermatitis mimicking mycosis Fungoides. *Am J Dermatopathol.* 2022;44(3):205–6.
110. Fernández-Aceñero MJ, Fernández-López P. Granulomatous contact dermatitis to palladium following ear piercing. *Indian J Dermatol Venereol Leprol.* 2008;74(6):651–2.
111. Matsudate Y, Yamashita M, Fujii Y, Urano Y. Contact granulomatous hypersensitivity to indium in a patient with orofacial granulomatosis. *Contact Dermatitis.* 2019;81(4):293–4.
112. Lugović-Mihić L, Pilipović K, Crnarić I, Šitum M, Duvančić T. Differential diagnosis of Cheilitis - how to classify Cheilitis? *Acta Clin Croat.* 2018;57(2):342–51.
113. Salah NB, Lahouel I, Belhadjali H, Machghoul S, Fadhel NB, Youssef M, et al. Pustular allergic contact dermatitis caused by a topical probiotic. *Dermatitis.* 2022;33(6):e66–e68.
114. Muñoz-Aceituno E, Reolid A, Chicharro P, Rodríguez-Jiménez P, Freih-Fraih A, Sánchez-Pérez JA. Occupational pustular allergic contact dermatitis. *Contact Dermatitis.* 2020;83(1):54–5.
115. Tracz ES, Sommerlund M, Bregnhøj A. Pustular allergic contact dermatitis caused by a sunscreen. *Contact Dermatitis.* 2020;83(4):328–9.
116. Merhy R, Weill A, Ingen-Housz-Oro S, Assier H. Pustular allergic contact dermatitis of the dorsum of hands related to rubber gloves additives. *Contact Dermatitis.* 2022;87(2):199–201.
117. Lahouel I, Thabouti M, Belhadjali H, Fadhel NB, Soua Y, Youssef M, et al. Pustular allergic contact dermatitis caused by colophonium in depilatory wax. *Contact Dermatitis.* 2021;84(2):130–2.
118. Lönngren V, Young E, Simanaitis M, Svedman C. Neutrophilic and eosinophilic dermatitis caused by contact allergic reaction to paraphenylenediamine in hair dye. *Arch Dermatol.* 2012;148(11):1299–301.
119. Webern RV, Mockenhaupt M, Schempp CM. Bullous irritant contact dermatitis caused by the spitting palm, *Euphorbia leuconeura* BOISSIER. *Contact Dermatitis.* 2022;87(5):460–2.
120. Soylu S. Bullous irritant contact dermatitis and discoloration due to Juglone: a frequently seen, rarely reported skin disorders. *Dermatitis.* 2023;34:571–2. <https://doi.org/10.1089/derm.2023.0038>
121. Soriano LF, Rasool S, Buckley DA. Bullous allergic contact dermatitis from disperse dyes in Lycra cycling shorts. *Contact Dermatitis.* 2020;82(2):122–4.
122. Naraghi ZS, Mansouri P, Mortazavi M. A clinicopathological study on acute cutaneous lesions induced by sulfur mustard gas (yperite). *Eur J Dermatol.* 2005;15(3):140–5.
123. Xará J, Matos A, Soares J, Teixeira J, Ramos L, Gonçalves M. Localized hypertrichosis as a manifestation of contact allergy to aluminium. *Contact Dermatitis.* 2023;89(1):58–9.
124. Liu KX, Zimarowski MJ, Wu PA. Contact dermatitis associated with alopecia and hyperpigmentation. *Pediatr Dermatol.* 2017;34(5):624–5.
125. Munoz CA, Gaspari A, Goldner R. Contact dermatitis from a prosthesis. *Dermatitis.* 2008;19(2):109–11.
126. Matos AL, Pacheco V, Lopes R, Cardoso JC, Carvalho L, Rustemeyer T, et al. Inflammatory lung pseudotumor possibly related to titanium hypersensitivity. *Contact Dermatitis.* 2023;88(5):419–22.
127. Wawrzynski J, Gil JA, Goodman AD, Waryasz GR. Hypersensitivity to orthopedic implants: a review of the literature. *Rheumatol Ther.* 2017;4(1):45–56.
128. Alves F, Ribeiro JC, Alves M, Gonçalves M. Titanium allergy as a possible cause of extrusion of a bone conduction ear implant. *Contact Dermatitis.* 2020;83(2):148–9.
129. Kerre S, Aerts O. Disfiguring angioedema-like airborne dermatitis from methylisothiazolinone in paints: about time to regulate? *Contact Dermatitis.* 2021;85(5):578–9.
130. Tukenmez Demirci G, Kivanc Altunay I, Atis G, Kucukunal A. Allergic contact dermatitis mimicking angioedema due to paraphenylenediamine hypersensitivity: a case report. *Cutan Ocul Toxicol.* 2012;31(3):250–2.
131. Sergoyne L, Mertens M, Dendooven E, Leysen J, Aerts O. Allergic contact dermatitis, mimicking atopic dermatitis, associated with the use of essential oils in "home-made" cosmetics and aromatherapy diffusers. *Contact Dermatitis.* 2020;83(4):311–3.
132. Oiso N, Ota T, Yoshinaga E, Endo H, Kawara S, Kawada A. Allergic contact dermatitis mimicking atopic dermatitis due to enoxolone in a topical medicament. *Contact Dermatitis.* 2006;54(6):351.
133. Raison-Peyron N, Litovsky J, Samaran Q, Dereure O. Severe allergic contact dermatitis to coumarin contained in a perfume mimicking skin drug reaction. *Contact Dermatitis.* 2023;88(5):416–7.
134. Casanova-Esquembre A, Lorca-Spröhnle J, Labrandero-Hoyos C, Peñuelas-Leal R, Zaragoza-Ninet V, Pérez-Ferriols A. Allergic contact dermatitis mimicking anti-interleukin 17 eczematous eruption. *Dermatitis.* 2022;34:464–5. <https://doi.org/10.1097/DER.0000000000000880>
135. Mestach L, Goossens A. Allergic contact dermatitis and nail damage mimicking psoriasis caused by nail hardeners. *Contact Dermatitis.* 2016;74(2):112–4.
136. Takeuchi S, Matsuzaki Y, Ikenaga S, Nishikawa Y, Kimura K, Nakano H, et al. Garlic-induced irritant contact dermatitis mimicking nail psoriasis. *J Dermatol.* 2011;38(3):280–2.

137. Coe J, Robinson R, Wilkinson SM. Nail dystrophy mimicking psoriatic disease caused by contact allergy to nail varnish allergens including copolymers. *Contact Dermatitis*. 2021;85(5):600–2.
138. Lapière K, Matthieu L, Meuleman L, Lambert J. Primula dermatitis mimicking lichen planus. *Contact Dermatitis*. 2001;44(3):199.
139. Panda S, Malakar S. Lichen nitidus-like allergic contact dermatitis - a clinicopathological study. *Contact Dermatitis*. 2003;49(1):43.
140. Lakshmi C, Srinivas CR, Pillai SB, Shanthakumari S. Parthenium dermatitis manifesting clinically as polymorphic light eruption and prurigo nodularis-like lesions with vasculitis-like picture on histopathology. *Indian Dermatol Online J*. 2011;2(2):97–100.
141. Aerts O, Van Dyck F, Dandelooy J, Mellaerts T, Lambert J. Contact dermatitis masquerading as folliculitis decalvans: methylisothiazolinone strikes again! *Dermatitis*. 2014;25(5):276.
142. Thomas P, Korting HC, Przybilla B. Propolis-induced allergic contact dermatitis mimicking pemphigus vulgaris. *Arch Dermatol*. 1998;134(4):511–3.
143. Navarro-Triviño FJ, Cuenca-Barrales C, Ruiz-Villaverde R. Eugenol allergy mimicking recurrent aphthous stomatitis and burning mouth syndrome. *Contact Dermatitis*. 2019;81(6):462–3.
144. Corazza M, Bencivelli D, Zedde P, Monti A, Zampino MR, Borghi A. Severe contact urticaria, mimicking allergic contact dermatitis, due to a surgical mask worn during the COVID-19 pandemic. *Contact Dermatitis*. 2021;84(6):466–7.
145. Ertugrul A, Hizli Demirkale Z, Bostanci I. Nickel allergy in an adolescent mimicking angioedema. *Contact Dermatitis*. 2021;84(4):263–5.
146. Opstrup MS, Garvey LH, Johansen JD, Bregnbak DK, Thyssen JP. A contact allergic reaction to budesonide mimicking immediate-type allergy. *Contact Dermatitis*. 2017;77(1):62–3.
147. Lee SX, Boontaveeyuwat E, Thaiwat S, Rustin M, McFadden J. Severe allergic contact dermatitis mimicking angioedema caused by propolis used as a traditional remedy. *Contact Dermatitis*. 2018;79(3):185–6.
148. Singh S, Khandpur S, Sharma VK. Allergic contact dermatitis to Parthenium hysterophorus mimicking actinic prurigo. *Indian J Dermatol Venereol Leprol*. 2015;81(1):82–4.
149. Feroze K, Manoj J, Venkitakrishnan S. Allergic contact dermatitis mimicking mammary paget's disease. *Indian J Dermatol*. 2008;53(3):154–5.
150. Orbaneja JG, Diez LI, Lozano JL, Salazar LC. Lymphomatoid contact dermatitis: a syndrome produced by epicutaneous hypersensitivity with clinical features and a histopathologic picture similar to that of mycosis fungoides. *Contact Dermatitis*. 1976;2(3):139–43.
151. Fisher AA. Allergic contact dermatitis mimicking mycosis fungoides. *Cutis*. 1987;40(1):19–21.
152. Hession MT, Scheinman PL. Lymphomatoid allergic contact dermatitis mimicking cutaneous T cell lymphoma. *Dermatitis*. 2010;21(4):220.
153. Pasolini G, Ghidini P, Arisi M, Pedretti A, Ungari M, Pinton PC. Pseudolymphoma tattoo-induced. *Dermatol Rep*. 2011;3(3):e47.
154. Hague J, Ilchyshyn A. Nickel allergy mimicking basal cell carcinoma. *Contact Dermatitis*. 2006;54(6):344–5.
155. Thomson KF, Charles-Holmes R, Beck MH. Primula dermatitis mimicking herpes simplex. *Contact Dermatitis*. 1997;37(4):185–6.
156. Du-Thanh A, Lalande M, Raison-Peyron N, Dereure O. Atypical and severe airborne isothiazolinone contact dermatitis mimicking Kaposi-Juliusberg syndrome. *Contact Dermatitis*. 2017;76(5):297–8.
157. Lu LY, Yen H, Chen WT. Zosteriform contact dermatitis mimicking herpes zoster. *Contact Dermatitis*. 2021;85(6):706–7.
158. Kim C, Craiglow BG, Watsky KL, Antaya RJ. Allergic contact dermatitis to benzoyl peroxide resembling impetigo. *Pediatr Dermatol*. 2015;32(4):e161–e162.
159. Gruson LM, Chang MW. Berloque dermatitis mimicking child abuse. *Arch Pediatr Adolesc Med*. 2002;156(11):1091–3.
160. Fanny SA, Burge L, Small A, Turbeville J, Donaruma-Kwoh M, Adekunle-Ojo A. When “the runs” also burns: case reports of severe diaper dermatitis in children with acute gastroenteritis mimicking scald burns. *Pediatr Dermatol*. 2023;40(2):341–4.
161. Tsai YC. Image gallery: contact dermatitis resulting from tarantula-hair injury mimicking ischaemia: a case with dermoscopy follow-up. *Br J Dermatol*. 2018;178(3):e227.
162. Elmas ÖF, Akdeniz N, Atasoy M, Karadag AS. Contact dermatitis: a great imitator. *Clin Dermatol*. 2020;38(2):176–92.
163. Salomon RJ, Briggaman RA, Wernikoff SY, Kayne AL. Localized bullous pemphigoid. A mimic of acute contact dermatitis. *Arch Dermatol*. 1987;123(3):389–92.
164. Soni BP, McAlvany JP, Fleischer AB, Sherertz EF. Bullous pemphigoid mimicking contact dermatitis. *Contact Dermatitis*. 1996;35(5):314.
165. Ajith C, Dogra S, Radotra BD, Handa S. Granulomatous rosacea mimicking eyelid dermatitis. *Indian J Dermatol Venereol Leprol*. 2005;71(5):365–6.
166. Johannemann D, Brunner M, Zouboulis CC. Facial sweet syndrome mimicking photoallergic contact dermatitis. *J Dtsch Dermatol Ges*. 2009;7(8):697–700.
167. Gambichler T, Doerler M, Scheel CH, Bastek K, Chatzipantazi M. Prurigo pigmentosa (Nagashima disease) in bathing suit distribution mimicking contact dermatitis. *J Eur Acad Dermatol Venereol*. 2023;37(4):e472–e473.
168. Sowerby L, Turchin I. Answer: can you identify this condition? *Can Fam Physician*. 2007;53(11):1902.
169. Sommer S, Wilkinson SM. Porphyria cutanea tarda masquerading as chronic hand eczema. *Acta Derm Venereol*. 2004;84(2):170–1.
170. Wolf R, Wolf D, Viskoper RJ, Sandbank M. Norwegian-type scabies mimicking contact dermatitis in an immunosuppressed patient. *Postgrad Med*. 1985;78(1):228–30.
171. Canan H, Altan-Yaycioglu R, Durdu M. Periocular Paederus dermatitis mimicking preseptal cellulitis. *Can J Ophthalmol*. 2013;48(2):121–5.
172. Johansen JD, Aalto-Korte K, Agner T, Andersen KE, Bircher A, Bruze M, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing - recommendations on best practice. *Contact Dermatitis*. 2015;73(4):195–221.
173. Wu PA. The importance of education when patch testing. *Dermatol Clin*. 2020;38(3):351–60.
174. Ramirez F, Chren MM, Botto N. A review of the impact of patch testing on quality of life in allergic contact dermatitis. *J Am Acad Dermatol*. 2017;76(5):1000–4.
175. Gonçalves M, Ferguson J, Boneville A, Bruynzeel DP, Giménez-Arnau A, Goossens A, et al. Photopatch testing: recommendations for a European photopatch test baseline series. *Contact Dermatitis*. 2013;68(4):239–43.
176. Lachapelle J-M, Marot L. Histopathological and immunohistochemical features of irritant and allergic contact dermatitis. 5th ed. Berlin-Heidelberg: Springer-Verlag; 2011.
177. Gru AA, Salavaggi AL. Common spongiotic dermatoses. *Semin Diagn Pathol*. 2017;34(3):226–36.
178. Bieber T. Atopic dermatitis. *Ann Dermatol*. 2010;22(2):125–37.
179. Rosa G. Allergic contact dermatitis. [PathologyOutlines.com](https://www.pathologyoutlines.com/topic/skinontumorallergiccontact.html) Available from: <https://www.pathologyoutlines.com/topic/skinontumorallergiccontact.html>. [Accessed 15th June 2023].
180. Rietschel RL, Fowler JF. Histology of contact dermatitis. In: Rietschel RL et al., editors. Hamilton, ON. Maidenhead: McGraw-Hill Education, distributor: BC Decker; 2008. p. 862.
181. Summey BT, Bowen SE, Allen HB. Lichen planus-like atopic dermatitis: expanding the differential diagnosis of spongiotic dermatitis. *J Cutan Pathol*. 2008;35(3):311–4.
182. Gonçalves M. Phototoxic and photoallergic reactions. In: Johansen J, Frosch P, Lepoittevin J, editors. Contact dermatitis. Springer International Publishing; Springer Nature Switzerland. 2020.
183. Yosipovitch G, Nedorost ST, Silverberg JI, Friedman AJ, Canosa JM, Cha A. Stasis dermatitis: an overview of its clinical presentation, pathogenesis, and management. *Am J Clin Dermatol*. 2023;24(2):275–86.
184. Rapini RP. Practical dermatopathology. 2nd ed. Edinburgh: Elsevier Saunders; 2012. p. 450.

185. Patterson JW. The spongiotic reaction pattern. In: Patterson JW, editor. *Weedon's skin pathology*. 4th ed. Amsterdam: Elsevier; 2015.
186. Crowson AN, Magro CM. The role of microvascular injury in the pathogenesis of cutaneous lesions of dermatomyositis. *Hum Pathol*. 1996;27(1):15–9.
187. Krathen MS, Fiorentino D, Werth VP. Dermatomyositis. *Curr Dir Autoimmun*. 2008;10:313–32.
188. Amsler E, Assier H, Soria A, Bara C, Ferrier le Bouëdec MC, Barbaud A, et al. What is the optimal duration for a ROAT? The experience of the French dermatology and Allergology group (DAG). *Contact Dermatitis*. 2022;87(2):170–5.
189. Goossens A. Alternatives aux patch-tests [alternatives to patch tests]. *Ann Dermatol Venereol*. 2009;136(8–9):623–5.

How to cite this article: Pesqué D, Aerts O, Bizjak M, Gonçalo M, Dugonik A, Simon D, et al. Differential diagnosis of contact dermatitis: A practical-approach review by the EADV Task Force on contact dermatitis. *J Eur Acad Dermatol Venereol*. 2024;00:1–19. <https://doi.org/10.1111/jdv.20052>