

## REVIEW ARTICLE

Cellular and immunological basis of the host-parasite relationship during infection with *Neospora caninum*

A. HEMPHILL\*, N. VONLAUFEN and A. NAGULESWARAN

*Institute of Parasitology, Vetsuisse Faculty, University of Berne, Länggass-Strasse 122, CH-3012 Berne, Switzerland**(Received 6 January 2006; revised 4 April 2006; accepted 8 April 2006; first published online 6 June 2006)*

## SUMMARY

*Neospora caninum* is an apicomplexan parasite that is closely related to *Toxoplasma gondii*, the causative agent of toxoplasmosis in humans and domestic animals. However, in contrast to *T. gondii*, *N. caninum* represents a major cause of abortion in cattle, pointing towards distinct differences in the biology of these two species. There are 3 distinct key features that represent potential targets for prevention of infection or intervention against disease caused by *N. caninum*. Firstly, tachyzoites are capable of infecting a large variety of host cells *in vitro* and *in vivo*. Secondly, the parasite exploits its ability to respond to alterations in living conditions by converting into another stage (tachyzoite-to-bradyzoite or *vice versa*). Thirdly, by analogy with *T. gondii*, this parasite has evolved mechanisms that modulate its host cells according to its own requirements, and these must, especially in the case of the bradyzoite stage, involve mechanisms that ensure long-term survival of not only the parasite but also of the host cell. In order to elucidate the molecular and cellular bases of these important features of *N. caninum*, cell culture-based approaches and laboratory animal models are being exploited. In this review, we will summarize the current achievements related to host cell and parasite cell biology, and will discuss potential applications for prevention of infection and/or disease by reviewing corresponding work performed in murine laboratory infection models and in cattle.

Key words: *Neospora caninum*, host cell invasion, cell surface receptors, host cell modulation, stage conversion, murine model, vaccination.

## NEOSPORA CANINUM, NEOSPOROSIS AND HOST-PARASITE INTERACTIONS

*Neospora caninum* (Apicomplexa: Eimeriina: Sarcocystidae) was initially reported as an unidentified protozoan in dogs with encephalomyelitis and myositis (Bjerkas *et al.* 1984). Dubey *et al.* (1988*a, b*) were the first to isolate and name the parasite. Meanwhile, *N. caninum* infection has been reported in various species of livestock, including sheep, goats, horses and deer (reviewed by Dubey and Lindsay, 1996; Hemphill, 1999; Dubey, 2003). However, most importantly, the current evidence strongly suggests that infection with *N. caninum* represents a major cause of reproductive failure in cattle worldwide (Hemphill and Gottstein, 2000; Dubey *et al.* 2002; Gondim *et al.* 2005; Innes *et al.* 2005). The exact phylogenetical relationship of *N. caninum* to other members of the Apicomplexa has been, and still is, under controversy (Tenter and Johnson, 1997; Dubey *et al.* 2002; Heydorn and Mehlhorn, 2002). McAllister *et al.* (1998) were the first to show that the dog is a definitive host for

*N. caninum*, and this was later confirmed by Lindsay *et al.* (1999*b*). Gondim *et al.* (2004) demonstrated that coyotes also shed oocysts, indicating that other final hosts cannot be ruled out.

There are different routes of transmission. One relevant route, but probably not the most important one, is through the oral uptake of sporozoite-containing oocysts (Trees *et al.* 2002; Gondim *et al.* 2005). Infection of an immunocompetent host with oocysts will not cause clinical disease, but liberated sporozoites, by analogy with other coccidians such as *Toxoplasma gondii*, are thought to infect the intestinal tissue, cross the epithelium, reach blood and lymphatic vessels, and infect other nucleated cells, including macrophages and lymphocytes. In the initial phases of the infection, parasites disseminate throughout the body, and transform to the rapidly proliferating tachyzoite stage. Subsequently, the stress caused by the host immune response is believed to be one of the factors that trigger stage conversion to the slowly proliferating *N. caninum* bradyzoites (reviewed by Buxton *et al.* 2002; Lyons *et al.* 2002). Bradyzoites represent a quiescent stage of the parasite, which forms intracellular tissue cysts, surrounded by a cyst wall that protects the parasite from immunological and physiological

\* Corresponding author. Tel: +41 31 6312384. Fax: +41 31 6312477. E-mail: andrew.hemphill@ipa.unibe.ch

reactions on part of the host. *N. caninum* tissue cysts have been identified almost exclusively within the central nervous system, with few exceptions (Peters *et al.* 2001). Bradyzoites can survive within a latently infected, but immunocompetent, animal for many years without causing any clinical signs. However, during pregnancy bradyzoites can get reactivated, and the partial immune-incompetence of a pregnant dam, namely the decrease of an efficient Th-1 response and shift to Th-2, leads to limited suppression of the cell-mediated immunity which normally keeps tachyzoite proliferation in check (Innes *et al.* 2000, 2002, 2005; Quinn *et al.* 2002). This can contribute to the reactivation of bradyzoites and re-conversion to tachyzoites, and to infection of the placenta and possibly the fetus with *N. caninum* tachyzoites. Transmission of *N. caninum* from mother to fetus *in utero* is highly efficient, and acute fetal neosporosis is a major cause for abortion, stillbirth, or at least clinical or subclinical disease in newborn calves (Williams *et al.* 2000; Trees *et al.* 2002; Gondim *et al.* 2005). Vertical transmission in cattle can occur over successive pregnancies, and in cases where congenitally infected but clinically healthy heifers are used for breeding, parasite spreading within a herd can take place very efficiently (Björkman *et al.* 1996). In the USA and the EU, neosporosis is reported as the leading cause of abortions in cattle (Hemphill and Gottstein, 2000; Dubey *et al.* 2002).

Since its discovery, the realization of the economical significance of this parasite has led to increased efforts in elucidating how *N. caninum* interacts with its host. These interactions occur on two levels. On the cellular level, *N. caninum* can only survive, proliferate and proceed during most stages of its life-cycle as an intracellular parasite. Thus the processes which lead to host cell invasion and intracellular development are of crucial importance (Hemphill, 1999; Hemphill *et al.* 2004). On the other hand, it is the complex relationship with the host immune system, which decides the fate of this parasite once it enters the host organism. In this review we will focus on the use of cell culture approaches and laboratory animals to elucidate how *N. caninum* interacts with its host on the cellular and immunological level.

#### NEOSPORA CANINUM AND ITS INTERACTIONS WITH THE HOST CELL

*N. caninum*, similar to *T. gondii*, is capable of actively invading a large variety of target cells. This process has been initially investigated *in vitro* using bovine aorta endothelial cell monolayers (Hemphill *et al.* 1996), and later for many other cell types (reviewed by Hemphill *et al.* 2004). While human cells or cell lines such as human foreskin fibroblasts, Hela cells and Caco2 (colon cancer) cells are readily

infected *in vitro*, there is no conclusive evidence that *N. caninum* actually infects humans (Peterson *et al.* 1999; Tranas *et al.* 1999; Graham *et al.* 1999; Dubey, 2003; Omata *et al.* 2005).

#### Surface constituents of *N. caninum* tachyzoites

The first step in the physical relationship between the parasite and the host cell is the establishment of a low-affinity contact between tachyzoite and host cell surface membrane, followed by the actual adhesion process, namely a more stable association between tachyzoites and the host cell surface (Fig. 1). In order to initiate host cell invasion, tachyzoites re-orientate themselves perpendicularly to the host cell surface membrane, and enter the host cell cytoplasm, by advancing anterior end first, until they are located in the cytoplasm, enclosed by a parasitophorous vacuole (PV) (Fig. 2A, D). Invasion is an active process requiring metabolic energy solely on the part of the parasite, but not on the part of the host cell. This is highlighted by earlier findings that *N. caninum* can even infect formaldehyde-fixed host cells (Hemphill *et al.* 1996). However, by far not all *N. caninum* tachyzoites adhering to the host cell surface will actually achieve host cell entry (Naguleswaran *et al.* 2003). Thus, specific signals and/or receptor-ligand interactions are required that enable tachyzoites to exploit their invasive capacity, and it will be a challenge in the future to elucidate those processes involved. Recently, comparative studies on host cell invasion by *T. gondii* and *N. caninum* using dog fibroblasts, cat kidney cells and Vero cells showed that *T. gondii* invaded all cell types with greater efficiency (Lei *et al.* 2005a).

On the molecular level, the initial low-affinity host-parasite contact is mediated, at least in part, through the 2 major immunodominant surface antigens of *N. caninum* tachyzoites, NcSAG1 and NcSRS2, which are both inserted into the plasma membrane by a GPI-anchor (Sonda *et al.* 1998; Howe *et al.* 1998; Schares *et al.* 2000). Polyclonal and monoclonal antibodies directed against these 2 surface antigens inhibit host cell adhesion and invasion (Hemphill, 1996; Nishikawa *et al.* 2000; Haldorson *et al.* 2005). Monoclonal antibodies directed against a 73 kDa *Neospora* surface antigen have also been shown to inhibit host cell invasion (Uchida *et al.* 2004). Specific *in vivo*-metabolic labelling of GPI-anchored proteins in *N. caninum* tachyzoites using radioactive <sup>3</sup>H-ethanolamine identified at least 7 protein bands carrying a GPI anchor, the most prominent of which largely co-migrated with NcSRS2 and NcSAG1, and the others were of 80 and 100 kDa (Fuchs *et al.* 1999). Thus there remain a number of surface antigens to be investigated. A possible approach to this has been undertaken by Lei *et al.* (2005b), who isolated pellicle and plasmalemma fractions of *N. caninum*




Fig. 1. Scanning electron microscopy of *Neospora caninum* tachyzoites in the process of invading Vero host cells. (A) Parasites are establishing contact with their host cells, and once dedicated to invade their host cells, they rearrange themselves perpendicularly to the host cell surface membrane (B), and move into the host cell with the apical end first (C).

tachyzoites using biochemical subcellular fractionation. They found that pellicle fractions contained several major proteins, and analyses of plasmalemma of *N. caninum* revealed the presence of 2 abundant proteins in addition to other lower abundance antigens, some of which were detectable by monoclonal antibodies (Lei *et al.* 2005b).

The surface of *N. caninum* tachyzoites exhibits considerable differences from *T. gondii* with regard to surface carbohydrate content. Baszler *et al.* (1996) were the first to demonstrate that the surface of *N. caninum* is glycosylated, and they showed that a monoclonal antibody directed against a periodate-sensitive epitope on a 65 kDa surface protein is useful for serological diagnosis of *N. caninum* infection by competitive ELISA (Baszler *et al.* 2001). Subsequently, Hemphill *et al.* (1997) reported that NcSAG1-epitopes recognized by polyclonal anti-SAG1 antibodies could be glycosylated. Fuchs *et al.* (1999) then demonstrated the presence of surface

carbohydrates on *N. caninum* tachyzoites by ruthenium red labelling, and the absence of glycans on the *T. gondii* surface. The same group showed that glycoproteins largely co-migrated with GPI-anchored proteins following SDS-PAGE separation. In addition, they used a panel of lectins for the identification of surface-associated glycoproteins in *N. caninum* tachyzoites, and showed that ConA stained the surface of *N. caninum*, but not of *T. gondii* tachyzoites. ConA binding sites were found to be localized to the surface and the dense granules in *N. caninum* tachyzoites. Therefore, an important feature that defines the differences between *N. caninum* and *T. gondii* could be post-translational modifications. The presence of carbohydrate modifications on antigenic proteins could potentially have implications regarding host-parasite interactions, especially with regard to masking of epitopes, and the immunological consequences thereof. Glycosylation could thus serve as a possible

explanation for the lack of extensive cross-antigenicity between the two closely related species.

#### *Secretory proteins and proteases*

Once a low-affinity contact between *N. caninum* and its host cell is achieved, *N. caninum* tachyzoites discharge secretory organelles named micronemes, rhoptries and dense granules, either prior, during or following host cell invasion. Using *T. gondii* as a model, there have been extensive advances during the last few years on the molecular composition of these secretory organelles and how these secretion processes are accomplished (for reviews refer to Soldati *et al.* 2001; Tomley and Soldati, 2001; Carruthers, 2002; Mercier *et al.* 2005). In this context, proteases have been shown to be critical for assembly and trafficking of microneme and rhoptry proteins, and parasite proteases are being considered as potential targets for chemotherapeutic intervention (Kim, 2004). Invasion of host cells by *T. gondii* takes place through the formation of a moving junction, which selectively excludes host cell plasma transmembrane proteins on the basis of their membrane anchoring (Mordue *et al.* 1999), and this mechanism most likely applies to all members of the apicomplexans. In addition, it was shown that *T. gondii* tachyzoite penetration of the host cell surface membrane is dependent on, and powered by, a parasite actin/myosin system, as determined by the use of specific inhibitors and parasite mutants (reviewed by Keeley and Soldati, 2004). More recently, it was shown that invasion itself is accompanied by proteolytic cleavage and shedding of secreted proteins during host cell invasion (Kim, 2004; Binder and Kim, 2004; Carruthers and Blackmann, 2005). Among the proteases involved in these processes, subtilisin-like serine proteases have essential roles in processing of secretory components. Other studies have demonstrated the role of cysteine proteases and shown that rhomboid proteases, a newly described class of serine proteases, are also important (Dowse and Soldati, 2005). It is conceivable that similar mechanisms would account for *N. caninum* invasive stages, although the respective key players have not been identified nor studied so far.

However, there is some experimental evidence that *N. caninum* and *T. gondii* differ with regard to their susceptibility to protease inhibitors (Naguleswaran *et al.* 2003). Comparative quantification of host cell invasion events has shown that *T. gondii* invasion of host cells is impaired by the serine protease inhibitor PMSF, while inhibitors affecting other protease classes (e.g. phenanthroline, E64, pepstatin) affect adhesion, but not invasion. In contrast, inhibition of serine-, metallo- and cysteine-proteases did not affect *N. caninum* adhesion nor invasion, but pepstatin, an inhibitor of aspartyl proteases, did

have a profound impact on *N. caninum* invasion (Naguleswaran *et al.* 2003). The molecular nature of the aspartyl protease activity has not been elucidated to date. The serine protease NcSUB1, formerly known as NC-p65 (Louie and Conrad, 1999) is the first proteolytic enzyme of *N. caninum* that was described at the molecular and functional level. Antibodies generated against an internal fragment of NcSUB1 (amino acids 649–783) labelled primarily the microneme organelles of the parasite. Analysis of secreted parasitic proteins indicated that a protein of 65 kDa (reduced) or 55 kDa (non-reduced) was recognized by the antibody, and the same secreted proteins were shown to contain major proteolytic activity by zymography (Louie *et al.* 2002). More recently, we have identified a metalloprotease activity present in a fetuin-binding fraction of *N. caninum* tachyzoite extracts, and antibody inhibition studies showed that this metalloprotease activity could also be implicated in host cell invasion (Vonlaufen *et al.*, manuscript submitted for publication). The importance and exact functional relevance of this fetuin-binding fraction needs to be further investigated.

#### *Micronemal proteins and host cell surface receptors*

The first organelles to be secreted by *N. caninum* tachyzoites at the onset of adhesion are the micronemes. Microneme proteins include potentially adhesive soluble components such as NcMIC1 (Keller *et al.* 2002), NcMIC2 (Lovett *et al.* 2000), NcMIC4 (Keller *et al.* 2004), and membrane-bound microneme proteins such as NcMIC3 (Sonda *et al.* 2000; Naguleswaran *et al.* 2001). While proteolytic processing of the soluble microneme proteins has been shown to occur, there is no indication so far that the membrane-bound NcMIC3 is undergoing modification. Secretion of microneme contents is initiated *in vitro* by simply incubating *N. caninum* tachyzoites in medium at 37 °C. This indicates that microneme secretory processes take place as soon as parasites egress from the host cells (Naguleswaran *et al.* 2001; Keller *et al.* 2002, 2004). Most likely, *Neospora* microneme proteins, by analogy with *T. gondii*, are deployed and function as protein complexes (Carruthers, 2002; Opitz and Soldati, 2002; Dowse and Soldati, 2004). Several *N. caninum* microneme proteins identified to date possess adhesive domains that could interact with receptors on the surface of target cells, similar to related domains found in vertebrate extracellular matrix proteins. These adhesive motifs include thrombospondin-(TSP-) like domains in NcMIC1 (Keller *et al.* 2002), integrin- and TSP-type I-like domains in NcMIC2 (Lovett *et al.* 2000), epidermal growth factor (EGF)-like domains in NcMIC3 (Sonda *et al.* 2000), and apple-domains in NcMIC4 (Keller *et al.* 2004).



Both, *N. caninum* and *T. gondii* have been shown to bind to their host cell surface via binding to sulphated host cell surface glycosaminoglycans (GAGs). However, despite the obvious similarities between the two species, there are distinct differences with regard to the actual host cell surface receptors. While *N. caninum* tachyzoites preferentially bind to chondroitin sulphate GAGs, it was shown that *T. gondii* tachyzoites preferentially interact with heparan sulfate residues (Naguleswaran *et al.* 2002). One of the microneme proteins suggested to be directly mediating the contact between *N. caninum* and the host cell surface is NcMIC3. NcMIC3 is secreted by tachyzoites at the apical tip, and remains bound to the tachyzoite surface for extended periods of time, with adhesive EGF-like domains exposed outwards (Naguleswaran *et al.* 2001). These NcMIC3-EGF-like domains, expressed in *E. coli* as a poly-his-recombinant protein, bind to host cell surface chondroitin sulfates (Naguleswaran *et al.* 2002). In contrast, neither recombinant NcSAG1 nor NcSRS2 bind to the host cell surface GAGs. Subsequent investigations showed that while chondroitin sulfate residues act as adhesion receptors, they do not mediate invasion of host cells (Naguleswaran *et al.* 2003). Thus, adhesion and invasion are distinct events.

Another protein recently demonstrated to be involved in parasite-host cell interaction is protein-disulfide isomerase (NcPDI; Naguleswaran *et al.* 2005). PDIs are generally involved in reduction, oxidation and isomerization of intra- and inter-molecular thiol-groups, and are thought to be responsible for maintaining the correct three-dimensional conformation of cysteine-rich proteins by modulating the formation of disulfide bridges. Our investigations showed that NcPDI is mainly found in the ER and in small vesiculated apical organelles resembling micronemes, but a fraction of NcPDI is also located on the surface of *N. caninum* tachyzoites (Naguleswaran *et al.* 2005). Pre-incubation of *N. caninum* tachyzoites with a panel of thioredoxin inhibitors, including the cell-impermeant PDI inhibitor bacitracin, have a negative impact on *N. caninum*, but not on *T. gondii*, tachyzoite host cell adhesion, despite the close PDI sequence similarity. The *Neospora* surface is largely composed of cysteine-rich proteins that are either constitutively or transiently expressed on the parasite surface, and the function of these proteins is highly dependent on their conformation. Therefore NcPDI could be an important factor mediating host cell interaction in *N. caninum* infection.

#### *Tachyzoite-bradyzoite stage conversion*

In the immunocompetent host such as cattle, *N. caninum* bradyzoites are found as a slowly proliferating and tissue cyst-forming stage. Bradyzoites

are orally infective. In addition, as immuno-competence gets impaired (such as during pregnancy), bradyzoites are reactivated, which is reflected by a raise of maternal antibodies in cattle (Guy *et al.* 2001). Parasites will resume proliferation and thus dissemination, and are then able to infect the placental tissue and possibly the unborn fetus, causing abortion, stillbirth or birth of weak offspring (Guy *et al.* 2001). Thus, *N. caninum* tissue cysts are in fact largely responsible for both horizontal (oral) transmission to the carnivorous final host, and vertical (transplacental) transmission to the fetus due to reactivation of quiescent bradyzoites, and are thus of prime epidemiological importance. Hence, an efficient chemotherapeutical treatment or any other means of intervention should target both tachyzoite and bradyzoite stages.

For *T. gondii* bradyzoite *in vitro* culture, several protocols had previously been developed, including modulation of culture conditions by altering the pH, increasing the temperature, applying chemical stress (Soete *et al.* 1994), or mitochondrial inhibitors (Bohne *et al.* 1994). In murine macrophages, interferon- $\gamma$  and SNP were shown to induce stage conversion of *T. gondii* by a mechanism related to nitric oxide (NO) release (Bohne *et al.* 1994). It was also shown that increased cyclic nucleotide levels in host or parasite seem to be linked to stage conversion (Kirkman *et al.* 2001).

*N. caninum* tissue cysts have been more difficult to obtain using *in vitro* culture. Protocols developed for *T. gondii*, have yielded only few parasites undergoing stage conversion, showing that the efficiency of the differentiation process *in vitro* is rather low compared to *Toxoplasma* (Weiss *et al.* 1999; Vonlaufen *et al.* 2002). Tunev *et al.* (2002) have further improved the *in vitro* procedure by using bovine macrophages and increasing the time of culture to 9 days, thus achieving a tachyzoite-to-bradyzoite conversion rate of 14% based on the immunofluorescent detection of BAG1 expression.

Vonlaufen *et al.* (2002) established an alternative model, which is based on the use of *N. caninum* Liverpool isolate and sodium nitroprusside (SNP)-treated murine epidermal keratinocytes as host cells (Fig. 2E, F). SNP releases nitric oxide (NO) that reacts with the iron sulphur centres of several proteins involved in electron transport of the respiratory chain and heme iron of cytochrome C oxidase. This results in a decrease of ATP formation and in a diminished binding of oxygen to cytochrome C oxidase (Cooper, 1999). The adaptation of the parasites to a decreased energy production and to an anaerobic environment may trigger the differentiation process, and thus results in the formation of slowly dividing bradyzoites, which consume less energy. More recently, *N. caninum* tachyzoite-to-bradyzoite *in vitro* cultivation has been adapted for the use of Vero host cells, a culture system that allows the separation of




Fig. 2. Transmission electron microscopy of the intracellular development of *Neospora caninum*. (A) Shortly following host cell invasion, tachyzoites are located in the cytoplasm, completely surrounded by a parasitophorous vacuole membrane. (B) This membrane becomes more evident once parasites release secretory products into the lumen of the vacuole (arrows). (C, D) As tachyzoites, *Neospora* proliferates rapidly by endodyogeny and forms large pseudocysts that are surrounded by a parasitophorous vacuole membrane only (arrowheads). During tachyzoite-to-bradyzoite *in vitro* stage conversion (E, F), proliferation is largely inhibited, and parasites form electron-dense deposits at the periphery of the vacuole, which are indicative of cyst wall formation. Co, conoid; mito, mitochondria; nu, nucleus; rho, rhoptries; mic, micronemes; dg, dense granules.

bradyzoites from host cells (Vonlaufen *et al.* 2004). Vero host cell invasion assays demonstrated significant differences between tachyzoites and bradyzoites with regard to invasive capacity and requirements of host cell surface receptor molecules. For instance, bradyzoites were shown to invade Vero cells much less efficiently than tachyzoites, and removal of sialic acid residues from the Vero cell surface impaired bradyzoite adhesion, but not tachyzoite host cell binding. In addition, a number of dense granule

proteins such as NcGRA1, NcGRA2 and NcGRA7 are secreted during *in vitro* stage conversion and are incorporated into the cyst wall (Vonlaufen *et al.* 2004). A similar procedure for obtaining bradyzoites in an efficient way in cell culture was established by Risco-Castillo *et al.* (2004), by using MARC145 monkey kidney epithelial cells as host cells. The same group recently reported on the identification and cloning of the NcSAG4 gene, an orthologue to the *T. gondii* TgSAG4 gene, which is

the first reported gene to be expressed specifically during the *N. caninum* bradyzoite stage (Fernandez-Garcia *et al.* 2005).

In *T. gondii*, *in vitro* stage conversion is readily achieved when using avirulent, or tissue cyst-forming strains (i.e. ME-49), while virulent strains such as RH hardly form tissue cysts nor express bradyzoite antigens under stress conditions. The situation appears different with *N. caninum*, where the more virulent Nc-Liverpool isolate (Barber *et al.* 1995) is useful for inducing tissue cyst formation *in vitro*, while the two less virulent isolates Nc-1 and NcSweB1 (Stenlund *et al.* 1997; Atkinson *et al.* 1999) were hardly able to produce bradyzoite stage parasites *in vitro* (Vonlaufen *et al.* 2002).

#### Host cell modulation and parasite-host cell cross-talk

Once inside the host cell, *N. caninum* resides within a parasitophorous vacuole (PV), surrounded by a parasitophorous vacuole membrane (PVM), which is essentially derived from the host cell surface membrane (Fig. 2A–D). There are many similarities with *T. gondii*-infected cells, where the PV resists acidification and phagolysosomal maturation (Mordue and Sibley, 1997; Mordue *et al.* 1999). Following invasion, the lumen of the vacuoles of both parasites, as well as its membrane, is extensively modified through secretory products, most likely originating from rhoptries and dense granules (Cesbron-Delauw, 1996; Hemphill *et al.* 1998; Vonlaufen *et al.*, 2004).

Host cell modulation by *N. caninum* has not yet been investigated extensively. It was demonstrated that infection of IFN- $\gamma$ -treated BABL/3T3 clone A31 fibroblasts with *N. caninum* tachyzoites *in vitro* causes apoptosis (Nishikawa *et al.* 2001*c*). Apoptosis of *N. caninum*-infected and IFN- $\gamma$ -treated cells was shown to be associated with increased DNA fragmentation, and increased caspase-3 and caspase-8 activity, and the administration of respective inhibitors inhibited cell death. The reduction in cell viability was prevented with the addition of anti-mouse FasL monoclonal antibody (Nishikawa *et al.* 2002). However, this aspect merits closer investigation. For sure, inhibition of apoptosis is a critical issue for the chronic phase of infection, during which *N. caninum* bradyzoites proliferate only slowly within a host cell, and form tissue cysts that can persist within infected tissue for years. These *N. caninum* bradyzoites, as well as their host cells, are pre-destined for long-term survival. It is conceivable, that host cells are modulated accordingly, by either utilizing the PVM as a signalling platform, or by actively secreting bioactive parasite-derived factors into the cytoplasm.

Host cell modulation has been much more extensively investigated in *T. gondii*-infected cells. Rhoptry proteins, discharged at the early phase of

vacuole formation, are involved in the biogenesis of the PV. *T. gondii* ROP2 has been reported to insert into the vacuole membrane and mediate anchoring to host cell mitochondria (Sinai and Joiner, 2001). Since in *T. gondii*-infected cells, the PVM actively recruits host cell mitochondria and ER (Speer *et al.* 1999), protocols have been developed for the enrichment of this organellar membrane complex (Sinai *et al.* 1997). There is accumulating evidence that the PVM, its constituents, as well as secretory parasite molecules passing through this membrane into the host cell cytoplasm, and are involved in cross-talk and manipulation of host cell functions. For instance, in *T. gondii*-infected macrophages, LPS-induced cytokine signalling is blocked by the presence of tachyzoites (Denkers *et al.* 2004). It has been shown that external addition of parasite lysates and secretion products do not mediate this suppression. Furthermore, killing of intracellular parasites by drug treatment relieved the blockage of LPS-induced secretion of TNF- $\alpha$  (Butcher and Denkers, 2002). Luder *et al.* (2001) showed that *T. gondii* down-regulates MHC class II gene expression and antigen presentation by murine macrophages via interference with nuclear translocation of STAT1 $\alpha$ . The same group has also demonstrated that *T. gondii* infection of macrophages results in reduced expression of inducible nitric oxide synthase and facilitates replication in activated macrophages (Luder *et al.* 2003*a*), and that infection in neural antigen-presenting cells inhibits MHC class II expression by down-regulating the class II transactivator CIITA (Luder *et al.* 2003*b*).

Other investigators found that *T. gondii* inhibits host cell apoptosis by inducing the activation of the transcription factor NF $\kappa$ B, which in turn regulates the expression of inhibitors of apoptosis in the host cell (reviewed by Sinai *et al.* 2004). Activation of the NF $\kappa$ B pathway by *T. gondii* correlates with the localization of phosphorylated I $\kappa$ B $\alpha$  at the PVM (Molestina *et al.* 2003) and, more recently, Molestina and Sinai (2005) detected a novel kinase activity at the *T. gondii* PVM capable of phosphorylating host I $\kappa$ B $\alpha$ . In this context, Goebel *et al.* (2001) reported that inhibition of host cell apoptosis by *T. gondii* is accompanied by reduced activation of the caspase cascade and alteration of poly(ADP-ribose) polymerase expression. However, the outcome of host cell modulation by *T. gondii* can differ, depending on the virulence of the infecting strain. Hisaeda *et al.* (1996, 1997) showed that the less-virulent Beverly strain inhibited apoptosis in infected macrophages, while the more virulent RH strain induced apoptosis. The authors suggested that expression of HSP65 in host cells was responsible for this difference. However, the timing of HSP expression appears to be critical. Nevertheless, expression of HSP65 seems to be another common mechanism for preventing apoptosis



during infection of mammalian cells with parasites (reviewed by Heussler *et al.* 2001).

Altogether, these findings indicate that messages to terminate host signalling may be delivered internally, possibly across the PVM. How this is achieved is not clear. There is only little information regarding the composition of the PVM proteome, and information on proteins that are secreted into the host cell cytoplasm across the PVM is sparse. In *T. gondii*, ROP2 and GRA5 are known to span the PVM, partially penetrating into the host cell cytosol (Beckers *et al.* 1994; Lecordier *et al.* 1999). Such PVM-spanning molecules could assist in modulating host cell signalling pathways. In addition, it has been hypothesized, that parasite molecules are specifically directed across the PVM into the cytoplasm, where they could target host-signalling cascades (Denkers *et al.* 2004).

#### *Immunological responses against N. caninum infection in mice*

*N. caninum* infection in mice has been associated with acute primary pneumonia, myositis, encephalitis, ganglioradiculoneuritis, and pancreatitis (Lindsay and Dubey, 1989), with particular mouse strains appearing more susceptible to infection of the CNS than others (Lindsay *et al.* 1995a; Long *et al.* 1998). A higher number of *N. caninum* tachyzoites is necessary for infection when compared to certain virulent strains of *T. gondii* such as RH, which are highly lethal at very small inoculum doses (Howe and Sibley, 1995). Both humoral and cellular immune responses are important to control infection.

The crucial role of the humoral immune response in *N. caninum* infection was demonstrated in experimental infection of  $\mu$ MT-antibody knock-out mice with *N. caninum* tachyzoites, as these were found to be much more susceptible to infection compared to wild-type C57BL/6 mice (Eperon *et al.* 1999). Other types of immunological deficiencies had been addressed to investigate the importance of specific components of the immune system to control the infection. Nude mice are very susceptible to *N. caninum* and develop acute neosporosis (Yamaga *et al.* 1996; Shibahara *et al.* 1999). Interferon- $\gamma$  knock-out mice developed acute and lethal neosporosis within 2 weeks, and IL-12 knock-out mice had exhibited a high susceptibility to acute neosporosis (Baszler *et al.* 1999; Ritter *et al.* 2002).

Khan *et al.* (1997) investigated the cellular immune response to *N. caninum* in inbred A/J mice during the first 14 days of infection. These mice exhibited no clinical signs of neosporosis and no significant histological evidence upon infection. Splenocytes obtained from infected mice proliferated *in vitro* in response to both *N. caninum* and *T. gondii* soluble antigens, suggesting the presence of

cross-reactive immune determinants. At day 7 after infection of mice with *N. caninum* tachyzoites, a transient (2–3 days) lymphocyte hypo-responsiveness was observed. A similar, although more prolonged, phenomenon had been previously demonstrated during acute *T. gondii* infection in mice with both virulent and avirulent strains (Howe and Sibley, 1995). In *N. caninum* infections, this immunosuppressive effect was restored at day 10, and this remained until day 14 p.i. when the study finished. The hypo-responsiveness to parasite antigen and mitogen was principally due to the induction of nitric oxide. Treatment of spleen cells with nitric oxide synthetase inhibitor partially restored this effect (Kahn *et al.* 1997). In acute murine toxoplasmosis, the cytokine IL-10 had earlier been found to be responsible for the downregulation of lymphocyte proliferation responses (Kahn *et al.* 1995). In *N. caninum* infections this is apparently not the case, since the principal mechanism for murine protection against *N. caninum* is likely to involve IL-12 and IFN- $\gamma$ , and mice treated with corresponding antibodies were rendered more susceptible to infection. Thus, the cellular mechanisms of host protection against *N. caninum* appear to be similar in many, but not all, aspects to the immune response elicited by *T. gondii* (reviewed by Darcy and Santoro, 1994; Sher *et al.* 1995; Gazzinelli *et al.* 1996; Alexander *et al.* 1997).

While the studies described above had focussed on the first 14 days of infection, Long *et al.* (1998) described a comparison of intracerebral parasite load, lesion development and systemic cytokines in different mouse strains (BALB/c, C57BL/6, B.10.D2) infected with *N. caninum* tachyzoites at 6 weeks post-infection. BALB/c and C57BL/6 mice were highly susceptible to the development of *N. caninum*-induced encephalitis, whereas B10.D2 mice were found to be resistant. Resistance in the latter was associated with high IFN- $\gamma$ :IL-4 ratio from antigen-stimulated splenocytes. More susceptible Balb/c and C57BL/6 mice generated a mixed Th1/Th2 type response (Long *et al.* 1998). Modulation of the normally mixed Th1/Th2-type cytokine response in susceptible Balb/c mice towards a Th1-type response by administration of recombinant IL12 resulted in a transient protective effect against *N. caninum* infection (Baszler *et al.* 1999). On the other hand, Nishikawa *et al.* (2001a) demonstrated in vaccination studies in mice employing recombinant NcSRS2-vaccinia virus, that high IgG1 levels are important for clearance of the parasite at the early stage of infection. Since Th1 cytokines, such as IL-12 and IFN- $\gamma$ , favour the production of IgG2a, and Th2 cytokines such as IL-4 and IL-10 are associated with the production of IgG1, the levels of antibody subclasses reflects the *in vivo* production of cytokines, and thus the type of immune response.



The role of immunity to control, or to permit, parasite recrudescence during pregnancy is crucial for the occurrence of fetal infection. In mice, the Th2 cytokine bias observed in pregnant animals (Athanasakis and Iconomidou, 1996) may favour the activation capacity of *N. caninum*. Thus, Long and Baszler (2000) demonstrated in *N. caninum*-infected mice that IL-4 neutralization before pregnancy – concomitant with the inoculation of an avirulent strain of *N. caninum* – decreased congenital transmission after a challenge during pregnancy. Quinn *et al.* (2004) compared the immune response following *N. caninum* infection in non-pregnant and pregnant mice. Spleen cells from both infected/non-pregnant and infected/pregnant mice produced IFN- $\gamma$ , interleukin-12 and TNF- $\alpha$ ; however, the levels of these Th1 cytokines were lower in infected/pregnant mice. Infected/non-pregnant and infected/pregnant mice also produced the Th2 cytokine interleukin-10, but there was no trend towards a decrease of this in pregnant mice. Interleukin-4 was exclusively produced at high levels by infected/pregnant mice and thus appeared responsible for the observed decline in Th1 cytokine production in pregnant mice. A bias towards Th2 cytokines such as IL-4 and IL-10 is normally associated with the maintenance of a viable pregnancy, and not with the control of protozoal infections (Quinn *et al.* 2004).

Kano *et al.* (2005) showed that mice infected during pregnancy may acquire a weaker immune response to the parasite than mice infected when they are not pregnant, and that mice infected during pregnancy show an enhanced type 2 immune response in the recrudescence of the infection. Rettigner *et al.* (2004) studied the immune response in mice chronically infected with *N. caninum* during successive pregnancies and in mice acutely infected during an ongoing pregnancy. Vertical transmission was demonstrated in chronically infected mice after the first pregnancy but the rate of fetal infection fell after further pregnancies. This was also in agreement with the findings of Cole *et al.* (1995) who observed, in mice inoculated during their first pregnancy, a transmission rate reduction of 25% after a second gestation and an absence of transfer after a third or a fourth gestation. In cattle, a reduction in transmission rates was also observed in successive pregnancies, indicating that immunity can protect against fetal infection (reviewed by Innes *et al.* 2005).

#### VACCINATION IN THE MURINE MODEL

A vaccine against neosporosis might be only successful if both humoral and cell-mediated immune responses are stimulated. Several authors have evaluated potential vaccines against neosporosis by employing experimental infections in

murine models. Some have considered the use of live vaccines, such as the temperature-sensitive strain of *N. caninum* described by Lindsay *et al.* (1999b). Kasper and Kahn (1998) demonstrated that vaccination of mice with intact *N. caninum* tachyzoites protects these mice against a lethal challenge from *T. gondii*, and that this protection is mediated by antigen cross-reactive CD8+ T-cells obtained from spleens of *N. caninum* vaccinated mice. Other authors suggested that vaccination with a killed vaccine, such as a crude *N. caninum* extract, induces protective immunity, both with regard to CNS infection (Lunden *et al.* 2002) and vertical transmission (Liddell *et al.* 1999). The notion that transplacental transmission of the parasite is preventable by vaccination was also demonstrated by Miller *et al.* (2005) who used outbred Qs mice as a model. Mice were immunized prior to pregnancy with live or a crude lysate of *N. caninum* (NC-Nowra isolate), and were then challenged with *N. caninum* (NC-Liverpool). They showed that injection of live NC-Nowra tachyzoites before pregnancy dramatically reduced transplacental transfer from 75 to 0.8% in one experiment and from 76 to 8% in a second experiment, while injection of a crude lysate of NC-Nowra tachyzoites was much less efficient and reduced transplacental transfer from 67 to 53% in one experiment and from 76 to 63% in a second experiment. Analysis of *N. caninum*-specific IgG1 and IgG2a antibody levels prior to pregnancy and challenge showed that NC-Nowra lysate induced a response skewed towards IgG1, whereas live parasites induced both IgG1 and IgG2a antibodies. (Miller *et al.* 2005).

Although a *Neospora* vaccine for cattle based on crude parasite extract is now commercialized (see below), there are efforts to develop a more defined vaccine. The preparation of crude *N. caninum*-antigen extracts depends on fresh parasite supply, and its composition may differ from one batch to another and can, therefore, not be accurately controlled and standardized. Crude parasite antigen extract may also contain factors which are non-protective or even immunosuppressive. In addition, its handling may be critical, mostly due to potential proteolytic activity. Thus, selected and more defined antigens should be considered as potential vaccine candidates. A number of different options have been investigated in the murine model.

Vaccination studies in mice showed that application of NcSRS2 and NcSAG1, expressed in the vaccinia virus system (Nishikawa *et al.* 2001a,b) protected against cerebral and fetal infection. Vaccination of mice with recombinant his-tagged NcSAG1 and NcSRS2 did not induce significant levels of protection against cerebral infection, but applied in combination with the corresponding DNA vaccine, high levels (75%) of protection were

achieved (Cannas *et al.* 2003a). Immunization of mice against neosporosis with recombinant NcSRS2 iscoms was reported to lead to induction of specific antibodies to native NcSRS2 and a significant reduction of cerebral parasite load in immunized mice (Pinitkiatisakul *et al.* 2005). Recently, immunization of mice with native NcSRS2, purified through monoclonal antibody affinity chromatography, also protected mice from transplacental infection, as reported by Haldorson *et al.* (2005). Native SRS2 antigen induced a Th-2 type cellular immune response, which would be compatible with successful pregnancy. These findings suggest that immunodominant surface antigens could be useful for vaccination, and confirm that these surface molecules play important roles during infection *in vivo*.

Other antigens have been investigated for protectivity. Among these, immunization of mice with DNA vaccines based on NcGRA7 and NcHSP33 conferred 54% and 47% protection, respectively, against vertical transmission of *N. caninum* in Balb/c mice (Liddell *et al.* 2003). Subsequently, Jenkins *et al.* (2004) found that inclusion of CpG adjuvant with NcGRA7 DNA vaccine improved the protective effect considerably. Vaccination of C57BL/6 mice with recombinant NcMIC3-poly-his-fusion proteins expressed and purified from *E. coli* resulted in profound (75%) protection of vaccinated animals against cerebral infection (Cannas *et al.* 2003b). Those animals which were infected exhibited a profoundly lower parasite burden compared to non-immunized mice. Serological analysis showed that immunization with recombinant NcMIC3 elicited a predominantly IgG1 antibody response against native NcMIC3. This is indicative for a Th-2 type immune response, which would be compatible with pregnancy (Cannas *et al.* 2003b). The protective potential of NcMIC1, has been assessed more recently, employing recombinant NcMIC1, NcMIC1-DNA vaccination, or combined DNA/recombinant NcMIC1 vaccination (Alaeddine *et al.* 2005). Upon vaccination with NcMIC1 and subsequent challenge infection, all mice were PCR positive in the brain, indicating that the overall protectivity of NcMIC1 was low. However, only those mice vaccinated with bacterially expressed recombinant NcMIC1 did not show any clinical signs of disease, while animals exhibiting clinical neosporosis were found in all other groups. In addition, quantitative PCR revealed that the cerebral parasite burden was significantly lower in the group vaccinated with recombinant NcMIC1 compared to all the other groups (Alaeddine *et al.* 2005). Therefore, recombinant NcMIC1 could be useful not on its own, but in a combination with other antigens, to minimize cerebral infection. Further vaccination studies in mice employing other antigens are ongoing.

#### HOST-PARASITE RELATIONSHIP DURING BOVINE NEOSPOROSIS

Excellent reviews documenting different aspects of the host-parasite relationship during *N. caninum* infection of cattle have recently been published by Innes *et al.* (2005) and Williams and Trees, (2006). In short, *N. caninum* is transmitted to cattle either through post-natal infection or through pre-natal infection. Most authors have presented data that strongly suggest that the latter represents the main route of transmission. Pre-natal infection (congenital or vertical transmission) has been found to be highly efficient, ranging from 81 to 95% in different studies (Paré *et al.* 1996; Schares *et al.* 1998; Davison *et al.* 1999). However, mathematical modelling suggested that, in order to maintain the infection within a herd, a limited degree of post-natal infection (horizontal transmission) must take place (French *et al.* 1999). This can be achieved either by dogs or other still yet unidentified final hosts shedding oocysts. In addition, experimental evidence suggested that calves could be orally infected by giving them colostrum spiked with *N. caninum* tachyzoites (Uggla *et al.* 1998). Abortion caused by *N. caninum* can occur at any time during gestation, but the majority of abortions occur at 5–6 months. Calves can be stillborn, or living calves with significant deformations can be born. In addition, early fetal death and resorption of the fetus can occur. However, the most common event is birth of chronically infected, but clinically asymptomatic calves (reviewed by Innes *et al.* 2005; Williams and Trees, 2006). Endemic abortions in a herd are usually found to be caused by vertical transmission, while abortion epidemics have been associated with point-source infections (Dijkstra *et al.* 2001), or could potentially also be caused by reactivation of latent infection due to factors causing immunosuppression (Wouda *et al.* 1999). Transplacental transmission can occur over consecutive pregnancies and congenitally infected heifers can transmit the parasite to their own offspring. However, there is evidence that cattle develop protective immunity against vertical transmission (Innes *et al.* 2001) and against abortion (Williams *et al.* 2003). Protection against vertical transmission has been achieved by experimental infection of naïve cattle with *N. caninum* tachyzoites prior to pregnancy, and subsequent challenge infection at mid-gestation. In none of the 6 sero-negative calves that were born, parasite DNA could be detected in any of the tissues, except in the spinal cord of 1 of the animals (Innes *et al.* 2001). Protection against abortion was observed in 5 naturally chronically infected cattle. These were experimentally challenged by *N. caninum* infection at 10 weeks of gestation. No abortion occurred, but 3 of these 5 live calves were infected. However, these transplacental infections were not

the result of the experimental infection, but occurred due to the recrudescence of the maternal infection. In a control group, 4 previously uninfected cows were challenged identically, and aborted (Williams *et al.* 2003). In addition, observations in the field showed that during point-source infection chronically infected cattle were less likely to abort, also indicating that protective immunity could occur (McAllister *et al.* 2000). This supports the notion that an effective vaccine against *N. caninum* could be developed.

Studies in non-pregnant cattle have shown, similar to the situation in mice, that following infection, a Th1-type immune response plays a crucial role. The expression of pro-inflammatory cytokines such as IFN $\gamma$  and IL-12 is important in limiting intracellular multiplication of the parasite. Several authors have shown that PBMC from experimentally infected cattle proliferate and produce IFN $\gamma$  when stimulated with crude *N. caninum* antigen extracts (Lunden *et al.* 1998; Andrianarivo *et al.* 1999; Williams *et al.* 2000). During pregnancy, the changes that occur in the immune system allow the dam to accept the fetus.

At this stage, Th2-type cellular immune responses secure the maintenance of the pregnancy and are crucial for regulating the potentially damaging effect of Th-1 responses (reviewed by Quinn *et al.* 2002; Innes *et al.* 2005; Williams and Trees, 2006). It was shown that in cattle PBMC proliferation and IFN $\gamma$  production were downregulated around mid-gestation and, as a consequence, this may mean that cattle are less able to control *N. caninum* infection and dissemination at this time, and thus the parasite is more likely to be transmitted to the fetus.

Another important determinant of transmission, and as a consequence abortion, is the gestational age when infection takes place. Experimental infection has shown that infection early in gestation is fatal for the fetus, whereas infection occurring in mid to late pregnancy may result in the birth of a congenitally infected, but otherwise healthy, calf (Williams *et al.* 2000). This is most likely related to the degree of immuno-competence of the fetus, as investigation on fetal immune responses have shown that at 14 weeks of gestation, lymphocytes only responded to mitogen, while by 24 weeks (mid-gestation), they responded to antigen by proliferating and releasing IFN $\gamma$  (reviewed by Innes *et al.* 2005).

A recent paper described the interaction of *N. caninum* tachyzoites with bovine natural killer (NK) cells (Boysen *et al.* 2006). NK cells represent key players in the early innate immune response, primarily indirectly by producing IFN $\gamma$  in response to cytokines such as IL-12. Live and heat-inactivated *N. caninum* tachyzoites, but not soluble parasite proteins, directly triggered production of IFN $\gamma$  in NK cells activated with IL-2, in a manner

independent of IL-12. Furthermore, *N. caninum*-infected autologous fibroblasts had increased susceptibility to NK cell cytotoxicity compared to uninfected fibroblasts. This cytotoxicity was largely mediated by a perforin-mediated mechanism. *N. caninum* tachyzoites were also able to infect and proliferate within cultured NK cells.

#### VACCINATION STRATEGIES AGAINST BOVINE NEOSPOROSIS

To date, the only commercially available *Neospora* vaccine, Neoguard<sup>TM</sup>, is marketed by Intervet. It is basically composed of a lysate of killed tachyzoites and a havlogen adjuvant. The immunogenicity of a killed whole *N. caninum* extract formulated with havlogen was compared with that of experimentally infected animals, and *N. caninum*-specific proliferation of PBMC was similar to proliferation of PBMC of infected animals (Andrianarivo *et al.* 1999). A field safety study on over 750 cattle demonstrated that the vaccine was safe to use in healthy animals (Choromanski, 2002). A standard field trial was carried out in Costa Rica to assess the effect of this vaccine on the abortion rate under field conditions (Romero *et al.* 2004). The study involved 876 cows, over 2.5 months into pregnancy, belonging to 25 Costa Rican dairy herds. For each vaccinated cow, another cow of the same herd, breed and age category, was selected as control. The treatments included 2 injections 1 month apart, the first dose given between days 75 and 90 of gestation. The treatment reduced the abortion rate from 20.8% in non-vaccinated to 11.2% in the vaccinated group, demonstrating that the killed whole tachyzoite preparation had a reasonable effect on the abortion rate in Costa Rican cattle.

However, as indicated above, somatic cellular extracts have inherent problems, and subunit vaccines are being searched for. Protective immunity is likely to involve both humoral and cell-mediated immune responses. Tuo *et al.* (2004) established *N. caninum* antigen-specific, short-term CD4+ T cell cultures from peripheral blood lymphocytes obtained from infected cattle. They separated *N. caninum* extracts by HPLC, and the assessment of different fractions by CD4 T cell proliferation assays showed that the antigenic properties could be attributed to defined fractions. Subsequently, Tuo *et al.* (2005) reported on a 16 kDa antigen interacting with CD4+ T cells that was also recognized by anti-*N. caninum* antibodies.

The presence of parasite-specific CD4+ and CD8+ cytotoxic T lymphocytes has been shown to be associated with immunity to natural *T. gondii* infection in humans (Purner *et al.* 1996), and it is likely that similar concepts would account for cellular immunity against neosporosis in cattle. Staska *et al.* (2003, 2005) focused on cytotoxic T cells. They



found that cattle experimentally infected with *N. caninum* develop parasite-specific CD4<sup>+</sup> cytotoxic T cells that lyse infected autologous target cells (Staska *et al.* 2003). More recently, they targeted the 2 major *N. caninum* surface antigens, NcSRS2 and NcSAG1, in order to identify specific epitopes inducing bovine cytotoxic T cells and helper T cell responses (Staska *et al.* 2005). This study showed that NcSRS2 induced potent memory CD4 and CD8 T cell activation, as evidenced by IFN $\gamma$  secretion and induction of proliferation, while NcSAG1 did not. They then went on to show that a specific immunodominant region on NcSRS2 spanning amino acids 133–155 was recognized by CD4<sup>+</sup> T cells from 4 experimentally infected cattle with 6 different major histocompatibility complex (MHC) class II haplotypes (Staska *et al.* 2005). This provides evidence that subunit vaccines incorporating NcSRS2 gene sequences or peptides should be further investigated for the development of a subunit vaccine against *N. caninum* in cattle.

#### CONCLUSIONS

Once inside its host cell, *N. caninum* will either undergo proliferation (tachyzoites), or tissue cyst formation (bradyzoites), and the decision on which way to proceed is dependent on the physiological status of the host. In any case, the survival and development of the parasite is ensured by a host cell, which is bound to do what the parasite wants it to do. Elucidating the mechanisms leading to host cell invasion and host cell manipulation is crucially important for the development of possible means of intervention, either by prevention of disease (e.g. by vaccination) or by treatment (e.g. chemotherapy). Studies on *N. caninum* and its interactions with the host have been immensely accelerated by corresponding investigations on *T. gondii*, which represents the best-characterized apicomplexan model organism. However, cell culture based approaches have demonstrated distinct differences between the two species, and will continue to provide important information on the cellular and molecular mechanisms defining the host-parasite relationship during neosporosis. In addition, laboratory *in vitro* and *in vivo* studies have already provided clues on potential targets for prevention and intervention. The main challenge in the future will be to set up a vaccination strategy that will prevent fetal infection by *N. caninum* tachyzoites (by e.g. limiting proliferation and dissemination of tachyzoites), and at the same time this vaccination strategy must result in an immune response that is compatible with pregnancy. Although laboratory mice represent valuable models, and will continue to provide important information on the host-parasite relationship during neosporosis, the true significance of the finding obtained in these models can

only be conclusively assessed in the canine and bovine hosts.

The authors acknowledge the financial support of the Swiss National Science Foundation (grant no. 3100A0-112532/1), the Swiss Secretariat of Education and Research (SBF; grant no. C03.0007) and the Foundation Research 3R. We also thank Bruno Gottstein and Norbert Müller for continuous support and helpful discussions.

#### REFERENCES

- Alaeddine, F., Keller, N., Leepin, A. and Hemphill, A.** (2005). Reduced infection and protection from clinical signs of cerebral neosporosis in C57BL/6 mice vaccinated with the recombinant microneme antigen NcMIC1. *Journal of Parasitology* **91**, 657–665.
- Alexander, J., Scharton-Kersten, T. M., Yap, G., Roberts, C. W., Liew, F. Y. and Sher, A.** (1997). Mechanisms of innate resistance to *Toxoplasma gondii*. *Philosophical Transactions of the Royal Society London* **352**, 1355–1359.
- Andrianarivo, A. G., Choromonaski, L., McDonough, S. P., Packham, A. E. and Conrad, P. A.** (1999). Immunogenicity of a killed whole *Neospora caninum* tachyzoite preparation formulated with different adjuvants. *International Journal for Parasitology* **29**, 1613–1625.
- Athanassakis, I. and Iconomidou, B.** (1996). Cytokine production in the serum and spleen of mice from day 6 to 14 of gestation: cytokines/placenta/spleen/serum. *Developmental Immunology* **4**, 247–255.
- Atkinson, R., Harper, P. A., Ryce, C., Morrison, D. A. and Ellis, J. T.** (1999). Comparison of the biological characteristics of two isolates of *Neospora caninum*. *Parasitology* **118**, 363–370.
- Barber, J. S., Homdahl, O. J., Owen, M. R., Guy, F., Uggla, A. and Trees, A. J.** (1995). Characterization of the first European isolate of *Neospora caninum* (Dubey, Carpenter, Speer, Topper and Uggla). *Parasitology* **111**, 563–568.
- Baszler, T. V., Adams, S., Vander-Schalie, J., Mathison, B. A. and Kostovic, M.** (2001). Validation of a commercially available monoclonal antibody-based competitive-inhibition enzyme-linked immunosorbent assay for detection of serum antibodies to *Neospora caninum* in cattle. *Journal of Clinical Microbiology* **39**, 3851–3857.
- Baszler, T. V., Knowles, D. P., Dubey, J. P., Gay, J. M., Mathison, B. A. and McElwain, T. F.** (1996). Serological diagnosis of bovine neosporosis by *Neospora caninum* monoclonal antibody-based competitive inhibition enzyme-linked immunosorbent assay. *Journal of Clinical Microbiology* **34**, 1423–1428.
- Baszler, T. V., Long, M. T., McElwain, T. F. and Mathison, B. A.** (1999). Interferon-gamma and interleukin-12 mediate protection to acute *Neospora caninum* infection in BALB/c mice. *International Journal for Parasitology* **29**, 1635–1646.
- Beckers, C. J., Dubremetz, J. F., Mercereau-Puijalon, O. and Joiner, K. A.** (1999). The *Toxoplasma gondii* rhoptry protein ROP 2 is inserted into the parasitophorous vacuole membrane, surrounding the intracellular parasite, and is exposed

- to the host cell cytoplasm. *Journal of Cell Biology* **27**, 947–961.
- Binder, E. M. and Kim, K.** (2004). Location, location, location: trafficking and function of secreted proteases of *Toxoplasma* and *Plasmodium*. *Traffic* **5**, 914–924.
- Bjerkås, I., Mohn, S. F. and Presthus, J.** (1984). Unidentified cyst-forming sporozoan causing encephalomyelitis and myositis in dogs. *Zeitschrift für Parasitenkunde* **70**, 271–274.
- Björkman, C., Johansson, O., Stenlund, S., Holmdahl, O. J. M. and Ugglä, A.** (1996). *Neospora* species infection in a herd of dairy cattle. *Journal of the American Veterinary Medical Association* **208**, 1441–1444.
- Bohne, W., Heesemann, J. and Gross, U.** (1994). Reduced replication of *Toxoplasma gondii* is necessary for induction of bradyzoite-specific antigens: a possible role for nitric oxide in triggering stage conversion. *Infection and Immunity* **62**, 1761–1767.
- Boysen, P., Klevar, S., Olsen, I., and Storset, A. K.** (2006). The protozoan *Neospora caninum* directly triggers bovine NK cells to produce gamma interferon and to kill infected fibroblasts. *Infection and Immunity* **74**, 953–960.
- Butcher, B. A. and Denkers, E. Y.** (2002). Mechanism of entry determines the ability of *Toxoplasma gondii* to inhibit macrophage proinflammatory cytokine production. *Infection and Immunity* **70**, 5216–5224.
- Buxton, D., McAllister, M. M. and Dubey, J. P.** (2002). The comparative pathogenesis of neosporosis. *Trends Parasitology* **18**, 546–552.
- Cannas, A., Naguleswaran, A., Müller, N., Gottstein, B., Eperon, S. and Hemphill, A.** (2003a). Vaccination of mice against experimental *N. caninum* infection using NcSAG1- and NcSRS2-based recombinant antigens and DNA-vaccines. *Parasitology* **126**, 303–312.
- Cannas, A., Naguleswaran, A., Müller, N., Gottstein, B. and Hemphill, A.** (2003b). Protective effect of vaccination with recombinant microneme protein NcMIC3 in mice following challenge infection with *Neospora caninum* tachyzoites. *Journal of Parasitology* **89**, 44–50.
- Carruthers, V. B.** (2002). Host cell invasion by the opportunistic pathogen *Toxoplasma gondii*. *Acta Tropica* **81**, 111–122.
- Carruthers, V. B. and Blackman, M. J.** (2005). A new release on life: emerging concepts in proteolysis and parasite invasion. *Molecular Microbiology* **55**, 1617–1630.
- Cesbron-Delauw, M. F., Lecordier, L. and Mercier, C.** (1996). Role of secretory dense granule organelles in the pathogenesis of toxoplasmosis. *Current Topics in Microbiology and Immunology* **219**, 59–65.
- Choromanski, L. J.** (2002). *Neospora caninum* vaccine field safety evaluation. *Intervet Bulletin* 1–3.
- Cole, R. A., Lindsay, D. S., Blagburn, B. L. and Dubey, J. P.** (1995). Vertical transmission of *Neospora caninum* in mice. *Journal of Parasitology* **81**, 730–732.
- Cooper, C. E.** (1999). Nitric oxide and iron proteins. *Biochimica et Biophysica Acta* **1411**, 290–309.
- Darcy, F. and Santoro, F.** (1994). Toxoplasmosis. In *Parasitic Infections and the Immune System* (ed. Kierszenbaum, F.), pp. 163–190. Academic Press, New York.
- Davison, H. C., Otter, A. and Trees, A. J.** (1999). Estimation of vertical and horizontal transmission parameters of *Neospora caninum* infections in dairy cattle. *International Journal for Parasitology* **29**, 1683–1689.
- Denkers, E. Y., Butcher, B. A., Del Rio, L. and Kim, L.** (2004). Manipulation of mitogen-activated protein kinase/nuclear factor-kappaB-signaling cascades during intracellular *Toxoplasma gondii* infection. *Immunology Review* **201**, 191–205.
- Dijkstra, T., Barkema, H. W., Eysker, M. and Wouda, W.** (2001). Evidence of post-natal transmission of *Neospora caninum* in Dutch dairy herds. *International Journal for Parasitology* **31**, 209–215.
- Dowse, T. J., Pascall, J. C., Brown, K. D. and Soldati, D.** (2005). Apicomplexan rhomboids have a potential role in microneme protein cleavage during host cell invasion. *International Journal for Parasitology* **35**, 747–756.
- Dowse, T. J. and Soldati, D.** (2004). Host cell invasion by the apicomplexans: the significance of microneme protein proteolysis. *Current Opinion in Microbiology* **7**, 388–396.
- Dubey, J. P.** (2003). Review of *Neospora caninum* and neosporosis in animals. *The Korean Journal of Parasitology* **41**, 1–16.
- Dubey, J. P., Barr, B. C., Barta, J. R., Bjerkas, I., Björkman, C., Blagburn, B. L., Bowman, D. D., Buxton, D., Ellis, J. T., Gottstein, B., Hemphill, A., Hill, D. E., Howe, D. K., Jenkins, M. C., Kobayashi, Y., Koudela, B., Marsh, A. E., Mattsson, J. G., McAllister, M. M., Modry, D., Omata, Y., Sibley, L. D., Speer, C. A., Trees, A. J., Ugglä, A., Upton, S. J., Williams, D. J. and Lindsay, D. S.** (2002). Redescription of *Neospora caninum* and its differentiation from related coccidia. *International Journal for Parasitology* **32**, 929–946.
- Dubey, J. P., Carpenter, J. L., Speer, C. A., Topper, M. J. and Ugglä, A.** (1988a). Newly recognized fatal protozoan disease of dogs. *Journal of the American Veterinary Medical Association* **192**, 1269–1285.
- Dubey, J. P., Hattel, A. L., Lindsay, D. S. and Topper, M. J.** (1988b). Neonatal *Neospora caninum* infection in dogs: isolation of the causative agent and experimental transmission. *Journal of the American Veterinary Medical Association* **193**, 1259–1263.
- Dubey, J. P. and Lindsay, D. S.** (1996). A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology* **67**, 1–59.
- Eperon, S., Broennimann, K., Hemphill, A. and Gottstein, B.** (1999). Susceptibility of B-cell deficient C57BL/6 ( $\mu$ MT) mice to *Neospora caninum* infection. *Parasite Immunology* **21**, 225–236.
- Fernandez-Garcia, A., Risco-Castillo, V., Zaballos, A., Alvarez-Garcia, G. and Ortega-Mora, L. M.** (2005). Identification and molecular cloning of the *Neospora caninum* SAG4 gene specifically expressed at bradyzoite stage. *Molecular and Biochemical Parasitology* [Epub ahead of print].
- French, N. P., Clancy, D., Davison, H. C. and Trees, A. J.** (1999). Mathematical models of *Neospora*

- caninum* infection in dairy cattle: transmission and options for control. *International Journal for Parasitology* **29**, 1691–1704.
- Fuchs, N., Bütikofer, P., Sonda, S. and Hemphill, A.** (1999). Detection of surface-associated and intracellular glycoconjugates and glycoproteins in *Neospora caninum* tachyzoites. *International Journal for Parasitology* **29**, 1597–1611.
- Gazzinelli, R. T., Amichay, D., Sharton-Kersten, T., Grunwald, E., Farber, J. M. and Sher, A.** (1996). Role of macrophage-derived cytokines in the induction and regulation of cell-mediated immunity to *Toxoplasma gondii*. *Current Topics in Microbiology and Immunology* **219**, 127–139.
- Goebel, S., Gross, U. and Luder, C. G.** (2001). Inhibition of host cell apoptosis by *Toxoplasma gondii* is accompanied by reduced activation of the caspase cascade and alterations of poly(ADP-ribose) polymerase expression. *Journal of Cell Science* **114**, 3495–3505.
- Gondim, M., McAllister, M. M., Anderson-Sprecher, R. C., Björkman, C., Locks, T. F., Firkins, L. D., Gao, L. and Fisxher, W. R.** (2005). Transplacental transmission and abortion in cows administered *Neospora caninum* oocysts. *Journal of Parasitology* **90**, 1394–1400.
- Gondim, M., McAllister, M. M., Pitt, W. C. and Zemlicka, D. E.** (2004). Coyotes (*Canis latrans*) are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **34**, 159–161.
- Graham, D. A., Calvert, V., Whyte, M. and Marks, J.** (1999). Absence of serological evidence for human *Neospora caninum* infection. *Veterinary Record* **144**, 672–673.
- Guy, C. S., Williams, D. J. L., Kelley, D. F., McGarry, J. W., Guy, F., Bjorkman, C., Smith, R. F. and Trees, A. J.** (2001). *Neospora caninum* in persistently infected, pregnant cows: spontaneous transplacental infection is associated with an acute increase in maternal antibody. *Veterinary Record* **149**, 443–449.
- Haldorson, G. J., Mathison, B. A., Wenberg, K., Conrad, P. A., Dubey, J. P., Trees, A. J., Yamane, I. and Baszler, T. V.** (2005). Immunization with native surface protein NcSRS2 induces a Th2 immune response and reduces congenital *Neospora caninum* transmission in mice. *International Journal for Parasitology* **35**, 1407–1415.
- Hemphill, A.** (1996). Subcellular localization and functional characterization of Nc-p43, a major *Neospora caninum* tachyzoite surface protein. *Infection and Immunity* **64**, 4279–4287.
- Hemphill, A.** (1999). The host-parasite relationship in neosporosis. *Advances in Parasitology* **43**, 47–104.
- Hemphill, A., Gajendran, N., Sonda, S., Fuchs, N., Gottstein, B., Hentrich, B. and Jenkins, M.** (1998). Identification and characterization of a dense granule-associated protein in *Neospora caninum* tachyzoites. *International Journal for Parasitology* **28**, 429–438.
- Hemphill, A. and Gottstein, B.** (2000). A European perspective on *Neospora caninum*. *International Journal for Parasitology* **30**, 877–924.
- Hemphill, A., Gottstein, B. and Kaufmann, H.** (1996). Adhesion and invasion of bovine endothelial cells by *Neospora caninum*. *Parasitology* **112**, 183–197.
- Hemphill, A., Fuchs, N., Sonda, S., Gottstein, B. and Hentrich, B.** (1997). Identification and partial characterization of a 36 kD surface protein on *Neospora caninum* tachyzoites. *Parasitology* **115**, 371–380.
- Hemphill, A., Vonlaufen, N., Naguleswaran, A., Keller, N., Riesen, M., Guetg, N., Srinivasan, S. and Alaeddine, F.** (2004). Tissue culture and explant approaches to studying and visualizing *Neospora caninum* and its interactions with the host cell. *Microscopy and Microanalysis* **10**, 602–620.
- Heussler, V., Küenzi, P. and Rottenberg, S.** (2001). Inhibition of apoptosis by intracellular protozoan parasites. *International Journal for Parasitology* **31**, 1166–1176.
- Heydorn, A. O. and Mehlhorn, H.** (2002). A re-evaluation of *Neospora* and *Hammondia* spp. *Trends in Parasitology* **18**, 246.
- Hisaeda, H., Sakai, T., Ishikawa, H., Maekawa, Y., Yasumoto, K., Good, R. A. and Himeno, K.** (1997). Heat shock protein 65 induced by gamma delta T cells prevents apoptosis of macrophages and contributes to host defense in mice infected with *Toxoplasma gondii*. *Journal of Immunology* **159**, 2375–2378.
- Hisaeda, H., Sakai, T., Nagasawa, H., Ishikawa, H., Yasutomo, K., Maekawa, Y. and Himeno, K.** (1996). Contribution of extrathymic interferon gamma delta T cells to the expression of heat shock protein and to protective immunity in mice infected with *Toxoplasma gondii*. *Immunology* **88**, 551–557.
- Howe, D. K. and Sibley, L. D.** (1995). *Toxoplasma gondii* comprises three clonal lineages: correlation of parasite genotype with human disease. *Journal of Infectious Diseases* **172**, 1561–1566.
- Howe, D. K., Crawford, A. C., Lindsay, D. and Sibley, L. D.** (1998). The p29 and p35 immunodominant antigens of *Neospora caninum* tachyzoites are homologous to the family of surface antigens of *Toxoplasma gondii*. *Infection and Immunity* **66**, 5322–5328.
- Innes, E. A., Andrianarivo, A. G., Björkman, C., Williams, D. J. and Conrad, P. A.** (2002). Immune responses to *Neospora caninum* and prospects for vaccination. *Trends in Parasitology* **18**, 497–504.
- Innes, E. A., Buxton, D., Eperon, S. and Gottstein, B.** (2000). Immunology of *Neospora caninum* infection in cattle and mice. *International Journal for Parasitology* **30**, 896–900.
- Innes, E. A., Wright, S. E., Maley, S., Rae, A., Schock, A., Kirvar, E., Bartley, P., Hamilton, C., Carey, I. M. and Buxton, D.** (2001). Protection against vertical transmission in bovine neosporosis. *International Journal for Parasitology* **31**, 1523–1534.
- Innes, E. A., Wright, S., Bartley, P., Maley, S., Macaldowie, C., Esteban Redondo, I. and Buxton, D.** (2005). The host-parasite relationship in bovine neosporosis. *Veterinary Immunology and Immunopathology* **108**, 29–36.
- Jenkins, M., Parker, C., Tuo, W., Vinyard, B. and Dubey, J. P.** (2004). Inclusion of Cpg adjuvant with plasmid DNA coding for NcGRA7 improves protection against congenital neosporosis. *Infection and Immunity* **72**, 1817–1819.
- Kano, R., Masukata, Y., Omata, Y., Kobayashi, Y., Maeda, R. and Saito, A.** (2005). Relationship



- between type 1/type 2 immune responses and occurrence of vertical transmission in Balb/c mice infected with *Neospora caninum*. *Veterinary Parasitology* **129**, 159–164.
- Kasper, L. H. and Kahn, I. A.** (1998). Antigen-specific CD8<sup>+</sup> T-cells protect against lethal toxoplasmosis in mice infected with *Neospora caninum*. *Infection and Immunity* **66**, 1554–1560.
- Keeley, A. and Soldati, D.** (2004). The glideosome: a molecular machine powering motility and host-cell invasion by Apicomplexa. *Trends in Cell Biology* **14**, 528–532.
- Keller, N., Naguleswaran, A., Cannas, A., Vonlaufen, N., Bienz, M., Björkman, C., Bohne, W. and Hemphill, A.** (2002). Identification of a *Neospora caninum* microneme protein (NcMIC1) which interacts with sulfated host cell surface glycosaminoglycans. *Infection and Immunity* **70**, 3187–3198.
- Keller, N., Riesen, M., Naguleswaran, A., Vonlaufen, N., Stettler, R., Leepin, A., Wastling, J. M. and Hemphill, A.** (2004). Identification and characterization of a *Neospora caninum* microneme-associated protein (NcMIC4) that exhibits unique lactose-binding properties. *Infection and Immunity* **72**, 4791–4800.
- Khan, I. A., Schwartzman, J. D., Fonseka, S. and Kasper, L. H.** (1997). *Neospora caninum*: role for immune cytokines in host immunity. *Experimental Parasitology* **85**, 24–34.
- Kim, K.** (2004). Role of proteases in host cell invasion by *Toxoplasma gondii* and other apicomplexa. *Acta Tropica* **91**, 69–81.
- Kirkman, L. A., Weiss, L. M. and Kim, K.** (2001). Cyclic nucleotide signaling in *Toxoplasma gondii* bradyzoite differentiation. *Infection and Immunity* **69**, 148–153.
- Lecordier, L., Mercier, C., Sibley, L. D. and Cesbron-Delauw, M. F.** (1999). Transmembrane insertion of the *Toxoplasma gondii* GRA5 protein occurs after soluble secretion into the host cell. *Molecular Biology of the Cell* **10**, 1277–1287.
- Lei, Y., Davey, M. and Ellis, J. T.** (2005a). Attachment and invasion of *Toxoplasma gondii* and *Neospora caninum* to epithelial and fibroblast cell lines *in vitro*. *Parasitology* **131**, 583–590.
- Lei, Y., Birch, D., Davey, M. and Ellis, J. T.** (2005b). Subcellular fractionation and molecular characterization of the pellicle and plasmalemma of *Neospora caninum*. *Parasitology* **131**, 467–475.
- Liddell, S., Jenkins, M., Collica, C. M. and Dubey, J. P.** (1999). Prevention of vertical transfer of *Neospora caninum* in BALB/c mice by vaccination. *Journal of Parasitology* **85**, 1072–1075.
- Liddell, S., Parker, C., Vinyard, B., Jenkins, M. and Dubey, J. P.** (2003). Immunization of mice with plasmid DNA coding for NcGRA7 and NcHSP33 confers partial protection against vertical transmission of *Neospora caninum*. *Journal of Parasitology* **89**, 496–500.
- Lindsay, D. S. and Dubey, J. P.** (1989). *Neospora caninum* (Protozoa: apicomplexa) infections in mice. *Journal of Parasitology* **75**, 772–779.
- Lindsay, D. S., Dubey, J. P. and Duncan, R. B.** (1999a). Confirmation that the dog is a definitive host for *Neospora caninum*. *Veterinary Parasitology* **82**, 327–333.
- Lindsay, S. D., Lenz, R., Blagburn, B. L. and Brake, D. A.** (1999b). Characterization of temperature-sensitive strains of *Neospora caninum* in mice. *Journal of Parasitology* **85**, 64–67.
- Lindsay, S. D., Lenz, R., Cole, R. A. J., Dubey, J. P. and Blagburn, B. L.** (1995). Mouse model for central nervous system *Neospora caninum* infections. *Journal of Parasitology* **81**, 313–315.
- Long, M. T. and Baszler, T. V.** (1996). Fetal loss in BALB/c mice infected with *Neospora caninum*. *Journal of Parasitology* **82**, 608–611.
- Long, M. T., Baszler, T. V. and Mathison, B. A.** (1998). Comparison of intracerebral parasite load, lesion development, and systemic cytokines in mouse strains infected with *Neospora caninum*. *Journal of Parasitology* **84**, 316–320.
- Long, M. T. and Baszler, T. V.** (2000). Neutralization of maternal IL-4 modulates congenital protozoal transmission: comparison of innate versus acquired immune responses. *Journal of Immunology* **164**, 4768–4774.
- Louie, K. and Conrad, P. A.** (1999). Characterization of a cDNA encoding a subtilisin-like serine protease (NC-p65) of *Neospora caninum*. *Molecular and Biochemical Parasitology* **107**, 211–223.
- Louie, K., Nordhausen, R., Robinson, T. W., Barr, B. C. and Conrad, P. A.** (2002). Characterization of a *Neospora caninum* protease, NcSUB1 (NC-P65), with rabbit anti-N54. *Journal of Parasitology* **88**, 1113–1119.
- Lovett, J. L., Howe, D. K. and Sibley, L. D.** (2000). Molecular characterization of a thrombospondin-related anonymous protein homologue in *Neospora caninum*. *Molecular and Biochemical Parasitology* **107**, 33–43.
- Luder, C. G., Algner, M., Lang, C., Bleicher, N. and Gross, U.** (2003a). Reduced expression of the inducible nitric oxide synthase after infection with *Toxoplasma gondii* facilitates parasite replication in activated murine macrophages. *International Journal for Parasitology* **33**, 833–844.
- Luder, C. G., Lang, C., Giraldo-Velasquez, M., Algner, M., Gerdes, J. and Gross, U.** (2003b). *Toxoplasma gondii* inhibits MHC class II expression in neural antigen-presenting cells by down-regulating the class II transactivator CIITA. *Journal of Neuroimmunology* **134**, 12–24.
- Luder, C. G., Walter, W., Beuerle, B., Maeurer, M. J. and Gross, U.** (2001). *Toxoplasma gondii* down-regulates MHC class II gene expression and antigen presentation by murine macrophages via interference with nuclear translocation of STAT1alpha. *European Journal of Immunology* **31**, 1475–1484.
- Lunden, A., Marks, J., Maley, S. W. and Innes, E. A.** (1998). Cellular immune responses in cattle experimentally infected with *Neospora caninum*. *Parasite Immunology* **20**, 519–526.
- Lunden, A., Wright, S., Allen, J. E. and Buxton, D.** (2002). Immunisation of mice against neosporosis. *International Journal for Parasitology* **32**, 867–876.
- Lyons, R. E., McLeod, R. and Roberts, C. W.** (2002). *Toxoplasma gondii* tachyzoite-bradyzoite interconversion. *Trends in Parasitology* **18**, 198–201.
- McAllister, M. M., Bjoerkman, C., Anderson-Sprecher, R. and Rogers, D. G.** (2000). Evidence

- of point-source exposure to *Neospora caninum* and protective immunity in a herd of beef cows. *Journal of the American Veterinary Medical Association* **217**, 881–887.
- McAllister, M. M., Dubey, J. P., Lindsay, D. S., Jolley, W. R., Wills, R. A. and McGuire, A. M.** (1998). Dogs are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* **28**, 1473–1478.
- Mercier, C., Adjogble, K. D., Daubener, W. and Delauw, M. F.** (2005). Dense granules: are they key organelles to help understand the parasitophorous vacuole of all apicomplexan parasites? *International Journal for Parasitology* **35**, 8298–8249.
- Miller, C. M., Quinn, H. E., Reichel, M. P. and Ellis, J. T.** (2005). Reduction in transplacental transmission of *Neospora caninum* in outbred mice by vaccination. *International Journal for Parasitology* **35**, 821–828.
- Molestina, R. E., Payne, T. M., Coppens, I. and Sinai, A. P.** (2003). Activation of NF-kappaB by *Toxoplasma gondii* correlates with increased expression of antiapoptotic genes and localization of phosphorylated IkappaB to the parasitophorous vacuole membrane. *Journal of Cell Science* **116**, 4359–4371.
- Molestina, R. E. and Sinai, A. P.** (2005). Detection of a novel parasite kinase activity at the *Toxoplasma gondii* parasitophorous vacuole membrane capable of phosphorylating host IkBa. *Cellular Microbiology* **7**, 351–362.
- Mordue, D. G., Desai, N., Dustin, M. and Sibley, L. D.** (1999). Invasion by *Toxoplasma gondii* establishes a moving junction that selectively excludes host cell plasma membrane proteins on the basis of their membrane anchoring. *Journal of Experimental Medicine* **190**, 1783–1792.
- Mordue, D. G. and Sibley, L. D.** (1997). Intracellular fate of vacuoles containing *Toxoplasma gondii* is determined at the time of formation and depends on the mechanism of entry. *Journal of Immunology* **159**, 4452–4459.
- Naguleswaran, A., Alaeddine, F., Guionaud, C., Vonlaufen, N., Sonda, S., Jenoe, P., Mevissen, M. and Hemphill, A.** (2005). *Neospora caninum* protein disulfide isomerase is involved in tachyzoite-host cell interaction. *International Journal for Parasitology* **35**, 1459–1472.
- Naguleswaran, A., Cannas, A., Keller, N., Vonlaufen, N., Björkman, C. and Hemphill, A.** (2002). Vero cell surface proteoglycan interaction with the microneme protein NcMIC3 mediates adhesion of *Neospora caninum* tachyzoites to host cells unlike that in *Toxoplasma gondii*. *International Journal for Parasitology* **32**, 695–704.
- Naguleswaran, A., Cannas, A., Keller, N., Vonlaufen, N., Schares, G., Conraths, F. J., Björkman, C. and Hemphill, A.** (2001). *Neospora caninum* microneme protein NcMIC3: Secretion, subcellular localization and functional involvement in host cell interaction. *Infection and Immunity* **69**, 6483–6494.
- Naguleswaran, A., Müller, N. and Hemphill, A.** (2003). *Neospora caninum* and *Toxoplasma gondii*: a novel adhesion/invasion assay reveals distinct differences in tachyzoite-host cell interactions. *Experimental Parasitology* **104**, 149–158.
- Nishikawa, Y., Inoue, N., Xuan, X., Nagasawa, H., Igarashi, I., Fujisaki, K., Otsuka, H. and Mikami, T.** (2001a). Protective vaccination by recombinant vaccinia virus against *Neospora caninum* infection. *Vaccine* **19**, 1381–1390.
- Nishikawa, Y., Xuan, X., Nagasawa, H., Igarashi, I., Fujisaki, K., Otsuka, H. and Mikami, T.** (2001b). Prevention of vertical transmission of *Neospora caninum* in BALB/c mice by recombinant vaccinia virus carrying NcSRS2 gene. *Vaccine* **19**, 1710–1716.
- Nishikawa, Y., Mishima, M., Nagasawa, H., Igarashi, I., Fujisaki, K., Otsuka, H. and Mikami, T.** (2001c). Interferon gamma induced apoptosis in host cells infected with *Neospora caninum*. *Parasitology* **123**, 25–31.
- Nishikawa, Y., Makala, L., Otsuka, H., Mikami, T. and Nagasawa, H.** (2002). Mechanisms of apoptosis in murine fibroblasts by two intracellular protozoan parasites, *Toxoplasma gondii* and *Neospora caninum*. *Parasite Immunology* **24**, 347–354.
- Nishikawa, Y., Xuan, X., Nagasawa, H., Igarashi, I., Fujisaki, K., Otsuka, H. and Mikami, T.** (2000). Monoclonal antibody inhibition of *Neospora caninum* tachyzoite invasion into host cells. *International Journal for Parasitology* **30**, 51–58.
- Omata, Y., Kano, R., Masukata, Y., Kobayashi, Y., Igarashi, M., Maeda, R. and Saito, R.** (2005). Development of *Neospora caninum* cultured with human serum *in vitro* and *in vivo*. *Journal of Parasitology* **91**, 222–225.
- Opitz, C. and Soldati, D.** (2002). ‘The glideosome’: a dynamic complex powering gliding motion and host cell invasion by *Toxoplasma gondii*. *Molecular Microbiology* **45**, 597–604.
- Paré, J., Thurmond, M. C. and Hietala, S. K.** (1996). Congenital *Neospora caninum* infection in dairy cattle and associated calfhood mortality. *Canadian Journal of Veterinary Research* **60**, 133–139.
- Peters, M., Lütkefals, E., Heckerroth, A. R. and Schares, G.** (2001). Immunohistochemical and ultrastructural evidence for *Neospora caninum* tissue cysts in skeletal muscles of naturally infected dogs and cattle. *International Journal for Parasitology* **31**, 1144–1148.
- Petersen, E., Lebech, M., Jensen, L., Lind, P., Rask, M., Bagger, P., Björkman, C. and Ugglä, A.** (1999). *Neospora caninum* infection and repeated abortions in humans. *Emerging Infectious Diseases* **5**, 278–280.
- Pinitkiatisakul, S., Mattsson, J. G., Wikman, M., Friedman, M., Bengtsson, K. L., Stahl, S. and Lunden, A.** (2005). Immunization of mice against neosporosis with recombinant NcSRS2 iscoms. *Veterinary Parasitology* **129**, 25–34.
- Purner, M. B., Berens, P. B., Nash, P. B., van Linden, A., Ross, E., Kruse, C., Krug, E. C. and Cureil, T. J.** (1996). CD4-mediated and CD8-mediated cytotoxic and proliferative immune responses to *Toxoplasma gondii* in seropositive humans. *Infection and Immunity* **64**, 4330–4338.
- Quinn, H. E., Ellis, J. T. and Smith, N. C.** (2002). *Neospora caninum*: a cause of immune-mediated failure of pregnancy? *Trends in Parasitology* **18**, 391–394.
- Quinn, H. E., Miller, C. M. and Ellis, J. T.** (2004). The cell-mediated immune response to *Neospora*

- caninum* during pregnancy in the mouse is associated with a bias towards production of interleukin-4. *International Journal for Parasitology* **34**, 723–732.
- Rettigner, C., De Meerschman, F., Focant, C., Vanderplasschen, A. and Losson, B.** (2004). The vertical transmission following the reactivation of a *Neospora caninum* chronic infection does not seem to be due to an alteration of the systemic immune response in pregnant CBA/Ca mice. *Parasitology* **128**, 149–160.
- Risco-Castillo, V., Fernandez-Garcia, A. and Ortega-Mora, L. M.** (2004). Comparative analysis of stress agents in a simplified *in vitro* system of *Neospora caninum* bradyzoite production. *Journal of Parasitology* **90**, 466–470.
- Ritter, D. M., Kerlin, R., Sibert, G. and Brake, D.** (2002). Immune factors influencing the course of infection with *Neospora caninum* in the murine host. *Journal of Parasitology* **88**, 271–280.
- Romero, J. J., Perez, E. and Frankena, K.** (2004). Effect of a killed whole *Neospora caninum* tachyzoite vaccine on the crude abortion rate of Costa Rican dairy cows under field conditions. *Veterinary Parasitology* **123**, 149–159.
- Schares, G., Peters, M., Wurm, R., Barwald, A. and Conraths, F. J.** (1998). The efficiency of vertical transmission of *Neospora caninum* in dairy cattle analysed by serological techniques. *Veterinary Parasitology* **80**, 87–98.
- Schares, G., Zinecker, C. F., Schmidt, J., Azzouz, N., Conraths, F. J., Gerold, P. and Schwarz, R. T.** (2000). Structural analysis of free and protein-bound glycosyl-phosphatidylinositols of *Neospora caninum*. *Molecular and Biochemical Parasitology* **105**, 155–161.
- Sher, A., Denkers, E. Y. and Gazzinelli, R. T.** (1995). Induction and regulation of host cell mediated immunity by *Toxoplasma gondii*. *Ciba Foundation Symposium* **195**, 95–104.
- Shibahara, T., Kokuho, T., Eto, M., Haritani, M., Hamaoka, T., Shimura, K., Nakamura, K., Yokomizo, Y. and Yamane, I.** (1999). Pathological and immunological findings of athymic nude and congenic wild type BALB/c mice experimentally infected with *Neospora caninum*. *Veterinary Pathology* **36**, 321–327.
- Sinai, A. P. and Joiner, K.** (2001). The *Toxoplasma gondii* protein ROP2 mediates host organelle association with the parasitophorous vacuole membrane. *Journal of Cell Biology* **154**, 95–108.
- Sinai, A. P., Payne, T. M., Carmen, J. C., Hardi, L., Watson, S. J. and Molestina, R. E.** (2004). Mechanisms underlying the manipulation of host apoptotic pathways by *Toxoplasma gondii*. *International Journal for Parasitology* **34**, 381–391.
- Sinai, A. P., Webster, P. and Joiner, K.** (1997). Association of host cell endoplasmic reticulum and mitochondria with the *Toxoplasma gondii* parasitophorous vacuole membrane: a high affinity interaction. *Journal of Cell Science* **110**, 2117–2128.
- Soete, M., Camus, D. and Dubremetz, J. F.** (1994). Experimental induction of bradyzoite-specific antigen expression and cyst formation by the RH strain of *Toxoplasma gondii* in vitro. *Experimental Parasitology* **78**, 361–370.
- Soldati, D., Dubremetz, J. F. and Lebrun, M.** (2001). Microneme proteins: structural and functional requirements to promote adhesion and invasion by the apicomplexan parasite *Toxoplasma gondii*. *International Journal for Parasitology* **31**, 1293–1302.
- Sonda, S., Fuchs, N., Connolly, B., Fernandez, P., Gottstein, B. and Hemphill, A.** (1998). The major 36 kDa *Neospora caninum* tachyzoite surface protein is closely related to the major *Toxoplasma gondii* surface antigen 1. *Molecular and Biochemical Parasitology* **97**, 97–108.
- Sonda, S., Fuchs, N., Gottstein, B. and Hemphill, A.** (2000). Molecular characterization of a novel microneme antigen in *Neospora caninum*. *Molecular and Biochemical Parasitology* **108**, 39–51.
- Speer, C. A., Dubey, J. P., McAllister, M. M. and Blixt, J. A.** (1999). Comparative ultrastructure of tachyzoites, bradyzoites, and tissue cysts of *Neospora caninum* and *Toxoplasma gondii*. *International Journal for Parasitology* **29**, 1509–1519.
- Staska, L. M., Dvies, C., Brown, W. C., McGuire, T. C., Suarez, C. E., Park, J. Y., Mathison, B. A., Abbott, J. R. and Baszler, T. V.** (2005). Identification of vaccine candidate peptides in the NcSRS2 surface protein of *Neospora caninum* by using CD4+ cytotoxic T lymphocytes and gamma interferon secreting T lymphocytes of infected Holstein cattle. *Infection and Immunity* **73**, 1321–1329.
- Staska, L. M., McGuire, T. C., Davies, C. J., Lewin, H. A. and Baszler, T. V.** (2003). *Neospora caninum* infected cattle develop parasite specific CD4+ cytotoxic T lymphocytes. *Infection and Immunity* **71**, 3272–3279.
- Stenlund, S., Bjorkman, C., Holmdahl, O. J., Kindahl, H. and Ugglä, A.** (1997). Characterization of a Swedish bovine isolate of *Neospora caninum*. *Parasitology Research* **83**, 214–219.
- Tenter, A. M. and Johnson, A. M.** (1997). Phylogeny of the tissue cyst-forming coccidian. *Advances in Parasitology* **39**, 169–139.
- Tomley, F. M. and Soldati, D. S.** (2001). Mix and match modules: structure and function of microneme proteins in apicomplexan parasites. *Trends in Parasitology* **17**, 81–88.
- Tranas, J., Heinzen, R. A., Weiss, L. M. and McAllister, M. M.** (1999). Serological evidence of human infection with the protozoan *Neospora caninum*. *Clinical and Diagnostic Laboratory Immunology* **6**, 765–777.
- Trees, A. J., McAllister, M. M., Guy, C. S., McGarry, J. W., Smith, R. F. and Williams, D. J.** (2002). *Neospora caninum*: oocyst challenge of pregnant cows. *Veterinary Parasitology* **109**, 147–154.
- Tunev, S. S., McAllister, M. M., Anderson-Sprecher, R. C. and Weiss, L. M.** (2002). *Neospora caninum* in vitro: evidence that the destiny of a parasitophorous vacuole depends on the phenotype of the progenitor zoite. *Journal of Parasitology* **88**, 1095–1099.
- Tuo, W., Davis, W. C., Fetterer, R., Jenkins, M., Boyd, P. C., Gasbarre, L. C. and Dubey, J. P.** (2004). Establishment of *Neospora* antigen-specific T cell lines of primarily CD4 T cells. *Parasite Immunology* **26**, 243–246.



- Tuo, W., Fetterer, R. H., Davis, W. C., Jenkins, M. C. and Dubey, J. P.** (2005). *Neospora* antigens defined by antigen-dependent bovine CD4+ T cells. *Journal of Parasitology* **91**, 564–568.
- Uchida, Y., Ike, K., Kurotaki, T., Ito, A. and Imai, S.** (2004). Monoclonal antibodies preventing invasion of *Neospora caninum* tachyzoites into host cells. *Journal of Veterinary Medical Science* **66**, 1355–1358.
- Uggla, A., Stenlund, S., Holmdahl, O. J. M., Jakubek, E. B., Thebo, P., Kindahl, H. and Björkman, C.** (1998). Oral *Neospora caninum* inoculation of neonatal calves. *International Journal for Parasitology* **28**, 1467–1472.
- Vonlaufen, N., Guetg, N., Naguleswaran, A., Müller, N., Bjorkman, C., Schares, G., von Blumroeder, D., Ellis, J. and Hemphill, A.** (2004). *In vitro* induction of *Neospora caninum* bradyzoites in Vero cells reveals differential antigen expression, localization, and host-cell recognition of tachyzoites and bradyzoites. *Infection and Immunity* **72**, 576–583.
- Vonlaufen, N., Müller, N., Keller, N., Naguleswaran, A., Bohne, W., McAllister, M., Björkman, C., Müller, E., Caldelari, R. and Hemphill, A.** (2002). Exogenous nitric oxide triggers *Neospora caninum* tachyzoite-to-bradyzoite stage conversion in murine epidermal keratinocyte cell cultures. *International Journal for Parasitology* **32**, 1253–1265.
- Weiss, L. M., Ma, Y. F., Halonen, S., McAllister, M. M. and Zhang, Y. W.** (1999). The *in vitro* development of *Neospora caninum* bradyzoites. *International Journal for Parasitology* **29**, 1713–1723.
- Williams, D. J., Guy, C. S., Smith, R. F., Guy, F., McGarry, J. W., McKay, J. S. and Trees, A. J.** (2003). First demonstration of protective immunity against foetopathy in cattle with latent *Neospora caninum* infection. *International Journal for Parasitology* **33**, 1059–1065.
- Williams, D. J., Guy, C. S., McGarry, J. W., Guy, F., Tasker, L., Smith, R. F., MacEachern, K., Cripps, P. J., Kelly, D. F. and Trees, A. J.** (2000). *Neospora caninum*-associated abortion in cattle: the time of experimentally-induced parasitaemia during gestation determines foetal survival. *Parasitology* **121**, 347–358.
- Williams, D. J. and Trees, A. J.** (2006). Protecting babies: vaccine strategies to prevent foetopathy in *Neospora*-infected cattle. *Parasite Immunology* **28**, 61–67.
- Wouda, W., Bartels, C. J. M. and Moen, A. R.** (1999). Characteristics of *Neospora caninum* associated abortion storms in dairy herds in the Netherlands (1995–1997). *Theriogenology* **52**, 233–245.
- Yamage, M., Flechtner, O. and Gottstein, B.** (1996). *Neospora caninum*: specific oligonucleotide primers for the detection of brain “cyst” DNA of experimentally infected nude mice by the polymerase chain reaction (PCR). *Journal of Parasitology* **82**, 272–279.