

In vitro efficacy of nitro- and bromo-thiazolyl-salicylamide compounds (thiazolides) against *Besnoitia besnoiti* infection in Vero cells

H. C. E. CORTES^{1*}, N. MUELLER², M. ESPOSITO², A. LEITÃO³, A. NAGULESWARAN² and A. HEMPHILL^{2*}

¹Laboratório de Parasitologia Victor Caeiro, Núcleo da Mitra, ICAM, Universidade de Évora, Apartado 94, 7000-554 Évora, Portugal

²Institute of Parasitology, University of Berne, Länggass-Strasse 122, CH-3012 Berne, Switzerland

³Instituto de Investigação Científica Tropical, CIISA, Faculdade de Medicina Veterinária, Av. da Universidade Técnica, 1300-447 Lisboa, Portugal

(Received 27 October 2006; revised 22 December 2006; accepted 2 January 2007; first published online 19 February 2007)

SUMMARY

Nitazoxanide (NTZ) and its deacetylated metabolite tizoxanide (TIZ) exhibit considerable *in vitro* activity against *Besnoitia besnoiti* tachyzoites grown in Vero cells. Real-time-PCR was used to assess *B. besnoiti* tachyzoite adhesion, invasion, and intracellular proliferation *in vitro*. A number of NTZ-derivatives, including Rm4822 and Rm4803, were generated, in which the thiazole-ring-associated nitro-group was replaced by a bromo-moiety. We here show that replacement of the nitro-group on the thiazole ring with a bromo (as it occurs in Rm4822) does not impair the efficacy of the drug, but methylation of the salicylate ring at the ortho-position in a bromo-derivative (Rm4803) results in complete abrogation of the antiparasitic activity. Treatment of extracellular *B. besnoiti* tachyzoites with NTZ has an inhibitory effect on host cell invasion, while treatments with TIZ, Rm4822 do not. TEM demonstrates that the effects of Rm4822 treatment upon the parasites are similar to the damage induced by NTZ. This includes increased vacuolization of the parasite cytoplasm, and loss of the structural integrity of the parasitophorous vacuole and its membrane. Thus, Rm4822, due to the absence of a potentially mutagenic nitro-group, may represent an important potential addition to the anti-parasitic arsenal for food animal production, especially in cattle.

Key words: thiazolides, *Besnoitia besnoiti*, *in vitro* drug treatment, besnoitiosis.

INTRODUCTION

Besnoitia besnoiti is the causative agent of bovine besnoitiosis, a parasitic disease of cattle widely distributed in subSaharian areas, which has recently received increased attention in Mediterranean countries (Cortes *et al.* 2003, 2005; Juste *et al.* 1990 and see P. J Bourdeau *et al.* (Abstract. IX European Multicolloquium of Parasitology, pp. 459–460, 2004)). Bovine besnoitiosis may have 2 clinical phases. During the initial phase, namely during the first 4 to 12 days of infection, a high parasitaemia occurs, and parasitaemia may be associated with fever and fever-associated signs. Subsequently, in the chronic stage, the parasite shows a high tropism to the skin, leading to dramatic thickening, hardening and folding or wrinkling of the skin, especially around

the neck, shoulders and rump, always accompanied by hyperkeratosis, hyperpigmentation and alopecia (Pols, 1960). Furthermore, abortion may occur during the acute phase (Juste *et al.* 1990) and irreversible aspermy is a common sequel developing during the chronic phase. The above-mentioned, together with the refractoriness to all anti-parasitic compounds tested so far, render bovine besnoitiosis an important re-emerging disease in the South of Portugal (Cortes *et al.* 2003). Despite considerable efforts to identify and develop effective agents to treat bovine besnoitiosis (Shkap *et al.* 1987), none has yet demonstrated any effect for treatment or prevention of disease.

In the United States, the thiazolide nitazoxanide [2-acetyloxy-*N*-(5-nitro-2-thiazolyl)benzamide] (NTZ; AliniaTM) is currently used in humans for the treatment of persistent diarrhoea caused by *Cryptosporidium parvum* and *Giardia intestinalis* in adults, adolescents, and children from 1 year of age. In addition, NTZ exhibits a broad spectrum of activities against a wide variety of enteric bacteria, protozoa, and intestinal and tissue-dwelling helminth parasites, infecting animals and humans (Fox and Saravolatz, 2005; Hemphill *et al.* 2006a;

* Corresponding authors: Laboratório de Parasitologia Victor Caeiro, Núcleo da Mitra, ICAM, Universidade de Évora, Apartado 94, 7000-554 Évora, Portugal. Tel: 00351 266760860. Fax: 00351 266 760912. E-mail: hcec@uevora.pt; Institute of Parasitology, Vetsuisse Faculty, University of Berne, Länggass-Strasse 122, CH-3012 Berne, Switzerland. Tel: +41 31 6312384. Fax: +41 31 6312477. E-mail: hemphill@ipa.unibe.ch




Fig. 1. Structures of the thiazolides investigated in this study.

Pankuch and Appelbaum, 2006). Against *Helicobacter pylori*, *Giardia intestinalis* and *Cryptosporidium parvum*, the drug has been postulated to act via reduction of its nitro-group by nitro-reductases, including pyruvate ferredoxin oxidoreductase (PFOR) (Coombs and Muller, 2002; Sisson *et al.* 2002). In terms of veterinary use, the drug is marketed for the treatment of equine myeloencephalitis caused by *Sarcocystis neurona* (NavigatorTM). However, its application can be related to severe gastrointestinal problems (NavigatorTM prescription information, IDDEX Pharmaceuticals Inc., Greensboro, NC, USA). It has been speculated that this is due to the action of PFOR (pyruvate ferredoxin oxidoreductase or other nitro-reductases, which could reduce the nitro-group and kill anaerobic intestinal bacteria through the production of free radicals (reviewed by Hemphill *et al.* 2006a). Thus, NTZ-derivatives have been synthesized in which the thiazole-associated nitro-group has been replaced by a bromo-group, and the salicylic acid moiety has been modified (Esposito *et al.* 2005, 2007; Muller *et al.* 2006).

Recently, *in vitro* studies demonstrated the efficacies of NTZ and several bromo-derivatives lacking the thiazole-associated nitro-group against *Neospora caninum* tachyzoites (Esposito *et al.* 2005). *Neospora caninum* is an apicomplexan parasite causing abortion and stillbirth in cattle (Dubey, 1999; Hemphill *et al.* 2006b). Since *Neospora*, *Cryptosporidium* and *Sarcocystis* all belong to the phylum Apicomplexa, we investigated the *in vitro* efficacy of NTZ and a limited number of NTZ-derivatives against *B. besnoiti* tachyzoites. For this we employed the Vero cell culture model (Cortes *et al.* 2006). *B. besnoiti* tachyzoite proliferation and adhesion/invasion was assessed by real-time PCR, and TEM was performed to document the drug-induced damage imposed upon the parasites. Our results suggest that thiazolides act against *B. besnoiti* tachyzoites, and compounds lacking the nitro-groups are equally effective,

suggesting that these could be potentially applied in food animals such as cattle.

MATERIALS AND METHODS

Vero cell culture and parasite purification

Besnoitia besnoiti tachyzoites (Bb1Evora03 isolate) were maintained in Vero cell cultures (culture medium: DMEM containing 2% fetal calf serum, 50 IU/ml penicillin, and 50 $\mu\text{g}/\text{ml}$ streptomycin). Parasites were harvested using a cell scraper when the cell monolayer started to be destroyed. Harvested material was repeatedly passaged through a 25-gauge needle and parasites were purified using cellulose CF11 columns.

Infection of Vero cells and in vitro drug treatment assays

Nitazoxanide (NTZ), tizoxanide (TIZ), and the NTZ-derivates Rm4822 and Rm4803 (see Fig. 1), were obtained from Romark Laboratories (Tampa, FL, USA), and were synthesized at the Department of Chemistry, University of Liverpool. Stock solutions were prepared at a concentration of 10 mg/ml in dimethyl sulfoxide (DMSO) and were kept at -20°C . Fresh dilutions to the appropriate concentration were prepared in culture media. DMEM with the same DMSO concentration was used as control. Vero Cell confluent monolayers in 24-well tissue culture plates (Sarstedt, Newton, MA) were inoculated with *B. besnoiti* tachyzoites (5×10^4) and incubated for 2 h at 37°C , 5% CO_2 . Unbound parasites were removed by washing with DMEM, and infected monolayers were maintained in culture medium containing the drugs, as indicated for the individual experiments. In order to assess selective toxicity, uninfected Vero cell monolayers were treated identically. Where indicated, NTZ was added to the Vero cells 24 h before infection. Samples

for monitoring parasite proliferation were taken at different times post-initiation of drug treatment. Each assay in a given experiment was carried out in triplicate.

Processing of DNA samples and light cycler-based quantitative PCR

DNA purification was performed using the DNAeasy kit (QIAGEN, Basel, Switzerland) as previously described for *N. caninum* infected cell cultures (Esposito *et al.* 2007). For quantitative PCR, primers directed to ITS1 region of *B. besnoiti* were used. Results shown correspond to mean values (plus/minus standard deviations) of triplicate determinations.

PDTC-based adhesion/invasion assay

This assay was performed as previously described for *N. caninum* and *Toxoplasma gondii* (Naguleswaran *et al.* 2003). Shortly, Vero cell monolayers were incubated with *B. besnoiti* tachyzoites either in the presence or absence of NTZ, TIZ, Rm4803 or Rm4820. Infected monolayers were incubated for 2 h at 37 °C in DMEM containing 100 µM pyrrolidine dithiocarbamate (PDTC), 0.2 µM CuSO₄ and rabbit anti-*B. besnoiti* antiserum (1 : 200), which killed and permeabilized extracellular tachyzoites. In parallel, wells were incubated in DMEM without any additives. The wells were then washed with DMEM and treated with 1 mg/ml DNase in DMEM for 1 h at 37 °C. Finally, the wells were washed with DMEM containing 1 mM EDTA and the cellular material was taken up in 180 µl of lysis buffer (DNAeasy kit; QIAGEN). The number of intracellular tachyzoites and the overall number of parasites in a given experiment were determined by real-time PCR as described above.

Transmission electron microscopy

Vero cell monolayers were grown in 6-well tissue culture plates, infected with *B. besnoiti* tachyzoites for 2–4 days, and treated with NTZ and Rm4822 (10 µg/ml) as described above. At different time-points, as indicated below, monolayers were washed with 100 mM sodium cacodylate buffer, pH 7.2, and fixed in 100 mM sodium cacodylate buffer containing 2.5% glutaraldehyde. Cells were scraped off with a rubber policeman, post-fixed in 1% OsO₄ (4 h, 4 °C), pre-stained in 1% uranyl acetate (1 h, 4 °C), dehydrated in ethanol, and embedded in Epon 820 resin. The resin was polymerized at 65 °C for 48 h. Ultrathin sections were cut and loaded onto 300-mesh copper grids (Plano GmbH). Staining with uranyl acetate and lead citrate was performed as described by Hemphill and Croft (1997). Finally, grids

were viewed on a Phillips 400 TEM operating at 80 kV.

Statistical analysis

For time-course and PDTC-based adhesion/invasion experiments, the significance of the differences between end-point values of the control and experimental assays was determined by Student's *t*-test, using the Microsoft Excel program. *P* values <0.05 were considered statistically significant.

RESULTS

The nitro-thiazolide compounds NTZ and TIZ inhibit the intracellular proliferation of B. besnoiti tachyzoites

In order to assess whether NTZ inhibits the proliferation of *B. besnoiti* tachyzoites, parasites were allowed to infect Vero cell monolayers for 1 h prior to addition of NTZ at concentrations from 1 to 10 µg/ml (Fig. 2A), and samples were collected at different time-points until day 6. Real-time PCR showed that NTZ readily inhibited proliferation of *Besnoitia* tachyzoites in a dose-dependent manner, with 5 and 10 µg/ml being most efficient (Fig. 1A). Similar results were obtained when NTZ was replaced by its deacetylated metabolite TIZ (data not shown). In parallel, light microscopical inspection of infected and non-infected Vero cell monolayers did not reveal any drug-induced alterations by these drugs, confirming the selective parasite-specific toxicity of NTZ and TIZ as previously reported by Esposito *et al.* (2005) (data not shown). Thus, further experiments were performed at a drug concentration of 10 µg/ml.

The bromo-compounds Rm4822 and Rm4803 exhibit differential effects

As shown in Fig. 1B, Rm4822, a thiazolide compound with a bromo-group substituting for the nitro-group on the thiazole-ring, plus as methylation at position 4 on the salicylic acid ring (Esposito *et al.* 2005), exhibited a proliferation inhibitory potential similar to NTZ. In contrast Rm4803, a bromo-derivative carrying the methylation on position 3, did not have any effect at all, and *Besnoitia* tachyzoite proliferation was comparable to controls where only the solvent was added (Fig. 2B). Thus, thiazolides act against *Besnoitia* using a mechanism which is independent of the nitro-group, and in which a defined position of the salicylic acid moiety plays a crucial role. Rm4822 inhibited *B. besnoiti* proliferation in a dose-dependent manner, with an efficiency similar to NTZ (Fig. 2C). The drug was most active at 5 and 10 µg/ml. In some experiments, Rm4822 treatment was discontinued at day 12 and the




Fig. 2. Effects of thiazolides on the proliferation of *Besnoitia besnoiti* tachyzoites in Vero cells *in vitro* as assessed by quantitative real time PCR. (A) Dose-dependent inhibition of tachyzoite proliferation by the nitrothiazole NTZ. Similar results were obtained for TIZ. (B) Inhibition of proliferation of *B. besnoiti* tachyzoites in the presence of NTZ and the bromo-derivative Rm4822, while Rm4803 did not affect proliferation. (C) Dose dependency of the inhibitory effect mediated by Rm4822. At 12 days of Rm4822 treatment, the drug was removed, and no further growth of parasites was noted until day 28, indicating the parasitocidal effect of Rm4822.




Fig. 3. Analysis of the effects of the nitrothiazoles NTZ and TIZ and the bromoderivatives Rm4822 and Rm4803 on host cell adhesion and invasion. Grey bars indicate values for the overall parasite numbers (adhered and invaded), and the dark bars indicate the numbers of invaded parasites only. Note that in the absence of drugs, around 80% of those parasites interacting with their host cells had actually invaded their host cells. In NTZ-treated cultures, a higher number of parasites adhered to, but only a small fraction of those actually had invaded their host cells. Values for TIZ and Rm4822 remained similar as for the controls. Rm4803 treatment resulted in higher numbers of parasites adhering to their host cells, but Rm4803 did not result in increased invasion.

cultures were maintained until day 28. In any case, no increase in parasite numbers was noted, indicating that *in vitro* treatment with both 5 and 10 µg/ml of Rm4822 was parasitocidal rather than only parasitostatic (Fig. 2C).

Effects of thiazolides on adhesion and invasion of Vero host cells by *B. besnoiti* tachyzoites

In order to study whether the impact of these drugs can be attributed to any interference in the adhesion/invasion process of *B. besnoiti* tachyzoites, freshly isolated parasites were incubated with NTZ, TIZ, Rm4822 and Rm4803 for 2 h prior to letting them interact with Vero cell monolayers. The overall number of parasites (adhered and invaded) interacting with Vero cells was quantified by real-time PCR. In a parallel experiment, the number of intracellular (invaded) tachyzoites was determined by PDTC/antiserum treatment, DNase digestion and real-time PCR, as previously described for *N. caninum* and *T. gondii* by Naguleswaran *et al.* (2003), (see Materials and Methods section). The invasion efficiency (percentage of invaded tachyzoites in relation to the overall parasite number) of control parasites (incubated in DMSO alone) was around 80% (Fig. 3). In NTZ-treated samples, the overall number of tachyzoites interacting with Vero cells was greatly increased by almost 2-fold, but the invasion efficiency reduced to roughly 10%, resulting in greatly decreased numbers of intracellular parasites. This shows that tachyzoites adhered more efficiently, but exhibited impaired invasion abilities (Fig. 3). In contrast, both overall parasite number and invasion rates for TIZ and Rm4822-treated tachyzoites were similar to the controls, indicating that neither

TIZ- nor Rm4822-treatment affected the initial host-parasite interaction. The overall number of tachyzoites interacting with Vero cells was increased by 2-fold in Rm4803-treated preparations, but nevertheless the number of invaded parasites was equal to the untreated control (Fig. 3).

Electron microscopy reveals ultrastructural effects of NTZ and Rm4822 in *B. Besnoiti*-infected HFF and Vero cell monolayers

The effects of NTZ in tachyzoite-infected Vero cell monolayers were studied in comparison to untreated cultures by TEM at different time-points p.i. In control cultures (Fig. 4), *B. besnoiti* tachyzoites were found within their host cells and localized within the parasitophorous vacuole, which was clearly separated from the host cell cytoplasm by the parasitophorous vacuole membrane (PVM, see large arrows in Fig. 4B,C). Tachyzoites exhibited the typical features of apicomplexan parasites such as conoid at the apical tip, micronemes and rhoptries (Fig. 4B,C), but only few dense granules could be seen. Rhoptry contents had an amorphous and rather electron-dense appearance (Fig. 4B,C). Tachyzoites were densely packed within the vacuole, and the vacuole membrane was often in tight contact with the parasite plasma membrane (large arrows in Fig. 4). The space between the tachyzoites was filled with granular material (small arrowheads in Fig. 4B,C).

In NTZ-treated parasites, clear alterations in parasite ultrastructure were visible already after 12 h of treatment. During the first 12 and 24 h following the initiation of drug treatment, the most prominent hallmark was the occurrence of numerous vacuoles




Fig. 4. TEM of Vero cells infected with *Besnoitia besnoiti* tachyzoites. (A) Low magnification view of a parasitophorous vacuole containing numerous tachyzoites. (B, C) Higher magnification views of parasitophorous vacuoles.

Rop = rhoptries, dg = dense granules, mic = micronemes, con = conoid, nuc = nucleus. Large arrows point towards the parasitophorous vacuole membrane, small arrows indicate the granular matrix filling the lumen of the vacuole. Note that parasites are extremely densely packed.

within the parasite cytoplasm (Fig. 5A, B). These vacuoles were often filled with membranous and filamentous material of unknown origin. While the

nucleus as such remained largely intact, other membranous structures within the parasites were dramatically affected (Fig. 5A). Overall, the PVM




Fig. 5. TEM of *Besnoitia besnoiti* in Vero cells treated with 10 $\mu\text{g/ml}$ of NTZ for 12 h (A), 24 h (B) and 48 h (C). Note the high degree of vacuolization of the parasite cytoplasm already at 12 h. Large arrows point towards the parasitophorous vacuole membrane which, in some instances, is already distorted at early time-points (A), small arrows indicate the granular matrix filling the lumen of the vacuole, which is partially distorted already after 12 h (A), but definitely non-existent at 48 h. Instead, the vacuole contains membranous debris of unknown origin. Nuc = nucleus, mito = host cell mitochondria in close vicinity of the parasites.

had started to disintegrate in some cells (large arrows in Fig. 5A), while in others (Fig. 5B) the PVM was partially still discernible. In addition, at 12 h of NTZ treatment, the granular material constituting the matrix of the parasitophorous vacuole was still visible in many instances. In some cases, (Fig. 5A), vacuoles were surrounded by numerous host cell mitochondria. At later time-points such as 24 h post-initiation of treatment, the damage became more dramatic, with even increased vacuolization of tachyzoites, distorted nuclei, and complete loss of the PVM (Fig. 5C). The granular material within the matrix of the parasitophorous vacuole disappeared, and in some cases was replaced with membranous and electron-dense clumps of unknown origin (small

arrow in Fig. 5C). Note that the parasite plasma membranes appeared still intact during this observation period. RM4822-treated cultures fixed at 12 h p.i., intracellular tachyzoites were also found to be extensively vacuolized. A clearly delineated parasitophorous vacuole and corresponding membrane was largely missing already at this time-point. (Fig. 6A, B). Only in very few cases, the granular parasitophorous vacuole matrix, or matrix remnants, were still partially visible (Fig. 6C, small arrows). Non-vacuolized parts of the tachyzoites started to become rather electron dense, and in longitudinal sections as in Fig 6C, membranous structures were distorted, while the nucleus and associated ER appeared unaffected. Rhoptries and micronemes and




Fig. 6. TEM of *Besnoitia besnoiti* in Vero cells treated with 10 $\mu\text{g}/\text{ml}$ of Rm4822 for 12 h (A–C) and 48 h (D, E). (B) Lower magnification view of (A). Note that parasites exhibit a high degree of vacuolization and aberrant membrane organization indicative for considerable metabolic stress. The parasitophorous vacuole membrane (large arrows) is considerably distorted already at 12 h (A–C), and this effect remains also at later time-points (D, E). In a few cases, matrix residues (small arrows) are still visible (small arrows). In (E), note the presence of highly condensed and electron-dense material within the nucleus (white arrow). However, the conoid as such (con) is still clearly discernable.

parasite mitochondrion could not be distinguished any more. At 24 h post-initiation of treatment (Fig. 6D, E), the situation remained the same. In some treated parasites, the nucleus contained electron-dense deposits, indicating that extensive chromatin condensation could have taken place (see Fig. 6E). Note that the parasite plasma membranes, and in many instances the conoid, were still clearly discernible and appeared still intact during the observation period.

DISCUSSION

In this study we demonstrate that the nitro-thiazole analogue NTZ, its metabolite TIZ, and a bromo-derivative, Rm4822, exhibit similar *in vitro* efficacies against *B. besnoiti* tachyzoites. This indicates, that the observed anti-parasitic activity is not dependent on the thiazole-ring-associated nitro-group. In addition, our findings have also shown that Rm4803, which structurally differs from Rm4822 with regard to the positioning of the methylation on the benzene ring of the salicylic acid moiety, exhibits a strongly impaired antiparasitic activity. This conforms to earlier *in vitro* findings obtained with *N. caninum* (Esposito *et al.* 2005, 2007), and *C. parvum* (Hemphill and Mathis, unpublished observations), which have also indicated that bromo-compounds exhibit similar anti-parasitic efficacy and that the activity of Rm4803 is reduced compared to other bromo-thiazolides (unpublished observations). Our experiments also demonstrate the importance of the benzene ring for drug efficacy, since exchanging the thiazole-nitro-group for a bromide does not notably impair the anti-parasitic activity of the molecule. This indicates that the nitro-group is not crucially instrumental in terms of parasitocidal activity, but that a crucial role is possibly mediated through the unmodified *ortho*-position on the benzene ring. On the other hand, a methyl substitution *ortho*-versus *para* on an aromatic ring is well known to critically influence reactivity of other substituents including the beta-ketoamide functionality.

While TIZ and Rm4822 appeared to exert their effect by inhibiting intracellular proliferation of tachyzoites, NTZ also inhibited the host cell invasion process. This differs from *N. caninum*, where NTZ-treatment of tachyzoites did not affect host cell entry (Esposito *et al.* 2005). Thus, our study indicates that not only intracellular *B. besnoiti*, but also extracellular tachyzoites, are susceptible to NTZ, but not to its metabolite TIZ, which represents the deacetylated compound. In addition, we also found that NTZ, and Rm4803-treated parasites exhibited increased adhesion to host cells. It is very likely that adhesion to host cells is, similar to other apicomplexans, dependent on the initial interaction between the host cell and tachyzoite plasma membrane and

the sequential release of proteins from secretory organelles including micronemes, rhoptries and dense granules (Sibley, 2004; Hemphill *et al.* 2006b), organelles that are also readily found in *B. besnoiti* tachyzoites. However, in *Besnoitia*, the major surface constituents have not been identified so far, and the secretory organelles have not been characterized on the molecular level. Currently, we do not know whether NTZ affects secretory events or has an impact on the functional activity of secretory proteins, and this needs to be investigated in the future. In any case, it is likely that the activity against the extracellular stages is not relevant to the killing mechanism of the drug, and that the decisive anti-parasitic action is taking place intracellularly, a suggestion earlier for *N. caninum* (Esposito *et al.* 2005).

TEM showed that treatment of intracellular tachyzoites with NTZ, TIZ and Rm4822 caused massive vacuolization and membranous vesiculation within the parasite cytoplasm after 12–48 h. These are signs of severely impaired metabolic activity and can be attributed to the action of the drugs. Other effects included distortion of the PVM, and degradation of the granular matrix of the parasitophorous vacuole. The effects appeared similar for all three drugs, and massive vacuolization of the cytoplasm of *N. caninum* and *T. gondii* tachyzoites was reported earlier also upon treatment of *N. caninum*- and *T. gondii*-infected cells with ponazuril (Darius *et al.* 2004; Mitchel *et al.* 2005). Darius *et al.* (2004) also indicated that ponazuril was targeting the *N. caninum* apicoplast. However, we found no clear evidence for thiazolide-induced damage specifically occurring at the apicoplast in *B. besnoiti*. In addition, similar findings, and especially interference in the maintenance of the structural integrity of the parasitophorous vacuole membrane, had been reported for *N. caninum* tachyzoites treated with NTZ (Esposito *et al.* 2005).

Detailed ultrastructural analysis indicated that distortion of the PVM and degradation of the granular matrix occurred earlier in Rm4822-treated cultures, namely shortly (12 h) after initiation of drug treatment, while similar effects in NTZ-treated cultures were seen only later. It has been shown that the parasitophorous vacuole represents an essential compartment for survival, development, and proliferation, not only of *B. besnoiti*, but also for other cyst forming coccidian parasites (Mordue *et al.* 1999). In *T. gondii* the membrane delineating the parasitophorous vacuole acts as a molecular sieve, allowing free diffusion of small molecules such as sugars, amino acids, nucleobases and cofactors from the host cytoplasm, to the vacuolar space (Schwab *et al.* 1994). In addition, it is conceivable that in *B. besnoiti*-infected cells, like for other apicomplexans, the PVM and its constituents, as well as parasite-derived secretory products passing through the

membrane into the host cell cytoplasm, are involved in manipulating host cell functions. Thus, the PVM participates in selective diffusion of molecules, and represent an important constituent involved in regulating import and export of molecules (Schwab *et al.* 1994), and as its structural integrity is distorted, parasites are deprived of their natural habitat, stop proliferation, and undergo distinct structural alterations and finally cell death. However, more detailed investigations are required in order to define the exact mechanism of the anti-parasitic activity of NTZ and Rm4822 against *B. besnoiti* tachyzoites. Most importantly, the respective drug targets need to be identified and characterized.

The fact that bromo-derivatives such as Rm4822 exhibit anti-parasitic activity against *B. besnoiti* is important. *B. besnoiti* is a parasite of cattle (Cortes *et al.* 2006). As NTZ is also active against both semi-aerobic and anaerobic bacteria, this will inevitably cause significant problems in the digestive system in ruminants and also any other mammals. The gut contains beneficial populations of microflora, principally comprised of anaerobic bacteria. Oral application of a broad spectrum compound such as NTZ will kill the bacterial flora, which may lead to secondary complications including severe gastrointestinal disturbances. This has been documented in horses, where administration of NTZ can disrupt the normal microbial flora of the gastrointestinal tract, which caused enterocolitis, and has even lead to death of individual horses (Navigator™ prescribing information, IDDEX Pharmaceuticals Inc., Greensboro, NC, USA). It has been speculated that this is due to the action of PFOR or other nitro-reductases, which reduce the nitro-group and kill the bacteria through the production of free radicals (reviewed by Hemphill *et al.* (2006a)). However, solid experimental evidence for this hypothesis is missing. Studies in *H. pylori* suggested that, in contrast to metronidazole, NTZ was largely non-mutagenic (Sisson *et al.* 2002). However, the prescribing information on NTZ (Alinia™) states that the drug was genotoxic in one tester strain (TA 100) in the Ames bacterial mutation assay. As potential mutagenicity would never be compatible with application of the drug in food and breeding animals bromo-compounds such as Rm4822 could represent a safe alternative. Rm4822 will probably have no effect against the intestinal microbial flora (Pankuch and Appelbaum, 2006), and computed toxicity suggested that Rm4822 is very unlikely to exhibit any mutagenicity (data not shown). Thus, bromo-derivatives represent a safer alternative to the parent compound NTZ, but with a similar efficacy against *B. besnoiti* and/or other apicomplexan parasites. Further studies are required to investigate whether Rm4822 will be useful for *in vivo* treatment of *B. besnoiti* infections, which may be of practical use in animal production.

The authors acknowledge financial support provided by the Swiss National Science Foundation (grant no. 3100A0-112532/1) and COST 854.

REFERENCES

- Coombs, G. H. and Muller, S.** (2002). Recent advances in the search for new anticoccidial drugs. *International Journal for Parasitology* **32**, 497–508.
- Cortes, H., Ferreira, M. L., Silva, J. F., Vidal, R., Serra, P. and Caeiro, V.** (2003). Contribuição para o estudo da besnoitiose bovina em Portugal. *Revista Portuguesa de Ciências Veterinárias* **98**, 43–46.
- Cortes, H., Leitão, A., Vidal, R., Vila-Vicosa, M. J., Ferreira, M. L., Caeiro, V. and Hjerpe, C. A.** (2005). Besnoitiosis in bulls in Portugal. *The Veterinary Record* **157**, 262–264.
- Cortes, H. C., Reis, Y., Waap, H., Vidal, R., Soares, H., Marques, I., Pereira da Fonseca, I., Fazendeiro, I., Ferreira, M. L., Caeiro, V., Shkap, V., Hemphill, A. and Leitao, A.** (2006). Isolation of *Besnoitia besnoiti* from infected cattle in Portugal. *Veterinary Parasitology* **141**, 226–233.
- Darius, A. K., Mehlhorn, H. and Heydorn, A. O.** (2004). Effects of toltrazuril and ponazuril on the fine structure and multiplication of tachyzoites of the NC-1 strain of *Neospora caninum* (a synonym of *Hammondia heydorni*) in cell cultures. *Parasitology Research* **92**, 453–458.
- Dubey, J. P.** (1999). Recent advances in *Neospora* and neosporosis. *Veterinary Parasitology* **84**, 349–367.
- Esposito, M., Stettler, R., Moores, S. L., Pidathala, C., Muller, N., Stachulski, A., Berry, N. G., Rossignol, J. F. and Hemphill, A.** (2005). In vitro efficacies of nitazoxanide and other thiazolides against *Neospora caninum* tachyzoites reveal antiparasitic activity independent of the nitro group. *Antimicrobial Agents and Chemotherapy* **49**, 3715–3723.
- Esposito, M., Mueller, N. and Hemphill, A.** (2007). Structure-activity relationships from *in vitro* efficacies of the thiazolide series against the intracellular apicomplexan protozoan *Neospora caninum*. *International Journal for Parasitology* **37**, 183–190. Epub ahead of print (doi:10.1016/j.ijpara.2006.10.009)
- Fox, L. M. and Saravolatz, L. D.** (2005). Nitazoxanide: a new thiazolide antiparasitic agent. *Clinical and Infectious Diseases* **40**, 1173–1180.
- Hemphill, A. and Croft, S. L.** (1997). Electron microscopy in parasitology. In *Analytical Parasitology* (ed. Rogan, M. T.), pp. 227–268. Springer Verlag, Heidelberg.
- Hemphill, A., Mueller, J. and Esposito, M.** (2006a). Nitazoxanide, a broad-spectrum thiazolide anti-infective agent for the treatment of gastrointestinal infections. *Expert Opinion on Pharmacotherapy* **7**, 953–964.
- Hemphill, A., Vonlaufen, N. and Naguleswaran, A.** (2006b). Cellular and immunological basis of the host-parasite relationship during infection with *Neospora caninum*. *Parasitology* **133**, 261–278.
- Juste, R. A., Cuervo, L. A., Marco, J. C. and Oregui, L. M.** (1990). La besnoitiosis bovina: desconocida en España? *Medicina Veterinaria* **7**, 613–618.

- Mitchell, S. M., Zajac, A. M., Davis, W. L., Kennedy, T. J. and Lindsay, D. S.** (2005). The effects of ponazuril on development of apicomplexans in vitro. *Journal of Eukaryotic Microbiology* **52**, 231–235.
- Mordue, D., Desai, N., Dustin, M. and Sibley, D. L.** (1999). Invasion by *Toxoplasma gondii* establishes a moving junction that selectively excludes host cell plasma membrane proteins on the basis of their membrane anchoring. *The Journal of Experimental Medicine*, **190**, 1783–1792.
- Muller, J., Ruhle, G., Muller, N., Rossignol, J. F. and Hemphill, A.** (2006). *In vitro* effects of thiazolides on *Giardia lamblia* WB clone C6 cultured axenically and in coculture with Caco2 cells. *Antimicrobial Agents and Chemotherapy* **50**, 162–170.
- Naguleswaran, A., Muller, N. and Hemphill, A.** (2003). *Neospora caninum* and *Toxoplasma gondii*: a novel adhesion/invasion assay reveals distinct differences in tachyzoite-host cell interactions. *Experimental Parasitology* **104**, 149–158.
- Pankuch, G. A. and Appelbaum, P. C.** (2006). Activities of tizoxanide and nitazoxanide compared to those of five other thiazolides and three other agents against anaerobic species. *Antimicrobial Agents and Chemotherapy* **50**, 1112–1117.
- Pols, J. W.** (1960). Studies on bovine besnoitiosis with special reference to the aetiology. *The Onderstepoort Journal of Veterinary Research* **28**, 265–356.
- Schwab, J. C., Beckers, C. J. and Joiner, K. A.** (1994). The parasitophorous vacuole membrane surrounding intracellular *Toxoplasma gondii* functions as a molecular sieve. *Proceedings of the National Academy of Sciences, USA* **91**, 509–513.
- Shkap, V., Pipano, E. and Ungar-Waron, H.** (1987). *Besnoitia besnoiti*: chemotherapeutic trials *in vivo* and *in vitro*. *Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux* **40**, 259–264.
- Sisson, G., Goodwin, A., Raudonikiene, A., Hughes, N. J., Mukhopadhyay, A. K., Berg, D. E. and Hoffman, P. S.** (2002). Enzymes associated with reductive activation and action of nitazoxanide, nitrofurans, and metronidazole in *Helicobacter pylori*. *Antimicrobial Agents and Chemotherapy* **46**, 2116–2123.