Characterization of the fetuin-binding fraction of *Neospora caninum* tachyzoites and its potential involvement in host-parasite interactions

N. VONLAUFEN, A. NAGULESWARAN†, C. GIANINAZZI and A. HEMPHILL*

Institute of Parasitology, Vetsuisse Faculty, University of Berne, Länggass-Strasse 122, CH-3012 Berne, Switzerland

(Resubmitted 11 October 2006; revised 18 November 2006; accepted 20 November 2006; first published online 12 February 2007)

SUMMARY

Terminal sialic acid residues on surface-associated glycoconjugates mediate host cell interactions of many pathogens. Addition of sialic acid-rich fetuin enhanced, and the presence of the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid reduced, the physical interaction of *Neospora caninum* tachyzoites and bradyzoites with Vero cell monolayers. Thus, *Neospora* extracts were subjected to fetuin-agarose affinity chromatography in order to isolate components potentially interacting with sialic acid residues. SDS-PAGE and silver staining of the fetuin binding fraction revealed the presence of a single protein band of ~65 kDa, subsequently named NcFBP (*Neospora caninum* fetuin-binding protein), which was localized at the apical tip of the tachyzoites and was continuously released into the surrounding medium in a temperature-independent manner. NcFBP readily interacted with Vero cells and bound to chondroitin sulfate A and C, and anti-NcFBP antibodies interfered in tachyzoite adhesion to host cell monolayers. In addition, analysis of the fetuin binding fraction by gelatin substrate zymography was performed, and demonstrated the presence of two bands of 96 and 140 kDa exhibiting metalloprotease-activity. The metalloprotease activity readily degraded glycosylated proteins such as fetuin and bovine immunoglobulin G heavy chain, whereas non-glycosylated proteins such as bovine serum albumin and immunoglobulin G light chain were not affected. These findings suggest that the fetuin-binding fraction of *Neospora caninum* tachyzoites contains components that could be potentially involved in host-parasite interactions.

Key words: *Neospora caninum*, tachyzoites, fetuin-binding protein, host-parasite interactions.

INTRODUCTION

Neospora caninum is an apicomplexan parasite, which causes neosporosis in cattle and dogs. The disease is characterized by abortion, stillbirth or calves born with congenital infection, and neuromuscular dysfunction in dogs (Dubey and Lindsay, 1996; Hemphill, 1999; Dubey *et al.* 2002). Although phylogenetically closely related to *Toxoplasma gondii*, *N. caninum* differs from *T. gondii* with regard to its ultrastructure (Hemphill *et al.* 2004), antigenic composition (Howe and Sibley, 1999; Hemphill *et al.* 2006), and host cell recognition (Naguleswaran *et al.* 2002, 2003), all most likely contributing to its host-range specificity and distinct pathology (Dubey *et al.* 2002). However, the infective stages of both species are capable of invading, and proliferating within, a wide range of tissues and cell types.

The physical interaction between host cells and members of the apicomplexa involves the sequential secretion from specialized organelles named micronemes, rhoptries and dense granules. Corresponding secreted proteins mediate adhesion to the host cell surface, host cell entry, formation of an intracellular parasitophorous vacuole in which the parasites reside, and modification and maturation of this vacuole (Mauel, 1996; Black and Boothroyd, 2000; Carruthers *et al.* 2000*a,b*; Tomley and Soldati, 2001; Binder and Kim, 2004; Mercier *et al.* 2005). An increasing body of literature demonstrates the important role of proteases in host cell adhesion and invasion by apicomplexan parasites. *Toxoplasma* and *Plasmodium* express serine and cysteine proteases required for attachment and penetration. These proteases are responsible for trimming of terminal peptides or segmentation of adhesive proteins, originating from micronemes, into multiple fragments. This eventually leads to activation of these adhesins and mediates improved binding activities to host cell surface receptors. The apically discharged adhesins then form a bridge connecting host receptors with the internal motility machinery comprised of a parasite-derived actin-myosin system (Dobrowolsky *et al.* 1996; Buscaglia *et al.* 2003;

* Corresponding author: Institute of Parasitology, Vetsuisse Faculty, University of Berne, Länggass-Strasse 122, CH-3012 Berne, Switzerland. Tel: +41 31 631 2384. Fax: +41 31 631 2477. E-mail: andrew.hemphill@ipa.unibe.ch

† Current address: Department of Pharmacology and Toxicology, Indiana University School of Medicine, 635 Barnhill Drive, MS A-515, Indianapolis, IN 46202, USA.

Jewett and Sibley, 2003; Carruthers and Blackman, 2005).

Although the mechanisms leading to host cell adhesion and invasion in apicomplexans are generally regarded as conserved across the phylum, we have previously reported on distinct differences between *Toxoplasma* and *Neospora* with regard to the utilization of host cell receptors and susceptibilities to protease inhibitors. For instance, pre-treatment of *Toxoplasma* tachyzoites with protease inhibitors such as phenanthroline, E64 and pepstatin severely affected host cell interactions, while treatment with PMSF, a serine protease inhibitor, lead to a significant reduction in host cell invasion (Conseil *et al.* 1999; Naguleswaran *et al.* 2003). Conversely, neither cysteine-, metallo- nor serine protease inhibitors affected the host cell interaction of *N. caninum* tachyzoites, but invasion was drastically reduced by the aspartic protease inhibitor pepstatin-A (Naguleswaran *et al.* 2003). Nevertheless, in *N. caninum* micronemes a subtilisin-like serine protease (NcSUB1; Louie and Conrad, 1999; Louie *et al.* 2002), homologous to the corresponding enzyme TgSUB1 in *T. gondii* (Miller *et al.* 2001), has been identified, and a putative role of this protease in host cell invasion has been discussed.

Investigations on host cell surface glycosaminoglycans as potential receptors demonstrated that *N. caninum* tachyzoites preferentially bound to host cell surface chondroitin sulfates, while *T. gondii* tachyzoites preferentially adhered to heparan sulfate glycosaminoglycans on the host cell surface (Naguleswaran *et al.* 2002, 2003). Although cell surface-associated terminal sialic acid residues are known to be important elements in host cell recognition, binding and invasion by viruses, bacteria and protozoan parasites (Vimr and Lichtensteiger, 2002), few reports exist on the role of sialic acid moieties during *Neospora* infection. Treatment of Vero cell monolayers with sialidase did not result in any notable alteration of the adhesive or invasive capacities of *N. caninum* tachyzoites (Naguleswaran *et al.* 2003). This was confirmed by Vonlaufen *et al.* (2004) who demonstrated that enzymatic removal of terminal sialic acid residues from either the host cell or the parasite surface increased the invasive capacities of bradyzoites, but not tachyzoites.

In this study, we found that pre-treatment of *N. caninum* tachyzoites with the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid reduced host cell adhesion of bradyzoites and to a lesser degree, also of tachyzoites, and incubation of tachyzoites with fetuin significantly increased the adhesive capacity of *N. caninum* tachyzoites to Vero cell monolayers. This pointed towards an as yet undefined role that sialic acid might play in the process of host cell recognition. Thus, *N. caninum* tachyzoite extracts were subjected to fetuin-agarose affinity chromatography, in order to isolate proteins

potentially interacting with fetuin and/or sialic acid residues, and which could play a role in the host-parasite relationship. We here report on the characterization of this fetuin-binding fraction.

MATERIALS AND METHODS

Buffers and reagents

Unless otherwise stated, all reagents, chemicals and tissue culture media were purchased from Sigma (St Louis, MO).

Tissue culture and parasite purification

Cultures of Vero cells were maintained in RPMI 1640 medium (Gibco-BRL, Basel, Switzerland) supplemented with 10% FCS, 2 mM glutamine, 50 U of penicillin/ml and 50 µg of streptomycin/ml at 37 °C with 5% CO₂ in tissue culture flasks. Cultures were trypsinized at least once a week. *Neospora caninum* tachyzoites of the Nc-Liverpool isolate (Barber *et al.* 1995) were maintained in Vero cell monolayer cultures during the time of which FCS was replaced with IgG-free horse serum. *N. caninum* tachyzoites were harvested as described previously (Hemphill, 1996).

Induction of *N. caninum* tachyzoite-to-bradyzoite stage conversion

In vitro stage conversion was carried out in sodium nitroprusside (SNP)-treated Vero cells according to the procedure described by Vonlaufen *et al.* (2004). Briefly, Vero cell monolayers grown on glass coverslips in 24-well tissue culture plates were infected with 10⁵/ml tachyzoites of the Nc-Liverpool isolate. The infected monolayers were cultured at 37 °C with 5% CO₂ for 8 days with daily addition of 17 µM SNP.

Protein extraction and fetuin-agarose affinity purification

Proteins were extracted from 1 × 10⁸/ml freshly isolated tachyzoites in an extraction buffer (100 mM phosphate, pH 6.8 containing 0.1% Triton X-100 and 1 mM PMSF) on ice by vortexing for 5 min. The protein suspension was centrifuged at 10 000 *g* for 30 min to remove the insoluble debris and the supernatant was used for fetuin agarose affinity chromatography. Fetuin-agarose was equilibrated with 100 mM phosphate buffer (pH 6.8) and the protein extract was loaded onto the affinity column. The column was washed extensively with 100 mM phosphate buffer (pH 6.8) containing 500 mM NaCl to remove non-specifically bound proteins, and the fetuin-binding protein fraction was eluted with

100 mM methylmannoside in 100 mM phosphate buffer (pH 6.8), aliquoted and stored at -80°C .

SDS-PAGE, immunoblotting and affinity purification of antibodies

The fractions eluted from fetuin-agarose affinity chromatography and *N. caninum* total extract were precipitated in methanol-chloroform (Wessel and Fluegge, 1984) and separated by SDS-PAGE under reducing conditions. Protein bands were visualized by silver staining. Immunoblots were prepared by electrophoretically transferring SDS-PAGE-separated proteins onto nitrocellulose, and non-specific binding sites were blocked in 3% bovine serum albumin (BSA) in TBS-Tween (20 mM Tris-HCl pH 7.6, 150 mM NaCl, 0.3% Tween 20) overnight at 4°C . Blots were labelled with anti-*N. caninum* hyperimmune serum (Hemphill *et al.* 1996) diluted in TBS-Tween/0.3% BSA for 2 h at room temperature. After extensive washing in TBS-Tween, bound antibodies were visualized using goat anti-rabbit-alkaline phosphatase conjugate (Promega) according to the instructions provided by the manufacturer. Affinity purification of mono-specific antibodies directed against the 65 kDa protein (NcFBP) out of a polyclonal anti-*N. caninum* antiserum was carried out as previously described (Sonda *et al.* 2000). Antibodies were supplemented with 0.3% BSA, aliquoted and stored at -20°C . For immunoblotting, affinity-purified antibodies were used at a dilution of 1:200.

Immunofluorescence localization of NcFBP

Immunofluorescence staining was performed as described by Vonlaufen *et al.* (2004) with tachyzoites and bradyzoites grown in Vero cell monolayers on glass cover-slips in 24-well tissue culture plates, either in the absence or presence of SNP. The following primary antibodies were diluted in PBS/0.3% BSA: (i) NcmAb-4, a monoclonal antibody directed against the immunodominant tachyzoite surface antigen NcSAG1 (Björkman and Hemphill, 1998) was used at $1\text{ }\mu\text{g/ml}$; (ii) affinity-purified polyclonal rabbit anti-NcFBP antibody was diluted at 1:1 dilution; (iii) mAbCC2, a rat mAb reacting with a *T. gondii* cyst wall protein (Gross *et al.* 1995), was diluted at 1:100. Incubation with primary antibodies was performed for 1 h, followed by 3 washes in PBS, 5 min each. Bound antibodies were detected with the appropriate tetramethylrhodamine isothiocyanate (TRITC)- or fluorescein isothiocyanate (FITC)-conjugated secondary antibodies. Finally, the preparations were washed in PBS 3 times for 5 min each and incubated in the fluorescent dye Hoechst 33258 ($1\text{ }\mu\text{g/ml}$ in PBS) for 2 min, rinsed again in PBS and mounted in

Fluoroprep (Biomerieux S.A.). All specimens were viewed on a Nikon Eclipse E 800 digital confocal fluorescence microscope. Processing of images was performed using the Openlab 2.0.7 software (Improvision).

Transmission electron microscopy (TEM) and immunogold labelling

For TEM, *N. caninum*-infected Vero cell cultures were fixed in 100 mM sodium cacodylate, pH 7.2, containing 2% paraformaldehyde for 30 min at 20°C . LR-White-embedding and on-section labelling was performed essentially as previously described (Vonlaufen *et al.* 2004). Sections were incubated in affinity-purified anti-FBP diluted 1:1 in PBS/0.3% BSA for 1 h. After washing in 5 changes of PBS, 2 min each, the goat anti-rabbit antibody conjugated to 10 nm gold particles (purchased from Amersham, Zürich, Switzerland) was applied (Vonlaufen *et al.* 2004). Finally, grids were stained with lead citrate and uranyl acetate and were subsequently viewed on a Phillips 400 transmission electron microscope operating at 80 kV.

Gelatin zymography

Protein fractions were precipitated with 80% pre-cooled acetone for 30 min on ice. The protein was resuspended in native sample buffer and separated on 10% SDS-PAGE containing 0.1% gelatin. After electrophoresis, gels were washed twice with 2.5% Triton X-100 for 30 min to remove the SDS and to re-nature the proteins. After rinsing with water, gels were incubated overnight at 37°C in an incubation buffer (50 mM Tris, pH 7.6, containing 50 mM NaCl and 10 mM CaCl_2) with gentle shaking. In some experiments, 1 mM 1,10-phenanthroline or 1 mM ZnCl_2 were added to the incubation buffer. Following overnight incubation, gelatinolytic activity was visualized with Coomassie brilliant blue G250 stain as clear bands on a blue background.

In vitro secretion assay

Freshly purified tachyzoites ($10^8/\text{ml}$) were resuspended in Earl's balanced salt solution (EBSS) and were incubated either on ice or at 37°C for 10 min. In some experiments, a cocktail of protease inhibitors (1 mM PMSF, 0.5 mM 1,10-phenanthroline, 0.02 mM E64 and 0.1 mM pepstatin-A) was added. After incubation, parasites were centrifuged at 2000 g and the supernatant was re-centrifuged at 10 000 g for 10 min at 4°C . Proteins arising from equal number of parasites from supernatants and pellets were separated on SDS-PAGE and immunoblots were performed using anti-NcFBP antibodies. Antibodies directed against NcMIC4, a

secretory *N. caninum* microneme protein (Keller *et al.* 2004) were included as a control.

Assessment of the effects of fetuin, anti-NcFBP antibodies and the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid on N. caninum tachyzoite and bradyzoite host cell interaction

Freshly purified tachyzoites (10^5) were resuspended in 200 μ l of RPMI 1640 medium. The suspension was supplemented either with fetuin (1 mg/ml) and incubated at 37 °C for 30 min, or with 50 μ l affinity-purified anti-NcFBP antibodies (diluted 1:1), or anti-beta-galactosidase (control) antibodies and incubated at 4 °C (Hemphill, 1996; Naguleswaran *et al.* 2005). In some experiments, parasites were pre-incubated with fetuin (1 mg/ml) for 30 min at 37 °C and subsequently re-incubated with affinity purified antibodies at 4 °C for 30 min. In other assays, the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid (10 μ M) was added for 30 min at 37 °C to tachyzoites and bradyzoites, followed by a centrifugation step to remove the sialidase inhibitor. The treated parasites were then allowed to interact with the Vero cell monolayer for 30 min at 37 °C. The unbound parasites were washed with medium and tachyzoites associated with host cells were quantified by LightCycler PCR according to the method of Müller *et al.* (2002). In some experiments, the PDTC-PCR-based adhesion/invasion assay according to Naguleswaran *et al.* (2003) was used to specifically quantify intracellular and extracellular tachyzoites and bradyzoites.

NcFBP – Vero cell co-sedimentation assays

These assays were done as previously described (Naguleswaran *et al.* 2002; Keller *et al.* 2004). Freshly trypsinized, non-adherent Vero cells were fixed in 2.5% glutaraldehyde in EBSS for 30 min, followed by post-fixation in 0.5% OsO₄ in 100 mM sodium phosphate buffer, pH 7.2. Subsequently, free aldehyde groups were blocked by incubation in 100 mM ethanolamine (pH 8) at 4 °C overnight. Triton X-100 extracts of *N. caninum* tachyzoites were prepared by incubating 5×10^8 freshly isolated *N. caninum* tachyzoites in 2 ml of PBS containing 1% Triton X-100 for 5 min at 4 °C, followed by centrifugation at 10 000 *g* for 30 min at 4 °C. Pre-fixed Vero cells (10^6) were then incubated in 250 μ l of the Triton X-100 extracts for 2 h at 4 °C. Subsequently the preparations were centrifuged at 10 000 *g* for 5 min, and the supernatants were collected and processed for SDS-PAGE and immunoblotting. The pellets were washed in PBS 3 times and were finally resuspended in SDS-PAGE sample buffer. Equal amounts of non-bound (supernatants) and bound

(pellets) proteins were loaded onto gels. Immunoblotting was performed as above.

Binding of NcFBP to defined glycosaminoglycans

Binding assays were carried out as described (Naguleswaran *et al.* 2003; Keller *et al.* 2004). Ninety-six-well ELISA plates were coated with heparin, dextran, dextran sulfate, or chondroitin sulfate A, B, or C (CSA, CSB, CSC) (5 mg/ml) for 12 h at 4 °C. Plates were washed and incubated with 100 μ l of the Triton X-100 extracts of *N. caninum* tachyzoites for 2 h at 4 °C. Specimens were then washed in PBS and nonspecific binding sites were blocked for 2 h with 1.5% BSA and 50 mM glycine in PBS. Bound NcFBP was detected using affinity-purified antibodies against NcFBP (diluted 1:50 in antibody dilution buffer), followed by the respective secondary antibodies conjugated to alkaline phosphatase. Antibody binding was visualized by using *p*-nitrophenylphosphate as a substrate and absorbance was measured at 405 nm on a Dynatech MR7000 enzyme-linked immunosorbent assay (ELISA) reader.

Proteolytic assays

For proteolysis of protein substrates, 200 μ l of the eluted *N. caninum* fetuin-binding protein fraction and 20 μ g of each fetuin, bovine immunoglobulin G and bovine serum albumin in 200 μ l of incubation buffer were incubated at 37 °C for 12 h. As a control, each protein substrate was incubated in incubation buffer and elution buffer. Protein degradation was assessed by SDS-PAGE followed by silver staining.

RESULTS

Involvement of sialic acid in the interaction between N. caninum and host cells

Pre-treatment of *N. caninum* with the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid resulted in a reduction in the number of parasites adhering/invading Vero cell monolayers, namely by 20% for tachyzoites and 40% for bradyzoites (Fig. 1A). The percentage of actually invaded parasites remained at similar levels independent of the treatments, i.e. 73% for the untreated tachyzoite control, 72% for the sialidase inhibitor-treated tachyzoites, and 28% and 30% for untreated bradyzoites and sialidase inhibitor-treated bradyzoites, respectively. This suggested that sialic acid could be involved in parasite attachment to host cells, but not necessarily in invasion. Fetuin, a liver-produced negative acute-phase protein rich in sialic acid content, was added to tachyzoites prior to letting them interact with Vero cell monolayers, and this resulted in a 2-fold increased number of parasites binding

Fig. 1. Sialic acid residues are involved in *Neospora caninum* host cell interaction. (A) Incubation of *N. caninum* tachyzoites and bradyzoites with the sialidase inhibitor 2-deoxy-2,3-dehydro-N-acetylneuraminic acid leads to a 20% and 40% reduction of parasites adhering to their host cells, respectively. However, the percentage of parasites invading the host cells remains the same. (B) The presence of fetuin leads to a massive increase (50%) of tachyzoites binding to host cells. The significance of the differences among the control and experimental assays was determined by Student's *t*-test using the Microsoft Excel program. *P* values < 0.05 were considered statistically significant and are marked with *.

and/or invading the host cells (Fig. 1B). Addition of desialylated asialofetuin did not have any effect (data not shown), suggesting that the increase in host cell associated tachyzoites could be attributed to the presence of sialic acid residues.

The N. caninum fetuin-binding fraction is constituted by a single protein of 65 kDa, which is expressed at the apical portion of N. caninum tachyzoites

Analysis of the fetuin-binding protein fraction of *N. caninum* tachyzoites by SDS-PAGE under reducing conditions and silver staining revealed the presence of a single fetuin binding protein band of 65 kDa, which was named NcFBP. (Fig. 2A, lane 2). When proteins were separated under non-reducing conditions, silver staining failed to detect any protein (Fig. 2A, lane 3). Immunoblotting showed that the 65 kDa band was the only protein recognized by a hyperimmune serum directed against *N. caninum* tachyzoites (Fig. 2B, lane 2), but no immunolabelling could be observed when SDS-PAGE was carried out

under non-reducing conditions (Fig. 2B, lane 3). Subsequently, affinity purification on the 65 kDa band yielded monospecific anti-NcFBP antibodies, and these specifically recognized the 65 kDa band on immunoblots of *Neospora* extracts (Fig. 2B, lane 4).

The affinity-purified antibodies localized the protein to the apical portion of both intracellular (Fig. 3A–C) and extracellular (Fig. 3D–F) tachyzoites, and closer inspection showed that at least a major portion of this protein expressed in these parasites was present on the tachyzoite surface (Fig. 3F). No cross-reaction of anti-NcFBP antibodies with host cells could be detected. Immunofluorescence labelling of parasites undergoing *in vitro* tachyzoite-to-bradyzoite stage conversion indicated that the expression of NcFBP during stage conversion was greatly reduced, or that the protein was modified in a way that rendered epitopes less accessible (Fig. 3G–I). Attempts to determine the exact subcellular localization of NcFBP by immunogold-TEM were hampered by the fact that anti-NcFBP

Fig. 2. Analysis of fetuin-binding fraction by SDS-PAGE and immunoblotting. (A) Fetuin-affinity chromatography of *Neospora caninum* extracts was analysed by reducing (lanes 1, 2) and non-reducing (lane 3) SDS-PAGE and stained by silver stain. Lane 1, *N. caninum* Triton X-100 soluble extract prior to affinity chromatography, lane 2, eluted fraction containing the fetuin-binding protein (NcFBP), lane 3, eluted fraction separated under non-reducing conditions. (B) Immunoblot of the same fractions as in panel A. Lanes 1–3 were labelled with anti-*N. caninum* antiserum and detected by goat anti-rabbit antibody conjugated to alkaline phosphatase. Lane 4 shows an immunoblot of *N. caninum* Triton X-100 soluble extract labelled with affinity-purified anti-NcFBP antibodies and detected by goat anti-rabbit antibody conjugated to alkaline phosphatase.

antibodies recognized fixation-sensitive epitope(s): immunogold labelling of paraformaldehyde/glutaraldehyde-fixed samples did not yield any staining (data not shown). However, TEM of sections of infected cells fixed in 2% paraformaldehyde confirmed the predominant apical localization of NcFBP in both intracellular and extracellular tachyzoites. Gold particles appeared to bind preferentially to vesiculated structures at the anterior end resembling micronemes (see Fig. 4A), while rhoptries were not labelled (Fig. 4B).

NcFBP is continuously released by N. caninum tachyzoites irrespective of the temperature

Experiments were undertaken to investigate whether NcFBP was part of the secretory fraction of *N. caninum* tachyzoites. Thus, purified tachyzoites were incubated either at 4 °C or 37 °C in EBSS, and parasite pellets and medium supernatants were collected and analysed by immunoblotting (Fig. 5A). The results showed that NcFBP was released or shed into the medium supernatant irrespective of the incubation temperature. As a control, the same samples were probed with anti-NcMIC4 antibodies, which detect a soluble microneme protein, NcMIC4 (Keller *et al.* 2004). NcMIC4 could only be detected in the supernatant sample previously incubated at 37 °C (Fig. 5B), indicating that active secretion has taken place. We incubated tachyzoites at 4 °C and 37 °C in the presence and absence of a protease

Fig. 3. Localization of the fetuin-binding protein. Immunofluorescence labelling of *Neospora caninum* tachyzoites grown in Vero cells (A–C) and purified parasites (D–F). Parasites were stained with anti-NcFBP antibodies and fluorescein isothiocyanate-conjugated anti-rabbit IgG (green) and with NcSAG1 antibody and tetramethyl rhodamine isothiocyanate-conjugated anti-mouse IgG (red). Nuclei were labelled with Hoechst 22358. (E–G) *In vitro*-generated bradyzoites exhibiting cyst wall staining with mAb CC2 (red) and down-regulated expression of NcFBP (green). Arrows in (C) and (D) highlight tachyzoites with NcFBP surface labelling.

inhibitor cocktail consisting of PMSF, pepstatin, E64 and phenanthroline, which would cover all 4 classes of proteases. Immunoblotting demonstrated that the release of NcFBP into the medium supernatant at both high and low temperature was strongly impaired (Fig. 5C), indicating that protease activity was required for release of NcFBP. However, when protease inhibitors were used individually, no effect was observed, indicating that more than one type of protease is involved in the release of NcFBP into the medium (data not shown).

NcFBP binds to Vero cells and influences tachyzoite-host cell interaction

Anti-NcFBP antibodies had a profound negative impact on host cell adhesion by reducing the adhesion/invasion of tachyzoites to Vero cell monolayers by 62% compared to the control anti-beta galactosidase antibodies (Fig. 6A). In a further experiment, pre-incubation of *N. caninum* tachyzoites with fetuin enhanced the adhesion to Vero cells by

Fig. 4. Immunogold labelling of *Neospora caninum*-infected Vero cells employing anti-NcFBP antibodies. Cells were fixed in 2% paraformaldehyde and embedded in LR-White resin. Note the arrows pointing towards gold particles mostly associated with the apical part of the tachyzoites. Nuc, nucleus; pv, parasitophorous vacuole; con, conoid; rop, rhoptries.

Fig. 5. Immunoblots of fractions secreted by tachyzoites incubated at 4 °C or 37 °C. Blots A and C were stained with anti-NcFBP antibodies and blot B was stained with anti-NcMIC4 antibodies. (S) Corresponds to the released fraction into the supernatant and the (P) corresponds to the pellet fraction. Blot C shows the release of NcFBP in the absence (–) or presence (+) of a cocktail of protease inhibitors (PI). Note the reduced release of NcFBP in the presence of a cocktail of protease inhibitors.

50% compared to the control, and this enhanced adhesive effect was reduced by 40%, when parasites pre-incubated with fetuin were re-incubated with anti-NcFBP antibodies prior to host cell interaction (Fig. 6B). Since anti-NcFBP antibodies reduced the adhesion of tachyzoites to Vero cell monolayers, we investigated whether NcFBP binds to the Vero cell surface. Triton X-100 extracts were incubated with pre-fixed Vero cells, and following centrifugation, co-sedimented and unbound proteins were subjected to SDS-PAGE and immunoblot analysis. Staining with anti-NcFBP antibodies showed that a substantial amount of NcFBP co-precipitated with the fraction bound to Vero cells (Fig. 6C). Moreover, solid-phase binding assays were performed to investigate whether

glycosaminoglycans could potentially be involved in binding of NcFBP. ELISA wells were coated with dextran, dextran sulfate, heparin and CSA, CSB, and CSC. As shown in Fig. 6D, NcFBP exhibited an increased binding to CSA, CSC and dextran, while CSB, heparin and dextran sulfate did not bind NcFBP.

The fetuin-binding fraction of N. caninum tachyzoites contains metalloprotease activity directed against glycosylated proteins

Since it was demonstrated earlier that human matrix metalloproteases bind to fetuin (Ochieng *et al.* 1995), we investigated whether metalloprotease activity was present in the fetuin-binding fraction of *N. caninum*

Fig. 6. NcFBP interacts with host cells, and antibodies inhibit tachyzoite adhesion. The parasites were quantified by real-time PCR, and parasite count per PCR-reaction is given in the y-axis. Black columns indicate the overall number of parasites interacting with their host cell. Percentages given above the black columns indicate the reduction in the overall parasite number with respect to the control. Error bars indicate standard deviation. (A) Pre-treatment of tachyzoites with anti-NcFBP antibodies resulted in an adhesion inhibition of 62% compared to a control comprised of an irrelevant anti-beta-galactosidase antibody. (B) Pre-treatment of tachyzoites with fetuin increased the adhesion by 50% compared to the control. Pre-treatment of tachyzoites with fetuin followed by incubation with anti-NcFBP antibodies resulted in an adhesion inhibition of 41%. (C) Co-precipitation assay demonstrating binding of NcFBP to pre-fixed Vero cells. The immunoblot was stained with anti-NcFBP antibodies. Lane P (pellet) represents the protein fraction bound to Vero cells; lane SN (supernatant) represents non-bound proteins. (D) Binding of NcFBP to immobilized heparin, dextran, dextran sulfate, CSA, CSB, and CSC. Binding of NcFBP was assessed by labelling with anti-NcFBP antibodies and alkaline phosphatase-conjugated secondary antibodies. Note that NcFBP binds preferentially to dextran, CSA and CSC but not to other glycosaminoglycans. OD 405 = optical density at 405 nm.

tachyzoites. Thus, total *Neospora* extract, secreted fraction of *N. caninum* tachyzoites, and fetuin-agarose eluted fraction, were examined by gelatin substrate zymography. The fetuin-agarose binding fraction exhibited 2 distinct protease bands at 96 kDa and 140 kDa, whereas the secreted fraction and the total *Neospora* extract revealed multiple proteolytic bands of different intensities, some of which co-migrated with the 2 bands of 96 kDa and 140 kDa in the fetuin-binding fraction (Fig. 7A). Fetuin-affinity chromatography of Vero cell extracts performed under identical conditions followed by zymography did not reveal any bands with protease activity (data not shown). In the presence of 1 mM of the zinc chelator 1,10-phenanthroline, the enzymatic activities in the fetuin-binding fraction were completely

abrogated. Further, addition of an excess of Zn^{2+} to the incubation buffer completely inhibited the proteolytic activities. Proteolytic activity was not affected by the presence of PMSF, pepstatin and E64, respectively (data not shown). This showed that the presence of the 2 bands was attributable to metalloprotease activity.

To address the question whether fetuin, besides binding metalloprotease, could also act as a substrate, fetuin was incubated at 37 °C, either in the presence or absence of the *N. caninum* fetuin-binding fraction, and the samples were analysed by SDS-PAGE and silver staining (Fig. 7B). When fetuin was incubated without the fetuin-binding fraction, the major protein band plus several other bands of lower *Mr* were visible. After incubation of fetuin in the presence of

Fig. 7. Proteolytic activity of fetuin-binding fractions. (A) Gelatin substrate zymography. Lane 1, *N. caninum* Triton X-100 soluble extract, lane 2, secreted fraction of *N. caninum* tachyzoites at 4 °C, lane 3, eluted fraction containing NcFBP, lanes 4 and 5, eluted fraction containing NcFBP incubated with 1,10-phenanthroline and 1 mM ZnCl₂, respectively were electrophoresed on a 0.1% gelatin containing 10% polyacrylamide gel and further processed as described in the Materials and Methods. (B) Proteolytic degradation of glycosylated and non-glycosylated substrates of NcFBP. All reaction mixtures were incubated at 37 °C for 12 h and degradation products were analysed by SDS-PAGE and visualized by silver stain. Lane 1, fetuin + NcFBP, lane 2, fetuin, lane 3, bovine immunoglobulin G + NcFBP, lane 4, bovine immunoglobulin G, lane 5, bovine serum albumin (BSA) + NcFBP, lane 6, BSA. Note the degradation of fetuin and bovine immunoglobulin G heavy chain (H). BSA and bovine immunoglobulin G light chain (L) remain undegraded.

N. caninum fetuin-binding proteins, the fetuin was largely digested, reflecting strong proteolytic activity. Identical experiments using bovine immunoglobulin G as a substrate revealed that the glycosylated heavy chain was completely digested, whereas the non-glycosylated light chain remained largely unaffected (Fig. 7B). In contrast, bovine

serum albumin, which represents a non-glycosylated substrate, was not affected by incubation with the fetuin-binding protein fraction (Fig. 7B). These experiments indicated that the metalloprotease activity was largely targeted towards glycoproteins, while non-glycosylated polypeptides remained largely unaffected.

DISCUSSION

Sialic acids are a structurally complex family of 9-carbon monosaccharides, that are involved in diverse activities related to glycoproteins and glycoconjugates, including host-pathogen interactions (Kelm and Schauer, 1997; Vimir and Lichtensteiger, 2002). The fact that functional inhibition of sialidase activity in *Neospora caninum* negatively affects tachyzoite host cell adhesion, but not invasion (see Fig. 1A), indicates, that sialic acid could also play a role in the physical relationship between *N. caninum* and its host cell. In the presence of fetuin, a highly sialylated glycoprotein with a triantennary glycan part that contains galactose, mannose, glucosamine and galactosamine as sugar molecules (Johnson and Heath, 1986), *N. caninum* tachyzoites bound much more efficiently to host cells compared to controls (Fig. 1B). It is likely that fetuin plays a similar role in adhesion as previously reported for laminin and *Toxoplasma gondii* (Furtado *et al.* 1992a) where laminin was proposed to act as a bridging molecule, interacting with both host and parasite cell surface constituents such as the integrin receptor $\alpha 6 \beta 1$ in human fibroblasts (Furtado *et al.* 1992b). Fetuin is regularly contained in FCS, which is used as an additive in culture medium, and in this context could contribute for efficient host cell interaction *in vitro*. However, fetuin is mainly a fetal protein in the sense that the highest concentrations are found in serum and body fluids of embryos and fetuses (Terkelsen, 1998), and this could be of relevance *in vivo* as neosporosis is mostly of concern in relation to fetal infection and disease in neonates.

In this study, we report on the characterization of a fetuin-binding fraction of *N. caninum* tachyzoites. In this fraction, silver staining revealed the presence of a single protein band of 65 kDa, and zymography demonstrated profound metalloprotease activity in 2 bands of 96 and 140 kDa, respectively. This is the first description of metalloproteases identified in *N. caninum*.

A 65 kDa fetuin-binding protein (NcFBP) is involved in host cell attachment of *N. caninum* tachyzoites

The purification of a single 65 kDa protein (NcFBP) was evident upon separation of the fetuin-binding fraction by SDS-PAGE under reducing conditions, while under non-reducing conditions this protein

could not be resolved by silver staining nor by immunoblotting. Several attempts to identify the 65 kDa NcFBP by tandem mass spectrometry have not been successful (data not shown), most likely due to the fact that the sequence of the *Neospora* genome is not fully available to date, and the expression level of this protein could be low and thus the corresponding sequence not well represented in the current EST-database. Further studies are required to elucidate the exact molecular identity of NcFBP. NcFBP was the only protein in the fetuin-binding fraction that was recognized by polyclonal anti-*N. caninum* antiserum, allowing the preparation of affinity-purified anti-NcFBP antibodies that exhibited high specificity on immunoblots. Immunofluorescence showed that the protein was concentrated at the apical part of intracellular, and the apical part as well as the surface of extracellular, tachyzoites. This was largely confirmed by immunogold TEM, which indicated that NcFBP was a component of microneme-like vesicles. This suggested that NcFBP could be transported onto the parasite surface and then released into the exterior medium. Indeed, secretion assays employing extracellular tachyzoites demonstrated that NcFBP was released into the medium supernatant independently of the metabolic state of the parasites, e.g. at high (37 °C) as well as at low (4 °C) temperature. Release of NcFBP did not simply reflect leakage by dead parasites, as another secretory and soluble protein, NcMIC4 (Keller *et al.* 2004), was not present in the 4 °C supernatants. Also other micronemal proteins such as NcMIC1, NcMIC2 and NcMIC3 had been reported earlier to be secreted in substantial amounts at 37 °C, but not at 4 °C (Lovett *et al.* 2000; Naguleswaran *et al.* 2001; Keller *et al.* 2004). This suggests that surface-associated NcFBP reaches the medium supernatant by a process distinct of the one described for other secretory components of apicomplexans, whereas the secretory events involving microneme and rhoptry proteins represent active processes requiring metabolic energy and protease activity on part of the parasite (Carruthers, 1999; Joiner and Roos, 2002; Binder and Kim, 2004). Nevertheless, release of NcFBP into the medium was also dependent on functional protease machinery, as the addition of a cocktail of protease inhibitors inhibited the release of the protein. In fact, release of NcFBP into the medium supernatant appears to be more efficient at 37 °C compared to 4 °C, but the mechanisms involved in NcFBP shedding remain to be elucidated.

NcFBP is likely to be one of the factors mediating tachyzoite host cell attachment, since (i) antibodies directed against NcFBP inhibited parasite adhesion, and (ii) also the enhanced binding of tachyzoites to host cells in the presence of fetuin was largely neutralized by anti-NcFBP antibodies (iii) NcFBP can bind to the surface of Vero cells, and (iv) also

binds to sulfated glycosaminoglycans, especially chondroitinsulfate A and C, which contain terminal sialic acid residues. Chondroitin sulfates, have earlier been shown to act as receptors for *N. caninum* tachyzoite adhesion to host cells (Naguleswaran *et al.* 2002, 2003). In this context it is interesting to note that earlier studies on *Toxoplasma* by Gross *et al.* (1993) had demonstrated the uptake of fetuin by *T. gondii*. Using fetuin-agarose chromatography of *T. gondii* tachyzoite extracts, they had identified a fetuin-binding protein of 15 kDa that specifically recognized fetuin glycan structures. In addition, Monteiro *et al.* (1998) demonstrated the involvement of host cell surface sialic acid residues in the process of host cell invasion by *Toxoplasma gondii*.

The N. caninum fetuin binding fraction contains metalloproteinase activity

The fetuin-binding fraction contains metalloprotease activity, which is evidenced by gelatin substrate zymography with the 2 bands of 96 and 140 kDa. The findings on metalloprotease activity are not overall surprising, as it has been shown previously that besides interacting with sialidases (Nok and Rivera, 2003), fetuin also binds metalloproteases. For instance, Ochieng *et al.* (1995) demonstrated that gelatinase A (MMP2) and B (MMP9) were both specifically bound to fetuin.

A profound role for metalloproteases in the interaction of protozoan parasites with host cells was reported in different studies. The gp63 surface metalloprotease in *Leishmania* promastigotes has been shown to be involved in the attachment of promastigotes to the fibronectin receptor of macrophages (Brittingham *et al.* 1999). Further, preincubation of *Trypanosoma cruzi* trypomastigotes with affinity-purified antibodies against Tcgp63 reduced the invasion rate of Vero cells by 50% (Cuevas *et al.* 2003). In a recent study a reduced invasion of *Plasmodium falciparum* merozoites into red blood cells was seen in the presence of 1,10-phenanthroline (Kitjaroenatham *et al.* 2005). Although phenanthroline was previously not found to be effective in influencing *N. caninum* tachyzoite host cell adhesion or invasion (Naguleswaran *et al.* 2003), it is still possible that *N. caninum* metalloproteinases play an important role in these processes, and additional investigations using other, less toxic and more specific, inhibitors need to be carried out.

Degradation of bovine immunoglobulin G by the metalloprotease activity found in fetuin-binding fractions could reflect a possible mechanism of *N. caninum* tachyzoites to evade the host immune response. In addition, digestion of glycosylated proteins such as fetuin and immunoglobulin G heavy chain suggests that the metalloproteases preferentially act on glycosylated proteins.

In conclusion, fetuin-binding fractions of *N. caninum* tachyzoites contain (i) a 65 kDa protein that could play a role in mediating the contact between the parasite and its host cell, and (ii) metalloproteinases that are targeted towards glycoproteins. It is presently not clear, whether the 2 metalloproteinase bands and NcFBP are identical or related proteins, or whether they represent completely different entities. Inspection of the *N. caninum* EST database for the occurrence of metalloproteases revealed only 4 entries coding for 3 proteins with similarities to metalloproteases, one of them coding for a putative sialoglycoprotease (Genbank No. CF274269; CF371306; CF597981). The corresponding genes are being cloned and respective proteins are currently investigated for their possible role in host-parasite interactions.

We acknowledge Alexander J. Trees and Diana Williams for providing the *Neospora caninum* Liverpool isolate for this work. We also acknowledge Jonathan Wastling and Sanya Sanderson for MS/MS analysis. This study was financially supported by the Foundation Research 3R, the Swiss National Science Foundation (Grant No. 3100A0-112532/1), and a grant obtained through the European Union from the Swiss Secretariat of Education and Research (SBF C03.0007). Many thanks are addressed to Norbert Müller for critically reading the manuscript and many helpful discussions.

REFERENCES

- Barber, J. S., Holmdahl, O. J., Owen, M. R., Guy, F., Uggla, A. and Trees, A. J. (1995). Characterization of the first European isolate of *Neospora caninum* (Dubey, Carpenter, Speer, Topper and Uggla). *Parasitology* **111**, 563–568.
- Binder, E. M. and Kim, K. (2004). Location, location, location: trafficking and function of secreted proteases of *Toxoplasma* and *Plasmodium*. *Traffic* **5**, 914–924.
- Björkman, C. and Hemphill, A. (1998). Characterization of *Neospora caninum* iscom antigens using monoclonal antibodies. *Parasite Immunology* **20**, 73–80.
- Black, M. W. and Boothroyd, J. C. (2000). Lytic cycle of *Toxoplasma gondii*. *Microbiology and Molecular Biology Reviews* **64**, 607–623.
- Brittingham, A., Chen, G., McGwire, B. S., Chang, K. P. and Mosser, D. M. (1999). Interaction of *Leishmania* gp63 with cellular receptors for fibronectin. *Infection and Immunity* **67**, 4477–4484.
- Buscaglia, C. A., Coppens, I., Hol, W. G. and Nussenzweig, V. (2003). Sites of interaction between aldolase and thrombospondin-related anonymous protein in plasmodium. *Molecular Biology of the Cell* **14**, 4947–4957.
- Carruthers, V. B. (1999). Armed and dangerous: *Toxoplasma gondii* uses an arsenal of secretory proteins to infect host cells. *Parasitology International* **48**, 1–10.
- Carruthers, V. B. and Blackman, M. J. (2005). A new release on life: emerging concepts in proteolysis and parasite invasion. *Molecular Microbiology* **55**, 1617–1630.
- Carruthers, V. B., Hakansson, S., Giddings, O. K. and Sibley, L. D. (2000a). *Toxoplasma gondii* uses sulfated proteoglycans for substrate and host cell attachment. *Infection and Immunity* **68**, 4005–4011.
- Carruthers, V. B., Sherman, G. D. and Sibley, L. D. (2000b). The *Toxoplasma* adhesive protein MIC2 is proteolytically processed at multiple sites by two different parasite-derived proteases. *Journal of Biological Chemistry* **275**, 14346–14353.
- Conseil, V., Soete, M. and Dubremetz, J. F. (1999). Serine protease inhibitors block invasion of host cell by *Toxoplasma gondii*. *Antimicrobial Agents and Chemotherapy* **43**, 1358–1361.
- Cuevas, I. C., Cazzulo, J. J. and Sanchez, D. O. (2003). gp63 homologues in *Trypanosoma cruzi*: surface antigens with metalloprotease activity and a possible role in host cell infection. *Infection and Immunity* **71**, 5739–5749.
- Dobrowolsky, J. M., Carruthers, V. B. and Sibley, L. D. (1996). Participation of myosin in gliding motility and host cell invasion by *Toxoplasma gondii*. *Molecular Microbiology* **26**, 163–173.
- Dubey, J. P., Barr, B. C., Barta, J. R., Bjerkas, I., Björkman, C., Blagburn, B. L., Bowman, D. D., Buxton, D., Ellis, J. T., Gottstein, B., Hemphill, A., Hill, D. E., Howe, D. K., Jenkins, M. C., Kobayashi, Y., Koudela, B., Marsh, A. E., Mattsson, J. G., McAllister, M. M., Modry, D., Omata, Y., Sibley, L. D., Speer, C. A., Trees, A. J., Uggla, A., Upton, S. J., Williams, D. J. and Lindsay, D. S. (2002). Redescription of *Neospora caninum* and its differentiation from related coccidia. *International Journal for Parasitology* **32**, 929–946.
- Dubey, J. P. and Lindsay, D. S. (1996). A review of *Neospora caninum* and neosporosis. *Veterinary Parasitology* **67**, 1–59.
- Furtado, G. C., Cao, Y. and Joiner, K. A. (1992a). Laminin on *Toxoplasma gondii* mediates parasite binding to the beta 1 integrin receptor alpha 6 beta 1 on human foreskin fibroblasts and Chinese hamster ovary cells. *Infection and Immunity* **60**, 4925–4931.
- Furtado, G. C., Slowik, M., Kleinman, H. K. and Joiner, K. A. (1992b). Laminin enhances binding of *Toxoplasma gondii* to JJ774 murine macrophage cells. *Infection and Immunity* **60**, 2337–2342.
- Gross, U., Bormuth, H., Gaissmaier, C., Dittrich, C., Krenn, V., Bohne, W. and Ferguson, D. J. (1995). Monoclonal rat antibodies directed against *Toxoplasma gondii* suitable for studying tachyzoite-bradyzoite interconversion in vivo. *Clinical and Diagnostic Laboratory Immunology* **2**, 542–548.
- Gross, U., Hambach, C., Windeck, T. and Heesemann, J. (1993). *Toxoplasma gondii*: uptake of fetuin and identification of a 14 kDa fetuin-binding protein. *Parasitology Research* **79**, 191–194.
- Hemphill, A. (1996). Subcellular localization and functional characterisation of Nc-p43, a major *Neospora caninum* tachyzoite surface protein. *Infection and Immunity* **64**, 4279–4287.
- Hemphill, A. (1999). The host-parasite relationship in neosporosis. *Advances in Parasitology* **43**, 48–105.
- Hemphill, A., Gottstein, B. and Kaufmann, H. (1996). Adhesion and invasion of bovine endothelial cells by

- Neospora caninum* tachyzoites. *Parasitology* **112**, 183–197.
- Hemphill, A., Vonlaufen, N., Naguleswaran, A., Guetg, N., Srinivasan, S. and Alaeddine, F.** (2004). Tissue culture and explant approaches to study and visualise *Neospora caninum* and its interactions with the host cell. *Microscopy and Microanalysis* **10**, 1–19.
- Hemphill, A., Vonalufen, N. and Naguleswaran, A.** (2006). Cellular and immunological basis of the host-parasite relationship during infection with *Neospora caninum*. *Parasitology* **133**, 261–278.
- Howe, D. K. and Sibley, L. D.** (1999). Comparison of the major antigens of *Neospora caninum* and *Toxoplasma gondii*. *International Journal for Parasitology* **29**, 1489–1496.
- Jewett, T. J. and Sibley, L. D.** (2003). Aldolase forms a bridge between cell surface adhesins and the actin cytoskeleton in apicomplexan parasites. *Molecular Cell* **11**, 885–894.
- Johnson, W. V. and Heath, E. C.** (1986). Structural features of bovine fetuin revealed from analysis of the primary translation product: anomalous behavior on sodium dodecyl sulfate-polyacrylamide gel electrophoresis is due largely to peptide and not solely to carbohydrate. *Archives in Biochemistry and Biophysics* **251**, 732–737.
- Joiner, K. A. and Roos, D.** (2002). Secretory traffic in the eukaryotic parasite *Toxoplasma gondii*: less is more. *Journal of Cell Biology* **157**, 557–563.
- Keller, N., Riesen, M., Naguleswaran, A., Vonlaufen, N., Stettler, R., Leepin, A., Wastling, J. M. and Hemphill, A.** (2004). Identification and characterization of a *Neospora caninum* microneme-associated protein (NcMIC) that exhibits unique lactose-binding properties. *Infection and Immunity* **72**, 4791–4800.
- Kelm, S. and Schauer, R.** (1997). Sialic acids in molecular and cellular interactions. *International Reviews in Cytology* **175**, 137–240.
- Kitjaroentharn, A., Suthiphongchai, T. and Wilairat, P.** (2005). Effect of metalloprotease inhibitors on invasion of red blood cell by *Plasmodium falciparum*. *Acta Tropica* **97**, 5–9.
- Louie, K. and Conrad, P. A.** (1999). Characterisation of a cDNA encoding a subtilisin-like serine protease (NC-p65) of *Neospora caninum*. *Molecular and Biochemical Parasitology* **103**, 211–223.
- Louie, K., Nordhausen, R., Robinson, T. W., Barr, B. C. and Conrad, P. A.** (2002). Characterisation of *Neospora caninum* protease, NcSUB1 (NC-P65), with rabbit anti-N54. *Journal of Parasitology* **88**, 1113–1119.
- Lovett, J. L., Howe, D. K. and Sibley, L. D.** (2000). Molecular characterization of a thrombospondin-related anonymous protein homologue in *Neospora caninum*. *Molecular and Biochemical Parasitology* **107**, 33–43.
- Mauel, J.** (1996). Intracellular survival of protozoan parasites with special reference to *Leishmania* spp., *Toxoplasma gondii* and *Trypanosoma cruzi*. *Advances in Parasitology* **38**, 1–51.
- Mercier, C., Adjogble, K. D., Daubener, W. and Delauw, M. F.** (2005). Dense granules: are they key organelles to help understand the parasitophorous vacuole of all apicomplexa parasites? *International Journal for Parasitology* **35**, 829–849.
- Miller, S. A., Binder, E. M., Blackman, M. J., Carruthers, V. B. and Kim, K.** (2001). A conserved subtilisin-like protein TgSUB1 in microneme organelles of *Toxoplasma gondii*. *Journal of Biological Chemistry* **276**, 45341–45348.
- Monteiro, V. G., Soares, C. P. and de Souza, W.** (1998). Host cell surface sialic acid residues are involved in the process of penetration of *Toxoplasma gondii* into mammalian cells. *FEMS Microbiological Letters* **164**, 323–327.
- Müller, N., Vonlaufen, N., Gianinazzi, C., Leib, S. and Hemphill, A.** (2002). Application of real-time fluorescent PCR for quantitative assessment of *Neospora caninum* infections in organotypic slice cultures of rat central nervous system tissue. *Journal of Clinical Microbiology* **40**, 252–255.
- Naguleswaran, A., Cannas, A., Keller, N., Vonlaufen, N., Schares, G., Conraths, F. J., Bjorkman, C. and Hemphill, A.** (2001). *Neospora caninum* microneme protein NcMIC3: secretion, subcellular localization, and functional involvement in host cell interaction. *Infection and Immunity* **69**, 6483–6494.
- Naguleswaran, A., Cannas, A., Keller, N., Vonlaufen, N., Björkman, C. and Hemphill, A.** (2002). Vero cell surface proteoglycan interaction with the microneme protein NcMIC3 mediates adhesion of *Neospora caninum* tachyzoites to host cells unlike that in *Toxoplasma gondii*. *International Journal for Parasitology* **32**, 695–704.
- Naguleswaran, A., Müller, N. and Hemphill, A.** (2003). *Neospora caninum* and *Toxoplasma gondii*: a novel adhesion/invasion assay reveals distinct differences in tachyzoite-host cell interactions. *Experimental Parasitology* **104**, 149–158.
- Naguleswaran, A., Alaeddine, F., Guionaud, C., Vonlaufen, N., Sonda, S., Jenoe, P., Mevissen, M. and Hemphill, A.** (2005). *Neospora caninum* protein disulfide isomerase is involved in tachyzoite-host cell interaction. *International Journal for Parasitology* **35**, 1459–1472.
- Nok, A. J. and Rivera, W.** (2003). Characterisation of sialidase from *Entamoeba histolytica* and possible pathogenic role in amebiasis. *Parasitology Research* **89**, 302–307.
- Ochieng, J., Warfield, P. and Green, B.** (1995). Interactions of gelatinases with soluble and immobilized fetuin and asialofetuin. *Archives in Biochemistry and Biophysics* **322**, 250–255.
- Sonda, S., Fuchs, N., Gottstein, B. and Hemphill, A.** (2000). Molecular characterisation of a novel microneme antigen in *Neospora caninum*. *Molecular and Biochemical Parasitology* **108**, 39–51.
- Terkelsen, O. B., Jahnen-Dechent, W., Nielsen, H., Moos, T., Fink, E., Nawratil, P., Muller-Esterl, W. and Mollgard, K.** (1998). Rat fetuin: distribution of protein and mRBA in embryonic and neonatal tissues. *Anatomy and Embryology* **197**, 125–133.
- Tomley, F. M. and Soldati, D.** (2001). Mix and match modules: structure and function of microneme proteins in apicomplexan parasites. *Trends in Parasitology* **17**, 81–88.

- Vimr, E. and Lichtensteiger, C.** (2002). To sialylate, or not to sialylate: that is the question. *Trends in Microbiology* **10**, 254–257.
- Vonlaufen, N., Guetg, N., Naguleswaran, A., Muller, N., Bjorkman, C., Schares, G., von Blumroeder, D., Ellis, J. and Hemphill, A.** (2004). *In vitro* induction of *Neospora caninum* bradyzoites in Vero cells reveals differential antigen expression, localization, and host-cell recognition of tachyzoites and bradyzoites. *Infection and Immunity* **72**, 576–583.
- Wessel, D. and Fluegge, U. I.** (1984). A method for the quantitative recovery of protein in dilute solution in the presence of detergents and lipids. *Analytical Biochemistry* **138**, 141–143.