

Heart rate variability does not discriminate between different levels of haemodynamic responsiveness during surgical anaesthesia[†]

M. Luginbühl^{1*}, H. Yppärilä-Wolters², M. Rüfenacht¹, S. Petersen-Felix¹ and I. Korhonen²

¹Department of Anaesthesiology, University Hospital of Bern, University of Bern, Bern, Switzerland.

²VTT Technical Research Centre of Finland, Tampere, Finland

*Corresponding author: Department of Anaesthesiology, University Hospital, CH-3010 Bern, Switzerland.

E-mail: martin.luginbuehl@dkf.unibe.ch

Background. Hypnotic depth but not haemodynamic responsiveness is measured with EEG-based monitors. In this study we compared heart rate variability (HRV) in unstimulated patients and stimulation-induced HRV at different levels of anaesthesia.

Methods. A total of 95 ASA I or II patients were randomly assigned to five groups (Group 1: BIS 45(5), remifentanyl 1 ng ml⁻¹; Group 2: BIS 45(5), remifentanyl 2 ng ml⁻¹; Group 3: BIS 45(5), remifentanyl 4 ng ml⁻¹; Group 4: BIS 30(5), remifentanyl 2 ng ml⁻¹; Group 5: BIS 60(5), remifentanyl 2 ng ml⁻¹). A time- and frequency-domain analysis of the RR interval (RRI) from the electrocardiogram was performed. HRV before induction, before and after a 5 s tetanic stimulus of the ulnar nerve, and before and after tracheal intubation was compared between groups, between stimuli, and between responders to intubation [systolic arterial pressure (SAP) increase >20 mm Hg, a maximal heart rate (HR) after intubation >90 min⁻¹ or both] and non-responders (ANOVA).

Results. Induction of anaesthesia significantly lowered HR and HRV. Mean RRI before stimulation was higher in G3 than in G1, G2, and G4 ($P<0.001$), whereas the other HRV parameters were similar. Intubation induced a greater HRV response than tetanic stimulation. The mean RRI after intubation was lower in G3 compared with the other groups and the SD of the RRI after tetanic stimulation was lower in G3 compared with G5. Otherwise, unstimulated HRV and stimulation-induced HRV were similar in responders and non-responders.

Conclusion. HRV parameters discriminate between awake and general anaesthesia, are different after tracheal intubation and a 5 s ulnar nerve stimulation, but do not discriminate between different levels of haemodynamic responsiveness during surgical anaesthesia.

Br J Anaesth 2007; **98**: 728–36

Keywords: anaesthetics i.v., propofol; analgesics opioid, remifentanyl; heart, heart rate variability; monitoring, bispectral index; Poincaré analysis

Accepted for publication: February 25, 2007

Although several types of hypnotic state monitors are now available,^{1,2} there is no monitor measuring haemodynamic responsiveness to noxious stimulation during general anaesthesia. The EEG-based hypnotic state monitors do not provide parameters predictive of haemodynamic reaction or movement,³ although the EEG response to noxious stimulation may reflect the analgesic drug concentration.⁴ Anaesthetic drug concentrations, although inversely correlated with the probability of a haemodynamic response, do not allow reliable prediction in every individual subject because of the variation in anaesthetic drug requirements.

A short electrical stimulation of the ulnar nerve elicits a short vasoconstriction, which is measurable with pulse plethysmography⁵ or laser Doppler,⁶ and is suppressed by increasing plasma concentrations of alfentanil⁵ or sevoflurane.⁶ This concept of applying a defined sub-maximal noxious stimulus during general anaesthesia and using the pulse plethysmography signal as response failed to

[†]The data were presented in part at the annual meeting of the European Society of Anaesthesiologists June 5–8, 2004, in Lisbon.

improve the prediction of the haemodynamic response to tracheal intubation based on hypnotic depth and predicted effect site remifentanyl concentration in a recent randomized double-blinded study.⁷ The same stimulus also elicits a short increase followed by a short decrease in the heart rate (HR), as measured with RR intervals (RRI). HR variability (HRV) is known to discriminate between awake and sedated or anaesthetized patients.⁸

The purpose of this study was to compare linear and non-linear parameters of HRV in patients at different levels of haemodynamic responsiveness during surgical anaesthesia before and after two standardized noxious stimuli. We hypothesized that some HRV parameters would discriminate between haemodynamic responsiveness at different levels of surgical anaesthesia, which were defined by different BIS levels and different effect-site remifentanyl concentrations in a previous study.⁷ HRV data recorded during a previous study on pulse plethysmography,⁷ which was recently published in this journal, were evaluated.

Methods

Patients

Data of the same, 95 ASA I or II patients enrolled in the pulse plethysmography study⁷ were evaluated. Patients with cardiovascular disease (e.g. arterial hypertension or antihypertensive treatment, cardiac, cerebrovascular, or peripheral vascular disease), any relevant pulmonary liver, kidney, or central nervous system disease, diabetes mellitus, alcohol or drug abuse, patients with a difficult airway (i.e. Mallampati class 3 or higher), or patients unable to give informed consent were excluded.

Study plan

The study plan has been described previously in detail.⁷ Briefly, the premedicated patients (midazolam 7.5 mg orally 30 min before induction) were monitored with ECG, non-invasive blood pressure cuff, and pulse oximeter (Datex AS3 monitor, Datex-Ohmeda, Instrumentarium Corporation, Helsinki, Finland). A venous cannula was inserted in the non-dominant arm. An A-2000 XP-BIS monitor (BIS software version 3.3, Aspect Medical Systems, Nattick, CA, USA) was installed with the sensor placed frontally, according to the manufacturer's instructions.

Skin electrodes for electrical stimulation of the ulnar nerve were placed on the dominant arm 15 and 23 cm distal to the cubital fold,⁶ and connected to a Digitimer DS7 constant current stimulator with a Digitimer DG2 trigger generator (Digitimer LTD, Hertfordshire, UK) and a timer device (constructed in our laboratory) with the positive pole attached proximally.

Before induction, the patients were randomly assigned to five treatment groups differing in target BIS level, and in remifentanyl bolus and infusion rates using a stratified

randomization protocol. In Groups 1–3, a propofol target-controlled infusion (Diprifusor®, AstraZeneca, Grafenau, Switzerland) was titrated to achieve a BIS level of 45(5), whereas in Groups 4 and 5 the target BIS levels were 30(5) and 60(5), respectively. In Groups 1–3, remifentanyl was administered with an initial bolus of 0.2, 0.4, and 0.8 $\mu\text{g kg}^{-1}$ immediately followed by an infusion of 0.04, 0.08, and 0.2 $\mu\text{g kg}^{-1} \text{min}^{-1}$, respectively. In Groups 4 and 5, the remifentanyl bolus was 0.4 $\mu\text{g kg}^{-1}$ and the infusion rate was 0.08 $\mu\text{g kg}^{-1} \text{min}^{-1}$.

After recording of baseline data for 5 min, the selected bolus of remifentanyl was given by i.v., and the related remifentanyl infusion was started immediately, followed by a target-controlled infusion of propofol. The target plasma propofol concentration was adjusted to achieve and maintain the selected BIS level. Muscle relaxation was achieved with vecuronium 0.15 mg kg^{-1} i.v., injected after loss of consciousness. After the requested effect of vecuronium was verified by train-of-four (suppression of at least three of four twitches), a 5 s, 60 mA, 50 Hz, and 0.25 ms square-wave electrical stimulus^{7,9} was applied to the ulnar nerve. After arterial pressure and HR had returned to the pre-stimulation level, an experienced anaesthetist blinded to the anaesthetic drug doses and the BIS level performed orotracheal intubation. During the study, arterial pressure was measured non-invasively (oscillometric method) at 1 min intervals (arterial pressure cuff on the arm opposite the pulse oximeter probe). The study was terminated 5 min after tracheal intubation.

Data recording

Arterial pressure, HR, end-tidal CO_2 , and SpO_2 were recorded on a laptop computer every 10 s. The BIS values from the A2000 monitor were recorded every 5 s. The quality of visualization of the vocal cords according to Wilson and colleagues¹⁰ and the duration of intubation were recorded. Patients with prolonged intubation (>45 s) were excluded from the study. The infusion rate of remifentanyl was also recorded on a laptop computer every 10 s. The target plasma propofol concentration before tracheal intubation was recorded manually. The ECG signal was digitized at 128 Hz (A/D conversion card, National Instruments Corporation, Austin, TX, USA) for off-line analysis.

Time of induction was defined as the start of the propofol infusion, time of stimulations as start of tetanic stimulation, and time of intubation as start of laryngoscopy. Heart rate (beats min^{-1}), systolic and diastolic arterial pressure (mm Hg), and BIS values were extracted from the files, and the mean (sd) value was calculated for the 120 s periods before induction, before stimulation (with stable anaesthetic drug concentrations), and before laryngoscopy. The maximal HR, arterial pressure, and BIS in the 300 s after intubation were determined. An increase in SAP by >20 mm Hg compared with pre-laryngoscopy, a maximal

HR after intubation >90 or both were defined as a response to tracheal intubation.^{7–11} The effect-site remifentanyl concentration at stimulation and intubation was computed from the recorded dosing history using the pharmacokinetic and pharmacodynamic parameter set of Minto and colleagues.¹²

Analysis of HRV

The R-waves of the ECG were detected automatically. This automatic detection was later verified visually, and separated arrhythmia, misclassified beats or both were manually corrected. A beat-to-beat RR interval (RRI) signal was constructed as a series of time differences between the successive R-waves.

The recorded signals from the pre-induction (120–60 s before the start of propofol), the pre-stimulation (60–0 s before tetanic stimulation), the post-stimulation (0–60 s after stimulation onset), the pre-intubation (60–0 s before start of laryngoscopy), and the post-intubation (0–180 s after intubation) periods were analysed off-line.

Heart rate variability was quantified with time- and frequency-domain parameters.^{13–14} The computed time-domain parameters were mean RRI, RRI standard deviation (RRI sd), and root of mean squared difference between successive RRI (RMSSD). The frequency-domain parameters—low-frequency (LF) power (0.04–0.15 Hz), high-frequency (HF) power (0.15–0.4 Hz), the LF/HF ratio, spectral entropy of RRI (0.04–0.4 Hz), HF spectral entropy (0.12–0.4 Hz)—and the amplitude of RRI variability (RRI amplitude) were computed. The quantitative Poincaré analysis was carried out as described by Tulppo and colleagues.¹⁴ In this analysis, RRI is plotted on an x - y plane (Fig. 1) so that the current RRI (on the y -axis) is related to the previous RRI (on the x -axis). This analysis provides a qualitative way of detecting deterministic patterns in complex data. For quantitative analysis of the plot, the standard deviations of the Poincaré plot against the axes $y=x$ (sd1) and $y'=-x+2m$ (sd2), where m is the mean RRI during the epoch of interest, and their ratio




Fig 1 Poincaré plot of the heart rate (HR) response to tetanic stimulation. RR intervals (RRI) of the pre- and post-stimulation window.

(sd1/sd2) were calculated. sd1 mainly describes the fast beat-to-beat HRV, while sd2 describes slower components of HRV.¹⁴

Statistics

The randomization procedure has been described previously.⁷ The HRV values after the events (induction, stimulation, and intubation) were normalized to the pre-event values by dividing the post-event values by the pre-event values. The post-induction window corresponded to the pre-stimulation window.

A two-way analysis of variance on ranks was performed to detect the induction effect and the group effect on the normalized HRV parameters. Repeated measurements of two-way analysis of variance on ranks with multiple pairwise comparisons among groups (Holm–Sidak test) was performed on the normalized HRV parameters to detect a stimulus effect (tetanic stimulation of ulnar nerve, tracheal intubation) and the group effect. $P < 0.05$ was considered statistically significant. SigmaStat for Windows Version 3.0 (SPSS Inc., Chicago, IL, USA) was used for the statistical analyses.

The different event-induced HRV variables in responders and non-responders to tracheal intubation were compared with a Mann–Whitney rank test. A logistic regression analysis was performed to determine the predictive value on the responder status of the following variables: the BIS value, the predicted effect-site remifentanyl concentration, the predicted plasma propofol concentration, and the HRV parameters related to tetanic stimulation, which were significantly different between responders and non-responders.

Results

Patients

A total of 95 patients (57 women and 38 men) were enrolled. The characteristics of the five treatment groups were similar (Table 1). Two patients of Group 1 were excluded from data analysis because of serious arrhythmias throughout the study. Various additional subjects were excluded from the analysis of induction and stimulation (see below).

Anaesthetic drug concentrations, BIS levels, and response to intubation

The computed effect-site remifentanyl concentrations, the BIS levels before laryngoscopy, the SAP and mean arterial pressure (MAP), HRs before induction and before laryngoscopy, the quality of vocal cord visualization, and duration of laryngoscopy and intubation are presented in Table 2. The increase in SAP, MAP, and HR in response to tracheal intubation and thus the probability of a response to tracheal intubation (SAP increase >20 mm Hg or maximal

Table 1 Characteristics of the study population. The data are given as mean (SD) or numbers of patients. Groups were compared with one-way ANOVA and χ^2 as appropriate. SAP, systolic arterial pressure; DAP, diastolic arterial pressure; HR, heart rate

	Group 1	Group 2	Group 3	Group 4	Group 5	P-values
Age	38.2(8.9)	40.9(10.5)	39.1(10.1)	35.8(9.0)	40.5(11.0)	0.639
Gender (M/F)	8/9	7/12	7/12	8/11	8/11	0.991
Weight (kg)	70.2(13.6)	68.4(11.6)	69.3(13.2)	69.1(14.9)	74.9(17.1)	0.654
BMI	24.1(3.5)	23.9(3.8)	23.8(3.2)	22.9(3.1)	25.4(4.2)	0.335
ASA I/II	14/3	14/5	16/3	14/5	16/3	0.797
SAP ward (mm Hg)	120(9)	118(12)	118(14)	122(12)	120(12)	0.889
DAP ward (mm Hg)	79(8)	74(8)	75(9)	74(8)	76(10)	0.661
HR ward (min ⁻¹)	72(10)	75(6)	75(8)	74(12)	73(10)	0.904

Table 2 BIS level, predicted effect-site remifentanil concentrations, predicted plasma propofol concentrations, and haemodynamic response to intubation.⁷ Numbers are mean (SD). BIS, BIS level before tracheal intubation; Remifentanil, predicted effect-site remifentanil concentration (ng ml⁻¹); Propofol, predicted plasma propofol concentration (μ g ml⁻¹) before laryngoscopy; P response to intubation, probability of haemodynamic response defined as increase of SAP >20 mm Hg compared with pre-laryngoscopy, a maximal heart rate after intubation or both >90 min⁻¹; SAP, systolic arterial pressure; MAP, mean arterial pressure; HR, heart rate; SAP-, MAP-, and HR-increase, maximum value after intubation—mean value during 120 s before intubation. One-way ANOVA on ranks with multiple pairwise comparisons (Dunn's test): ¹ $P < 0.05$ compared with all other treatment groups. ² $P < 0.05$ compared with Group 5, ³ $P < 0.05$ compared with Groups 3 and 5, ⁴ $P < 0.05$ compared with Groups 2 and 3, ⁵ $P < 0.05$ compared with Group 3, ⁶ $P < 0.05$ compared with Groups 3 and 4, ⁷ $P < 0.05$ compared with Groups 2 and 4, ⁸ $P < 0.05$ compared with Group 3

	Group 1	Group 2	Group 3	Group 4	Group 5	P-values
Measured BIS	44(5)	41(6)	43(4)	31(6)	57(4) ¹	<0.001
Remifentanil effect site (ng ml ⁻¹)	1.1(0.1) ¹	2.1(0.3)	4.7(0.7) ¹	2.0(0.2)	2.1(0.4)	<0.001
Propofol	3.4(0.9) ²	3.2(0.9) ²	2.9(0.7)	4.0(0.8) ³	2.2(0.5)	<0.001
Laryngoscopy (Wilson ¹⁰ I/II/III)	11/2/0	15/4/0	12/6/1	14/3/0	13/5/1	0.795
Duration of intubation (s)	18.5(11.0)	16.3(4.9)	16.9(4.4)	16.4(4.9)	17.2(3.7)	0.840
SAP before induction (mm Hg)	129(18)	124(12)	116(31)	127(14)	124(12)	0.622
SAP before laryngoscopy (mm Hg)	105(7)	97(7)	96(8)	100(8)	98(14)	0.064
SAP increase (mm Hg)	30(19) ⁴	17(11)	11(11)	11(8)	22(13) ⁵	<0.001
MAP before induction (mm Hg)	95(10)	91(7)	91(10)	94(11)	93(8)	0.638
MAP before laryngoscopy (mm Hg)	75(7)	72(6)	71(7)	74(8)	74(11)	0.148
MAP increase (mm Hg)	26(12) ⁶	18(9)	11(11)	12(6)	21(13)	<0.001
HR before induction (min ⁻¹)	79(19)	79(17)	68(21)	78(16)	78(13)	0.234
HR before laryngoscopy (min ⁻¹)	63(7)	66(10)	57(7) ⁷	65(11)	61(7)	0.023
HR increase (min ⁻¹)	21(11) ⁸	20(10)	8(8)	9(7)	15(7)	<0.001
P response to intubation	0.77	0.47	0.05	0.18	0.52	0.03

HR >90 min⁻¹) were significantly different among groups (one-way ANOVA and χ^2 test $P < 0.001$, respectively).

HRV before and after induction: comparison among treatment groups

Because of bad data quality in the pre-induction window, eight additional patients of Groups 2–5 were excluded, leaving data of 85 patients for analysis (Tables 3 and 4). All HRV parameters (time domain and frequency domain) except for the ratio of Poincaré SD1/SD2 and LF/HF power were significantly different before and after induction (stable anaesthetic conditions, Tables 3 and 4). Except for mean RRI, which was significantly higher in Group 3 than in Groups 1, 2, and 4, reflecting the lower HR in the subjects with the highest remifentanil concentration, the HRV parameters after induction were similar among the groups.

Stimulation-induced HRV: stimulation and group effect

For the analysis of the effect of tetanic stimulation, no further subjects were excluded, so that data of 93 patients

were analysed. For the analysis of the effect of tracheal intubation, four additional subjects of Group 1 were excluded because of prolonged intubation (>45 s) and a further seven subjects were excluded because of insufficient data quality in the pre-intubation window. The effect of tracheal intubation on HRV was therefore analysed in 82 subjects (Tables 5 and 6).

Tetanic stimulation induced a smaller HRV response (RRI SD and Poincaré SD2) in Group 3 compared with Group 5, whereas spectral entropy was higher in Group 2 than in all other groups after tetanic stimulation (Tables 5 and 6).

Tracheal intubation induced a stronger HRV response than the 5 s tetanic stimulation of the ulnar nerve, in all parameters of HRV except for root mean squared SD of RRI and Poincaré SD1 (Tables 5 and 6). There was a significant interaction between group effect and stimulation effect, implying that the group effect was stimulation dependent. Mean RRI after tracheal intubation was significantly higher in Group 3 compared with Groups 1, 2, and 5, indicating that intubation induced less tachycardia in subjects with the highest remifentanil concentrations.

Table 3 Heart rate variability before and after induction: induction effect and group effect measured with time-domain parameters. Heart rate variability before induction of anaesthesia (Pre) and before tetanic stimulation of the ulnar nerve (after induction, under stable anaesthetic conditions, Post). Numbers are median (inter-quartile range). Mean RRI, mean RR interval; RRI sd, standard deviation of RRI; RMSSD, root mean-squared standard deviation of RRI; Poincaré sd1 and Poincaré sd2 (see Fig. 2 for details). Two-way ANOVA on ranks with multiple pairwise comparison among groups to detect induction effect and group effect (Holm–Sidak test). *Group effect: $P < 0.05$ compared with Groups 1, 2, and 4

	Induction effect	Group effect	Group 1 (n=16)		Group 2 (n=17)		Group 3 (n=18)		Group 4 (n=18)		Group 5 (n=16)	
			Pre	Post	Post	Post	Pre	Post	Pre	Post	Pre	Post
Mean	<0.001	0.25	814.1 (740.87, 872.58)	932.48 (877.61, 970.89)	837.95 (706.37, 903.19)	933.32 (813.85, 1012.27)	876.82 (810.34, 928.35)	1031.05 (954.06, 1113.95)*	774.37 (726.19, 858.76)	920.39 (818.79, 1012.2)	858.67 (714.95, 920.67)	981.65 (902.98, 1034.3)
RRI sd	<0.001	0.896	42.93 (24.07, 62.49)	21.69 (16.24, 32.88)	38.96 (30.31, 67.42)	20.37 (17.5, 30.26)	51.2 (26.45, 74.84)	21.08 (18.93, 25.64)	58.36 (37.09, 65.2)	23.26 (15.14, 25.23)	60.04 (32.76, 77.63)	18.74 (15.03, 25.6)
RMSSD	<0.001	0.722	29.86 (16.48, 35.4)	10.64 (7.74, 13.94)	29.08 (17.08, 42.31)	12.35 (6.97, 19.31)	31.95 (21.19, 48.01)	15.23 (10.07, 19.17)	38.2 (20.12, 60.37)	13.49 (8.95, 23.51)	33.15 (16.47, 53.41)	12.86 (9.39, 16.64)
Poincaré sd1	<0.001	0.724	21.26 (11.73, 25.2)	7.58 (5.52, 9.93)	20.7 (12.14, 30.11)	8.78 (4.96, 13.76)	22.74 (15.09, 34.22)	10.84 (7.15, 13.69)	27.13 (14.28, 42.94)	9.6 (6.36, 16.75)	23.6 (11.72, 38.02)	9.15 (6.69, 11.84)
Poincaré sd2	<0.001	0.924	55.93 (32.87, 85.42)	29.6 (22.15, 42.09)	50.15 (40.97, 90.8)	27.35 (23.76, 39.67)	64.5 (35.24, 100.7)	26.32 (24.85, 33.55)	73.55 (50.32, 82.77)	26 (20.65, 32.8)	80.18 (43.31, 105.83)	24.81 (20.05, 33.27)

Table 4 HRV before and after induction: induction effect and group effect measured with frequency-domain parameters. Heart rate variability before induction of anaesthesia (Pre) and before tetanic stimulation of the ulnar nerve (after induction, under stable anaesthetic conditions, Post). Numbers are median (inter-quartile range). LF power, low-frequency power of RRI variability (0.04–0.15 Hz); HF power, high-frequency power of RRI variability (0.15–0.4 Hz); LF/HF ratio, ratio of low- to high-frequency power; entropy, spectral entropy of RRI variability (0.04–0.4 Hz); HF entropy, high-frequency spectral entropy (0.12–0.4 Hz). Two-way ANOVA on ranks with multiple pairwise comparisons among groups to detect induction effect and group effect (Holm–Sidak test)

	Induction effect	Group effect	Group 1 (n=16)		Group 2 (n=17)		Group 3 (n=18)		Group 4 (n=18)		Group 5 (n=16)	
			Pre	Post	Pre	Post	Pre	Post	Pre	Post	Pre	Post
LF power	<0.001	0.464	57.54 (10.6, 89.1)	6.16 (2.26, 19.32)	44.69 (16.37, 72.62)	8.62 (3.36, 27.85)	47.32 (8.83, 214.01)	5.17 (3.37, 7.46)	85.09 (32.21, 144.5)	7.31 (3.28, 10.31)	120.57 (35.45, 228.51)	4.75 (3.94, 12.33)
HF power	<0.001	0.641	26.51 (2.38, 40.02)	1.98 (0.84, 3.49)	26.31 (3.08, 54.17)	3.88 (1.18, 7.54)	18.64 (7.72, 47.77)	4.61 (1.78, 9.41)	29.66 (6.65, 120.25)	5.77 (1.5, 7.86)	29.36 (7.73, 121.44)	2.17 (1.68, 3.89)
LF/HF ratio	0.068	0.462	2.91 (1.58, 6.99)	2.77 (0.95, 7.52)	1.97 (0.92, 4.98)	2.7 (1.45, 5.95)	2.0 (0.83, 8.77)	0.98 (0.43, 2.6)	3.55 (1.54, 7.67)	1.99 (1.07, 2.48)	6.64 (1.64, 9.82)	2.71 (1.28, 4.31)
Entropy	<0.001	0.679	0.78 (0.71, 0.87)	0.83 (0.79, 0.88)	0.77 (0.69, 0.85)	0.81 (0.71, 0.87)	0.79 (0.67, 0.89)	0.84 (0.77, 0.87)	0.76 (0.72, 0.81)	0.85 (0.81, 0.88)	0.73 (0.68, 0.8)	0.85 (0.79, 0.87)
HF entropy	<0.001	0.502	0.83 (0.75, 0.86)	0.86 (0.81, 0.9)	0.8 (0.73, 0.84)	0.84 (0.81, 0.88)	0.81 (0.75, 0.9)	0.84 (0.8, 0.89)	0.82 (0.76, 0.88)	0.87 (0.8, 0.92)	0.79 (0.77, 0.84)	0.86 (0.81, 0.88)

Table 5 Stimulation-induced HRV: stimulus and group effects, group–stimulus interaction in time-domain parameters. Numbers are median (inter-quartile range) of post-stimulation values normalized to pre-stimulation values (post-stimulation/pre-stimulation). TET, tetanic stimulation of the ulnar nerve ($n=94$); INT, tracheal intubation ($n=82$). Two-way ANOVA on ranks with stimulus type, group, and group–stimulus interaction as factors and normalized post-stimulation HRV variables as dependant variables. † P -values of two-way ANOVA on ranks for group effects, stimulus effects, and group–stimulus interaction. One-way ANOVA on ranks: group effects at each stimulus: * $P<0.05$ compared with Groups 1, 2, and 5; ** $P<0.05$ compared with Group 5. Significant values are highlighted bold-italic

	Group effect†	Stimulus effect†	G–S interaction†	Group 1 (TET, $n=17$) (INT, $n=12$)		Group 2 (TET, $n=19$) (INT, $n=18$)		Group 3 (TET, $n=19$) (INT, $n=18$)		Group 4 (TET, $n=9$) (INT, $n=16$)		Group 5 (TET, $n=19$) (INT, $n=18$)	
				TET	INT	TET	INT	TET	INT	TET	INT	TET	INT
Mean RRI	0.133	<0.001	0.038	1.015 (0.996, 1.035)	0.894 (0.854, 0.962)	1.017 (0.995, 1.025)	0.909 (0.832, 0.968)	1.007 (1.001, 1.02)	0.961 (0.946, 1.021)*	1.012 (1.004, 1.023)	0.938 (0.901, 0.997)	1.011 (0.996, 1.021)	0.913 (0.871, 0.947)
RRI SD	0.053	<0.001	0.042	1.671 (1.266, 2.281)	2.41 (1.647, 3.826)	1.566 (1.187, 2.107)	3.531 (2.148, 4.445)	1.354 (1.088, 1.704)**	2.278 (1.823, 3.589)	1.615 (1.459, 2.025)	1.943 (1.389, 2.593)	2.314 (1.702, 3.087)	2.448 (1.729, 2.873)
RMSSD	0.865	0.132	0.200	1.148 (0.948, 1.623)	1.128 (0.879, 1.428)	1.23 (1.008, 1.376)	1.018 (0.679, 1.665)	1.094 (0.995, 1.266)	1.233 (1.13, 1.729)	1.2 (1.126, 1.584)	0.985 (0.811, 1.163)	1.402 (0.99, 1.952)	1.082 (0.924, 1.854)
Poincaré SD1	0.876	0.123	0.200	1.148 (0.954, 1.63)	1.122 (0.875, 1.42)	1.23 (1.009, 1.375)	1.019 (0.674, 1.646)	1.091 (0.995, 1.263)	1.225 (1.124, 1.722)	1.197 (1.126, 1.58)	0.994 (0.806, 1.156)	1.39 (0.988, 1.954)	1.082 (0.919, 1.843)
Poincaré SD2	0.082	<0.001	0.034	1.776 (1.254, 2.357)	2.53 (1.653, 4.046)	1.629 (1.234, 2.217)	3.669 (2.276, 4.94)	1.373 (1.119, 1.835)**	2.63 (2.007, 3.884)	1.702 (1.521, 2.23)	2.026 (1.537, 2.868)	2.353 (1.903, 3.26)	2.581 (1.803, 3.036)

Table 6 Stimulation-induced HRV: stimulus and group effects, group–stimulus interaction in frequency-domain parameters. Numbers are median (inter-quartile range) of post-stimulation values normalized to pre-stimulation values (post-stimulation/pre-stimulation). TET, tetanic stimulation of the ulnar nerve ($n=94$); INT, tracheal intubation ($n=82$). Two-way ANOVA on ranks with stimulus type, group, and group–stimulus interaction as factors and normalized post-stimulation HRV variables as dependant variables. † P -values of two-way ANOVA on ranks for group effects, stimulus effects, and group–stimulus interaction. One-way ANOVA on ranks: group effects at each stimulus: * $P < 0.05$ compared with Groups 1, and 3–5. Significant values are highlighted bold-italic

	Group effect†	Stimulus effect†	G–S interaction†	Group 1 (TET, $n=17$) (INT, $n=12$)		Group 2 ((TET, $n=19$) (INT, $n=18$)		Group 3 (TET $n=19$) (INT, $n=18$)		Group 4 (TET, $n=19$) (INT, $n=16$)		Group 5 (TET, $n=19$) (INT, $n=18$)	
				TET	INT	TET	INT	TET	INT	TET	INT	TET	INT
LF power	0.695	<0.001	0.223	1.291 (0.606, 2.373)	19.118 (4.449, 225.848)	0.717 (0.277, 1.256)	23.188 (4.825, 56.846)	1.465 (0.589, 2.36)	12.059 (6.476, 31.412)	1.322 (0.683, 3.402)	7.232 (4.319, 20.284)	1.135 (0.644, 2.372)	11.118 (4.01, 34.3)
HF power	0.118	<0.001	0.274	1.001 (0.773, 1.518)	10.328 (4.052, 41.056)	0.772 (0.575, 1.082)	3.19 (0.93, 16.923)	0.689 (0.473, 1.498)	7.945 (3.728, 13.789)	1.393 (0.634, 1.935)	5.791 (2.316, 13.873)	1.222 (0.667, 1.908)	9.013 (1.874, 15.953)
LF/HF ratio	0.931	0.030	0.053	1.671 (0.399, 3.437)	2.502 (0.972, 7.426)	0.777 (0.304, 1.632)	3.583 (1.307, 8.267)	1.609 (0.625, 4.725)	1.491 (0.678, 2.964)	1.286 (0.575, 2.341)	1.094 (0.359, 8.643)	1.546 (0.514, 2.248)	1.516 (0.68, 3.251)
Entropy	0.984	0.017	0.025	0.952 (0.885, 1.095)	0.997 (0.868, 1.033)	1.056 (0.978, 1.183)*	0.892 (0.857, 0.933)	0.967 (0.923, 1.016)	0.942 (0.868, 1.004)	0.952 (0.911, 1.008)	0.941 (0.852, 0.994)	0.987 (0.903, 1.018)	0.945 (0.895, 1.019)
HF entropy	0.565	0.002	0.794	1.008 (0.932, 1.098)	0.998 (0.906, 1.023)	0.996 (0.955, 1.076)	0.927 (0.819, 1.005)	1.017 (0.911, 1.064)	0.927 (0.868, 1.011)	0.943 (0.89, 1.06)	0.923 (0.812, 0.969)	0.988 (0.958, 1.017)	0.957 (0.877, 1.026)




Fig 2 The grand average values of RRI before and after tetanic stimulation of the ulnar nerve are plotted for each group. The RRI are normalized to the mean RRI before stimulation (measured RRI/mean RRI before stimulation). The stimulation induced first a short decrease of RRI (=increase of heart rate), which was equal in all groups, followed by an increase (=decrease of heart rate). The difference in increase among the groups visible on the plot was not statistically significant.

The grand averages of RRI before and after tetanic stimulation and tracheal intubation in the five treatment groups are presented in Figures 2 and 3.

HRV: comparison between responders and non-responders to tracheal intubation

Of the 91 patients with uneventful intubation, 35 were responders and 56 were non-responders. Of 82 patients




Fig 3 The grand average values of RRI before and after tracheal intubation. The stimulation-induced RRI is similar among the five treatment groups. The RRI are normalized to the mean RRI before intubation (measured RRI/mean RRI before intubation). Intubation was followed by a decrease of RRI (=tachycardia), which was significantly smaller in Group 3 ($P < 0.05$).

included in the analysis of intubation-induced HRV, 31 were responders and 51 were non-responders. A total of 17 patients showed purely an arterial pressure response, 6 purely a HR response, and 8 a combined response.

HRV before ulnar nerve stimulation and before the start of laryngoscopy (i.e. unstimulated HRV) were similar in responders and non-responders to tracheal intubation (data not shown). The mean (inter-quartile range) normalized RRI after intubation in responders and non-responders was 0.867 (0.825–0.912) and 0.959 (0.914–1.001), respectively ($P < 0.001$; i.e. responders became more tachycardic than non-responders). The other parameters of HRV response were similar (data not shown).

The HRV induced by ulnar nerve stimulation was also similar in responders and non-responders (data not shown). In the logistic regression analysis the BIS level, the predicted effect-site remifentanyl concentration and the predicted plasma propofol concentration but not HRV variables were kept as independent parameters. The BIS level and the predicted effect-site remifentanyl concentration, but not the propofol concentration, were significant predictors of the responder status. The logistic regression equation to calculate the probability of response was therefore similar to a previously published equation.⁷

Discussion

Induction of general anaesthesia significantly and substantially changed HRV. Different levels of haemodynamic responsiveness during general anaesthesia expressed by the *post-hoc* probability of blood pressure and HR response were not reflected by any parameter of HRV, except that patients with high remifentanyl concentrations were more bradycardic. The sd of the RRI after induction, the Poincaré sd1 and sd2, and the RMSSD of RRI were lower and the entropy of RRI was higher after induction, reflecting a reduced HRV compared with the awake state. The LF and the HF power significantly decreased, with the LF/HF power ratio decreasing in Groups 3–5 and increasing in Groups 1 and 2. The stimulation-induced HRV response was similar among the groups, but dependent on the stimulus intensity; intubation induced a stronger HR response (reflected by most of the parameters) than the 5 s ulnar nerve stimulation. Responders and non-responders to tracheal intubation had similar HRV parameters except for a lower post-intubation HR in non-responders. The prediction of the response to tracheal intubation estimated from the pre-intubation BIS level and effect-site remifentanyl concentration was not improved by adding HRV parameters.

HRV reflects autonomic nervous system activity and is affected by anaesthesia. Several linear and non-linear methods have been used to assess HRV. The most frequently used linear method is the (frequency domain)

power spectral analysis of RRI, defining a LF band between 0.04 and 0.15 Hz, and a HF band between 0.15 and 0.4 or 0.5 Hz.¹⁵ Another linear method calculates RRI SD and RMSSD of HRV (time-domain parameters). Non-linear methods such as spectral entropy or Poincaré analysis, in contrast to linear methods, are better suited for non-stationary signals as may occur in patients under noxious stimulation.¹⁶

Our results on the induction effect confirm a previous study using power spectral analysis of HRV. In that study, total power, LF power (representing sympathetic tone) and HF power (representing parasympathetic tone), the LF/HF ratio, and the HR entropy were significantly reduced in propofol sedated spontaneously breathing patients (OAA/S 3) compared with awake patients.¹⁷

Other studies have reported partially conflicting results on induction-induced HRV. Induction with propofol (to a BIS level of 30) reduced HF but not LF power and LF/HF ratio, induction with sevoflurane reduced LF but not HF power, whereas entropy was reduced with both induction methods in patients under assisted spontaneous respiration.⁸ Thiopental or etomidate induction markedly reduced both LF power and HF power of HRV,¹⁸ whereas induction with midazolam and intermediate to high-dose fentanyl in cardiac surgery patients only decreased LF power independent of the dose.¹⁹ Conversely, midazolam sedation in ICU patients dose-dependently reduced HRV (frequency-domain parameters) correlating with the Ramsay score.²⁰ The clinical meaning of a change in HRV at induction of anaesthesia is limited, because the effect of induction agents is monitored clinically and eventually by EEG-derived parameters.

Because opioids predominantly reduce LF power,^{19 21} reflecting an increased parasympathetic tone, and because opioids are known to reduce HR increase after noxious stimulation, one might expect that a relative reduction in the sympathetic tone by opioids would reduce haemodynamic responsiveness to noxious stimulations. Our data show that HR increase after intubation was significantly lower in patients with predicted effect-site remifentanyl concentration of 4.7 ng ml⁻¹ (Group 3) compared with the other groups. Nevertheless, the stimulation-induced HRV response did not discriminate between the treatment groups or between responders and non-responders to tracheal intubation, except that RRI SD and Poincaré RRI SD² responses after tetanic stimulation were lower in Group 3 (highest remifentanyl concentration) compared with Group 5 (highest BIS level, intermediate remifentanyl concentration). In another study, intraoperative noxious stimulation did not affect power spectral HRV, and HRV did not correlate with the humoral stress response in adults.²² Conversely, painful stimulation induced a significant increase in RRI SD, LF power, and LF/HF ratio, and a decrease in approximate entropy in propofol-anaesthetized children.²³ Tachycardia induced by surgical skin incision in adults was reflected by a decrease in mean RRI, an

increase in RRI SD, and a decrease in HF power in patients under sevoflurane anaesthesia.¹⁶

Not surprisingly, we found a significant difference of HRV response induced by two stimuli of different intensity. Tracheal intubation, which is one of the strongest stimuli, elicited a significantly stronger HRV response than 5 s tetanic stimulation on the ulnar nerve, which is weaker than surgical skin incision. In a recent study, the RRI response to a 30 s tetanic stimulus discriminated between patients with a plasma remifentanyl concentration of 1 ng ml⁻¹²⁴ and those with 3 or 5 ng ml⁻¹. In our study, the RRI response induced by our 5 s tetanic stimulation was detectable, but did not discriminate between predicted effect-site remifentanyl concentrations of 1.1, 2.1, and 4.7 ng ml⁻¹ (Fig. 2). The RRI response to tracheal intubation (Fig. 3), however, discriminated between 4.7 ng ml⁻¹ and the lower remifentanyl concentrations. This suggests that only strong or even very strong stimuli allow discrimination between different remifentanyl concentrations or different states of haemodynamic responsiveness during general anaesthesia.

The prediction of the arterial pressure and HR response to tracheal intubation based on the BIS level and the predicted remifentanyl concentration was therefore not improved by HRV response induced by the tetanic stimulation. At present, the state of haemodynamic responsiveness during general anaesthesia is best assessed by the predicted plasma or effect-site opioid concentrations and using a depth of sedation monitor (e.g. BIS).

The definition of the responder status in this study may be a limitation. The pre-intubation arterial pressure was lower than the pre-induction arterial pressure (Table 2). The maximal SAP after intubation therefore was generally <160 mm Hg, also including blood pressure responders. Only an SAP >160 mm Hg was associated with an increased risk of stroke in heart surgery patients.²⁵ In the same study, only an HR >100 min⁻¹ was associated with an increased incidence of postoperative myocardial infarction, and an HR >120 min⁻¹ was even associated with increased mortality.²⁵ Only two patients in our study had a maximal HR >100 min⁻¹, and another two had a maximal SAP >160 mm Hg after intubation (all in Groups 1 and 2). The HRV parameters of these 'heavy responders' did not differ from the other responders, however (data not shown). Although our responder definition potentially limits the clinical relevance of the response in the current study, we could not change it, because our current study re-evaluates data from our previous study, and thus the same definition had to be used.⁷

We conclude that HRV in the unstimulated patient discriminates between awake and anaesthetized subjects, and stimulation-induced HRV response is affected by the stimulus intensity. The different HRV parameters are similar in a wide range of surgical anaesthesia with controlled ventilation and are therefore of limited value in assessing haemodynamic responsiveness.

Acknowledgement

This research was funded by the Department of Anaesthesiology, University Hospital of Bern, Switzerland

References

- 1 Rampil IJ. A primer for EEG signal processing in anesthesia. *Anesthesiology* 1998; **89**: 980–1002
- 2 Dutton RC, Smith WD, Rampil IJ, Chortkoff BS, Eger EI. Forty-hertz midlatency auditory evoked potential activity predicts wakeful response during desflurane and propofol anesthesia in volunteers. *Anesthesiology* 1999; **91**: 1209–20
- 3 Kalkman CJ, Drummond JC. Monitors of depth of anesthesia, quo vadis? *Anesthesiology* 2002; **96**: 784–7
- 4 Guignard B, Menigaux C, Dupont X, Fletcher D, Chauvin M. The effect of remifentanyl on the bispectral index change and hemodynamic responses after orotracheal intubation. *Anesth Analg* 2000; **90**: 161–7
- 5 Luginbühl M, Reichlin F, Sigurdsson GH, Zbinden AM, Petersen-Felix S. Prediction of the haemodynamic response to tracheal intubation: comparison of laser-Doppler skin vasomotor reflex and pulse wave reflex. *Br J Anaesth* 2002; **89**: 389–97
- 6 Shimoda O, Ikuta Y, Sakamoto M, Terasaki H. Skin vasomotor reflex predicts circulatory responses to laryngoscopy and intubation. *Anesthesiology* 1998; **88**: 297–304
- 7 Luginbühl M, Rufenacht M, Korhonen I, *et al.* Stimulation induced variability of pulse plethysmography does not discriminate responsiveness to intubation. *Br J Anaesth* 2006; **93**: 323–9
- 8 Kanaya N, Hirata N, Kurosawa S, Nakayama M, Namiki A. Differential effects of propofol and sevoflurane on heart rate variability. *Anesthesiology* 2003; **98**: 34–40
- 9 Petersen-Felix S, Zbinden AM, Fischer M, Thomson DA. Isoflurane minimum alveolar concentration decreases during anesthesia and surgery. *Anesthesiology* 1993; **79**: 959–65
- 10 Wilson ME, Spiegelhalter D, Robertson JA, Lesser P. Predicting difficult intubation. *Br J Anaesth* 1988; **61**: 211–6
- 11 Gan TJ, Glass PS, Windsor A, *et al.* Bispectral index monitoring allows faster emergence and improved recovery from propofol, alfentanil, and nitrous oxide anesthesia. BIS Utility Study Group. *Anesthesiology* 1997; **87**: 808–15
- 12 Minto CF, Schnider TW, Egan TD, *et al.* The influence of age and gender on the pharmacokinetics and pharmacodynamics of remifentanyl I. Model development. *Anesthesiology* 1997; **86**: 10–23
- 13 Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability: standards of measurement, physiological interpretation and clinical use. *Circulation* 1996; **93**: 1043–65
- 14 Tulppo MP, Makikallio TH, Takala TE, Seppanen T, Huikuri HV. Quantitative beat-to-beat analysis of heart rate dynamics during exercise. *Am J Physiol* 1996; **271**: H244–252.
- 15 Pomfrett CJ. Heart rate variability, BIS and 'depth of anaesthesia'. *Br J Anaesth* 1999; **82**: 659–62
- 16 Seitsonen ER, Korhonen IK, van Gils MJ, *et al.* EEG spectral entropy, heart rate, photoplethysmography and motor responses to skin incision during sevoflurane anaesthesia. *Acta Anaesthesiol Scand* 2005; **49**: 284–92
- 17 Win NN, Fukayama H, Kohase H, Umino M. The different effects of intravenous propofol and midazolam sedation on hemodynamic and heart rate variability. *Anesth Analg* 2005; **101**: 97–102
- 18 Latson TW, McCarroll SM, Mirhej MA, *et al.* Effects of three anesthetic induction techniques on heart rate variability. *J Clin Anesth* 1992; **4**: 265–76
- 19 Zickmann B, Hofmann HC, Pottkamper C, *et al.* Changes in heart rate variability during induction of anesthesia with fentanyl and midazolam. *J Cardiothorac Vasc Anesth* 1996; **10**: 609–13
- 20 Haberthur C, Lehmann F, Ritz R. Assessment of depth of midazolam sedation using objective parameters. *Intensive Care Med* 1996; **22**: 1385–90
- 21 Michaloudis D, Kochiadakis G, Georgopoulou G, *et al.* The influence of premedication on heart rate variability. *Anaesthesia* 1998; **53**: 446–53
- 22 Ledowski T, Bein B, Hanss R, *et al.* Neuroendocrine stress response and heart rate variability: a comparison of total intravenous versus balanced anesthesia. *Anesth Analg* 2005; **101**: 1700–5
- 23 Toweill DL, Kovarik WD, Carr R, *et al.* Linear and nonlinear analysis of heart rate variability during propofol anesthesia for short-duration procedures in children. *Pediatr Crit Care Med* 2003; **4**: 308–14
- 24 Rantanen M, Yppärilä-Wolters H, Yli-Hankala A, Kymäläinen M, Korhonen I. Tetanic stimulus of the ulnar nerve as a predictor of heart rate response to skin incision in propofol-remifentanyl anaesthesia. *Eur J Anaesthesiol* 2006; **23**(Suppl 37): pA95
- 25 Reich DL, Bodian CA, Krol M, *et al.* Intraoperative hemodynamic predictors of mortality, stroke, and myocardial infarction after coronary artery bypass surgery. *Anesth Analg* 1999; **89**: 814–22