

Marcel Arnold
Michael Halpern
Niklaus Meier
Urs Fischer
Tobias Haefeli
Liliane Kappeler
Caspar Brekenfeld
Heinrich P. Mattle
Krassen Nedeltchev

Age-dependent differences in demographics, risk factors, co-morbidity, etiology, management, and clinical outcome of acute ischemic stroke

Received: 2 September 2007
Received in revised form: 26 February 2008
Accepted: 26 March 2008
Published online: 28 July 2008

M. Arnold, MD (✉) · M. Halpern · N. Meier
· U. Fischer · T. Haefeli · L. Kappeler ·
H. P. Mattle · K. Nedeltchev
Dept. of Neurology
University Hospital
Inselspital
Freiburgstrasse
3010 Berne, Switzerland
Tel.: +41-31/632-3332
Fax: +41-31/632-9679
E-Mail: marcel.arnold@insel.ch

C. Brekenfeld
Dept. of Diagnostic and Interventional
Neuroradiology
Inselspital
University of Bern
Bern, Switzerland

Abstract *Background* Comparisons between younger and older stroke patients including co-morbidities are limited. *Methods* Prospective data of consecutive patients with first ever acute ischemic stroke were compared between younger (≤ 45 years) and older patients (> 45 years). *Results* Among 1004 patients, 137 (14%) were ≤ 45 years. Younger patients were more commonly female (57% versus 34%; $p < 0.0001$), had a lower frequency of diabetes (1% versus 15%; $p < 0.0001$), hypercholesterolemia (26% versus 56%; $p < 0.0001$), hypertension (19% versus 65%; $p < 0.0001$), coronary heart disease (14% versus 40%; $p < 0.0001$), and a lower mean Charlson co-morbidity index (CCI), (0.18 versus 0.84; $p < 0.0001$). Tobacco use was more prevalent in the young (39% versus 26%; $P < 0.0001$). Large artery disease (2% versus 21%; $p < 0.0001$), small artery disease (3% versus 12%; $p = 0.0019$) and atrial fibrillation (1% versus 17%; $p = 0.001$) were less common in young pa-

tients, while other etiologies (31% versus 9%; $p < 0.0001$), patent foramen ovale or atrial septal defect (44% versus 26%; $p < 0.0001$), and cervical artery dissection (26% versus 7%; $p < 0.0001$) were more frequent. A favorable outcome (mRS 0 or 1) was more common (57.4% versus 46.9%; $p = 0.023$), and mortality (5.1% versus 12%; $p = 0.009$) was lower in the young. After regression analysis, there was no independent association between age and outcome ($p = 0.206$) or mortality ($p = 0.073$). Baseline NIHSS score ($p < 0.0001$), diabetes ($p = 0.041$), and CCI ($p = 0.002$) independently predicted an unfavorable outcome. *Conclusions* Younger patients were more likely to be female, had different risk factors and etiologies and fewer co-morbidities. There was no independent association between age and clinical outcome or mortality.

Key words stroke · outcome · mortality

Introduction

Stroke in young adults is rare. The majority of strokes in the younger age group are cerebral infarcts, and their etiologies are manifold and often remain undetermined [1–3].

Clinical functional outcome has been considered

more favorable and survival rates are higher in younger than in older individuals, but severe and fatal strokes may occur at any age and co-morbidities have not been systematically analyzed in previous studies [1].

In recent years, new insights in stroke epidemiology, aetiology and prognosis in young stroke patients have been gained [4–7]. However, comparisons between younger and older patients with ischemic stroke includ-

ing risk factors, extensive diagnostic investigations to determine etiology, management, co-morbidities, and outcomes are still limited. Therefore we aimed to compare younger and older patients with first-ever ischemic strokes in a large cohort of consecutive patients.

Methods

The study sample included consecutive patients older than 16 years who suffered a first ever acute ischemic stroke and had been admitted to The Berne Stroke Unit from January 2000 to July 2006. Our University Hospital-based Stroke Unit has a catchment population of about 1.5 million.

Data were collected prospectively and entered continuously in the hospital-based Bernese stroke database.

Stroke was defined according to the World Health Organization criteria as an acute focal neurological deficit with symptoms lasting longer than 24 hours [8]. Clinical data of some patients were reported previously [3].

All patients underwent a standard investigation protocol immediately upon admission, including blood tests (determination of red cell, white cell and platelet counts, prothrombin time, activated partial thromboplastin time, serum electrolytes, glucose, urea, creatinine, transaminases, and cholesterol levels), 12-lead ECG, cranial computed tomography (CT) and/or conventional magnetic resonance imaging (MRI) (1.5 Tesla system providing axial T1, T2, and proton density weighted images, and gadolinium-enhanced T1-weighted images). All patients were investigated by neurovascular ultrasound or magnetic resonance angiography (MRA) or digital subtraction angiography (DSA) of the brain supplying arteries upon admission or during their hospital stay. The following ancillary investigations were carried out at the discretion of the treating stroke neurologist: a search for vasculitis (including C reactive protein, erythrocyte sedimentation rate, antinuclear antibodies, and antiphospholipid antibodies), transthoracic echocardiography, transesophageal echocardiography, 24-hour ECG monitoring, cervical MRI with fat suppression technique, diffusion-weighted and perfusion-weighted MRI. We assessed etiologic stroke subtype according to the TOAST criteria (trial of ORG 10172) [9]. The frequency of atrial fibrillation, cervical artery dissection, foramen ovale or atrial septal defect and atrial septal aneurysm (in patients undergoing TEE), coagulopathy and vasculitis were analyzed separately.

In addition infarctions were categorized according to the Oxfordshire Community Stroke Project (OCSP) classification [10]. The following risk factors were assessed: sex, hypertension (defined by pre-admission history and medical records), diabetes mellitus (fasting venous plasma glucose values ≥ 7.0 mmol/l on at least two separate occasions, and/or ≥ 11.1 mmol/l 2 hours after intake of 75 g of oral glucose and on one other occasion during the 2 hour test, or pre-admission history of diabetes), current cigarette smoking, hypercholesterolemia (total venous plasma cholesterol concentration > 5 mmol/l), and coronary heart disease.

Pre-existing co-morbid conditions were assessed using the Charlson Co-morbidity Index (CCI) [11]. The term "myocardial infarction" was replaced by "coronary heart disease" proved by coronary angiography or electrocardiography [12]. The severity of the neurological deficit was assessed at admission by a neurologist using the National Institutes of Health Stroke Scale (NIHSS) [13].

Antithrombotic therapy and secondary prevention was performed according to a standardized protocol similar to international guidelines [14, 15].

Clinical follow-up information of ischemic stroke patients was obtained by neurological examination (n = 307) or a structured telephone interview (n = 680) by different neurologists 3 months after stroke onset. The modified Rankin scale score (mRS) was recorded [16].

Seventeen patients (1.7%) were lost to follow-up because they had moved or were living in foreign countries.

Statistical analysis

Statistical analysis was performed with SPSS 10 for Macintosh statistical software (2001; SPSS Inc). The following variables were analyzed and compared between younger (≤ 45 years) and older patients (> 45 years): gender; smoking; hypertension; diabetes mellitus; hypercholesterolemia; family history of stroke; NIHSS score on admission; Charlson Co-morbidity Index; stroke etiology; intravenous and intra-arterial thrombolysis; antithrombotic treatment; overall frequency of transesophageal echocardiography (TEE), transthoracic echocardiography (TTE), and 24-hour ECG; duration of hospital stay; discharge to home; 3 months functional clinical outcome; and mortality.

For comparison of outcome we divided patients into two groups with favorable (mRS score 0 to 1) and non-favorable (mRS score 2 to 6) outcome. The χ^2 test was performed for cross tabulation. Mann-Whitney U test was used for non-categorical variables. Logistic regression analysis with a forward stepwise method was performed including the NIHSS score on admission and the variables that showed significant associations or trends ($p < 0.20$) with age when tested with univariate analyses, to determine an independent association between age and functional outcome or mortality.

Results

Among 1004 patients with acute ischemic stroke, 137 (14%) younger patients (≤ 45 years) and 867 (86%) older patients (> 45 years) were identified.

The mean age was 35.8 years (SD 7.3) in the younger and 64.7 years (SD 9.9) in the older patients.

The percentage of women was higher in younger than in older patients (57% versus 34%; $p < 0.0001$). Risk factors, stroke severity on admission and co-morbidities are shown in Table 1: Younger patients had a lower overall frequency of diabetes mellitus (1% versus 15%; $p < 0.0001$), hypercholesterolemia (26% versus 56%; $p < 0.0001$), hypertension (19% versus 65%; $p < 0.0001$), and coronary heart disease (14% versus 40%; $p < 0.0001$) but were more often current smokers (39% versus 26%;

Table 1 Demographics, risk factors, stroke severity, and co-morbidity in younger (< 45 years) and older (> 45 years) patients

| Characteristic | Younger patients n (%) | Older patients n (%) | P-value |
|---------------------------------------|---------------------------|-------------------------|--------------|
| No. of patients | 137 (14) | 867 (86) | |
| Female sex | 78 (57) | 291 (34) | $p < 0.0001$ |
| Family history of stroke | 24 (22) | 183 (28) | $p = 0.155$ |
| Diabetes mellitus | 2 (1) | 128 (15) | $p < 0.0001$ |
| Current smoking | 52 (39) | 221 (26) | $p = 0.002$ |
| Hypertension | 26 (19) | 562 (65) | $p < 0.0001$ |
| Hypercholesterolemia | 34 (26) | 474 (56) | $p < 0.0001$ |
| Coronary heart disease | 19 (14) | 347 (40) | $p < 0.0001$ |
| Mean NISS score on admission (SD) | 8.5 (7.7) | 8.6 (7.2) | $p = 0.845$ |
| Mean Charlson Co-morbidity Index (SD) | 0.18 (0.45) | 0.84 (1.36) | $p < 0.0001$ |

$P < 0.0001$) than older patients. The NIHSS score on admission did not differ between younger and older patients (mean 8.5 (SD 7.7) versus 8.6 (SD 7.2; $p = 0.845$). Mean CCI was lower in the younger age group (0.18 versus 0.84; $p < 0.0001$).

Younger patients more often underwent TEE [(98 patients (72 %) versus 442 (56 %); $p < 0.0001$)] and less frequently TTE [15 (11 %) versus 174 (20 %) patients; $p = 0.011$]. A 24-hour ECG was performed in 105 (77 %) younger and in 704 (81 %) older patients ($p = 0.21$).

Strokes due to large artery disease (2 % versus 21 %; $p < 0.0001$) and small artery disease (3 % versus 12 %; $p = 0.0019$) were less frequent, and strokes due to other etiologies (31 % versus 9 %; $p < 0.0001$) were more common in the younger age group. Patent foramen ovale or atrial septal defect (44 % versus 26 %; $p < 0.0001$), and cervical artery dissection were more common (26 % versus 7 %; $p < 0.0001$), and atrial fibrillation less frequent (1 % versus 17 %; $p = 0.001$) among younger patients. The stroke etiology remained undetermined in 27 % of the younger and in 25 % of the older patients (Table 2).

There was no significant association between age and OCSF categories ($p = 0.120$). However, a trend towards a higher frequency of lacunar syndromes (16 % versus 12 %; 0.250) and total anterior circulation syndromes (16 % versus 9 %; $p = 0.062$) and a lower frequency of partial anterior circulation syndromes (45 % versus 53 %; $p = 0.098$) in the older patients was observed (Table 2).

Twenty-eight patients (20 %) ≤ 45 years and 235 patients (27 %) > 45 years were treated with local intra-

Table 3 Three-month clinical outcome in 136 younger (<45 years) and 851 older (>45 years) patients with ischemic stroke

| Modified Rankin Scale Score | Younger patients n (%) | Elderly patients n (%) | P-value univariate | P-value Regression analyses |
|-----------------------------|------------------------|------------------------|--------------------|-----------------------------|
| 0 | 45 (33.1) | 195 (22.9) | 0.023 | 0.258 |
| 1 | 33 (24.3) | 204 (24.0) | | |
| 2 | 28 (20.6) | 168 (19.7) | | |
| 3 | 12 (8.8) | 102 (12.0) | | |
| 4 | 5 (3.7) | 74 (8.7) | | |
| 5 | 6 (4.4) | 6 (0.7) | | |
| 6 | 7 (5.1) | 102 (12.0) | 0.009 | 0.051 |

arterial thrombolysis ($p = 0.099$), and 6 (4 %) younger and 20 (2 %) older patients with intravenous thrombolysis ($p = 0.156$).

Stroke treatment and prevention included aspirin in 84 younger (61 %) and 572 (66 %) older patients ($p = 0.29$). Twenty-six older patients received clopidogrel (3 %). Heparin in therapeutic dose and/or warfarin was given to 53 (39 %) younger and 269 (31 %) older patients ($p = 0.074$).

Statins were given to 28 (20 %) younger and 395 (46 %) older patients ($p < 0.0001$).

The duration of hospital stay (10.3 (SD 7.5) days in the young versus 9.9 (SD 6.5) days in the elderly; $p = 0.154$) and the frequency of discharge at home (41 % in the young versus 35 % in the elderly; $p = 0.21$) were similar in both groups

At three months 107 (78 %) younger and 631 (73 %) older patients were living at home ($p = 0.19$). At univari-

Table 2 Stroke etiology according to TOAST classification and Oxfordshire Community Stroke Project Subtype in younger (<45 years) and older (>45 years) patients

| Characteristic | Younger patients n (%) | Older Patients n (%) | P-value |
|---------------------------------------|------------------------|----------------------|----------|
| Atrial fibrillation | 2 (1) | 147 (17) | < 0.0001 |
| PFO or atrial septal defect | 43 (44) | 129 (26) | < 0.0001 |
| Atrial septal aneurysm | 5 (5) | 27 (6) | 0.44 |
| Cervical artery dissection | 35 (26) | 62 (7) | < 0.0001 |
| Coagulopathy | 1 (1) | 2 (0.2) | 0.99 |
| Vasculitis | 2 (1.5) | 4 (0.5) | 0.16 |
| TOAST criteria | | | |
| Large artery disease | 3 (2) | 181 (21) | < 0.0001 |
| Cardioembolic | 51 (37) | 284 (33) | 0.30 |
| Small artery disease | 4 (3) | 101 (12) | 0.0019 |
| Other determined etiology | 42 (31) | 78 (9) | < 0.0001 |
| Undetermined, complete examination | 37 (27) | 214 (25) | 0.56 |
| Undetermined, incomplete examination | 0 (0) | 3 (0.3) | 0.33 |
| More than one cause | 0 (0) | 6 (0.7) | 0.33 |
| OCSF criteria | | | |
| Total anterior circulation syndrome | 13 (9) | 135 (16) | 0.062 |
| Partial anterior circulation syndrome | 72 (53) | 390 (45) | 0.098 |
| Lacunar syndrome | 17 (12) | 141 (16) | 0.250 |
| Posterior circulation syndrome | 35 (26) | 201 (23) | 0.544 |

PFO patent foramen ovale; p the difference between subgroups by χ^2 test or Mann-Whitney-test; TOAST trial of ORG 10172 in acute stroke treatment; OCSF Oxfordshire Community Stroke Project

ate analysis, younger patients were more likely to have a favorable outcome (modified Rankin scale Score ≤ 1) after 3 months (57% versus 47%; $p=0.023$) and had a lower mortality (5% versus 12%; $p=0.009$).

After regression analysis, there was no independent association between age and outcome ($p=0.206$) or mortality ($p=0.073$). Higher baseline NIHSS score ($p<0.0001$), higher CCI ($p=0.002$) and diabetes ($p=0.041$) were the only independent predictors of an unfavorable clinical functional outcome. Mortality was independently associated with higher baseline NIHSS score ($p<0.0001$), higher CCI ($p=0.001$), diabetes ($p=0.008$), cardioembolic etiology ($p=0.001$) or stroke due to other etiology ($p=0.015$).

Discussion

In this hospital-based study the overall percentage of younger patients ≤ 45 years of age was 14%. The higher percentage of younger patients compared with population-based studies reporting less than 5% young stroke rates probably reflects a selection bias [1, 17]. In Switzerland, younger patients are more likely to be admitted to a stroke unit.

The percentage of women was higher in younger than in older patients (57% versus 34%).

These age-dependent gender differences are difficult to explain and may be partly due to higher stroke incidences in men in the sixth and seventh decade [18]. Another reason may be that elderly women are less likely to be admitted to a stroke unit than elderly men because of a gender-based selection bias [19]. A sex-based selection bias in the young is unlikely because in Switzerland young adults with ischemic stroke are usually referred to a stroke center independent of their sex.

Hypertension, diabetes, hypercholesterolemia, and coronary heart disease were more common among older patients. This reflects the higher prevalence of these risk factors among older subjects in the general population. Only smoking was more frequent among young stroke patients. The fact that more than one third of the younger group were current smokers indicates that smoking is a particularly important modifiable risk factor in younger subjects. This emphasizes the need to enforce preventive strategies to target behavioral risk factors in this age group.

One of the strengths of this study is a complete evaluation including echocardiography and 24-hour ECG to determine stroke etiology in almost all patients. Three fourths of the younger and more than half of the older patients underwent TEE. This explains the lower frequency of stroke with undetermined etiology compared with other series [20–22]. The etiology of stroke differed greatly between younger and older patients. Cardiac em-

bolism and other determined etiologies such as cervical internal carotid and vertebral artery dissections were the most common causes in younger patients. These results are in agreement with previous studies [2, 23]. Atrial fibrillation was more frequent in older patients as reported previously [24]. PFO or atrial septal defect were detected by TEE in 44% among younger and in only 26% among older patients. These findings support the current hypothesis that a right to left shunt may play a more important role in the pathogenesis of stroke in the young, whereas other etiologies such as atrial fibrillation, large and small vessel disease are the principal causes of stroke in older patients.

Previous reports on clinical functional outcome in ischemic stroke in the young are contradictory. In a French study, 86.8% of young adults were independent 3 years after ischemic stroke [2]. In contrast, functional outcome was worse in other studies reporting 37% to 47% moderate or severe disabilities in young ischemic stroke patients [1, 25]. In our study, a favorable outcome (mRS 0 or 1) was more common (57.4% versus 46.9%), and case fatality (5.1% versus 12%) was lower in the young compared with older patients. The low mortality rates in our young adults is in agreement with previous studies reporting case fatality rates ranging from 4.9% to 7.7% in young stroke patients [1, 2]. However, after regression analysis, there was no independent association between age and clinical outcome or mortality in our patients. Higher baseline NIHSS score, higher CCI and diabetes were the only independent predictors of an unfavorable outcome. These results emphasize the need to analyze co-morbidities such as the CCI in stroke studies aiming to predict clinical outcome or mortality.

The main limitation of this hospital-based study is a selection bias. Our cohort had a lower mean age than patients in population-based studies. However, in contrast to most population-based studies the strength of this study is a more extensive diagnostic investigation including neurovascular ultrasound or MRA of the brain supplying arteries, echocardiography and 24-hour ECG in most patients.

In conclusion, in this hospital-based study with a potential selection bias, young adult stroke patients were more likely to be female than older patients and had different stroke etiologies and vascular risk factors. More than one third of the young stroke patients were current smokers. Co-morbidities were less common in the young. After correction for other variables including CCI, there was no independent association between age and clinical outcome or mortality.

■ **Acknowledgments** We thank Pietro Ballinari, PhD, for statistical advice. We wish to thank Dr. Andros Tofield, MD and Mr. Fred Buerki for their proofreading and editing.

References

1. Rozenthul-Sorokin N, Ronen R, Tamir A, et al. (1996) Stroke in the young in Israel. Incidence and outcomes. *Stroke* 27:838–841
2. Leys D, Bandu L, Henon H, Lucas C, Mounier-Vehier F, Rondepierre P, Godefroy O (2002) Clinical outcome in 287 consecutive young adults (15 to 45 years) with ischemic stroke. *Neurology* 59:26–33
3. Nedeltchev K, Auf der Maur T, Georgiadis D, Arnold M, Caso V, Mattle HP, Schroth G, Remonda L, Sturzenegger M, Fischer U, Baumgartner RW (2005) Ischemic stroke in young adults: predictors of outcome and recurrence. *J Neurol Neurosurg Psychiatry* 76: 2191–2195
4. Neau JP, Ingrand P, Mouille-Brachet C, et al. (1998) Functional recovery and social outcome after cerebral infarction in young adults. *Cerebrovasc Dis* 8:296–302
5. Marini C, Totaro R, Carolei A (1999) Long-term prognosis of cerebral ischemia in young adults. National research council study group on stroke in the young. *Stroke* 30:2320–2325
6. Naess H, Nyland HI, Thomassen L, et al. (2002) Incidence and short-term outcome of cerebral infarction in young adults in western Norway. *Stroke* 33:2105–2108
7. Jacobs BS, Boden-Albala B, Lin IF, et al. (2002) Stroke in the young in the northern Manhattan stroke study. *Stroke* 33:2789–2793
8. Aho K, Harmsen P, Hatano S, et al. (1980) Cerebrovascular disease in the community: results of a WHO collaborative study. *Bull WHO* 58:113–130
9. Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE 3rd, and the TOAST investigators (1993) Classification of subtype of acute ischemic stroke: definitions for use in a multicenter clinical trial. TOAST. Trial of Org 10172 in Acute Stroke Treatment. *Stroke* 24:35–41
10. Bamford J, Sandercock P, Dennis M, Burn J, Warlow C (1991) Classification and natural history of clinically identifiable subtypes of cerebral infarction. *Lancet* 22(337):1521–1526
11. Charlson ME, Pompei P, Ales KL, MacKenzie CR (1987) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 40: 373–383
12. Fischer U, Arnold M, Nedeltchev K, Schoenenberger RA, Kappeler L, Hollinger P, Schroth G, Ballinari P, Mattle HP (2006) Impact of comorbidity on ischemic stroke outcome. *Acta Neurol Scand* 113:108–113
13. Brott T, Adams HP Jr, Olinger CP, Marler JR, Barsan WG, Biller J, Spilker J, Holleran R, Eberle R, Hertzberg V, Roorick M, Moomaw CJ, Walker M (1989) Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 20:846–870
14. Toni D, Chamorro A, Kaste M, Lees K, Wahlgren NG, Hacke W, for the EUSI Executive Committee and the EUSI Writing Committee (2004) Acute Treatment of Ischaemic Stroke. European Stroke Initiative. *Cerebrovasc Dis* 17(Suppl 2):30–46
15. Leys D, Kwicinski H, Bogousslavsky J, Bath P, Brainin M, Diener HC, Kaste M, Sivenius J, Hennerici MG, Hacke W, for the EUSI Executive Committee and the EUSI Writing Committee (2004) Prevention. European Stroke Initiative. *Cerebrovasc Dis* 17(Suppl 2):15–29
16. Van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, von Gijn J (1988) Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 19:604–607
17. Nencini P, Inzitari D, Baruffi MC, et al. (1988) Incidence of stroke in young adults in Florence, Italy. *Stroke* 19: 977–981
18. Johansson B, Norrving B, Lindgren A (2000) Increased Stroke Incidence in Lund-Orup, Sweden, Between 1983 to 1985 and 1993 to 1995. *Stroke* 31: 481–486
19. Kapral MK, Fang J, Hill MD, Silver F, Richards J, Jaigobin C, Cheung AM (2005) Investigators of the Registry of the Canadian Stroke Network. Sex differences in stroke care and outcomes: Results from the registry of the Canadian Stroke Network. *Stroke* 36: 809–814
20. Adams HP Jr, Kapelle LJ, Biller J, Gordon DL, Love BB, Gomez F, Heffner RN (1995) Ischemic stroke in young adults: experience in 329 patients in the Iowa Registry of Stroke in Young Adults. *Arch Neurol* 52:491–495
21. Pohjasvaara T, Erkinjuntti T, Vataja R, Kaste M (1997) Comparison of stroke features and disability in daily life in patients with ischemic stroke aged 55 to 70 and 71 to 85 years. *Stroke* 28: 729–735
22. Lipska K, Sylaja PN, Sarma SP, Kutty VR, Thankappan KR, Vasan RS, Radhakrishnan K (2007) Risk factors for acute ischaemic stroke in young adults in South India. *J Neurol Neurosurg Psychiatry* 78:959–963
23. Kristensen B, Malm J, Carlberg B, et al. (1997) Epidemiology and Etiology of Ischemic Stroke in Young Adults Aged 18 to 44 Years in Northern Sweden. *Stroke* 28:1702–1709
24. Zeiler K, Siostrzonek P, Lang W, et al. (1992) Different risk factor profiles in young and elderly stroke patients with special reference to cardiac disorders. *J Clin Epidemiol* 45:1383–1389
25. Marini C, Totaro R, De Santis F, Ciancarelli I, Baldassarre M, Carolei A (2001) Stroke in Young Adults in the Community-Based L'Aquila Registry. Incidence and Prognosis. *Stroke* 32: 52–56