

# Index of Complexity, Outcome and Need scored on plaster and digital models

A. C. Veenema\*, C. Katsaros\*\*\*, S. C. Boxum\*\*, E. M. Bronkhorst\* and A. M. Kuijpers-Jagtman\*

\*Departments of Orthodontics and Oral Biology and Community and Restorative Dentistry, Radboud University Nijmegen Medical Centre, \*\*Department of Orthodontics, University Medical Centre Groningen, The Netherlands and \*\*\*Department of Orthodontics and Dentofacial Orthopedics, University of Bern, Switzerland

**SUMMARY** The aim of this study was to compare standard plaster models with their digital counterparts for the applicability of the Index of Complexity, Outcome, and Need (ICON). Generated study models of 30 randomly selected patients: 30 pre- ( $T_0$ ) and 30 post- ( $T_1$ ) treatment. Two examiners, calibrated in the ICON, scored the digital and plaster models. The overall ICON scores were evaluated for reliability and reproducibility using kappa statistics and reliability coefficients.

The values for reliability of the total and weighted ICON scores were generally high for the  $T_0$  sample (range 0.83–0.95) but less high for the  $T_1$  sample (range 0.55–0.85). Differences in total ICON score between plaster and digital models resulted in mostly statistically insignificant values ( $P$  values ranging from 0.07 to 0.19), except for observer 1 in the  $T_1$  sample. No statistically different values were found for the total ICON score on either plaster or digital models.

ICON scores performed on computer-based models appear to be as accurate and reliable as ICON scores on plaster models.

## Introduction

Digital, computer-based three-dimensional (3D) study models are an alternative to plaster models. Users of digital models can retrieve and store their records electronically, reducing the chance of loss or damage and reducing storage space. A virtual 3D set of models can be manipulated in all planes of space, sectioned in any plane and measured along any plane. The virtual images can be sent worldwide for instant referral or consultation as needed, for Internet study groups or research purposes. Because computer-based 3D study models are a component of the digital orthodontic record, they contribute to a paperless office. Digital models have been shown to be a valid tool for undertaking simple diagnostic measurements such as tooth size, arch width, overjet, overbite, arch length, and Bolton ratio (Tomassetti *et al.*, 2001; Santoro *et al.*, 2003; Zilberman *et al.*, 2003; Quimby *et al.*, 2004; Stevens *et al.*, 2006). For direct measurements, the above authors did not find clinically significant differences between measurements made on digital or plaster models. Statistical differences were found for tooth width by Santoro *et al.* (2003) and Zilberman *et al.* (2003). Shrinkage of alginate and difficulties in identifying landmarks of a 3D image on a two-dimensional (2D) screen are mentioned explanations. One of the greatest sources of random error is the difficulty in identifying landmarks (Houston, 1983). Zilberman *et al.* (2003) stated that this is a particular concern for digital models because a 3D structure is viewed as a 2D image and identifying landmarks becomes more difficult. Quimby *et al.* (2004) and Stevens *et al.* (2006) found no statistical differences between measurements made on plaster or digital models.

Besides direct measurements, other methods are used to quantify malocclusion and treatment results, such as occlusal indices.

Numerous indices have been developed since the 1960s either to rank or score the severity of malocclusion relative to a pre-conceived orthodontic ideal, or in terms of treatment need (Draker, 1960; Salzmann, 1968; Summers, 1971; Linder-Aronson, 1974; Lundström, 1977; Brook and Shaw, 1989; Buchanan, 1991; Shaw *et al.*, 1991; Richmond *et al.*, 1992; Casco *et al.*, 1998; Daniels and Richmond, 2000). Examples of these indices are the Dental Aesthetic Index (DAI), the Peer Assessment Rating (PAR) Index, the Index of Orthodontic Treatment Need (IOTN), American Board of Orthodontics Objective grading system (ABO-OGS), and the Index of Complexity Outcome, and Need (ICON). From the various indices only two, PAR and ABO-OGS, have been compared for digital and plaster study models. For the PAR Index, two studies did not find significant differences (Mayers *et al.*, 2005; Stevens *et al.*, 2006). For the ABO-OGS, statistically significant differences were found (Costalos *et al.*, 2005; Okunami *et al.*, 2007) for the components ‘alignment’ and ‘buccolingual inclination’ (Costalos *et al.*, 2005) and ‘occlusal contacts and relationships’ (Okunami *et al.*, 2007). According to those authors, possible explanations for these differences might be the difficulty in identifying the same landmarks on plaster and digital models, a need for adequate calibration to achieve repeatability in both methods and a difference in angulation of the models while measurements were taken. Although some findings showed statistical differences, clinically the differences were too small to be noticed during

the grading process. Although these indices are widely used, they are not validated for determining treatment need and do not take aesthetics into consideration.

The ICON is an index of treatment need, severity of malocclusion, and treatment outcome and as such offers significant advantages over other indices of treatment need (Daniels and Richmond, 2000; Louwse *et al.*, 2006; Onyeaso and BeGole, 2007). A single set of weightings is used to score four separate facets of orthodontic care. The ICON weightings are based on the opinion of an international panel of 97 orthodontists from nine countries (Richmond and Daniels, 1998a,b). The aesthetic component (AC; Brook and Shaw, 1989) of the IOTN, crossbite, upper arch crowding or spacing, overbite or open bite, and buccal segment antero-posterior relationship are used to determine treatment need, treatment outcome, complexity, and degree of improvement (Daniels and Richmond, 2000). The ICON shows good overall agreement with the DAI, PAR, and ABO-OGS (Onyeaso and BeGole, 2007). Fox *et al.* (2002) found significant correlations between the IOTN and ICON with respect to need and PAR and ICON with respect to outcome.

A review of the literature did not identify any studies that investigated the clinical applicability of the ICON on digital models. Therefore, the aim of this study was to determine the reliability and reproducibility of ICON scores derived from digital study models compared with scores from plaster models of the same patients. This was carried out by comparing the values scored on plaster models with those scored on digital models and comparing the reproducibility of scoring on digital models with that on plaster models.

**Material and methods**

*Sample*

Pre- (T<sub>0</sub>) and post- (T<sub>1</sub>) treatment dental casts of 30 patients were randomly selected from the patient archive of the Department of Orthodontics and Oral Biology, Radboud University Nijmegen Medical Centre. Random numbers were

generated by the computer. The study sample met the following inclusion criteria: (1) permanent dentition, (2) apparently normal crown morphology (casts showing gross abnormalities were excluded), and (3) no features that would alter the natural mesiodistal or buccolingual crown diameter, such as restorations, caries, attrition, or fracture. The dental casts represented a spectrum of malocclusion types and severity before treatment. Patients with clefts and other craniofacial deformities were excluded. Five subjects had a Class I molar occlusion at T<sub>0</sub>, 19 were Class II division 1, five Class II division 2, and one Class III. Five dental casts displayed a crossbite or scissor bite and five showed an anterior open bite. Five patients were treated with extractions. The distribution of the Dental Health Component (DHC) grades of the IOTN is shown in Table 1.

To process digital models, the models were sent to Orthoproof®, Nieuwegein, The Netherlands. The plaster casts were scanned with a Flash CT scanner (Hytec Inc, Los Alamos, Oklahoma, USA). The same plaster casts were returned and used for direct measurements. The corresponding digital models were returned via electronic mail within 48 hours. The file size of one set of digital models is approximately 10 Mb.

*Measurements*

The measurements used were those described in computing the ICON, as shown in Table 2 (Daniels and Richmond, 2000). The components and the scoring range of the ICON are shown in Table 3. Two observers (SCB and MAB),

**Table 1** Distribution of the Dental Health Component (DHC) grades of the Index of Orthodontic Treatment Need (IOTN) of the sample (total *n* = 60).

IOTN DHC grade	Treatment need	<i>N</i> pre-treatment	<i>N</i> post-treatment
1	None	0	6
2	Little	0	23
3	Moderate	11	1
4	High	16	0
5	Very high	3	0

**Table 2** Protocol for occlusal trait scoring (reproduced from Daniels C, Richmond S 2000. The development of the Index of Complexity, Outcome and Need (ICON). *Journal of Orthodontics* 27:149–162, with kind permission of Maney Publishing).

Score	0	1	2	3	4	5
Aesthetic	1–10 as judged using IOTN-AC					
Upper arch crowding	Score only the highest trait either spacing or crowding	Less than 2 mm	2.1–5 mm	5.1 to 9 mm	9.1 to 13 mm	13.1 to 17 mm >17 mm or impacted teeth
Upper arch spacing		Up to 2 mm	2.1–5 mm	5.1 to 9 mm	>9 mm	
Crossbite	Transverse relationship of cusp to cusp or worse	No crossbite	Crossbite present			
Incisor overbite	Lower incisor coverage	Up to 1/3 tooth	1/3–2/3 coverage	2/3 up to full coverage	Full coverage	
Sagittal relationship of the buccal segment	Left and right added together	Cusp to embrasure relationship only, Class I, II or III	Any cusp relation up to but not including cusp to cusp	Cusp to cusp relationship		

calibrated in the use for the ICON, scored the models. The casts were displayed in a fixed order on tables. There was no time limit. After 1 week the digital models were scored. The digital models were viewed by using the proprietary software (Digimodel®, version 2.2.1, Nieuwegein, The Netherlands). The digital models were displayed on screen with four views of one set of the dental casts of one patient (Figure 1). A 15.4 inch LCD laptop screen with a resolution of 1280 × 800 pixels with 32 bit colour was used. The observer could manipulate the position of the models in view. For ease and accuracy of measurements, the images were enlarged on screen as required with the magnifying feature. Overbite was assessed by making the maxilla transparent. Posterior displacement from ideal interdigitation was determined with a view perpendicular to the posterior quadrant.

The second scoring of the digital models was carried out 2 weeks after the first scoring and 1 week thereafter the second scoring of the plaster models was undertaken.

Reliability was considered as the extent to which a measurement was repeatable under identical conditions for the new diagnostic test (digital models) and the gold standard (plaster).

*Statistical analysis*


The various components of the ICON have different scales. Some components result in ordinal data (e.g. crossbite), while other components are recorded on a metric scale. For the metric variables (AC, incisor overbite, and sagittal relationship of the buccal segment, total score, and weighted score), the observer performance is expressed in the reliability index (calculated by Pearson correlation coefficient) and the results of a paired *t*-test (*P* value and mean difference). For the ordinal components (upper arch crowding/spacing and crossbite), kappa values were calculated to analyse observer agreement. Statistical analysis was performed with the Statistical Package for Social Sciences version 12 (SPSS Inc., Chicago, Illinois, USA). Statistical significance was set at a *P* < 0.05.

**Results**

Descriptive analysis of the total and weighted total scores for the samples at T<sub>0</sub> and T<sub>1</sub> are shown in Table 4 and statistical comparison between digital and plaster models in Table 5.

**Table 3** Components of the Index of Complexity, Outcome, and Need (ICON) with their scoring range and weights.

ICON components	Scoring range	Weight
Aesthetic Component of the Index of Orthodontic Treatment Need	1–10	7
Upper arch crowding/spacing	0–5	5
Crossbite	0–1	5
Incisor overbite	0–4	4
Sagittal relationship of the buccal segment	0–2	3



**Figure 1** Digital model used in the study (Digimodel®).

Differences in total ICON score between the plaster and digital models resulted in mostly statistically insignificant values (*P* values ranging from 0.07 to 0.19), except for observer 1 in the T<sub>1</sub> sample. Paired *t*-tests showed that the total ICON score between plaster and digital models for observer 1 differed, with a plaster score being on average 0.73 points lower than the score on digital models (*P* < 0.01). This difference was found for the components, sagittal relationship of the buccal segment (*P* < 0.01), and crossbite (kappa = 0).

Significant differences between the digital and plaster models in the weighted ICON scores were found. Observer 1 showed a difference (*P* = 0.01) in the T<sub>1</sub> sample. This difference was also seen in the total ICON score for that sample. Observer 2 showed a difference in the weighted total score for both the T<sub>0</sub> and T<sub>1</sub> sample (*P* = 0.03 and *P* < 0.01, respectively).

**Table 4** Descriptive statistics for the pre- (T<sub>0</sub>) and post- (T<sub>1</sub>) treatment sample. Mean, standard deviation (SD), and (range) are given in Index of Complexity, Outcome, and Need (ICON) points.

	Time	Plaster						Digital					
		Observer 1			Observer 2			Observer 1			Observer 2		
		Mean	SD	Range	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range
Total ICON score	T <sub>0</sub>	11.47	2.37	(5–16)	11.53	2.64	(5–16)	10.97	2.47	(6–16)	11.13	2.39	(5–17)
	T <sub>1</sub>	3.4	1.07	(1–6)	4.3	1.05	(2–6)	4.13	1.31	(1–7)	4.6	1.13	(2–6)
Weighted Total ICON score	T <sub>0</sub>	59.7	14.37	(28–84)	60.03	15.39	(28–98)	56.73	14.19	(31–84)	57.03	13.14	(28–87)
	T <sub>1</sub>	17.23	4.85	(7–28)	20.9	4.47	(10–30)	19.93	5.87	(7–33)	21.13	4.76	(10–30)

**Table 5** Index of Complexity, Outcome, and Need (ICON) scores of plaster versus digital models compared for both observers.

	ICON component	Pre-treatment (T <sub>0</sub> )				Post-treatment (T <sub>1</sub> )			
		Pearson correlation coefficient	P value	Diff	Kappa	Pearson correlation coefficient	P value	Diff	Kappa
Observer 1	Aesthetic component	0.68	0.29	0.30		0.43	0.13	-0.17	
	U-arch crowding/spacing				0.75				1
	Crossbite				0.86				0
	Incisor overbite	0.99	0.33	0.03		0.70	0.33	-0.03	
	Sagittal buccal segment	0.66	0.79	0.03		0.61	<0.01	-0.63	
	Total ICON	0.64	0.19	0.5		0.60	0.01	-0.73	
	Weighted total ICON	0.62	0.20	3		0.56	0.01	-2.70	
Observer 2	Aesthetic component	0.89	0.01	0.47		0.57	0.06	0.17	
	U-arch crowding/spacing				0.64				1
	Crossbite				0.9				1
	Incisor overbite	*	*	*	*	*	*	*	*
	Sagittal buccal segment	0.85	0.66	-0.03		0.71	<0.01	-0.47	
	Total ICON	0.89	0.08	0.4		0.68	0.07	-0.30	
	Weighted total ICON	0.89	0.03	3		0.63	<0.01	-3.67	

Diff, mean difference plaster minus digital.

\*No values possible due to complete agreement between the values for plaster and digital models.

A difference was found for the components sagittal relationship of the buccal segment ( $P < 0.01$ ) and AC ( $P = 0.01$ ).

Intraobserver reproducibility for the plaster and digital models are shown in Tables 6 and 7. The values for reliability of the total and weighted total ICON scores were generally high for the T<sub>0</sub> sample (range 0.83–0.95) but lower for the T<sub>1</sub> sample (range 0.55–0.85). No statistical differences were found for the total ICON score on either plaster or digital models. Intraobserver differences in repeated weighted total ICON scoring of digital models resulted in statistically significant values for observer 2 ( $P = 0.04$ ) for the T<sub>0</sub> sample. This difference was mainly due to the AC.

**Discussion**

The present study assessed the reliability and reproducibility of ICON scores derived from 3D digital study models using plaster models as the gold standard. The number of models to be compared (both for T<sub>0</sub> and T<sub>1</sub>) had to be large enough to achieve sufficient power. For metric variables, the standard

error (SE) in the difference of two variables with approximately the same standard deviation (SD) can be calculated as:  $SE = \sqrt{(2/N)} \times SD$ . By setting  $N$  at 30, the standard error of the mean is close to a factor of 4 smaller than the SD. This implies that differences between observers, or between digital and plaster models, would reach statistical significance at a level of 0.5 SD or more. For those cases where the kappa statistic was used, the power depends both on the expected and the observed level of agreement, so the power calculation is more arbitrary. Setting the expected level of agreement at 0.25 (as can be seen in an equally distributed four-point scale or approximately in a skewed distributed five-point scale), then a level of agreement of 0.75 would result in a kappa of 0.67. Obviously, the same  $N$  as for metric variables has to be used. Setting  $N$  at 30 with these levels of agreement gives a SE for the kappa value of 0.11. This was considered to be sufficiently precise.

Two samples were used: 30 pre- (T<sub>0</sub>) and 30 post- (T<sub>1</sub>) treatment models. Whereas the T<sub>0</sub> sample contained a variety of malocclusions, the T<sub>1</sub> sample was more homogeneous.

**Table 6** Intraobserver reproducibility for the pre-treatment models.

		Pearson correlation coefficient		P value		Mean Difference		Kappa	
		Plaster	Digital	Plaster	Digital	Plaster	Digital	Plaster	Digital
Observer 1	Aesthetic component	0.92	0.84	0.66	0.38	-0.07	0.17	0.81 1	0.25 0.86
	Upper-arch crowding/spacing								
	Crossbite								
	Incisor overbite	0.95	0.95	1	0.33	0	-0.07		
	Sagittal buccal segment	0.87	0.81	0.42	0.71	-0.07	-0.03		
Observer 2	Total ICON	0.87	0.84	0.32	0.36	-0.23	-0.23		
	Weighted total ICON	0.88	0.83	0.39	0.65	-1.17	-0.7		
	Aesthetic component	0.95	0.95	0.54	0.17	0.07	-0.17	0.37 1	0.60 1
	Upper-arch crowding/spacing								
	Crossbite								
	Incisor overbite	0.94	0.99	0.66	0.33	0.03	-0.03		
	Sagittal buccal segment	0.84	0.84	1	0.42	0	0.07		
Total ICON	0.95	0.93	0.38	0.07	-0.13	-0.33			
Weighted total ICON	0.95	0.94	0.51	0.04*	-0.57	-2.1			

\*Significant at  $P < 0.05$ .

**Table 7** Intraobserver reproducibility for the post-treatment models.

		Pearson correlation coefficient		P value		Difference		Kappa	
		Plaster	Digital	Plaster	Digital	Plaster	Digital	Plaster	Digital
Observer 1	Aesthetic component	0.43	0.26	0.54	1	0.07	0	1 0.78	1 1
	Upper-arch crowding/spacing								
	Crossbite								
	Incisor overbite	0.42	0.80	0.04	0.33	-0.13	-0.03		
	Sagittal buccal segment	0.85	0.81	0.26	0.38	-0.1	-0.1		
Observer 2	Total ICON	0.79	0.68	0.38	0.46	-0.13	-0.13		
	Weighted total ICON	0.67	0.55	0.81	0.63	-0.2	-0.04		
	Aesthetic component	0.51	0.83	0.42	0.08	0.07	-0.1	0 0	1 1
	Upper-arch crowding/spacing								
	Crossbite								
	Incisor overbite	—	—	0.33	0.33	-0.03	-0.03		
	Sagittal buccal segment	0.90	0.77	0.10	0.29	-0.13	0.13		
Total ICON	0.85	0.75	0.13	1	-0.17	0			
Weighted total ICON	0.74	0.78	0.49	0.48	-0.4	0.43			

Therefore, the pre- and post-treatment samples were analysed separately. Low intrasample variation in the T<sub>1</sub> sample, which is inherent post-treatment (Table 4), is an important factor for the statistical differences found in total and weighted total score for the comparison between plaster and digital models (Table 5). A much lower interobserver agreement in decisions of treatment acceptability and lower predictive accuracy for treatment outcome for the ICON compared with treatment need have been reported previously (Richmond and Daniels, 1998b). The difference for weighted total score (3 points) in the T<sub>0</sub> sample for observer 2 does not appear to be clinically relevant, the SD in this sample being 15.39 points. The digital models analysed by observer 2 in the T<sub>0</sub> sample (Table 6) had a weighted total score difference of 2.1, which is very small. The different components of the ICON show low kappa values for the ordinal components. This is because kappa can drop dramatically based

on the prevalence of the variable involved (Altman, 1991). The components, upper arch crowding/spacing and crossbite, show this phenomenon. Stevens *et al.* (2006) reported difficulties in observing crossbites. The posterior teeth can falsely appear in crossbite on screen or they will seem to have a positive overjet in the posterior segment when they do not. With the cross-section function of the program, this can be checked.

The AC gave a statistically different value for observer 2 (Table 5). This difference (0.47) is very small and it should be stressed that in the calibrating process, the AC of the ICON proved difficult to learn. Indeed, studies assessing the IOTN-AC demonstrated moderate validity (Richmond *et al.*, 1995; Beglin *et al.*, 2001). When the plaster and digital models were compared in the present study, differences were found for the sagittal relationship of the buccal segment (range 0.47–0.63). A slight rotation of the digital model around the vertical

axis affected assessment of molar and canine relationships and this could explain these differences. Other studies (Mayers *et al.*, 2005; Rheude *et al.*, 2005; Stevens *et al.*, 2006) also found a slightly lower reliability for digital models for buccal occlusion. However, overall, the differences between plaster and digital models do not seem to be clinically important.

In all studies so far undertaken, statistical differences might be found, but they do not seem to be clinically relevant. It is likely that these small differences do not have an influence on diagnosis and treatment planning, as confirmed by the studies of Rheude *et al.* (2005) and Whetten *et al.* (2006) who showed that digital models are a valid alternative to conventional plaster models in treatment planning.

### Conclusion

Despite some minor differences between ICON scores on plaster and digital models, it can be concluded that ICON scoring can be performed reliably on digital models.

### Address for correspondence

Professor A. M. Kuijpers-Jagtman  
Department of Orthodontics and Oral Biology  
Radboud University Nijmegen Medical Centre  
P.O. Box 9101  
NL-6500 HB Nijmegen  
The Netherlands  
E-mail: orthodontics@dent.umcn.nl

### Acknowledgement

Special thanks to Manon A. Borsje, University Medical Centre Groningen, for her assistance in this research.

### References

- Altman D G 1991 Practical statistics for medical research. Chapman & Hall, London
- Beglin F M, Firestone A R, Vig K W, Beck F M, Kuthy R A, Wade D 2001 A comparison of the reliability and validity of 3 occlusal indexes of orthodontic treatment need. *American Journal of Orthodontics and Dentofacial Orthopedics* 120: 240–246
- Brook P H, Shaw W C 1989 The development of an index of orthodontic treatment priority. *European Journal of Orthodontics* 11: 309–320
- Buchanan I B 1991 The evaluation and initial testing of an index to assess orthodontic treatment standards: the PAR index. Thesis, University of Manchester
- Casko J, Vaden J, Kokich V, Damone J, James R, Cangialosi T 1998 Objective grading system for dental casts and panoramic radiographs. *American Journal of Orthodontics and Dentofacial Orthopedics* 114: 589–599
- Costalos P A, Sarraf K, Cangialosi T J, Efstratiadis S 2005 Evaluation of the accuracy of digital model analysis for the American Board of Orthodontics objective grading system for dental casts. *American Journal of Orthodontics and Dentofacial Orthopedics* 128: 624–629
- Daniels C, Richmond S 2000 The development of the Index of Complexity, Outcome and Need (ICON). *Journal of Orthodontics* 27: 149–162
- Draker H L 1960 Handicapping labio-lingual deviations: a proposed index for public health purposes. *American Journal of Orthodontics* 46: 295–315
- Fox N A, Daniels C, Gilgrass T 2002 A comparison of the Index of Complexity Outcome and Need (ICON) with the peer assessment rating (PAR) and the Index of Orthodontic Treatment Need (IOTN). *British Dental Journal* 193: 225–230
- Houston W J B 1983 The analysis of errors in orthodontic measurements. *American Journal of Orthodontics* 83: 382–390
- Linder-Aronson S 1974 Orthodontics in the Swedish public dental health system. *Transactions of the European Orthodontic Society*, pp.233–240
- Louwerse T J, Aartman I H A, Kramer G J C, Prahll-Andersen B 2006 The reliability and validity of the Index of Complexity, Outcome and Need for determining treatment need in Dutch orthodontic practice. *European Journal of Orthodontics* 28: 58–64
- Lundström A 1977 Need for treatment in cases of malocclusion. *Transactions of the European Orthodontic Society*, pp.111–123
- Mayers M, Firestone A R, Rashid R, Vig K W L 2005 Comparison of Peer Assessment Rating (PAR) Index scores of plaster and computer-based digital models. *American Journal of Orthodontics and Dentofacial Orthopedics* 128: 431–434
- Okunami T R, Kusnoto B, BeGole E, Evans C A, Sadowsky C, Shahrbanoo F 2007 Assessing the American Board of Orthodontics Objective Grading System: digital vs plaster dental casts. *American Journal of Orthodontics and Dentofacial Orthopedics* 131: 51–56
- Onyeaso C O, BeGole E 2007 Relationship between Index of Complexity, Outcome and Need, Dental Aesthetic Index, Peer Assessment Rating Index, and American Board of Orthodontics Objective Grading System. *American Journal of Orthodontics and Dentofacial Orthopedics* 131: 248–252
- Quimby M, Vig K, Rashid R, Firestone A, Mayers M 2004 The accuracy and reliability of measurements made on computer-based digital models. *Angle Orthodontist* 74: 298–303
- Rheude B, Sadowsky P L, Ferreira A, Jacobson A 2005 An evaluation of the use of digital study models in orthodontic diagnosis and treatment planning. *Angle Orthodontist* 75: 300–304
- Richmond S, Daniels C P 1998a International comparisons of professional assessments in orthodontics: part 1—treatment need. *American Journal of Orthodontics and Dentofacial Orthopedics* 113: 180–185
- Richmond S, Daniels C P 1998b International comparisons of professional assessments in orthodontics: part 2—treatment outcome. *American Journal of Orthodontics and Dentofacial Orthopedics* 113: 324–328
- Richmond S *et al.* 1992 The development of the PAR index (Peer Assessment Rating): reliability and validity. *European Journal of Orthodontics* 14: 125–139
- Richmond S *et al.* 1995 Calibration of dentists in the use of occlusal indices. *Community Dentistry and Oral Epidemiology* 23: 173–176
- Salzmann J A 1968 Handicapping malocclusion assessment to establish treatment priority. *American Journal of Orthodontics* 54: 749–765
- Santoro M, Galkin S, Teredesai M, Nicolay O, Cangialosi T 2003 Comparison of measurements made on digital and plaster models. *American Journal of Orthodontics and Dentofacial Orthopedics* 124: 101–105
- Shaw W C, Richmond S, O'Brien K D, Brook P, Stephens C D 1991 Quality control in orthodontics: indices of treatment need and treatment standards. *British Dental Journal* 170: 107–112
- Stevens R, Flores-Mir C, Nebbe B, Raboud D W, Heo G, Major P W 2006 Validity, reliability, and reproducibility of plaster vs digital study models: comparison of Peer Assessment Rating and Bolton analysis and their constituent measurements. *American Journal of Orthodontics and Dentofacial Orthopedics* 129: 794–803
- Summers C J 1971 The occlusal index: a system for identifying and scoring occlusal disorders. *American Journal of Orthodontics* 59: 552–567
- Tomassetti J, Taloumis L, Denny J, Fisher J 2001 A comparison of 3 computerized Bolton tooth-size analysis with a commonly used method. *Angle Orthodontist* 71: 351–357
- Whetten J L, Williamson P C, Heo G, Varnhagen C, Major P W 2006 Variations in orthodontic treatment planning decisions of Class II patients between virtual 3-dimensional models and traditional plaster study models. *American Journal of Orthodontics and Dentofacial Orthopedics* 130: 485–491
- Zilberman O, Huggare J, Parikakis K 2003 Evaluation of the validity of tooth size and arch width measurements using conventional and three-dimensional virtual orthodontic models. *Angle Orthodontist* 73: 301–306