

Analyticity and Riesz basis property of semigroups associated to damped vibrations

BIRGIT JACOB, CARSTEN TRUNK and MONIKA WINKLMEIER

Abstract. Second order equations of the form $\ddot{z}(t) + A_0 z(t) + D\dot{z}(t) = 0$ are considered. Such equations are often used as a model for transverse motions of thin beams in the presence of damping. We derive various properties of the operator matrix $\mathcal{A} = \begin{bmatrix} 0 & I \\ -A_0 & -D \end{bmatrix}$ associated with the second order problem above. We develop sufficient conditions for analyticity of the associated semigroup and for the existence of a Riesz basis consisting of eigenvectors and associated vectors of \mathcal{A} in the phase space.

1. Introduction

A linear equation describing transverse motions of a thin beam can be written in the form

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2}{\partial r^2} \left[E \frac{\partial^2 u}{\partial r^2} + C_d \frac{\partial^3 u}{\partial r^2 \partial t} \right] = 0, \quad r \in (0, 1), t > 0,$$

where $u(r, t)$ is the transverse displacement of the beam at time t and position r . The existence and behaviour of solutions u depend also on boundary and initial conditions. In the example above we are interested in solutions having finite energy, i.e. solutions such that $\|u(\cdot, t)\|^2 + \|u''(\cdot, t)\|^2 < \infty$ for all $t > 0$ where $\|\cdot\|$ denotes the usual norm in the Hilbert space $L^2(0, 1)$. Identifying the function $u(\cdot, t)$ with an element $z(t) \in L^2(0, 1)$ by $z(t)(r) = u(r, t)$ we obtain from the partial differential equation above a second order equation in $L^2(0, 1)$ of the form

$$\ddot{z}(t) + A_0 z(t) + D\dot{z}(t) = 0, \quad (1)$$

where $A_0 = E \frac{\partial^4}{\partial r^4}$, $D = \frac{\partial^2}{\partial r^2} C_d \frac{\partial^2}{\partial r^2}$ acting in $L^2(0, 1)$ with appropriate domains encoding the boundary conditions under consideration. We will come back to this example in Section 6.

In this paper we study second order equations of type (1) in an abstract Hilbert space H where the stiffness operator A_0 is a possibly unbounded positive operator on H and

Mathematics Subject Classifications (2000): 47A10, 47A70, 34G10, 47D06.

Key words: Operator matrices, second order equations, spectrum, Riesz basis, analytic semigroup

is assumed to be boundedly invertible, and D , the damping operator, is an unbounded operator such that $A_0^{-1/2}DA_0^{-1/2}$ is a bounded non-negative operator on H . This second order equation is equivalent to the standard first-order equation $\dot{x}(t) = \mathcal{A}x(t)$ where $\mathcal{A} : \mathcal{D}(\mathcal{A}) \subset \mathcal{D}(A_0^{1/2}) \times H \rightarrow \mathcal{D}(A_0^{1/2}) \times H$ is given by

$$\mathcal{A} = \begin{bmatrix} 0 & I \\ -A_0 & -D \end{bmatrix},$$

$$\mathcal{D}(\mathcal{A}) = \left\{ \begin{bmatrix} z \\ w \end{bmatrix} \in \mathcal{D}(A_0^{1/2}) \times \mathcal{D}(A_0^{1/2}) \mid A_0z + Dw \in H \right\}.$$

This operator matrix has been studied in the literature for more than 20 years. Interest in this particular model is motivated by various problems such as stabilization, see for example [8], [35], [36], [39], solvability of Riccati equations [18], minimum-phase property [23] and compensator problems with partial observations [19].

It is well-known that \mathcal{A} generates a C_0 -semigroup of contractions in $H_{1/2} \times H$, where $H_{1/2} = \mathcal{D}(A_0^{1/2})$ is equipped with the norm $\|x\|_{1/2} = \|A_0^{1/2}x\|_H$, and thus the spectrum of \mathcal{A} is located in the closed left half plane. This goes back to [4] and [34], see also [5], [11]. Several authors have proved independently of each other that the condition $\langle A_0^{-1/2}Dz, A_0^{1/2}z \rangle_H \geq \beta \|z\|_H^2$ is sufficient for exponential stability of the C_0 -semigroup generated by \mathcal{A} , see for example [4], [5], [7], [11], [17], [20], [41], [42].

In this paper we focus on two properties of the operator \mathcal{A} : Analyticity of the generated semigroup and the Riesz basis property in the phase space $H_{1/2} \times H$. Analyticity of the C_0 -semigroup generated by \mathcal{A} has been studied in many papers, see [10], [11], [12], [13], [16], [21], [22], and [30]. Most of the papers require that the damping operator D is comparable with A^ρ for some $\rho \in [1/2, 1]$. In [30] the damping operator D is of the form

$$D = \alpha A_0 + B, \tag{2}$$

where $\alpha > 0$ is a constant, A_0^{-1} is compact and B is symmetric and A_0 -compact. If $-1/\alpha \notin \sigma_p(\mathcal{A})$, then it is shown in [30] that \mathcal{A} generates an analytic semigroup. In this case the essential spectrum of the operator $A_0^{-1}D$, considered as an operator acting in $H_{1/2}$ consists of the point $-1/\alpha$ only. We extend the result of [30] to more general damping operators D : If A_0^{-1} is compact in H and $0 \notin \sigma_{ess}(A_0^{-1}D)$, then \mathcal{A} generates an analytic semigroup on $H_{1/2} \times H$ (cf. Theorem 4.1 below). In particular, this implies that the above mentioned result from [30] holds even if $-1/\alpha \in \sigma_p(\mathcal{A})$.

Note that analyticity of the semigroup generated by \mathcal{A} already implies that the semigroup satisfies the spectral mapping theorem

$$\sigma(T(t)) \setminus \{0\} = e^{t\sigma(\mathcal{A})}, \quad t \geq 0,$$

where $(T(t))_{t \geq 0}$ is the semigroup generated by \mathcal{A} , see [15, Chapter IV, Section 3.10].

We further develop conditions guaranteeing that the space $H_{1/2} \times H$ possesses a Riesz basis consisting of eigenvectors and finitely many associated vectors of \mathcal{A} . The existence of such a system has many important implications for the operator \mathcal{A} ; for instance, it implies that \mathcal{A} satisfies the weak spectral mapping theorem, that is,

$$\sigma(T(t)) = \overline{e^{t\sigma(\mathcal{A})}}, \quad t \geq 0.$$

In particular, it follows that the semigroup is exponentially stable if and only if the spectrum of \mathcal{A} is contained in the open left half plane and uniformly bounded away from the imaginary axis.

The Riesz basis property has been shown in [30] in the situation where A_0^{-1} is a compact operator, D is of the form (2) for some $\alpha \geq 0$ with a symmetric operator B and $-1/\alpha \notin \sigma_p(\mathcal{A})$, if $\alpha \neq 0$ (and with some additional assumptions in the case $\alpha = 0$). Similar results were obtained [12, Appendix A] in a more special situation. All these assumptions guarantee that the essential spectrum of \mathcal{A} consists at most of one point.

In this paper we also assume that A_0^{-1} is a compact operator, but we allow a more general damping operators D . In particular, the essential spectrum of \mathcal{A} may contain infinitely many points. For most of our results we need the assumption that $0 \notin \sigma_{ess}(A_0^{-1}D)$, where $A_0^{-1}D$ is seen as an operator acting on $H_{1/2} = \mathcal{D}(A_0^{1/2})$. This, however, implies that we cannot handle the case where $A_0^{-1}D$ is a compact operator in $H_{1/2}$ unless H is of finite dimension. Together with some rather weak conditions the above mentioned imply that there exists a Riesz basis in the phase space $H_{1/2} \times H$ consisting of eigenvalues and finitely many associated vectors of \mathcal{A} (cf. Theorem 5.1 below). For results involving compact $A_0^{-1}D$ we refer the reader to [30].

Throughout this paper we assume that all Hilbert and Krein spaces are infinite dimensional. Since we are interested in applications to partial differential equations, this is no major restriction.

We proceed as follows. In Section 2 we provide some useful results on the spectrum of operators in Krein spaces. In particular, we recall the notion of spectral points of positive and negative type and of type π_+ and type π_- . One main tool of this paper is to show that certain spectral points of \mathcal{A} are of positive or negative type or of type π_+ or π_- . In Section 3 we give the precise definition of the operator \mathcal{A} and prove some of its properties. The main results of this paper are contained in Sections 4 and 5 where we always assume that that A_0^{-1} is a compact operator and that $0 \notin \sigma_{ess}(A_0^{-1}D)$. The main result of Section 4 is that \mathcal{A} generates an analytic strongly continuous semigroup. Further, we show that ∞ is a spectral point of negative type and that every real spectral point is of type π_+ . As a consequence we obtain that \mathcal{A} is definitizable and that the non-real spectrum of \mathcal{A} consists of at most finitely many points belonging to the point spectrum of \mathcal{A} . Further, the operator \mathcal{A} can be written as a direct sum of a self-adjoint operator in a Hilbert space and a bounded self-adjoint operator in a Pontryagin space. Section 5 is devoted to the Riesz basis property of the operator \mathcal{A} , that is, it is shown that under additional weak conditions there exists a

Riesz basis of $\mathcal{D}(A_0^{1/2}) \times H$ consisting of eigenvalues and finitely many associated vectors of A . Finally, in Section 6 the results are illustrated by an example: the Euler-Bernoulli beam with distributed Kelvin-Voigt damping.

2. Spectrum of operators in Krein spaces

Let $(\mathcal{H}, [\cdot, \cdot])$ be a Krein space. We briefly recall that a complex linear space \mathcal{H} with a hermitian nondegenerate sesquilinear form $[\cdot, \cdot]$ is called a *Krein space* if there exists a decomposition $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ with subspaces \mathcal{H}_\pm being orthogonal to each other with respect to $[\cdot, \cdot]$ such that $(\mathcal{H}_\pm, \pm[\cdot, \cdot])$ are Hilbert spaces. In the following all topological notions are understood with respect to some Hilbert space norm $\|\cdot\|$ on \mathcal{H} such that $[\cdot, \cdot]$ is $\|\cdot\|$ -continuous. Any two such norms are equivalent. For the basic theory of Krein space and operators acting therein we refer to [9] and [2].

Let A be a closed operator in \mathcal{H} . Analogously to the Hilbert space case we define the extended spectrum $\sigma_e(A)$ of A by $\sigma_e(A) := \sigma(A)$ if A is bounded and $\sigma_e(A) := \sigma(A) \cup \{\infty\}$ if A is unbounded. The resolvent set of A is denoted by $\rho(A)$. By $\sigma_{p,norm}(A)$ we denote the set of all $\lambda \in \mathbb{C}$ which are isolated points of $\sigma(A)$ and normal eigenvalues of A , that is, the corresponding Riesz-Dunford projection is of finite rank. A point $\lambda_0 \in \mathbb{C}$ is said to belong to the *approximative point spectrum* $\sigma_{ap}(A)$ of A if there exists a sequence $(x_n) \subset \mathcal{D}(A)$ with $\|x_n\| = 1$, $n = 1, 2, \dots$, and $\|(A - \lambda_0 I)x_n\| \rightarrow 0$ if $n \rightarrow \infty$. For a self-adjoint operator A in \mathcal{H} all real spectral points of A belong to $\sigma_{ap}(A)$ (see e.g. [9, Corollary VI.6.2]). The operator A is called *Fredholm* if the dimension of the kernel of A and the codimension of the range of A are finite. The set

$$\sigma_{ess}(A) := \{\lambda \in \mathbb{C} \mid A - \lambda I \text{ is not Fredholm}\}$$

is called the *essential spectrum* of A .

Using the indefiniteness of the scalar product on \mathcal{H} we have the notions of spectral points of positive and negative type and of type π_+ and type π_- . The following definition was given in [29] and [33] for bounded self-adjoint operators.

DEFINITION 2.1. For a self-adjoint operator A in \mathcal{H} a point $\lambda_0 \in \sigma(A)$ is called a spectral point of *positive (negative) type of A* if $\lambda_0 \in \sigma_{ap}(A)$ and for every sequence $(x_n) \subset \mathcal{D}(A)$ with $\|x_n\| = 1$ and $\|(A - \lambda_0 I)x_n\| \rightarrow 0$ as $n \rightarrow \infty$ we have

$$\liminf_{n \rightarrow \infty} [x_n, x_n] > 0 \quad (\text{resp. } \limsup_{n \rightarrow \infty} [x_n, x_n] < 0).$$

The point ∞ is said to be of *positive (negative) type of A* if A is unbounded and for every sequence $(x_n) \subset \mathcal{D}(A)$ with $\lim_{n \rightarrow \infty} \|x_n\| = 0$ and $\|Ax_n\| = 1$ we have

$$\liminf_{n \rightarrow \infty} [Ax_n, Ax_n] > 0 \quad (\text{resp. } \limsup_{n \rightarrow \infty} [Ax_n, Ax_n] < 0).$$

We denote the set of all points of $\sigma_e(A)$ of positive (negative) type by $\sigma_{++}(A)$ (resp. $\sigma_{--}(A)$).

It is not difficult to see that the sets $\sigma_{++}(A)$ and $\sigma_{--}(A)$ are contained in $\overline{\mathbb{R}}$. Moreover the non-real spectrum of A cannot accumulate to $\sigma_{++}(A) \cup \sigma_{--}(A)$.

In a similar way as above we define subsets $\sigma_{\pi_+}(A)$ and $\sigma_{\pi_-}(A)$ of $\sigma_e(A)$ containing $\sigma_{++}(A)$ and $\sigma_{--}(A)$, respectively (cf. [3, Definition 5]).

DEFINITION 2.2. For a self-adjoint operator A in \mathcal{H} a point $\lambda_0 \in \sigma(A)$ is called a spectral point of *type π_+* (*type π_-*) of A if $\lambda_0 \in \sigma_{ap}(A)$ and if there exists a linear submanifold $\mathcal{H}_0 \subset \mathcal{H}$ with $\text{codim } \mathcal{H}_0 < \infty$ such that for every sequence $(x_n) \subset \mathcal{H}_0 \cap \mathcal{D}(A)$ with $\|x_n\| = 1$ and $\|(A - \lambda_0 I)x_n\| \rightarrow 0$ as $n \rightarrow \infty$ we have

$$\liminf_{n \rightarrow \infty} [x_n, x_n] > 0 \quad (\text{resp. } \limsup_{n \rightarrow \infty} [x_n, x_n] < 0). \tag{3}$$

The point ∞ is said to be of *type π_+* (*type π_-*) of A if A is unbounded and if there exists a linear submanifold $\mathcal{H}_0 \subset \mathcal{H}$ with $\text{codim } \mathcal{H}_0 < \infty$ such that for every sequence $(x_n) \subset \mathcal{H}_0 \cap \mathcal{D}(A)$ with $\lim_{n \rightarrow \infty} \|x_n\| = 0$ and $\|Ax_n\| = 1$ we have

$$\liminf_{n \rightarrow \infty} [Ax_n, Ax_n] > 0 \quad (\text{resp. } \limsup_{n \rightarrow \infty} [Ax_n, Ax_n] < 0).$$

We denote the set of all points of $\sigma_e(A)$ of type π_+ (type π_-) of A by $\sigma_{\pi_+}(A)$ (resp. $\sigma_{\pi_-}(A)$).

Recall that a self-adjoint operator A in a Krein space $(\mathcal{H}, [\cdot, \cdot])$ is called *definitizable* if $\rho(A) \neq \emptyset$ and if there exists a rational function $p \neq 0$ having poles only in $\rho(A)$ such that $[p(A)x, x] \geq 0$ for all $x \in \mathcal{H}$. Then the spectrum of A is real or its non-real part consists of a finite number of points. Moreover, A has a spectral function $E(\cdot)$ defined on the ring generated by all connected subsets of $\overline{\mathbb{R}}$ whose endpoints do not belong to some finite set which is contained in $\{t \in \mathbb{R} : p(t) = 0\} \cup \{\infty\}$ (see [32]).

For a definitizable operator A a spectral point $t \in \overline{\mathbb{R}}$ is called a *critical point* of A if there is no open subset Δ with $t \in \Delta$ such that either $\Delta \subset \sigma_{++}(A)$ or $\Delta \subset \sigma_{--}(A)$. A critical point t is called *regular* if there exists an open deleted neighbourhood δ_0 of t such that the set of the projections $E(\delta)$ is bounded where δ runs through all intervals δ with $\bar{\delta} \subset \delta_0$, see [32].

THEOREM 2.3. *Let A be a self-adjoint operator in \mathcal{H} satisfying*

$$\sigma_{ess}(A) \subset \mathbb{R}, \quad \infty \in \sigma_{++}(A) \cup \sigma_{--}(A) \quad \text{and} \quad \sigma(A) \cap \mathbb{R} \subset \sigma_{\pi_+}(A) \cup \sigma_{\pi_-}(A).$$

Then A is a definitizable operator, and the non-real spectrum of A consists of at most finitely many points which belong to $\sigma_{p,norm}(A)$.

Proof. Assume $\infty \in \sigma_{--}(A)$. By [3, Lemma 2] there exists a neighbourhood \mathcal{U} of ∞ in $\overline{\mathbb{C}}$ with

$$\mathcal{U} \setminus \overline{\mathbb{R}} \subset \rho(A) \quad \text{and} \quad \mathcal{U} \cap \mathbb{R} \subset \sigma_{--}(A) \cup \rho(A).$$

From this and [3, Theorem 18] we conclude that the non-real spectrum of A consists of at most finitely many points which belong to $\sigma_{p,norm}(A)$. Then, by [3, Theorem 23] and [25, Theorem 4.7], the operator A is definitizable. A similar reasoning applies to the case $\infty \in \sigma_{++}(A)$. □

3. Framework and preliminary results

Throughout this paper we make the following assumptions.

(A1) The stiffness operator $A_0 : \mathcal{D}(A_0) \subset H \rightarrow H$ is a self-adjoint uniformly positive operator on a Hilbert space H . We define $H_{\frac{1}{2}} = \mathcal{D}(A_0^{1/2})$ equipped with the norm $\|\cdot\|_{H_{\frac{1}{2}}} := \|A_0^{1/2} \cdot\|_H$ and $H_{-\frac{1}{2}} = H_{\frac{1}{2}}^*$. Here the duality is taken with respect to the pivot space H , that is, equivalently $H_{-\frac{1}{2}}$ is the completion of H with respect to the norm $\|z\|_{H_{-\frac{1}{2}}} = \|A_0^{-1/2}z\|_H$. Thus A_0 restricts to a bounded operator $A_0 : H_{\frac{1}{2}} \rightarrow H_{-\frac{1}{2}}$. We use the same notation A_0 to denote this restriction.

We denote the inner product on H by $\langle \cdot, \cdot \rangle_H$ or $\langle \cdot, \cdot \rangle$, and the duality pairing on $H_{-\frac{1}{2}} \times H_{\frac{1}{2}}$ by $\langle \cdot, \cdot \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}}$. Note that for $(z', z) \in H \times H_{\frac{1}{2}}$, we have

$$\langle z', z \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} = \langle z', z \rangle_H.$$

(A2) The damping operator $D : H_{\frac{1}{2}} \rightarrow H_{-\frac{1}{2}}$ is a bounded operator such that $A_0^{-1/2}DA_0^{-1/2}$ is a bounded self-adjoint operator in H and satisfies

$$\langle Dz, z \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} \geq 0, \quad z \in H_{\frac{1}{2}}.$$

The equation (1) is equivalent to the following standard first-order equation

$$\dot{x}(t) = \mathcal{A}x(t) \tag{4}$$

where $\mathcal{A} : \mathcal{D}(\mathcal{A}) \subset H_{\frac{1}{2}} \times H \rightarrow H_{\frac{1}{2}} \times H$, is given by

$$\mathcal{A} = \begin{bmatrix} 0 & I \\ -A_0 & -D \end{bmatrix},$$

$$\mathcal{D}(\mathcal{A}) = \left\{ \begin{pmatrix} z \\ w \end{pmatrix} \in H_{\frac{1}{2}} \times H_{\frac{1}{2}} \mid A_0z + Dw \in H \right\}.$$

The operator \mathcal{A} itself is not self-adjoint in the Hilbert space $H_{\frac{1}{2}} \times H$. It is easy to see (e.g. [42]) that \mathcal{A} has a bounded inverse in $H_{\frac{1}{2}} \times H$ given by

$$\mathcal{A}^{-1} = \begin{bmatrix} -A_0^{-1}D & -A_0^{-1} \\ I & 0 \end{bmatrix}, \quad (5)$$

where $A_0^{-1}D$ is considered as an operator acting in $H_{\frac{1}{2}}$. This together with the fact that

$$J\mathcal{A}, \quad \text{where } J = \begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix},$$

is a symmetric operator in the Hilbert space $H_{\frac{1}{2}} \times H$, imply the self-adjointness of $J\mathcal{A}$ in $H_{1/2} \times H$. Therefore, (compare also [40, Proof of Lemma 4.5])

$$\mathcal{A}^* = J\mathcal{A}J, \quad \text{with } \mathcal{D}(\mathcal{A}^*) = J\mathcal{D}(\mathcal{A})$$

and

$$\operatorname{Re} \langle \mathcal{A}x, x \rangle \leq 0 \quad \text{for } x \in \mathcal{D}(\mathcal{A}) \quad \text{and} \quad \operatorname{Re} \langle \mathcal{A}^*x, x \rangle \leq 0 \quad \text{for } x \in \mathcal{D}(\mathcal{A}^*).$$

Hence, \mathcal{A} is the generator of a strongly continuous semigroup of contractions on the state space $H_{\frac{1}{2}} \times H$. This fact is well-known, see e.g. [4], [5], [11], [17], [34] or [42].

For $\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \in H_{\frac{1}{2}} \times H$ we define an indefinite inner product on $H_{\frac{1}{2}} \times H$ by

$$\left[\begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \right] := \left\langle J \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_2 \\ y_2 \end{pmatrix} \right\rangle = \langle x_1, x_2 \rangle_{H_{\frac{1}{2}}} - \langle y_1, y_2 \rangle. \quad (6)$$

Then $(H_{\frac{1}{2}} \times H, [\cdot, \cdot])$ is a Krein space and \mathcal{A} is a self-adjoint operator with respect to $[\cdot, \cdot]$.

In the following proposition we collect the above considerations.

PROPOSITION 3.1. *The operator \mathcal{A} is self-adjoint in the Krein space $(H_{\frac{1}{2}} \times H, [\cdot, \cdot])$, its spectrum is contained in the closed left half plane and lies symmetric with respect to the real line. The operator \mathcal{A} has a bounded inverse and is the generator of a strongly continuous semigroup of contractions on $H_{\frac{1}{2}} \times H$.*

This implies that the spectrum of \mathcal{A} is a subset of the closed left half plane without the origin and symmetric with respect to the real axis. However, otherwise the spectrum of \mathcal{A} is quite arbitrary. For an example with $\sigma(\mathcal{A}) = \{s \in \mathbb{C} \mid \operatorname{Re} s \leq 0, |s| \geq \varepsilon\}$, $\varepsilon > 0$, we refer to [24].

In the following theorem we give an estimate for the neighbourhood of the origin which lies in the resolvent set of \mathcal{A} and for the modulus of the eigenvalues of \mathcal{A} .

THEOREM 3.2. *We have $\lambda \in \rho(\mathcal{A})$ if and only if the operator $I + \lambda A_0^{-1}(D + \lambda I)$, considered as an operator in $\mathcal{L}(H_{\frac{1}{2}})$, is boundedly invertible. In particular*

$$\left\{ \lambda \in \mathbb{C} \mid \|\lambda A_0^{-1}D + \lambda^2 A_0^{-1}\|_{\mathcal{L}(H_{\frac{1}{2}})} < 1 \right\} \subset \rho(\mathcal{A}). \tag{7}$$

Moreover, each $\lambda \in \sigma_p(\mathcal{A})$ satisfies

$$|\lambda| \geq \frac{2}{\sqrt{\|A_0^{-1}D\|_{\mathcal{L}(H_{\frac{1}{2}})}^2 + 4\|A_0^{-1}\|_{\mathcal{L}(H)} + \|A_0^{-1}D\|_{\mathcal{L}(H_{\frac{1}{2}})}}}. \tag{8}$$

Proof. Let $\lambda \in \rho(\mathcal{A})$. Then by [24, Proposition 2.2] the operator

$$\lambda^2 A_0^{-1} + \lambda A_0^{-1/2} D A_0^{-1/2} + I$$

is bounded and boundedly invertible in H , hence

$$A_0^{-1/2} \left(\lambda^2 A_0^{-1} + \lambda A_0^{-1/2} D A_0^{-1/2} + I \right) A_0^{1/2} = I + \lambda A_0^{-1}(D + \lambda I)$$

is boundedly invertible in $H_{\frac{1}{2}}$. For the contrary choose $\lambda \in \mathbb{C}$, $\begin{pmatrix} u \\ v \end{pmatrix} \in \mathcal{D}(\mathcal{A})$ and $\begin{pmatrix} x \\ y \end{pmatrix} \in H_{\frac{1}{2}} \times H$. Then we have

$$(\mathcal{A} - \lambda I) \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$$

if and only if the following equations hold

$$v = x + \lambda u$$

$$-A_0(I + \lambda A_0^{-1}(D + \lambda I))u = y + Dx + \lambda x.$$

This implies the first assertion of Theorem 3.2. Let $\lambda \in \sigma_p(\mathcal{A})$. Then the above calculations imply that the operator $I + \lambda A_0^{-1}(D + \lambda I)$ in $H_{\frac{1}{2}}$ is not injective. Therefore, there exists a non-zero vector $f \in H_{\frac{1}{2}}$ with

$$f = -\lambda A_0^{-1}(D + \lambda I)f.$$

Hence,

$$\|f\|_{H_{\frac{1}{2}}} \leq |\lambda| \left(\|A_0^{-1}D\|_{\mathcal{L}(H_{\frac{1}{2}})} + |\lambda| \|A_0^{-1}\|_{\mathcal{L}(H_{\frac{1}{2}})} \right) \|f\|_{H_{\frac{1}{2}}}$$

and, as $\|A_0^{-1}\|_{\mathcal{L}(H_{\frac{1}{2}})} = \|A_0^{-1}\|_{\mathcal{L}(H)}$, we conclude

$$\left(|\lambda| + \frac{\|A_0^{-1}D\|_{\mathcal{L}(H_{\frac{1}{2}})}}{2\|A_0^{-1}\|_{\mathcal{L}(H)}} \right)^2 - \frac{1}{\|A_0^{-1}\|_{\mathcal{L}(H)}} - \frac{\|A_0^{-1}D\|_{\mathcal{L}(H_{\frac{1}{2}})}^2}{4\|A_0^{-1}\|_{\mathcal{L}(H)}^2} \geq 0$$

and Theorem 3.2 is proved. □

REMARK 3.3. The estimate (8) for the eigenvalues is optimal since in the case $D = 0$ it follows that μ is an eigenvalue of A_0 if and only if $\pm i\sqrt{\mu}$ are eigenvalues of \mathcal{A} . If the uniformly positive operator A_0 has a compact resolvent, then the smallest eigenvalue of A_0 equals $\|A_0^{-1}\|^{-1}$ and the eigenvalue λ_{\min} of \mathcal{A} with smallest absolute value is given by

$$|\lambda_{\min}| = \sqrt{\min\{\mu \mid \mu \text{ eigenvalue of } A_0\}} = \sqrt{\|A_0^{-1}\|^{-1}}$$

which is equal to the right hand side of (8) if D is set to be 0.

4. Analyticity

Throughout this section we assume that A_0^{-1} is a compact operator. Note that $A_0^{-1}D$, considered as an operator acting in $H_{\frac{1}{2}}$, is a bounded non-negative operator. In [24, Theorem 4.1] it is shown that under this assumption for $\lambda \in \mathbb{C} \setminus \{0\}$ we have

$$\lambda \in \sigma_{ess}(-A_0^{-1}D) \quad \text{if and only if} \quad 1/\lambda \in \sigma_{ess}(\mathcal{A}). \tag{9}$$

If not explicitly stated otherwise, the operator $A_0^{-1}D$ is always considered as an operator acting on $H_{1/2}$.

We obtain the following main result concerning analyticity.

THEOREM 4.1. *Assume that A_0^{-1} is compact in H and that $0 \notin \sigma_{ess}(A_0^{-1}D)$. Then \mathcal{A} generates an analytic semigroup on $H_{1/2} \times H$.*

The proof of this theorem will be given at the end of this section. We first prove some properties of the point infinity. The following theorem shows in particular that $\infty \in \sigma_{--}(\mathcal{A})$ if $0 \notin \sigma_{ess}(A_0^{-1}D)$.

THEOREM 4.2. *Assume that the operator A_0^{-1} is a compact operator in H and that $0 \notin \sigma_{ess}(A_0^{-1}D)$. Then*

$$\infty \in \sigma_{--}(\mathcal{A}) \quad \text{and} \quad \mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A}) \cup \rho(\mathcal{A}).$$

Moreover, the operator \mathcal{A} is definitizable and there exists a neighbourhood \mathcal{U} of ∞ in $\overline{\mathbb{C}}$ and constants $M > 0$, $m \in \mathbb{N}$ and $\eta > 0$ such that

$$\mathcal{U} \setminus \overline{\mathbb{R}} \subset \rho(\mathcal{A}) \quad \text{and} \quad \mathcal{U} \cap \mathbb{R} \subset \sigma_{--}(\mathcal{A}) \cup \rho(\mathcal{A}) \tag{10}$$

and

$$\|(\mathcal{A} - \lambda I)^{-1}\| \leq \frac{M}{|\operatorname{Im} \lambda|} \quad \text{for all } \lambda \in \mathcal{U} \setminus \overline{\mathbb{R}}, \tag{11}$$

$$\|(\mathcal{A} - \lambda I)^{-1}\| \leq \frac{M}{|\operatorname{Im} \lambda|^m} \quad \text{for all } \lambda \in \rho(\mathcal{A}) \setminus \mathbb{R} \text{ with } |\operatorname{Im} \lambda| \leq \eta. \tag{12}$$

Further, the non-real spectrum of \mathcal{A} consists of at most finitely many points which belong to $\sigma_{p,norm}(\mathcal{A})$.

Proof. The proof is divided into two steps. First we will prove that $\infty \in \sigma_{--}(\mathcal{A})$. In the second step we will show that $\mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A}) \cup \rho(\mathcal{A})$. Since by (9) the essential spectrum of \mathcal{A} is real, Theorem 2.3 yields that \mathcal{A} is a definitizable operator and the non-real spectrum of \mathcal{A} consists of at most finitely many points which belong to $\sigma_{p,norm}(\mathcal{A})$. Further, (10), (11) and (12) follow from [3, Lemma 2 and Proposition 3] and from [32, Proposition II.2.1].

Step 1. By [3, Lemma 10], ∞ belongs to $\sigma_{--}(\mathcal{A})$ if and only if ∞ belongs to $\sigma_{\pi_-}(\mathcal{A})$. It is easily seen (see e.g. [1]) that this is the case if and only if $0 \in \sigma_{\pi_-}(\mathcal{A}^{-1})$. Assume $0 \notin \sigma_{\pi_-}(\mathcal{A}^{-1})$. Then there exists a sequence $((\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})) \subset H_{\frac{1}{2}} \times H$ with $\|(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})\|_{H_{\frac{1}{2}} \times H}^2 = \|x_n\|_{H_{\frac{1}{2}}}^2 + \|y_n\|^2 = 1$ and $\mathcal{A}^{-1}(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}) \rightarrow 0$ as $n \rightarrow \infty$ and

$$\limsup_{n \rightarrow \infty} [(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}), (\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})] = \limsup_{n \rightarrow \infty} \left(\|x_n\|_{H_{\frac{1}{2}}}^2 - \|y_n\|^2 \right) \geq 0. \tag{13}$$

By [3, Theorem 14] this sequence can be chosen to converge to zero weakly. This gives

$$\|A_0^{-1}Dx_n + A_0^{-1}y_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{and} \quad \|x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{14}$$

The sequence $(A_0^{-1/2}y_n)$ converges weakly to zero in $H_{\frac{1}{2}}$. As A_0^{-1} is a compact operator in H , $A_0^{-1/2}$ is a compact operator in $H_{\frac{1}{2}}$. It follows that $(A_0^{-1}y_n)$ converges to zero in $H_{\frac{1}{2}}$. Then, by (14), we have

$$\|A_0^{-1}Dx_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Moreover, the sequence (x_n) converges weakly to zero in $H_{\frac{1}{2}}$, hence the assumption $0 \notin \sigma_{ess}(A_0^{-1}D)$ implies

$$\|x_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then $\|y_n\| \rightarrow 1$ as $n \rightarrow \infty$, in contradiction to (13) and $0 \in \sigma_{\pi_-}(\mathcal{A}^{-1})$ follows.

Step 2. We now choose $\mu \in (-\infty, 0)$ and

$$G_\mu := \text{span} \{x \in H_{\frac{1}{2}} \mid A_0x = \mu x, \mu \leq \mu^2\}.$$

Then G_μ is a finite dimensional subspace of $H_{\frac{1}{2}}$. For every sequence $((\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}))$ in $\mathcal{D}(\mathcal{A}) \cap (G_\mu \times G_\mu)^\perp$ with $\|(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})\|_{H_{\frac{1}{2}} \times H}^2 = 1$ and $(\mathcal{A} - \mu I)(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}) \rightarrow 0$ as $n \rightarrow \infty$ we have

$$\|y_n - \mu x_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{and} \quad \|A_0x_n + Dy_n + \mu y_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This gives

$$\begin{aligned} \liminf_{n \rightarrow \infty} [(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}), (\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})] &= \liminf_{n \rightarrow \infty} \left(\langle x_n, x_n \rangle_{H_{\frac{1}{2}}} - \langle y_n, y_n \rangle \right) \\ &= \liminf_{n \rightarrow \infty} \left(\langle A_0x_n, x_n \rangle - \mu^2 \langle x_n, x_n \rangle \right) > 0, \end{aligned}$$

where the last inequality follows from the fact that $x_n \in G_\mu^\perp, n \in \mathbb{N}$. Therefore $\mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A})$ and Theorem 4.2 is proved. \square

REMARK 4.3. The stronger assumption $0 \notin \sigma(A_0^{-1}D)$ implies that there exist constants $\alpha, \gamma > 0$ with

$$\gamma \langle A_0 x, x \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} \leq \langle Dx, x \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} \leq \alpha \langle A_0 x, x \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} \text{ for } x \in H_{\frac{1}{2}}.$$

Proof of Theorem 4.1. Since \mathcal{A} is the generator of a strongly continuous semigroup, estimate (11) shows immediately that \mathcal{A} generates an analytic semigroup, see [15, Chapter II, Section 4.5]. \square

The following corollary shows that under the assumptions of Theorem 4.2 the operator \mathcal{A} can be written as a direct sum of a self-adjoint operator on a Hilbert space and a bounded self-adjoint operator on a Pontryagin space. In the situation of $D = \rho A_0^\alpha, \rho > 0$ and $\alpha \in (0, 1], \mathcal{A}$ is the direct sum of two normal operators [12].

COROLLARY 4.4. Assume that the operator A_0^{-1} is a compact operator in H and that $0 \notin \sigma_{ess}(A_0^{-1}D)$. Then the space $(H_{\frac{1}{2}} \times H, [\cdot, \cdot])$ decomposes into the direct sum of two \mathcal{A} -invariant closed subspaces H' and H'' , which are orthogonal with respect to $[\cdot, \cdot]$, such that:

1. The space $(H', -[\cdot, \cdot])$ is a Hilbert space, $\mathcal{A}|_{H'}$ is a self-adjoint operator in this Hilbert space and

$$\sigma(\mathcal{A}|_{H'}) \subset \overline{\mathbb{R}} \setminus (-M, \infty),$$

where M is as in (15).

2. The space $(H'', [\cdot, \cdot])$ is a Pontryagin space, $\mathcal{A}|_{H''}$ is a bounded self-adjoint operator in this Pontryagin space with

$$\sigma(\mathcal{A}|_{H''}) \subset [-M, 0) \cup \Theta \text{ and } \sigma(\mathcal{A}|_{H''}) \subset \sigma_{++}(\mathcal{A}|_{H''}) \cup \Xi$$

where $\Xi, \Theta \subset \mathbb{C}$ are empty or consist of finitely many points and $\Theta \subset \sigma_{p,norm}(\mathcal{A}|_{H''})$.

Proof. By Theorem 4.2. the operator \mathcal{A} is definitizable with $\infty \in \sigma_{--}(\mathcal{A})$ and $\mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A}) \cup \rho(\mathcal{A})$. Denote by E the spectral function of \mathcal{A} . Since $\infty \in \sigma_{--}(\mathcal{A})$, there exists $M > 0$ with

$$\overline{\mathbb{R}} \setminus (-M, M) \subset \sigma_{--}(\mathcal{A}) \cup \rho(\mathcal{A}), \tag{15}$$

see [3, Lemma 2]. Set $\Delta_0 := \overline{\mathbb{R}} \setminus [-M, M], H' := E(\Delta_0)(H_{\frac{1}{2}} \times H)$ and $H'' := (I - E(\Delta_0))(H_{\frac{1}{2}} \times H)$. Then the assertions above follow from [25, Theorem 3.18] and Theorem 3.1. \square

5. Expansion in eigenfunctions

In the sequel we always assume the Hilbert space H to be separable. An at most countable set \mathcal{M} of elements of a Hilbert space is said to be a *Riesz basis* if there exists an isomorphic mapping \mathcal{M} onto an orthonormal basis, cf. [38, Lecture VI].

Condition (17) of Theorem 5.1 below appears already in the celebrated works [27, 28], where the case of a bounded self-adjoint operator D and a positive compact operator A_0 was discussed. We will use this approach in the proof of Theorem 5.1.

THEOREM 5.1. *Assume that the operator A_0^{-1} is compact in H and that*

$$0 \notin \sigma_{ess}(A_0^{-1}D), \quad (16)$$

where $A_0^{-1}D$ is considered as an operator acting in $H_{\frac{1}{2}}$. Assume that the set $\sigma_{ess}(A_0^{-1}D)$ is countable and has at most countably many accumulation points. Moreover, let at least one of the following conditions be satisfied.

(a) *There exists a $\delta > 0$ such that for all $f \in H_{\frac{1}{2}}$ with $\|f\|_{H_{\frac{1}{2}}} = 1$ we have*

$$\left\langle A_0^{-1}Df, f \right\rangle_{H_{\frac{1}{2}}}^2 - 4 \left\langle A_0^{-1}f, f \right\rangle_{H_{\frac{1}{2}}} > \delta. \quad (17)$$

(b) *For all $\mu \in \sigma_{ess}(-A_0^{-1}D)$ we have either $\frac{1}{\mu} \notin \sigma_p(\mathcal{A})$ or, if $\frac{1}{\mu} \in \sigma_p(\mathcal{A})$, there exists no non-zero $\begin{pmatrix} y \\ \mu^{-1}y \end{pmatrix} \in \ker(\mathcal{A} - \mu^{-1}I)$ such that*

$$\mu^2 \langle y, w \rangle_{H_{\frac{1}{2}}} = \langle y, w \rangle \quad \text{for all } \begin{pmatrix} w \\ \mu^{-1}w \end{pmatrix} \in \ker(\mathcal{A} - \mu^{-1}I). \quad (18)$$

(c) $\|A_0^{-1/2}\| < \inf\{\lambda > 0 \mid \lambda \in \sigma_{ess}(A_0^{-1}D)\}$.

Then the following assertions hold.

1. *There exists a subspace of $H_{\frac{1}{2}} \times H$ of at most finite codimension which has a Riesz basis consisting of eigenvectors of \mathcal{A} .*
2. *There exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors and finitely many associated vectors of \mathcal{A} .*
3. *Moreover, if (a) holds, then \mathcal{A} has no associated vectors, i.e. there are no Jordan chains of length greater than one, the spectrum of \mathcal{A} is real and there exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors of \mathcal{A} .*

As a corollary we obtain the following result

COROLLARY 5.2. *Assume that the operator D has the form*

$$D = \alpha A_0 + B,$$

where $\alpha > 0$ is a constant and B is a symmetric A_0 -compact operator. If $-1/\alpha \notin \sigma_p(\mathcal{A})$, then there exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors and finitely many associated vectors of \mathcal{A} , and \mathcal{A} generates an analytic semigroup.

Proof. We define $H_1 = \mathcal{D}(A_0)$ equipped with the norm $\|\cdot\|_{H_1} := \|A_0 \cdot\|$ and H_{-1} is the completion of H with respect to the norm $\|\cdot\|_{H_{-1}} := \|A_0^{-1} \cdot\|$. Then, by assumption, B , restricted from H_1 to H , is a compact operator, cf. [26, IV 1.12]. By the symmetry of B , B^* is an extension of B and a compact operator acting from H into H_{-1} . Thus, by interpolation, the operator B considered as an operator from $H_{\frac{1}{2}}$ to $H_{-\frac{1}{2}}$ is compact, hence $\sigma_{ess}(A_0^{-1}D) = \alpha$ and Corollary 5.2 follows from Theorem 5.1. \square

In [30] it is shown that under the assumption of the Corollary 5.2 there exists a Riesz basis of $H_{\frac{1}{2}} \times H_{\frac{1}{2}}$ consisting of eigenvectors and finitely many associated vectors of \mathcal{A} , and \mathcal{A} generates an analytic semigroup.

Proof of Theorem 5.1. It suffices to prove Part 1 and 3 of the theorem, as Part 2 follows immediately from Part 1.

Assume that condition (a) holds. By Theorem 4.2 we have $\mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A}) \cup \rho(A)$. Let $\lambda \in \sigma_{\pi_+}(\mathcal{A})$. Then there exists a sequence $((\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}))$ in $\mathcal{D}(\mathcal{A})$ with $\|(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})\|_{H_{\frac{1}{2}} \times H}^2 = 1$ and $(\mathcal{A} - \lambda I)(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}) \rightarrow 0$ as $n \rightarrow \infty$. Hence

$$\|y_n - \lambda x_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{and} \quad \|A_0 x_n + \lambda D x_n + \lambda^2 x_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

This implies

$$\|x_n + \lambda A_0^{-1} D x_n + \lambda^2 A_0^{-1} x_n\|_{H_{\frac{1}{2}}} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{19}$$

and

$$\begin{aligned} \liminf_{n \rightarrow \infty} [(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}), (\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})] &= \liminf_{n \rightarrow \infty} \left(\langle x_n, x_n \rangle_{H_{\frac{1}{2}}} - \langle y_n, y_n \rangle \right) \\ &= \liminf_{n \rightarrow \infty} \left(\langle x_n, x_n \rangle_{H_{\frac{1}{2}}} - \lambda^2 \langle x_n, x_n \rangle \right) \\ &= \liminf_{n \rightarrow \infty} -\lambda \left(\langle A_0^{-1} D x_n, x_n \rangle_{H_{\frac{1}{2}}} + 2\lambda \langle A_0^{-1} x_n, x_n \rangle_{H_{\frac{1}{2}}} \right). \end{aligned}$$

Similarly, we have

$$\limsup_{n \rightarrow \infty} [(\begin{smallmatrix} x_n \\ y_n \end{smallmatrix}), (\begin{smallmatrix} x_n \\ y_n \end{smallmatrix})] = \limsup_{n \rightarrow \infty} -\lambda \left(\langle A_0^{-1} D x_n, x_n \rangle_{H_{\frac{1}{2}}} + 2\lambda \langle A_0^{-1} x_n, x_n \rangle_{H_{\frac{1}{2}}} \right).$$

Now (a) implies $\sigma(\mathcal{A}) \subset \mathbb{R}$ (see, e.g. [24, Theorem 3.3]), hence, by Proposition 3.1, we have $\lambda \in (-\infty, 0)$. Moreover, by (a), the operator pencil

$$L(s) := s^2I + sA_0^{-1}D + A_0^{-1}, \quad s \in \mathbb{C},$$

considered as a pencil with values in the bounded operators acting on $H_{\frac{1}{2}}$, is strongly hyperbolic, see e.g. [37, Lemma 31.23]. Therefore, see e.g. [31], we have

$$\liminf_{n \rightarrow \infty} \left(\langle A_0^{-1}Dx_n, x_n \rangle_{H_{\frac{1}{2}}} + 2\lambda \langle A_0^{-1}x_n, x_n \rangle_{H_{\frac{1}{2}}} \right) > 0$$

or

$$\limsup_{n \rightarrow \infty} \left(\langle A_0^{-1}Dx_n, x_n \rangle_{H_{\frac{1}{2}}} + 2\lambda \langle A_0^{-1}x_n, x_n \rangle_{H_{\frac{1}{2}}} \right) < 0.$$

This gives

$$\sigma(\mathcal{A}) \subset \sigma_{++}(\mathcal{A}) \cup \sigma_{--}(\mathcal{A}) \subset \mathbb{R}$$

and \mathcal{A} has no associated vectors and there exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors of \mathcal{A} .

Assume that condition (b) holds and let $\mu \in \sigma_{ess}(-A_0^{-1}D)$ such that $\frac{1}{\mu} \in \sigma_p(\mathcal{A})$. Now, (b) implies that there are no Jordan chains of \mathcal{A} corresponding to the eigenvalue $\frac{1}{\mu}$ of length greater than one, and, moreover, that $\ker(\mathcal{A} - \mu^{-1}I)$ is a non-degenerate subspace of $(H_{\frac{1}{2}} \times H, [\cdot, \cdot])$, that is,

$$\ker(\mathcal{A} - \mu^{-1}I) \cap \left(\ker(\mathcal{A} - \mu^{-1}I) \right)^{[\perp]} = \{0\},$$

where $\left(\ker(\mathcal{A} - \mu^{-1}I) \right)^{[\perp]}$ is the orthogonal companion of $\ker(\mathcal{A} - \mu^{-1}I)$ with respect to $[\cdot, \cdot]$. Moreover, (18) implies that μ^{-1} is a regular critical point of \mathcal{A} , see [32] or [14, Proposition 1.4]. As all points from $\sigma(\mathcal{A}) \setminus \sigma_{ess}(\mathcal{A})$ belong to $\sigma_{p,norm}(\mathcal{A})$, it turns out that \mathcal{A} has only regular critical points and the eigenvectors of \mathcal{A} form a Riesz basis of a subspace of $H_{\frac{1}{2}} \times H$ of an at most finite codimension. The eigenvectors and associated vectors of \mathcal{A} form a Riesz basis of $H_{\frac{1}{2}} \times H$.

Condition (c) implies (b), hence Theorem 5.1 is proved. \square

Theorem 5.1 implies that the operator \mathcal{A} is the direct sum of an operator similar to a self-adjoint operator in a Hilbert space and a bounded operator in a finite-dimensional space.

REMARK 5.3. Assume that the operator A_0^{-1} is compact in H and that (16) holds. Then it was shown in the proof of Theorem 5.1 that if (i) holds or if for all $\mu^{-1} \in \sigma_p(\mathcal{A})$ we have that (18) holds, all critical points of \mathcal{A} are regular and there are no associated vectors. Hence, \mathcal{A} is similar to a self-adjoint operator in the Hilbert space $H_{\frac{1}{2}} \times H$.

REMARK 5.4. We mention that Theorem 5.1 can be obtained also by methods from [2]. For this, one has to show that the operator \mathcal{A}^{-1} belongs to the class **(H)**, cf. [2, Chapter 3, §5], and then apply [2, Theorem 4.2.12].

6. Example: Euler-Bernoulli Beam with distributed Kelvin-Voigt damping

We consider a beam of length 1 with a thin film of piezoelectric polymer applied to one side and we study transverse vibrations only. Let $u(r, t)$ denote the deflection of the beam from its rigid body motion at time t and position r . Use of the Euler-Bernoulli model for the beam deflection and the Kelvin-Voigt damping model leads to the following description of the vibrations [6], [43]:

$$\frac{\partial^2 u}{\partial t^2} + \frac{\partial^2}{\partial r^2} \left[E \frac{\partial^2 u}{\partial r^2} + C_d \frac{\partial^3 u}{\partial r^2 \partial t} \right] = 0, \quad r \in (0, 1), t > 0. \quad (20)$$

Here the flexural rigidity E is a positive physical constant which is determined by the beam's area momentum of inertia and its modulus of elasticity and $C_d \in L^\infty(0, 1)$ with $C_d(t) \geq c > 0$ a.e. describes the damping properties of the piezoelectric film. Assuming that the beam is pinned at point 0 and sliding at 1, we have for all $t > 0$ the following boundary conditions:

$$u|_{r=0} = 0, \quad \frac{\partial u}{\partial r}|_{r=1} = 0, \quad \frac{\partial^2 u}{\partial r^2}|_{r=0} = 0, \quad \frac{\partial^3 u}{\partial r^3}|_{r=1} = 0. \quad (21)$$

We consider the partial differential equation (20)-(21) as a second order problem in the Hilbert space $H = L^2(0, 1)$. In H we define the operator A_0 by

$$A_0 = E \frac{d^4}{dr^4}, \quad \mathcal{D}(A_0) = \left\{ z \in H^4(0, 1) \mid z(0) = z'(1) = z''(0) = z'''(1) = 0 \right\}.$$

It is easy to see that the operator A_0 satisfies assumption (A1) and that A_0^{-1} is a compact operator. We have

$$H_{\frac{1}{2}} = \left\{ z \in H^2(0, 1) \mid z(0) = z'(1) = 0 \right\}$$

with inner product $\langle z, v \rangle_{H_{\frac{1}{2}}} = E \langle z'', v'' \rangle$. The operator $A_0^{1/2}$ is given by

$$A_0^{1/2} = E^{1/2} \frac{d^2}{dr^2} \quad \text{and} \quad \|z\|_{H_{\frac{1}{2}}}^2 \geq \frac{\pi^4 E}{16} \|z\|^2 \quad \text{for } z \in H_{\frac{1}{2}}. \quad (22)$$

Let $x(t) = (u(\cdot, t), \dot{u}(\cdot, t))$. Then $\|x(t)\|_{H_{\frac{1}{2}} \times H}^2 = \|u''(\cdot, t)\|^2 + \|\dot{u}(\cdot, t)\|^2$ corresponds to the energy of the beam which justifies the choice of $L^2(0, 1)$ as the Hilbert space for the analysis of the boundary value problem (20)-(21).

By $M_{C_d} \in \mathcal{L}(H)$ we denote the multiplication operator

$$(M_{C_d} f)(x) = C_d(x) f(x)$$

and we define the damping operator as

$$D = \frac{1}{E} A_0^{1/2} M_{C_d} A_0^{1/2}.$$

For $z \in H_{\frac{1}{2}}$ we have

$$\langle Dz, z \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}} = \langle C_d z'', z'' \rangle_H \geq c \|z''\|^2 \geq \frac{\pi^4}{16} c \|z\|^2, \tag{23}$$

and thus the assumption (A2) holds as well. Furthermore, each solution of the abstract problem $\ddot{z}(t) + A_0 z(t) + D\dot{z}(t) = 0$ corresponds to a solution of the boundary value problem (20)-(21). We have the following lemma.

LEMMA 6.1. *The operators $A_0^{-1/2} M_{C_d} A_0^{-1/2}$ and $A_0^{-1} D$ are bounded self-adjoint operators in H and $H_{\frac{1}{2}}$, respectively, with*

$$\sigma_{ess}(A_0^{-1} D) = \sigma_{ess}(A_0^{-1/2} M_{C_d} A_0^{-1/2}) = \sigma(E^{-1} M_{C_d}).$$

Proof. A vector $x \in H_{\frac{1}{2}}$ belongs to $\ker(A_0^{-1} D - \lambda I)$ if and only if $A_0^{1/2} x$ belongs to $\ker(A_0^{-1/2} D A_0^{-1/2} - \lambda I)$. The operator $A_0^{1/2}$ maps $H_{\frac{1}{2}}$ isometrically onto H , therefore

$$\dim \ker (A_0^{-1} D - \lambda I) = \dim \ker (A_0^{-1/2} D A_0^{-1/2} - \lambda I).$$

Obviously, $A_0^{-1/2} D A_0^{-1/2} = M_{C_d}$ and $A_0^{-1} D$ are bounded self-adjoint operators in H and $H_{\frac{1}{2}}$, respectively, and therefore we have for real λ

$$\begin{aligned} \operatorname{codim} \operatorname{ran} (A_0^{-1} D - \lambda I) &= \dim \ker (A_0^{-1} D - \lambda I) \\ &= \dim \ker (A_0^{-1/2} D A_0^{-1/2} - \lambda I) \\ &= \operatorname{codim} \operatorname{ran} (A_0^{-1/2} D A_0^{-1/2} - \lambda I). \quad \square \end{aligned}$$

Lemma 6.1 implies $0 \notin \sigma_{ess}(A_0^{-1} D)$, as the function C_d satisfies $C_d(t) \geq c > 0$ a.e. By (23), the corresponding operator \mathcal{A} generates an exponentially stable semigroup on $H_{\frac{1}{2}} \times L^2(0, 1)$ (see the introduction). Moreover, the assumptions of Theorem 4.1 are satisfied and thus \mathcal{A} generates an analytic semigroup. By Theorem 4.2, \mathcal{A} is definitizable, $\infty \in \sigma_{--}(\mathcal{A})$, $\mathbb{R} \subset \sigma_{\pi_+}(\mathcal{A}) \cup \rho(\mathcal{A})$ and the non-real spectrum of \mathcal{A} consists of at most finitely many points which belong to $\sigma_{p,norm}(\mathcal{A})$.

In addition, we now assume that the film on the beam consists of several patches, that is,

$$C_d(x) = \sum_{k=1}^n a_k \chi_{A_k}(x),$$

where $n \in \mathbb{N}$, $a_k > 0$, $k = 1, \dots, n$, A_k are measurable disjoint subsets of $(0, 1)$ and

$$\overline{\bigcup_{k=1}^n A_k} = [0, 1].$$

THEOREM 6.2. *If*

$$a_k > \frac{8}{\pi^2 \sqrt{E}} \quad (24)$$

holds for $k = 1, \dots, n$, then (a) from Theorem 5.1 is satisfied, that is, the spectrum of \mathcal{A} is real and there exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors of \mathcal{A} . If for all a_k , $k = 1, \dots, n$, with

$$a_k \leq \frac{4}{\pi^2} \sqrt{E}$$

we have $-E/a_k \notin \sigma_p(\mathcal{A})$, then (b) of Theorem 5.1 is satisfied, that is, there exists a Riesz basis of $H_{\frac{1}{2}} \times H$ consisting of eigenvectors and finitely many associated vectors of \mathcal{A} . In particular, this holds true if we have

$$a_k > \frac{4}{\pi^2} \sqrt{E} \quad \text{for } k = 1, \dots, n.$$

Proof. By (22) and (24), we have for $f \in H_{\frac{1}{2}}$

$$\begin{aligned} \langle A_0^{-1} Df, f \rangle_{H_{\frac{1}{2}}}^2 &= \langle Df, f \rangle_{H_{-\frac{1}{2}} \times H_{\frac{1}{2}}}^2 \geq (\min\{a_k \mid k = 1, \dots, n\})^2 \|f\|_{H_{\frac{1}{2}}}^4 \\ &> 4\|f\|^2 \|f\|_{H_{\frac{1}{2}}}^2 + \delta \|f\|_{H_{\frac{1}{2}}}^4 \end{aligned}$$

for some sufficiently small $\delta > 0$ and the first assertion of Theorem 6.2 is proved.

The second and third assertion follow from the fact that for $a_k > \frac{4}{\pi^2} \sqrt{E}$ we have

$$a_k \langle y, y \rangle_{H_{\frac{1}{2}}} > \|y\|^2$$

and (b) (resp. (c)) of Theorem 5.1 is satisfied. \square

There is an obvious generalization of Theorem 6.2 for the case of countably many patches which we do not give here in detail.

REFERENCES

- [1] AZIZOV, T. YA., BEHRNDT, J., JONAS, P. and TRUNK, C., Spectral points of type π_+ and π_- for closed linear relations in Krein spaces. submitted.
- [2] AZIZOV, T. YA. and IOKHVIDOV, I. S., *Linear operators in spaces with an indefinite metric*, Pure and Applied Mathematics (New York). John Wiley & Sons Ltd., Chichester, 1989.

- [3] AZIZOV, T. YA., JONAS, P. and TRUNK, C., *Spectral points of type π_+ and π_- of self-adjoint operators in Krein spaces*, J. Funct. Anal., 226(1) (2005), 114–137.
- [4] BANKS, H. T. and ITO, K., *A unified framework for approximation in inverse problems for distributed parameter systems*, Control Theory Adv. Tech., 4(1) (1988) 73–90.
- [5] BANKS, H. T., ITO, K. and WANG, Y., *Well posedness for damped second-order systems with unbounded input operators*, Differential Integral Equations, 8(3) (1995) 587–606.
- [6] BANKS, H. T., SMITH, R. C. and WANG, Y., *The modeling of piezoceramic patch interactions with shells, plates, and beams*, Quart. Appl. Math., 53(2) (1995), 353–381.
- [7] BÁTKAI, A. and ENGEL, K., *Exponential decay of 2×2 operator matrix semigroups*, J. Comput. Anal. Appl., 6(2) (2004), 153–163.
- [8] BENCHIMOL, C. D., *A note on weak stabilizability of contraction semigroups*, SIAM J. Control Optimization, 16(3) (1978), 373–379.
- [9] BOGNÁR, J., *Indefinite inner product spaces*, Springer-Verlag, New York, 1974. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 78.
- [10] CHEN, G. and RUSSELL, D. L., *A mathematical model for linear elastic systems with structural damping*, Q. Appl. Math., 39 (1982) 433–454.
- [11] CHEN, S., LIU, K. and LIU, Z., *Spectrum and stability for elastic systems with global or local Kelvin-Voigt damping*, SIAM J. Appl. Math., 59(2) (electronic), (1999), 651–668.
- [12] CHEN, S. and TRIGGIANI, R., *Proof of extensions of two conjectures on structural damping for elastic systems*, Pacific J. Math., 136(1) (1989), 15–55.
- [13] CHEN, S. and TRIGGIANI, R., *Characterization of domains of fractional powers of certain operators arising in elastic systems, and applications*, J. Differ. Equations, 88(2) (1990), 279–293.
- [14] DIJKSMA, A. and LANGER, H., *Operator theory and ordinary differential operators*. In Lectures on operator theory and its applications (Waterloo, ON, 1994), volume 3 of Fields Inst. Monogr., pages 73–139. Amer. Math. Soc., Providence, RI, 1996.
- [15] ENGEL, K. and NAGEL, R., *One-parameter semigroups for linear evolution equations*, volume 194 of Graduate Texts in Mathematics. Springer-Verlag, New York, 2000.
- [16] GRINIV, R. O. and SHKALIKOV, A. A., *Operator models in elasticity theory and hydromechanics and the associated analytic semigroups*, Mosc. Univ. Math. Bull., 54(5) (1999), 1–10.
- [17] GRINIV, R. O. and SHKALIKOV, A. A., *Exponential stability of semigroups related to operator models in mechanics*, Mat. Zametki, 73(5) (2003), 657–664.
- [18] HENDRICKSON, E. and LASIECKA, I., *Numerical approximations and regularizations of Riccati equations arising in hyperbolic dynamics with unbounded control operators*, Comput. Optim. Appl., 2(4) (1993), 343–390.
- [19] HENDRICKSON, E. and LASIECKA, I., *Finite-dimensional approximations of boundary control problems arising in partially observed hyperbolic systems*, Dynam. Contin. Discrete Impuls. Systems, 1(1) (1995), 101–142.
- [20] HRYNIV, R. O. and SHKALIKOV, A. A., *Exponential decay of solution energy for equations associated with some operator models of mechanics*, Funct. Anal. Appl., 38(3) (2004), 163–172.
- [21] HUANG, F., *On the mathematical model for linear elastic systems with analytic damping*, SIAM J. Control Optimization, 26(3) (1988), 714–724.
- [22] HUANG, F., *Some problems for linear elastic systems with damping*, Acta Math. Sci., 10(3) (1990), 319–326.
- [23] JACOB, B., MORRIS, K. and TRUNK, C., *Minimum-phase infinite-dimensional second-order systems*, IEEE Transactions on Automatic Control, 52 (2007), 1654–1665.
- [24] JACOB, B. AND TRUNK, C., *Location of the spectrum of operator matrices which are associated to second order equations*, Operators and Matrices, 1 (2007), 45–60.
- [25] JONAS, P., *On locally definite operators in Krein spaces*, In Spectral analysis and its applications, volume 2 of Theta Ser. Adv. Math., pages 95–127. Theta, Bucharest, 2003.
- [26] KATO, T., *Perturbation theory for linear operators*, Springer-Verlag, Berlin, second edition, 1976. Grundlehren der Mathematischen Wissenschaften, Band 132.
- [27] KREIN, M. G. AND LANGER, H., *On some mathematical principles in the linear theory of damped oscillations of continua I*, Integral Equations Oper. Theory, 1 (1978), 364–399.

- [28] KREIN, M. G. and LANGER, H., *On some mathematical principles in the linear theory of damped oscillations of continua II*, Integral Equations Oper. Theory, 1 (1978), 539–566.
- [29] LANCASTER, P., MARKUS, A. S. and MATSAEV, V. I., *Definitizable operators and quasihyperbolic operator polynomials*, J. Funct. Anal., 131(1) (1995), 1–28.
- [30] LANCASTER, P. and SHKALIKOV, A. A., *Damped vibrations of beams and related spectral problems*, Canad. Appl. Math. Quart., 2(1) (1994), 45–90.
- [31] LANGER, H., *Über stark gedämpfte Scharen im Hilbertraum*, J. Math. Mech., 17 (1967/1968), 685–705.
- [32] LANGER, H., *Spectral functions of definitizable operators in Kreĭn spaces*, In *Functional analysis (Dubrovnik, 1981)*, volume 948 of *Lecture Notes in Math.*, pages 1–46. Springer, Berlin, 1982.
- [33] LANGER, H., MARKUS, A. S. and MATSAEV, V. I., *Locally definite operators in indefinite inner product spaces*, Math. Ann., 308(3) (1997), 405–424.
- [34] LASIECKA, I., *Stabilization of wave and plate-like equations with nonlinear dissipation on the boundary*, J. Differential Equations, 79(2) (1989), 340–381.
- [35] LASIECKA, I. and TRIGGIANI, R., *Uniform exponential energy decay of wave equations in a bounded region with $L_2(0, \infty; L_2(\Gamma))$ -feedback control in the Dirichlet boundary conditions*, J. Differential Equations, 66(3) (1987), 340–390.
- [36] LEVAN, N., *The stabilizability problem: a Hilbert space operator decomposition approach*, IEEE Trans. Circuits and Systems, 25(9) (1978), 721–727. Special issue on the mathematical foundations of system theory.
- [37] MARKUS, A. S., *Introduction to the spectral theory of polynomial operator pencils*, volume 71 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1988.
- [38] NIKOL'SKIĬ, N. K., *Treatise on the shift operator*, volume 273 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1986.
- [39] SLEMROD, M., *Stabilization of boundary control systems*, J. Differential Equations, 22(2) (1976), 402–415.
- [40] TUCSNAK, M. and WEISS, G., *How to get a conservative well-posed linear system out of thin air*, II. Controllability and stability. SIAM J. Control Optim., 42(3) (2003), (electronic), 907–935.
- [41] VESELIĆ, K., *Energy decay of damped systems*, ZAMM Z. Angew. Math. Mech., 84(12) (2004), 856–863.
- [42] WEISS, G. AND TUCSNAK, M., *How to get a conservative well-posed linear system out of thin air*, I. Well-posedness and energy balance. ESAIM Control Optim. Calc. Var., 9 (2003), (electronic), 247–274.
- [43] YOU, P. H., *Boundary feedback control of elastic beam equation with structural damping and stability*, Acta Math. Appl. Sinica (English Ser.), 6(4) (1990), 373–382.

Birgit Jacob
Department of Applied Mathematics
Delft University of Technology
P.O. Box 5031, 2600 GA Delft
The Netherlands
e-mail: b.jacob@tudelft.nl

Carsten Trunk
Institut für Mathematik
Technische Universität Ilmenau
Postfach 100565, D-98684 Ilmenau,
Germany
e-mail: carsten.trunk@tu-ilmenau.de

Monika Winklmeier
Mathematisches Institut, Universität Bern
Sidlerstrasse 5
CH-3012 Bern
Switzerland
e-mail: monika.winklmeier@math.unibe.ch