Cytogenetic and Genome Research

Cytogenet Genome Res 2009;126:63–76 DOI: 10.1159/000245907 Accepted after revision: June 15, 2009 by M. Schmid

Molecular Cytogenetics and Gene Mapping in Sheep (*Ovis aries,* 2n = 54)

T. Goldammer^a G.P. Di Meo^c G. Lühken^b C. Drögemüller^d C.H. Wu^e J. Kijas^f B.P. Dalrymple^f F.W. Nicholas^g J.F. Maddox^h L. lannuzzi^c N.E. Cockett^e

^aForschungsbereich Molekularbiologie, Forschungsinstitut für die Biologie landwirtschaftlicher Nutztiere (FBN), Dummerstorf, ^bInstitut für Tierzucht und Haustiergenetik, Justus-Liebig-Universität Giessen, Giessen, Germany; ^cNational Research Council, Laboratory of Animal Cytogenetics and Gene Mapping, Naples, Italy; ^dInstitut für Genetik, Universität Bern, Bern, Switzerland; ^eADVS Department, College of Agriculture, Utah State University, Logan, Utah, USA; ^fCSIRO Livestock Industries, St. Lucia, Qld., ^gFaculty of Veterinary Science, University of Sydney, Sydney, N.S.W., ^hDepartment of Veterinary Science, The University of Melbourne, Melbourne, Vic., Australia

Key Words

Chromosome structure · Genome analysis · Linkage mapping · QTL · Sheep cytogenetic map

Abstract

The development of a completely annotated sheep genome sequence is a key need for understanding the phylogenetic relationships and genetic diversity among the many different sheep breeds worldwide and for identifying genes controlling economically and physiologically important traits. The ovine genome sequence assembly will be crucial for developing optimized breeding programs based on highly productive, healthy sheep phenotypes that are adapted to modern breeding and production conditions. Scientists and breeders around the globe have been contributing to this goal by generating genomic and cDNA libraries, performing genome-wide and trait-associated analyses of polymorphism, expression analysis, genome sequencing, and by developing virtual and physical comparative maps. The International Sheep Genomics Consortium (ISGC), an informal network of sheep genomics researchers, is playing a major role in coordinating many of these activities. In addition to serving as an essential tool for monitoring chromosome abnormalities in specific sheep populations, ovine molecular cytogenetics provides physical anchors which link and order genome regions, such as sequence contigs, genes and polymorphic DNA markers to ovine chromosomes. Likewise, molecular cytogenetics can contribute to the process of defining evolutionary breakpoints between related species. The selective expansion of the sheep cytogenetic map, using loci to connect maps and identify chromosome bands, can substantially contribute to improving the guality of the annotated sheep genome sequence and will also accelerate its assembly. Furthermore, identifying major morphological chromosome anomalies and micro-rearrangements, such as gene duplications or deletions, that might occur between different sheep breeds and other Ovis species will also be important to understand the diversity of sheep chromosome structure and its implications for cross-breeding. To date, 566 loci have been assigned to specific chromosome regions in sheep and the new cytogenetic map is presented as part of this review. This review will also summarize the current cytogenomic status of the sheep genome, describe current activities in the sheep cytogenomics research sector, and will discuss the cytogenomics data in context with other major sheep genomics projects.

Copyright © 2009 S. Karger AG, Basel

Dr. Tom Goldammer, Forschungsbereich Molekularbiologie Forschungsinstitut für die Biologie landwirtschaftlicher Nutztiere (FBN) Wilhelm-Stahl-Allee 2, DE–18196 Dummerstorf (Germany) Tel. +49 38208 68708, Fax +49 38208 68702 E-Mail tomgoldammer@fbn-dummerstorf.de

KARGER

Fax +41 61 306 12 34 E-Mail karger@karger.ch www.karger.com © 2009 S. Karger AG, Basel 1424–8581/09/1262–0063\$26.00/0

Accessible online at: www.karger.com/cgr

The number of domestic sheep (*Ovis aries*, 2n = 54) in the world is steadily growing and currently amounts to about 1.3 billion animals. Extremely large numbers of sheep can be found in China, Australia, India, Iran, Sudan, and New Zealand [Wint and Robinson, 2007]. Sheep production plays an important economic and agricultural role in developed and developing countries. Several products are obtained from sheep, such as wool, milk, meat, leather, and fertilizer. In less developed and poor countries, sheep often provide the only economic basis for survival of many rural people. Although nearly 60 sheep breeds are known to be extinct, and an additional 120 breeds are at risk of extinction, the existence of more than 1,400 known breeds make sheep an ideal model for studying biodiversity or genome variation [Scherf, 1995, 2000]. The relatively close phylogenetic relationship between sheep and humans means that sheep also serve as biomedical models [e.g. Goetz et al., 2008; Neddermann et al., 2009].

Over the past 10 years, the interests of breeder associations, whose goal is to produce healthy, highly productive sheep in a cost-efficient manner, and of scientists studying ovine genomics have merged, leading to a powerful symbiotic alliance that is focusing on the complete analysis of the sheep genome. In addition, automated cytogenetic and molecular genetic tools have become more affordable, which has tremendously accelerated sheep genome analysis. The International Sheep Genomics Consortium (ISGC, www.sheephapmap.org) is an informal network of sheep genomics researchers that is playing a major role in coordinating many of the sheep genomics activities and ensuring that the results from these initiatives are placed in the public domain.

Cytogenetic Analysis of the Sheep Genome

Many cytogenomics resources have provided information about chromosome structure in sheep, in the context of related species. These resources include the following. A comparative Zoo-FISH map has been created and gives an overview of evolutionarily conserved syntenic chromosome segments and breakpoints between ovine and human chromosomes [Burkin et al., 1997; Iannuzzi et al., 1999]. Comprehensive cytogenetic maps corresponding to the sheep ISCNDB 2000 chromosome standard [Di Berardino et al., 2001] contain about 450 loci [Di Meo et al., 2003, 2007]. High-resolution or targeted-cytogenetic mapping has also linked trait-associated genome regions and candidate genes to chromosome bands for traits such as scrapie susceptibility [Cosseddu et al., 2002; presenilin 1 (PSEN1) and dihydrolipoamide S-succinvltransferase (DLST), in this review], and apoptosis [Lyahyai et al., 2005]. Several studies have used modern cytogenetic techniques such as fiber-FISH or regionspecific chromosome painting probes, generated by chromosome microdissection, for sex chromosome analysis in sheep [e.g. Di Berardino et al., 2004; Chaves et al., 2005; Bugno et al., 2008]. Zoo-FISH using bovine painting probes has also been used to investigate the differences in chromosome abnormalities of in vitro and in vivo generated ovine embryos [Coppola et al., 2007] which in the long term will help with improving the production of in vitro embryos. Ovine chromosome-specific painting probes have been produced and used for comparative studies between sheep, pig, and javelina (Pecari tajacu) [Adega et al., 2006]. Research has also been performed to analyze distinctive features of sheep chromosome morphology, such as translocations or fragile sites [e.g. Chaves et al., 2003; Ali et al., 2008].

Genetic and Physical Analysis of the Sheep Genome

A range of other genomics resources have been developed for sheep and are summarized schematically in figure 1. Sheep genetic maps of increasing resolution have been assembled [http://rubens.its.unimelb.edu.au/ ~jillm/jill.htm; Crawford et al., 1995; de Gortari et al., 1998; Maddox et al., 2001, 2008; Maddox, 2005]. Several studies have been published on the use of polymorphic markers in the discovery of genetic associations and chromosomal locations associated with quantitative trait loci (QTL). A QTL database for the sheep genome, Sheep QTLdb, has been developed in association with the AnimalQTLdb project that summarizes mapped QTL and displays QTL chromosome linkages based on map information of a suitable format given in published manuscripts [http://sphinx.vet.unimelb.edu.au/cgi-bin/QTLdb/ OA/viewmap; Hu et al., 2007]. Numerous new unlisted QTL for milk, growth or health parameters have recently been identified in several sheep breeds, e.g. milk fat content [Calvo et al., 2006], carcass [Johnson et al., 2005], muscle depth [Hadjipavlou et al., 2008], facial eczema [Phua et al., 2009], scrapie incubation time [Moreno et al., 2008], or resistance to internal parasites such as Haemonchus contortus [Marshall et al., 2009].

An ovine bacterial artificial chromosome (BAC) library of 10–12-fold genome coverage, comprising more than 200,000 clones, was developed from a Texel ram



Fig. 1. Sheep genome analyses at different resolutions using different genomic resources.

[Osoegawa et al., 1998] as an ISGC project and is distributed as the CHORI-243 library through BACAPAC Resources (http://bacpac.chori.org/library.php?id=162). End sequencing from both ends of approximately 180,000 BAC clones from the CHORI-243 library allowed generation of comparative genome contigs (BAC-CGCs) covering 91.2% of the human genome. The BAC end sequences were combined with a comparative mapping approach using human, dog and cattle genome data as a framework and used to construct a virtual sheep genome (VSG) [www.livestockgenomics.csiro.au/perl/gbrowse.cgi/ vsheep2/; Dalrymple et al., 2007]. Two publicly available cell lines have also been made from the Texel ram [Chitko-McKown et al., 2008] which will ensure both a ready supply of DNA from this animal and be useful for functional studies.

Two ovine whole-genome radiation hybrid panels have been produced allowing different resolutions for physi-cal mapping of the sheep genome. A 5,000-rad panel, USUoRH5000 has been used to generate a comparative and comprehensive RH map with 3,227 loci that spans the sheep genome by mapping markers that were a mix of monomorphic markers such as BAC end sequences, genes, and polymorphic markers [Cockett, 2006; Wu et al., 2007, 2008a, b, 2009]. A 12,000-rad panel was produced at the Institut National de la Recherche Agrono-mique (INRA) and has been used to generate a high-resolution map in a scrapie-associated chromosome region from *Ovis aries* (OAR) chromosome 18 [Laurent et al., 2007].

More than 200,000 ovine ESTs from different sources have been deposited in public genome databases (http:// compbio.dfci.harvard.edu/tgi/cgi-bin/tgi/T_release. pl?gudb=sheep) and several high-density sheep cDNA and oligo microarrays have been developed [e.g. Keane et al., 2006; Pariset et al., 2008; Sgorlon et al., 2008]. In addition, the close evolutionary relationship between sheep and cattle has allowed several groups to use bovine microarrays for functional studies in sheep with some success [e.g. Tao et al., 2004; Diez-Tascón et al., 2005; Cao et al., 2006].



Fig. 2. Assignment of *PSEN1* and *DLST* to OAR7q33–q34, a chromosome region known to be associated with scrapie susceptibility. The FISH experiment was done on G-banded ovine metaphase chromosomes. DNA sequencing with gene-specific primer sets confirmed gene specificity within ovine BAC probes. Primer sets for PCR and DNA sequencing: *DLST* (5'-GCTCCTGCTAAGGCCAG-3'/5'-CCGA-TGTTCTGAACGCAGAC-3') and *PSEN1* (5'-CATTATTGCTCCTCGCCATT3'/AAGATGAAGAGTCCTGTTTGGT-TT-3'). Arrows indicate the position of hybridization signals.

Recently, 1.5K and 50K SNP chips for sheep genotyping have been created under the auspices of the ISGC [Kijas et al., 2009]. The 50K SNP chip is being used for generating the most comprehensive map of haplotypes (Hap-Map) for livestock species from 2,890 domestic sheep representing 64 breeds (http://www.sheephapmap.org/ hapmap.php).

A first draft of the sheep genome sequence assembly (version 1.0) that corresponds to 2,784.6 Mbp of DNA sequence generated from 6 diverse animals has been produced in another ISGC project and has been made available at http://www.livestockgenomics.csiro.au/perl/ gbrowse.cgi/oar1.0/.

Most of the mapping and sequencing data generated for sheep have been made publically available and can be found at the listed websites. It can be suggested that the sheep genome will soon become a reference genome for the annotation of genomes of phylogenetically related species, such as goats and other *Ovis* species.

Although the international cooperative research within the last few years has made significant progress in the field of sheep genomics, more work remains to be done. Challenges such as limited funding and personnel have impeded the development of a high-quality annotated sheep genome sequence. However, once completed, the reference sequence will serve as a useful tool for the development of modern breeding plans, or to answer phylogenetic questions, or for the identification of causal mutations for specific traits. Another obstacle, which is the focus of this review, is the relatively low number of cytogenetic anchors that can be used to bridge between the different types of maps and chromosome morphology in sheep. We present here the current status of the cytogenetic map and discuss these data in the light of completing a sheep genome sequence assembly.

The 2009 Cytogenetic Map of the Sheep Genome

For sheep, the average number of cytogenetic loci per sheep chromosome is 20, which is similar to the number of loci that have been assigned to specific chromosome bands or segments in other farm animals, such as cattle with an average of 21 loci per chromosome or goat with about 14 loci (http://locus.jouy.inra.fr/). However, the resolution of the cytogenetic sheep map is low compared Downloaded from http://karger.com/cgr/article-pdf/126/1-2/63/2378136/000245907.pdf by UniversitA¤tsbibliothek Bern user on 31 May 2023



Fig. 3. Assignment of *QDPR* to OAR6q36. This FISH experiment was done on RBPI-banded (R-banding by late BrdU incorporation and propidium iodide staining) ovine metaphase chromosomes with simultaneous visualization of FITC signals (arrows). The majority of loci shown in figure 4 has been mapped to the sheep genome using this approach. Iannuzzi and Di Berardino [2008] have described this mapping approach in more detail.

to more completely analyzed genomes, such as the human genome with more than 11,000 cytogenetically assigned loci [http://www.ncbi.nlm.nih.gov/genome/cyto/ hbrc.shtml; The BAC Resource Consortium et al., 2001] or the mouse genome with more than 1,400 mapped loci [http://www.informatics.jax.org/; Bult et al., 2008]. One reason for the low number in sheep is that cytogenetic mapping approaches require appropriate expertise, have lower throughput, and are seen as relatively expensive technologies when compared to sequence and bioinformatics approaches so that cytogenetic research in sheep is limited to a very small number of labs. Recent contributions to the sheep cytogenetic map have mainly focused on mapping trait-associated candidate genes [e.g. Cosseddu et al., 2002; Lyahyai et al., 2005; shown for PSEN1 and DLST in fig. 2], or on the analysis of evolutionary breakpoint regions between sheep and other bovids or human [e.g. Perucatti et al., 2007, 2009; Chessa et al., 2009]. The most current ovine cytogenetic map (table 1) contains 566 loci, which is 112 more than for

earlier cytogenetic maps [e.g. Di Meo et al., 2007]. 375 are type I loci (known genes) and 191 are type II loci (SSRs – simple sequence repeats, microsatellite markers, STSs – sequence tagged sites). All of these loci have been assigned to specific chromosome regions or bands. Most data have been obtained by using simultaneous visualization of FITC-signal hybridization and RBPIbanding (fig. 3). A new cytogenetic map for the whole ovine genome is shown in figure 4 and presents the actual number of cytogenetic loci assigned to specific chromosome regions on the sheep R-band ideogram according to the ISCNDB 2000 standard [Di Berardino et al., 2001].

The cytogenetic data presented in figure 4 have been analyzed in context with other recent and relevant data sets from the sheep genome analysis. From this, we conclude that various genomic maps, such as the SNP, genetic or RH maps, of the sheep genome have reached an appropriate density, and a first build of the sheep genome sequence assembly has been produced, but only a relatively small number of cytogenetic anchors link these maps to specific chromosome bands. This clearly impedes a deeper analysis of morphological and structural chromosome rearrangements.

The 566 loci on the cytogenetic map result in an average distance of 5.1 Mbp per locus based on the first draft of the ovine genome sequence assembly. In order to visualize the correspondence between cytogenetic location and DNA sequence position, we have constructed diagrams for OAR7 and OARX, 2 chromosomes having 2.6 and 2.1 mapped loci in average per chromosome band, respectively. Both sheep chromosomes have assignments of more than 40 DNA sequences and the average distance between cytogenetic loci was calculated as 2.5 and 2.9 Mbp, respectively (fig. 5). In general, the relative positions derived from the FISH mapping data and sequencing data are consistent. However, OAR7 and OARX contain chromosome bands where physical anchors are missing and there are order discrepancies be-

(For figure 4 see next pages.)

Fig. 4. The ovine cytogenetic map as of May 2009. 566 loci assigned to specific chromosome regions and chromosome bands are presented on the sheep R-band standard ideogram of the ISCNDB 2000 [Di Berardino et al., 2001]. Standard characters indicate type I loci (375) and italicized characters indicate type II loci (191). If more than 1 locus has been found for a gene (e.g. *CYP19A1* or *IL2*), the most precise position has been included in the drawing.

Sheep Genome Analysis



Goldammer et al.



Locus symbol	Locus name	Localization OAR	Reference
BMS963	DNA segment D3S73	1p21	Tabet-Aoul et al., 2000
BMS2321	DNA segment D1S91	1q12	
BMS1341	DNA segment D8S42	2p17	
BMS356	DNA segment D2S71	2q44	
BMS460	DNA segment D11S87	3p25	
BMS1172	DNA segment D0S044	4q15	
BMS522	DNA segment D7S47	5q26	
BMS360	DNA segment D6S48	6q22-q24	
BMS1290	DNA segment D9S47	8q22-q23	
BMS1678	DNA segment D14S38	9q15-q16	
BMS1304	DNA segment D14S44	9q23	
BMS1316	DNA segment D12S50	10q26	
ETH3	DNA segment D19S2	11q22	
BMS538	DNA segment D16S45	12q15	
BMS1669	DNA segment D13S40	13q22	
BMS2355	DNA segment D18S35	14q13	
BMS820	DNA segment D15S42	15q29	
MCM150	DNA segment L39829	16q22-q23	
TGLA231	DNA segment D17S19	17q21	
BMS2815	DNA segment D21S49	18q17-q21	
BMS517	DNA segment D22S34	19q13-q21	
BMS1948	DNA segment D29S38	21q24	
BMS882	DNA segment D26S31	22q22-q23	
ILSTS65	DNA segment L37269	23q21	
BMS1232	DNA segment D25S32	24q13-q14	
OARFCB4	DNA segment OARFCB4	25q17	
BMS2104	DNA segment D27S28	26q12-q14	
BMS1008	DNA segment D0S025	Xq45	
SPG21	spastic paraplegia 21 (autosomal recessive, Mast syndrome)	7q13	Cosseddu et al., 2002
RABIIA	CNAD (il l 2	/q15	
SMAD3	SMAD family member 3	/q13 7-24	
BAHDI	bromo adjacent nomology domain containing 1	/q24 7-24 -25	
GATM	giveine amiginotransferase (L-arginine:giveine amiginotransferase)	/q24-q25 7a25_a26	
MIUSC DEL24D1	myosin vC	/q25-q26 7-25	
KSL24DI	ribosomai L24 domain containing 1	7q25 7a24_a25	
ADAMIU MVO1C	ADAM metalopeptidase domain 10	7q24-q25 7a24_a25	
WICIC VDS12C	IIIyosiii IC	7q24-q25 7a24_a25	
VP313C DD\$271	ribosomal protein S27 like	7q24-q25	
NF327L SNV1	sorting pavin 1	7q24-q25	
ZNEA10	zinc finger protein 410	7q24-q23	
ZNF410 EIE2D2	aukarwatic translation initiation factor 2P, subunit 2 bata 20 kDa	7q34	
UIF2D2 VASH1	vasobibin 1	7q34	
SNIW1	SNW domain containing 1	7q34	
TSHD	thyroid stimulating hormone receptor	7q34 7a34	
CALM1	calmodulin 1 (phosphorylase kinase delta)	7q34	
EIE3I	eukarvotic translation initiation factor 3 subunit I	7q34	
NDN	necdin homolog (mouse)	18012	
IIRE3A	ubiquitin protein ligase F3A	18012	
CARRS3	gamma-aminobutyric acid (GABA) A recentor beta 3	18a12_a13	
OCA2	oculocutaneous albinism II	18a12-a13	
A VEN	apontosis, caspase activation inhibitor	18012-015	
SMFK1	SMEK homolog 1 suppressor of mek1 (Dictuactelium)	18023	
DDX24	DFAD (Asp-Glu-Ala-Asp) hox polypertide 24	18023-024	
DICER1	dicer 1 ribonuclease type III	18g23-q27	
DIO1	deiodinase iodothyronine type III	18024	
CINP	cyclin_dependent kingse 2_interacting protein	18024	
UTIVI	cychin-acpendent Kinase 2-interacting protein	10424	

Table 1. A list of 113 loci (83 type I, 30 type II) added to the comprehensive cytogenetic sheep map of Di Meo et al. [2007], arranged by year of publication

Table 1 (continued)

Locus symbol	Locus name	Localization OAR	Reference		
BCL2L2	BCL2-like 2	7q15-q21	Lyahyai et al., 2005		
BCL2L1	BCL2-like 1	13q22			
BCL2	B-cell CLL/lymphoma 2	23q27			
DCTD	dCMP deaminase	26q15	Perucatti et al., 2007		
UFSP2	UFM1-specific peptidase 2	26q15			
CASP3	caspase 3, apoptosis-related cysteine peptidase	26q15			
TLR3	Toll-like receptor 3	26q15			
MSR1	macrophage scavenger receptor 1	26q15			
FAT1	FAT tumor suppressor homolog 1 (Drosophila)	26q15			
LONRF1	LON peptidase N-terminal domain and ring finger 1	26q17			
DLC1	deleted in liver cancer 1	26q17			
C8orf41	chromosome 8 open reading frame 41	26q17			
CSSM036	DNA segment U03827	26q17			
LSM1	LSM1 homolog, U6 small nuclear RNA associated (S. cerevisiae)	26q17			
EIF4EBP1	eukaryotic translation initiation factor 4E binding protein 1	26q17			
ATP9B	ATPase, class II, type 9B	23q12	Tetens et al., 2007		
LOC643448	hypothetical protein LOC643448	23q13			
SYT4	synaptotagmin IV	23q13			
ROCK1	Rho-associated, coiled-coil containing protein kinase 1	23q23-q24			
COLEC12	collectin sub-family member 12	23q23-q24			
SPIRE1	spire homolog 1 (Drosophila)	23q25			
SETBP1	SET-binding protein 1	23q25			
PHLPP	PH domain and leucine-rich repeat protein phosphatase	23q27			
enJSRV-8	endogenous retroviruses	3q21	Chessa et al., 2009		
enJSRV-15	endogenous retroviruses	6q13			
enJSRV-16	endogenous retroviruses	10q24			
enJSRV-18	endogenous retroviruses	11q17			
enJS5F-16	endogenous retroviruses	15q23			
QRFPR	pyroglutamylated RFamide peptide receptor	6q13	Perucatti et al., 2009		
TRAM1L1	translocation-associated membrane protein 1-like 1	6q13dist			
PPP3CA	protein phosphatase 3 (formerly 2B), catalytic subunit, alpha isoform	6q15			
SNCA	synuclein, alpha (non-A4 component of amyloid precursor)	6q17			
PPARGC1A	peroxisome proliferator-activated receptor gamma, coactivator 1α	6q17			
UGDH	UDP-glucose dehydrogenase	6q22prox			
KDR	kinase insert domain receptor (a type III receptor tyrosine kinase)	6q22			
CNO16L	CCR4-NO1 transcription complex, subunit 6-like	6q32prox			
NUP54	nucleoporin 54 kDa	6q32			
DMPI	dentin matrix acidic phosphoprotein 1	6q34dist-q36prox			
	quinoia ainyaropteriaine reductase	6436			
ASMTL	acetylserotonin O-methyltransferase-like	Xp12	Goldammer et al., 2009		
CTPS2	CTP synthase II	Xq11-q12			
NHS	Nance-Horan syndrome (congenital cataracts and dental anomalies)	Xq12			
CNKSR2	connector enhancer of kinase suppressor of Ras 2	Xq12			
DDX53	DEAD (Asp-Glu-Ala-Asp) box polypeptide 53	Xq12			
ILIKAPLI CV auf 50	interleukin 1 receptor accessory protein-like 1	Xq12-q21			
CAOTJ59	chromosome A open reading frame 59	Xq21 X-22			
MED14	Cdc42 guarina nucleotida avehanga factor (CEE) 0	Xq22 Xq22			
IADID1C	lycine (K) specific demethylase 5C	Xq22 Xq23			
PHF8	PHD finger protein 8	Xq23 Xq23			
DCAT216	diacylglycerol () acyltransferase 2 like 6	Xq25 q26			
CXorf26	chromosome X open reading frame 26	Xq25-q26			
LRCH2	leucine-rich repeats calponin homology (CH) domain containing ?	Xa36-a42			
ACSL4	acyl-CoA synthetase long-chain family member 4	Xq30 q=2 Xq43			
NXT2	nuclear transport factor 2-like export factor 2	Xq43			
BMS1820	DNA segment D0S108	Xq43			
	0	1 .			





Table 2. Comp	orehensive	data set	of ovine	genome	analysis
---------------	------------	----------	----------	--------	----------

OAR Chr	Zoo-FISH [Iannuzzi et al., 1999] HSA Chr	Recent Cyto Map loci No.	Average loci number per band [399 band level]	Bands with no cyto loci [399 band level]	Chr length i (OGA v1.0, May 2009) Mbp	Average cyto locus distance in OGA v1.0 Mbp	RH Map (USUo 5,000 panel, May 2009) loci No.	Microsatellite/SNP Map sex averaged (Maddox v5)		SNP50 (pre-	QTL (data base
								loci No.	сM	dicted) SNP No.	plus new QTL) QTL No.
1	1, 3, 21	49	1.2	22	299.8	6.1	297	251	340.0	6,016	2
2	1, 2, 9	42	1.2	13	263.2	6.3	141	240	333.0	5,553	4
3	2, 9, 12, 22	33	1.1	8	242.9	7.4	130	227	313.4	5,071	8
4	4	19	1.1	4	128.1	6.7	64	95	167.7	2,723	0
5	5, 19	19	1.1	11	117.2	6.2	68	91	168.2	2,385	3
6	7	26	1.4	8	129.1	5.0	181	103	154.6	2,624	3
7	14, 15	44	2.6	6	109.0	2.5	73	98	147.9	2,280	0
8	8	15	0.9	7	98.0	6.5	148	61	131.5	2,084	2
9	6	15	1.1	8	100.8	6,7	102	73	138.0	2,168	0
10	13	13	1.2	2	94.2	7.3	78	58	118.3	1,871	2
11	17	22	2.2	3	67.1	3.0	45	81	117.7	1,202	2
12	1	13	1.1	5	86.5	6.7	58	64	120.5	1,738	1
13	10, 20	20	2.2	4	89.1	4.4	57	78	135.2	1,718	2
14	16, 19	20	1.7	7	69.3	3.4	51	77	126.8	1,193	5
15	8	13	1.0	5	90.1	6.9	61	91	130.7	1,716	1
16	5	11	0.8	7	77.2	7.0	54	55	100.2	1,596	0
17	4, 12, 22	14	1.3	3	78.6	5.6	98	80	130.4	1,439	0
18	14, 15	22	2.0	2	72.5	3.3	46	88	130.8	1,434	4
19	3	10	1.4	2	65.0	6.5	42	64	84.7	1,257	0
20	6	18	2.0	5	55.9	3.1	123	63	104.9	1,166	6
21	11	9	1.3	4	55.5	6.2	35	56	78.4	912	0
22	10	10	1.2	4	55.9	5.6	42	33	104.6	1,107	0
23	18	16	1.6	3	66.8	4.2	110	79	93.3	1,140	2
24	8	8	1.0	1	45.3	5.7	49	60	81.6	750	2
25	10	8	0.9	2	48.3	6.0	37	63	84.0	1,016	8
26	7,16	22	2.4	3	50.1	2.3	30	42	81.8	933	0
Х	Х	45	2.1	2	129.1	2.9	130 ^a	147	93.7	1,502	1
Y	Y	10	2.0	0	-	-	0	-	-	1	0
Total Avera	age	566	1.4	151 2	,784.6	5.1	2,350 ^a 2	2,518 3	,811.9	54,595 ^b	58 ^c

OAR Chr = Ovine chromosome number; OGA = ovine genome assembly. QTL database, http://sphinx.vet.unimelb.edu.au/cgi-bin/QTLdb/OA/viewmap.

^a Data in Wu et al., 2008b together with data in Goldammer et al., 2009a.

^b Additional 382 unpositioned SNPs. New QTL (not in QTL database): OAR1 – milk fat content [Calvo et al., 2006]; OAR2 – muscle depth [Hadjipavlou et al., 2008]; carcass [Johnson et al., 2005]; OAR3 – facial eczema [Phua et al., 2009]; OAR6, OAR18 – scrapie incubation time [Moreno et al., 2008].

^c A QTL for host resistance to internal parasite *Haemonchus contortus* [Marshall et al., 2009] is not included.

tween the 2 maps (fig. 5). Given the precision of the ovine FISH mapping process, including a step whereby probes are sequenced to confirm gene content, it is likely that most of the FISH positions are correct (e.g. *HEXA* and *TGM1* loci on OAR7) and that the sequence assembly positions, largely derived from chromosome orders for species other than sheep, are incorrect. The FISH map-

ping method offers a way for positioning more loci to the sheep cytogenetic map and contributes to a sheep sequence assembly process of higher quality. In summary, the generation of many more cytogenetic loci is important to complete the accurate assignment between ovine DNA sequence and metaphase chromosome band morphology.

Sheep Genome Analysis

The relatively low density of cytogenetically mapped loci as compared to other mapped chromosome features is shown in table 2, as well as the gross correspondence between sheep and human chromosomes. Zoo-FISH has been used to identify several rearrangements between ovine and human chromosomes. The description of the regions flanking the breakpoints at the cytogenetic level in sheep is in most cases very imprecise. The same conclusion can also be made when comparing the number of loci with chromosome bands, as on average only 1.42 DNA probes per band have been assigned to date. The ISCNDB 2000 publication [Di Berardino et al., 2001] describes 399 distinct single chromosome bands for the sheep genome. Only 62% of the bands (248 bands) have one or more loci that have been cytogenetically mapped to them (fig. 4). No physical anchors are available for definition or differentiation of the remaining 151 bands. Comparing the number of DNA sequences obtainable from radiation hybrid or genetic maps with the cytogenetic map indicates again that the link between these maps to the metaphase chromosome morphology is only of appropriate density for a small number of genome regions. Consequently, the current cytogenetic map provides several useful physical anchors for sheep genome analysis or comparative studies, but it is inadequate for the completion of the sheep genome assembly or for studies that depend on a high resolution of loci.

Conclusions and Future Aspects

Major efforts based on close collaboration of the laboratories have contributed to the development of the cytogenetic sheep map. However, an increase in the number of mapped loci per chromosome is needed. In addition to the targeted assignment of trait-associated candidate genes or informative markers, which link a QTL region to a chromosome segment, future cytogenetic mapping in sheep should focus on the chromosomal assignment of selected DNA probes which anchor the different types of maps to banded metaphase chromosomes. Targeted comparative mapping is suggested based on chromosome morphology and DNA sequence information between human and sheep and will be necessary to assign genes to ovine chromosome bands that currently have no locus assignments. Likewise, the ovine DNA probes that are used should be selected according to their use in comparative mapping approaches, such as cytogenetically mapped loci in other domestic Bovi*dae* such as cattle, goat or river buffalo. Building these bridges between the different maps and chromosome morphology has already been reported for OAR23 [Tetens et al., 2007], OAR10 [Drögemüller et al., 2008], OAR6 [Perucatti et al., 2009] and OARX [Goldammer et al., 2009a, b]. Other map comparisons are being performed for other sheep chromosomes to demonstrate the evolutionary rearrangements in other Bovidae and human.

Procedures for the identification of chromosomal polymorphisms such as whole-comparative genomic hybridization (W-CGH) or DNA breakage detection FISH (DBD-FISH) have also proven useful in phylogenetic studies between domestic sheep and the mouflon, a direct ancestor [Dávila-Rodríguez et al., 2009]. Expanding such studies to other Ovis species should improve information about the genetic diversity and evolution of the genus Ovis and complement other phylogenetics studies such as the recent study that used retroviral insertion to delineate different sheep lineages and mitochondrial lineages studies [e.g. Tapio et al., 2006; Meadows et al., 2007; Chessa et al., 2009]. For Hominidae, studies have shown that the process of chromosome evolution is the result of numerous small segment duplications and deletions [Locke et al., 2003; Fortna et al., 2004], and this is likely to be the case also for Caprinae and other ruminants. The human studies used array-based methods, with the first being an array-comparative genomic hybridization (A-CGH) approach that used a human BAC array and showed about 60 small segment mutations. The second approach was a higher-resolution approach that looked at differences in gene copy number in Hominidae by using a cDNA-array that included almost the complete gene coding genome. Generation of analogous genome-wide DNA arrays based on the well-characterized ovine CHO-RI-243 BAC library or cDNA libraries from sheep, would be powerful tools for analysis of chromosome evolution in sheep. Moreover, such tools would also be useful for comparative analysis in other closely related Ruminantia and for clinical cytogenetics. In addition to their use for investigating the evolution of sheep, such information would contribute to the completion of the sheep genome DNA sequence and could likewise be linked with information about quantitative traits. In this way, ovine cytogenomic data would be a valuable contribution to the study of sheep phenotypes that are optimally adapted to a specific environment, as well as modern breeding and production conditions.

The authors thank Domenico Incarnato, CNR-ISPAAM of Naples for construction of the sheep cytogenetic map on an Rbanded standard ideogram, as well as the listed organizations for funding sheep genome analysis: German Research Foundation – DFG-Grant GO-896/6-3; USDA National Research Initiative Grants no. 2005-591-0649-01; National Research Council of Rome, Italy, DG.RSTL.083.001 and DG.RSTL.083.002; LR5 of Campania Region, Italy; co-authors funding bodies e.g. sheep-GENOMICS (a joint initiative of Meat and Livestock Australia (MLA) and Australian Wool Innovation (AWI).

References

- Adega F, Chaves R, Kofler A, Krausman PR, Masabanda J, et al: High-resolution comparative chromosome painting in the Arizona collared peccary (*Pecari tajacu*, Tayassuidae): a comparison with the karyotype of pig and sheep. Chromosome Res 14:243–251 (2006).
- Ali A, Abdullah M, Babar ME, Javed K, Nadeem A: Expression and identification of folatesensitive fragile sites in British Suffolk sheep (*Ovis aries*). J Genet 87:219–227 (2008).
- Bugno M, Zyga A, Słota E: Microdissected bovine X chromosome segment delineates homologous chromosomal regions in sheep and goat. Folia Biol (Krakow) 56:149–151 (2008).
- Bult CJ, Eppig JT, Kadin JA, Richardson JE, Blake JA; the Mouse Genome Database Group: The Mouse Genome Database (MGD): mouse biology and model systems. Nucleic Acids Res 36:D724–D728 (2008).
- Burkin DJ, Yang F, Broad TE, Wienberg J, Hill DF, Ferguson-Smith MA: Use of the Indian muntjac idiogram to align conserved chromosomal segments in sheep and human genomes by chromosome painting. Genomics 46:143–147 (1997).
- Calvo JH, Martínez-Royo A, Beattie AE, Dodds KG, Marcos-Carcavilla A, Serrano M: Fine mapping of genes on sheep chromosome 1 and their association with milk traits. Anim Genet 37:205–210 (2006).
- Cao H, Kabaroff LC, You Q, Rodriguez A, Boermans HJ, Karrow NA: Characterization of ovine hepatic gene expression profiles in response to *Escherichia coli* lipopolysaccharide using a bovine cDNA microarray. BMC Vet Res 2:34 (2006).
- Chaves R, Adega F, Wienberg J, Guedes-Pinto H, Heslop-Harrison JS: Molecular cytogenetic analysis and centromeric satellite organization of a novel 8;11 translocation in sheep: a possible intermediate in biarmed chromosome evolution. Mamm Genome 14:706–710 (2003).
- Chaves R, Guedes-Pinto H, Heslop-Harrison JS: Phylogenetic relationships and the primitive X chromosome inferred from chromosomal and satellite DNA analysis in Bovidae. Proc Biol Sci 272:2009–2016 (2005).
- Chessa B, Pereira F, Arnaud F, Amorim A, Goyache F, et al: Revealing the history of sheep domestication using retrovirus integrations. Science 324:532–536 (2009).

- Chitko-McKown CG, Leonard JC, Moscatello D, Miller LC, Freking BA: Development of cell lines from the sheep used to construct the CHORI-243 Ovine BAC library. Anim Biotechnol 19:84–88 (2008).
- Cockett NE: The sheep genome. Genome Dyn 2: 79–85 (2006).
- Coppola G, Alexander B, Di Berardino D, St John E, Basrur PK, King WA: Use of crossspecies in-situ hybridization (ZOO-FISH) to assess chromosome abnormalities in day-6 in-vivo- or in-vitro-produced sheep embryos. Chromosome Res 15:399–408 (2007).
- Cosseddu GM, Oustry-Vaiman A, Jego B, Moreno C, Taourit S, et al: Sheep/human comparative map in a chromosome region involved in scrapie incubation time shows multiple breakpoints between human chromosomes 14 and 15 and sheep chromosomes 7 and 18. Chromosome Res 10:369–378 (2002).
- Crawford AM, Dodds KG, Pierson CA, Ede AJ, Montgomery GW, et al: An autosomal genetic linkage map of the sheep genome. Genetics 140:703–724 (1995).
- Dalrymple BP, Kirkness EF, Nefedov M, McWilliam S, Ratnakumar A, et al: International Sheep Genomics Consortium: Using comparative genomics to reorder the human genome sequence into a virtual sheep genome. Genome Biol 8:R152 (2007).
- Dávila-Rodríguez MI, Cortés-Gutiérrez EI, López-Fernández C, Pita M, Mezzanotte R, Gosálvez J: Whole-comparative genomic hybridization in domestic sheep (*Ovis aries*) breeds. Cytogenet Genome Res 124:19–26 (2009).
- de Gortari MJ, Freking BA, Cuthbertson RP, Kappes SM, Keele JW, et al: A second-generation linkage map of the sheep genome. Mamm Genome 9:204–209 (1998).
- Di Berardino D, Di Meo GP, Gallagher DS, Hayes H, Iannuzzi L (co-ordinator) (eds): ISCNDB 2000: International System for Chromosome Nomenclature of Domestic Bovids. Cytogenet Cell Genet 92:283–299 (2001).
- Di Berardino D, Vozdova M, Kubickova S, Cernohorska H, Coppola G, et al: Sexing river buffalo (*Bubalus bubalis* L.), sheep (*Ovis aries* L.), goat (*Capra hircus* L.), and cattle spermatozoa by double color FISH using bovine (*Bos taurus* L.) X- and Y-painting probes. Mol Reprod Dev 67:108–115 (2004).

- Diez-Tascón C, Keane OM, Wilson T, Zadissa A, Hyndman DL, et al: Microarray analysis of selection lines from outbred populations to identify genes involved with nematode parasite resistance in sheep. Physiol Genomics 21: 59–69 (2005).
- Di Meo GP, Perucatti A, Gautier M, Hayes H, Incarnato D, et al: Chromosome localization of the 31 type I Texas bovine markers in sheep and goat chromosomes by comparative FISH-mapping and R-banding. Anim Genet 34:294–296 (2003).
- Di Meo GP, Perucatti A, Floriot S, Hayes H, Schibler L, et al: An advanced sheep (*Ovis aries*, 2n = 54) cytogenetic map and assignment of 88 new autosomal loci by fluorescence in situ hybridization and R-banding. Anim Genet 38:233–240 (2007).
- Drögemüller M, Tetens J, Dalrymple B, Goldammer T, Wu CH, et al: A comparative radiation hybrid map of sheep chromosome 10. Cytogenet Genome Res 121:35–40 (2008).
- Fortna A, Kim Y, MacLaren E, Marshall K, Hahn G, et al: Lineage-specific gene duplication and loss in human and great ape evolution. PLoS Biol 2:E207 (2004).
- Goetz WA, Tan TE, Lim KH, Salgues Sle H, Grousson N, et al: Truly stentless molded autologous pericardial aortic valve prosthesis with single point attached commissures in a sheep model. Eur J Cardiothorac Surg 33: 548–553 (2008).
- Goldammer T, Brunner RM, Rebl A, Wu CH, Nomura K, et al: A high-resolution radiation hybrid map of sheep chromosome X and comparison with human and cattle. Cytogenet Genome Res 125:40–45 (2009a).
- Goldammer T, Brunner RM, Rebl A, Wu CH, Nomura K, et al: Cytogenetic anchoring of radiation hybrid and virtual maps of sheep chromosome X and comparison of X chromosomes in sheep, cattle, and human. Chromosome Res 17:497–506 (2009b).
- Hadjipavlou G, Matika O, Clop A, Bishop SC: Two single nucleotide polymorphisms in the myostatin (*GDF8*) gene have significant association with muscle depth of commercial Charollais sheep. Anim Genet 39:346–352 (2008).
- Hu ZL, Fritz ER, Reecy JM: AnimalQTLdb: a livestock QTL database tool set for positional QTL information mining and beyond. Nucleic Acids Res 35(Database issue):D604– D609 (2007).

- Iannuzzi L, Di Berardino D: Tool of the trade: diagnostic and research in domestic animal cytogenetics. J Appl Genet 49:357–356 (2008).
- Iannuzzi L, Di Meo GP, Perucatti A, Incarnato D: Comparison of the human with the sheep genomes by use of human chromosome-specific painting probes. Mamm Genome 10: 719–723 (1999).
- Johnson PL, McEwan JC, Dodds KG, Purchas RW, Blair HT: A directed search in the region of *GDF8* for quantitative trait loci affecting carcass traits in Texel sheep. J Anim Sci 83: 1988–2000 (2005).
- Keane OM, Zadissa A, Wilson T, Hyndman DL, Greer GJ, et al: Gene expression profiling of naïve sheep genetically resistant and susceptible to gastrointestinal nematodes. BMC Genomics 7:42 (2006).
- Kijas JW, Townley D, Dalrymple BP, Heaton MP, Maddox JF, et al; International Sheep Genomics Consortium: A genome wide survey of SNP variation reveals the genetic structure of sheep breeds. PLoS ONE 4:e4668 (2009).
- Laurent P, Schibler L, Vaiman A, Laubier J, Delcros C, et al: A 12 000-rad whole-genome radiation hybrid panel in sheep: application to the study of the ovine chromosome 18 region containing a QTL for scrapie susceptibility. Anim Genet 38:358–363 (2007).
- Locke DP, Segraves R, Carbone L, Archidiacono N, Albertson DG, et al: Large-scale variation among human and great ape genomes determined by array comparative genomic hybridization. Genome Res 13:347–357 (2003).
- Lyahyai J, Goldammer T, Beattie AE, Zaragoza P, Martín-Burriel I: Positional and functional characterisation of apoptosis related genes belonging to the BCL2 family in sheep. Cytogenet Genome Res 109:519–526 (2005).
- Maddox JF: A presentation of the differences between the sheep and goat genetic maps. Genet Sel Evol 37 Suppl 1:S1–S10 (2005).
- Maddox JF, Davies KP, Crawford AM, Hulme DJ, Vaiman D, et al: An enhanced linkage map of the sheep genome comprising more than 1000 loci. Genome Res 11:1275–1289 (2001).
- Maddox JF, Crawford AM, Ingersoll R, Kijas J, Dalrymple BP, et al, and the International Sheep Genomics Consortium: Using a 1.5K ovine SNP array to expand the sheep linkage map, p 2160. Proc 31st ISAG conference, Amsterdam, The Netherlands (2008).
- Marshall K, Maddox JF, Lee SH, Zhang Y, Kahn L, et al: Genetic mapping of quantitative trait loci for resistance to *Haemonchus contortus* in sheep. Anim Genet 40:262–272 (2009).
- Meadows JR, Cemal I, Karaca O, Gootwine E, Kijas JW: Five ovine mitochondrial lineages identified from sheep breeds of the near East. Genetics 175:1371–1379 (2007).

- Moreno CR, Cosseddu GM, Schibler L, Roig A, Moazami-Goudarzi K, et al: Identification of new quantitative trait loci (other than the *PRNP* gene) modulating the scrapie incubation period in sheep. Genetics 179:723–726 (2008).
- Neddermann A, Willbold E, Witte F, Hurschler C, Hankemeier S, et al: Tunnel widening after anterior cruciate ligament reconstruction: an experimental study in sheep. Am J Sports Med 37:1609–1617 (2009).
- Osoegawa K, Woon PY, Zhao B, Frengen E, Tateno M, et al: An improved approach for construction of bacterial artificial chromosome libraries. Genomics 52:1–8 (1998).
- Pariset L, Chillemi G, Bueno S, Prosperini G, Bongiorni S, Valentini A: From public sequences to custom microarrays: an easy and affordable tool, p 1007, Proc 31st ISAG conference, Amsterdam, The Netherlands (2008).
- Perucatti A, Di Meo GP, Goldammer T, Incarnato D, Brunner R, Iannuzzi L: Comparative FISH-mapping of twelve loci in river buffalo and sheep chromosomes: comparison with HSA8p and HSA4q. Cytogenet Genome Res 119:242–244 (2007).
- Perucatti A, Di Meo GP, Goldammer T, Incarnato D, Nicolae I, et al: FISH-mapping comparison between river buffalo chromosome 7 and sheep chromosome 6: assignment of new loci and comparison with HSA4. Cytogenet Genome Res 124:106–111 (2009).
- Phua SH, Dodds KG, Morris CA, Henry HM, Beattie AE, et al: A genome-screen experiment to detect quantitative trait loci affecting resistance to facial eczema disease in sheep. Anim Genet 40:73–79 (2009).
- Scherf DB (ed): World Watch List for Domestic Animal Diversity, ed. 2. (Food and Agriculture Organization of the United Nations, Rome, Italy 1995).
- Scherf DB (ed): World Watch List for Domestic Animal Diversity, ed. 3. (Food and Agriculture Organization of the United Nations, Rome, Italy 2000).
- Sgorlon S, Ferrarini A, Asquini E, Dreos R, Bruno S: Nutrigenomics studies in sheep using a custom microarray platform, p 2208, Proc 31st ISAG conference, Amsterdam, The Netherlands (2008).
- Tabet-Aoul K, Oustry-Vaiman A, Vaiman D, Saïdi-Mehtar, Cribiu ED, Lantier F: Cytogenetical anchoring of sheep linkage map and syntenic groups using a sheep BAC library. Genet Sel Evol 32:441–457 (2000).
- Tao W, Mallard B, Karrow N, Bridle B: Construction and application of a bovine immune-endocrine cDNA microarray. Vet Immunol Immunopathol 101:1–17 (2004).
- Tapio M, Marzanov N, Ozerov M, Cinkulov M, Gonzarenko G, et al: Sheep mitochondrial DNA variation in European, Caucasian, and Central Asian areas. Mol Biol Evol 23:1776– 1783 (2006).

- Tetens J, Goldammer T, Maddox JF, Cockett NE, Leeb T, Drögemüller C: A radiation hybrid map of sheep chromosome 23 based on ovine BAC-end sequences. Anim Genet 38:132– 140 (2007).
- The BAC Resource Consortium, Cheung VG, Nowak N, Jang W, Kirsch IR, et al: Integration of cytogenetic landmarks into the draft sequence of the human genome. Nature 409: 953–958 (2001).
- Wint GRW, Robinson TP (eds): Gridded Live Stock of the World 2007, Appendix B: 99–107, p 131 (Food and Agriculture Organization of the United Nations, Rome, Italy 2007).
- Wu CH, Nomura K, Goldammer T, Hadfield T, Womack JE, Cockett NE: An ovine wholegenome radiation hybrid panel used to construct an RH map of ovine chromosome 9. Anim Genet 38:534–536 (2007).
- Wu CH, Nomura K, Goldammer T, Hadfield T, Dalrymple BP, et al: A high-resolution comparative radiation hybrid map of ovine chromosomal regions that are homologous to human chromosome 6 (HSA6). Anim Genet 39: 459–467 (2008a).
- Wu CH, Nomura K, Goldammer T, Tetens J, Hadfield T, et al: A first-generation ovine whole genome radiation hybrid map using the USUoRH5000 panel, p 2228, Proc 31st ISAG conference, Amsterdam, The Netherlands (2008b).
- Wu CH, Jin W, Hadfield TS, Moore SS, Dalrymple BP, et al: Expanding the ovine radiation hybrid map with loci from the ovine 1.5K and bovine 3.0K SNP chips, p 497, Proc 27th Plant & Animal Genomes Conference (PAG), San Diego, CA, USA (2009).

Referenced web pages:

- BACAPAC Resources CHORI-243 library: http://bacpac.chori.org/library.php?id=162
- Human BAC Resources: www.ncbi.nlm.nih.gov/ genome/cyto/hbrc.shtml
- International Sheep Genomics Constortium (ISGC): www.sheephapmap.org
- Mouse genome database: www.informatics.jax. org/
- NCBI genome database: www.ncbi.nlm.nih.gov/ Ovine 50K SNP HapMap: www.sheephapmap. org/hapmap.php
- Ovine Cytogenetic Map: http://locus.jouy.inra.fr/
- Ovine genome assembly version 1.0: www.livestockgenomics.csiro.au/perl/gbrowse.cgi/ oar1.0/
- Ovine Linkage Map v4.2: http://rubens.its.unimelb.edu.au/~jillm/jill.htm
- Sheep QTL map: http://sphinx.vet.unimelb.edu. au/cgi-bin/QTLdb/OA/viewmap
- The Gene Index database: http://compbio.dfci. harvard.edu/tgi/cgi-bin/tgi/T_release.pl? gudb=sheep
- Virtual sheep genome (VSG): www.livestockgenomics.csiro.au/perl/gbrowse.cgi/vsheep2/