

SLR-derived terrestrial reference frame using observations to LAGEOS-1 / 2, Starlette, Stella, and AJISAI

Krzysztof Sośnica (1), Adrian Jäggi (1),
Daniela Thaller (2), Gerhard Beutler (1),
Rolf Dach (1), Christian Baumann (1)

(1) *Astronomical Institute, University of Bern, Switzerland*

(2) *Bundesamt für Kartographie und Geodäsie, Frankfurt am Main, Germany*

18th International Workshop on Laser Ranging

11–15 November 2013 Fujiyoshida, Japan


Motivation





Etalon-1/2

Current ILRS products:

- LAGEOS-1/2 & Etalon-1/2 solutions only,
- On average ~3000 normal points to LAGEOS-1/2 and ~300 normal points to Etalon-1/2 per week,
- The impact of Etalon-1/2 on the solution is virtually negligible



Motivation



	LAGEOS-1	LAGEOS-2	AJISAI	Starlette	Stella
No of normal points per week	1500	1500	3000	1600	800
	LAGEOS-1/2	LEO	Combined		
No of stations per week	19.8	21.1	22.4		

SLR solutions in the Bernese GNSS Software

Parameter	LAGEOS	LEO
Station Coordinates	Weekly	Weekly
Earth Rotation Parameters	PWL daily	PWL daily
Geocenter Coordinates	Weekly	Weekly
Gravity field	Up to d/o 4	Up to d/o 4
Range Biases	Selected stations	All stations
Orbit	Osculating Elements	Weekly
	Constant along-track S0	Weekly
	Air Drag Scaling Factor	-
	Once-per-rev SS, SC	Weekly
	Once-per-rev WS, WC	Weekly (when not estimating gravity field)
	Pseudo-Stochastic Pulses	-
		Once-per-rev in along-track

Bernese GNSS Software, v.5.3

10 years of processed data (2002-2012)

Scale



TRF scale estimated from the Helmert 7-parameter transformation of weekly SLR solutions

Orbit modeling deficiencies related to non-gravitational forces appear as the periods of the draconitic year.



Draconitic year is a time interval between two consecutive passes of the Sun through the orbital plane of a satellite (in the same direction).

Draconitic years of geodetic satellites:

- 222 days: LAGEOS-2,
- 560 days: LAGEOS-1,
- 89 days: AJISAI,
- 73 days: Starlette,
- 182 days: Stella.



Geocenter coordinates



The origin of the reference frame (geocenter coordinates) are best defined by the SLR technique. The X and Y components can be also defined by other techniques, e.g., DORIS, GNSS, but the Z component is strongly affected by the deficiencies in the solar radiation pressure modeling, and thus, can only be established by the SLR solutions*.


The LAGEOS-1/2 solutions or the Star+Ste+Aji solutions show very small orbit modeling deficiencies (draconitic year of LAGEOS-2 and Ajisai).

All amplitudes related to draconitic years are substantially reduced in the combined solutions.

* see Meindl et al. (2013) Geocenter coordinates estimated from GNSS data as viewed by perturbation theory. Advances in Space Research, Volume 51, Issue 7, p. 1047–1064


Weekly repeatabilities of station coordinates

Station coordinate repeatability in LAGEOS-1/2 and the **combined** solutions



The station repeatability is improved in the combined solutions for East and North components of non-core SLR station

Correlations




Correlation coefficients between selected parameters.

Estimated parameters (ERPs, geopotential, station coordinates) can be substantially decorrelated when using many SLR satellites, due to:

- better observation geometry,
- larger number of SLR observations,
- different orbital characteristics (altitudes, inclination angles, eccentricities).



Earth Rotation Parameters (w.r.t. IERS-08-C04)



Solution type		X pole [μas]		Y pole [μas]		LoD [μs]		Repeatability [mm]		
		bias	WRMS	bias	WRMS	bias	WRMS	Up	North	East
LAGEOS-1/2	gravity up to 4/4	4.1	160.0	-8.0	155.2	6.1	57.0	11.1	10.2	12.3
LAGEOS-1/2	no gravity	45.8	168.5	-54.1	153.5	77.3	120.5	10.9	10.0	12.4
SLR-LEO	gravity up to 4/4	38.3	267.9	-7.8	217.6	-38.5	105.6	15.3	15.4	15.2
SLR-LEO	no gravity	190.1	437.5	-61.1	315.9	189.6	359.3	15.8	15.6	16.8
multi-SLR	gravity up to 4/4	6.4	148.9	8.5	140.3	6.3	56.3	11.3	11.2	11.7
multi-SLR	no gravity	83.7	153.1	63.3	156.7	75.8	121.7	11.1	11.3	11.8

Simultaneous estimation of all parameters
(gravity field+station coord+ERPs+orbits)
is beneficial for SLR solutions
(in particular for combined L1/L2/Sta/Ste/Aji solutions).



Gravity field parameters



C_{20} can be well-established from LAGEOS-1/2 solutions, but the amplitude of annual signal is by 20% larger than in the multi-SLR solutions.

C_{30} from LAGEOS-1/2 shows a clear alias period with draconitic year of LA-2.

Gravity field parameters



Gravity field parameters (besides C_{20}) can be much better established from the multi-SLR solutions (in particular for degree higher than 2). LAGEOS-1/2 solution reveals variations related to the draconitic years or their harmonics.

Summary

Advantages of the multi-SLR solutions :

- The estimated **parameters** can be substantially **decorrelated**.
- The observation **geometry** is **improved**.
- **Orbit modeling** deficiencies of **non-gravitational** forces (solar radiation pressure, albedo, Earth's infrared radiation, the Yarkovsky effect, the Yarkovsky–Schach effect, light aberration, etc.) can be **reduced**.
- As a result, the terrestrial reference frame parameters (**scale**, **geocenter coordinates**, **ERPs**) can be **better** established.
- The repeatability of some SLR station **coordinates** can be **improved**.


Disadvantages of the multi-SLR solutions:

- The repeatability of some of best-performing SLR station coordinates is slightly worse w.r.t. the LAGEOS-1/2 solutions,
- High-quality solutions of low orbiting satellites require more user's attention (screening process and the orbit modeling optimization).




Etalon-1/2

Thank you for your attention





Single-satellite solutions

Stella leads to worse
Coordinates, but LoD is
improved



	Aposteriori sigma of unit weight [mm]	Mean number of observation per week	X pole bias [μas]	X pole WRMS [μas]	Y pole bias [μas]	Y pole WRMS [μas]	LoD bias [μs]	LoD WRMS [μs]
AJISAI	6.31	3011	36.4	266.3	3.6	233.9	-17.3	108.5
Starlette	6.45	1697	21.8	339.5	-6.5	290.5	-18.0	133.0
Stella	6.03	813	120.0	901.6	-11.8	829.0	9.6	110.7
AJISAI+Starlette	6.85	4708	32.0	207.3	-3.0	184.4	-35.2	136.9
AJISAI+Stella	7.24	3824	71.8	304.4	-3.8	256.6	-1.0	93.1
Starlette+Stella	7.62	2510	75.2	365.2	-19.1	291.5	-3.7	99.4
All satellites	7.78	5521	57.7	269.8	-8.7	218.1	-3.6	106.5

Geocenter coordinates



The Y geocenter component from the multi-SLR solution shows large annual variations and a secular drift, whose origin remains unclear. Nevertheless, the combined solutions are slightly less noisy than the LAGEOS-1/2 solutions.

References

References:

- Beutler G (2005) Methods of Celestial Mechanics. Volume II: Application to Planetary System, Geodynamics and Satellite Geodesy. Springer Verlag. ISBN 978-3-540-26512-2.
- Dach R, Hugentobler U, Fridez P, Meindl M (2007) Bernese GPS Software Version 5.0. Astronomical Institute, University of Bern, Switzerland.
- Jäggi A, Sośnica K, Thaller D, Beutler G (2012) Validation and estimation of low-degree gravity field coefficients using LAGEOS. Proceedings of 17th ILRS Workshop, Bundesamt für Kartographie und Geodäsie, 48, Frankfurt, 2012.
- Michael M, Beutler G, Thaller D, Dach R, Jäggi A (2013) Geocenter coordinates estimated from GNSS data as viewed by perturbation theory. *Advances in Space Research*, Volume 51, Issue 7, p. 1047–1064.
- Rodriguez-Solano CJ, Hugentobler U, Steigenberger P, Lutz S (2012) Impact of Earth radiation pressure on GPS position estimates. *J Geod* 86(5):309–317. doi: 10.1007/s00190-011-0517-4.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Impact of atmospheric pressure loading on SLR-derived products and on the consistency between GNSS and SLR results. *Journal of Geodesy* 87 (8), 751–769. DOI: 10.1007/s00190-013-0644-1.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Contribution of Starlette, Stella, and AJISAI to the SLR-derived global reference frame. *Journal of Geodesy* (submitted manuscript).
- Sośnica K, Thaller D, Jäggi A, Dach R, Beutler G (2012) Sensitivity of Lageos Orbits to Global Gravity Field Models. *Art Sat*, 47(2), pp. 35–79. doi:10.2478/v10018-012-0013-y.
- Sośnica K, Thaller D, Jäggi A, Dach R, Beutler G (2012) Can we improve LAGEOS solutions by combining with LEO satellites? *Proceedings of the International Technical Laser Workshop 2012 (ITLW-12)*, Frascati (Rome), Italy, November 5–9, 2012.
- Sośnica K, Thaller D, Dach R, Jäggi A, Beutler G (2013) Time variable Earth's gravity field from SLR data. *Advances in Space Research* (to be submitted).
- Sośnica K, Jäggi A, Thaller D, Meyer U, Beutler G, Dach R (2013) Earth gravity field recovery using GPS, GLONASS, and SLR satellites. *18th International Workshop on Laser Ranging*, 11–15 November 2013 Fujiyoshida, Japan.
- Thaller D, Sośnica K, Dach R, Jäggi A, Beutler G (2011) LAGEOS-ETALON solutions using the Bernese Software. *Mitteilungen des Bundesamtes fuer Kartographie und Geodäsie*, Proceedings of the 17th International Workshop on Laser Ranging, Extending the Range, Bad Kötzting, Germany, May 16–20, 2011, vol. 48, pp.333–336, Frankfurt.
- Thaller D, Sośnica K, Mareyen M , Dach R, Jäggi A, Beutler G (2013) Geodetic parameters estimated from LAGEOS and Etalon data and comparison to GNSS-estimates. *Journal of Geodesy* (submitted manuscript).
- Thaller D, Sośnica K, Dach R, Jäggi A, Beutler G, Mareyen M, Richter B (2013) Geocenter coordinates from GNSS and combined GNSS-SLR solutions using satellite co-locations. *IAG Symposia Series* 139.