

# Patterns of millennial variability over the last 500 ka

M. Siddall<sup>1</sup>, E. J. Rohling<sup>2</sup>, T. Blunier<sup>3</sup>, and R. Spahni<sup>4</sup>

<sup>1</sup>Department of Earth Science, University of Bristol, Bristol, UK

<sup>2</sup>National Oceanography Centre, Southampton, Southampton, UK

<sup>3</sup>Centre for Ice & Climate, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

<sup>4</sup>Climate and Environmental Physics and Oeschger Centre for Climate Change Research, University of Bern,

Bern, Switzerland

Received: 14 January 2010 – Published in Clim. Past Discuss.: 29 January 2010 Revised: 13 April 2010 – Accepted: 22 April 2010 – Published: 5 May 2010

Abstract. Millennial variability is a robust feature of many paleoclimate records, at least throughout the last several glacial cycles. Here we use the mean signal from Antarctic climate events 1 to 4 to probe the EPICA Dome C temperature proxy reconstruction through the last 500 ka for similar millennial-scale events. We find that clusters of millennial events occurred in a regular fashion over half of the time during this with a mean recurrence interval of 21 kyr. We find that there is no consistent link between ice-rafted debris deposition and millennial variability. Instead we speculate that changes in the zonality of atmospheric circulation over the North Atlantic form a viable alternative to freshwater release from icebergs as a trigger for millennial variability. We suggest that millennial changes in the zonality of atmospheric circulation over the North Atlantic are linked to precession via sea-ice feedbacks and that this relationship is modified by the presence of the large, Northern Hemisphere ice sheets during glacial periods.

# 1 Introduction

(Background) Ice-core proxy records of high latitude Northern Hemisphere temperature reveal a distinctive pattern of repeated rapid warming events of 8–15 °C during Marine Isotope Stage 3 (MIS 3), known as Dansgaard-Oeschger (D-O) events (Dansgaard et al., 1984, 1993; Oeschger et al., 1984). These rapid warmings are interspersed with cold periods such that MIS 3 is a period of substantial millennialscale climate variability. This variability is found throughout much of the Northern Hemisphere in marine sediments and also continental records (Shackleton et al., 2000; Wang et al., 2001; Voelker et al., 2002; Rohling et al., 2003; Denton



Correspondence to: M. Siddall mark.siddall@bristol.ac.uk

et al., 2005). D-O events in MIS 3 appear in groups with a longer warm period followed by several shorter warm periods, interspersed with cold periods (Bond and Lotti, 1995). These D-O groups, sometimes called Bond cycles, end in a cold culmination, expressed as a so-called Heinrich event of massive ice-rafted debris (IRD) deposition in the North Atlantic between about 40 and 50° N (see overview in Hemming, 2004). Such indications of substantial iceberg release have motivated modelling studies which consider freshwater pulses into the North Atlantic as a trigger for rapid cooling during D-O variability (Stocker and Wright, 1992).

Blunier et al. (1998) and Blunier and Brook (2001) used variations in the concentration of atmospheric methane (a globally well-mixed gas) in air bubbles enclosed within the ice, to synchronise ice-core records from Antarctica and Greenland. This work showed that D-O events correspond to slower, smaller changes in Antarctica. Longer D-O stadials in Greenland correspond to the largest variations in Antarctic temperature (Antarctic events A1 to A4) (Stocker and Johnsen, 2003). High-resolution temperature reconstructions from Antarctic ice cores have confirmed that shorter D-O events are also associated with smaller Antarctic warming events and that the magnitude of the Antarctic warming is proportional to the duration of D-O stadial events (EPICA Community members, 2006). This paper is concerned with the larger Antarctic climate events, corresponding to the longer D-O stadials during MIS 3.

The detailed phase response between Antarctic and Greenland records during MIS 3 has been described by Stocker and Johnsen (2003) – in response to rapid warmings in Greenland, Antarctica starts to gradually cool and in response to rapid coolings in Greenland, Antarctica starts to gradually warm. The physical mechanism responsible for this "bipolar seesaw" is linked to an active meridional overturning circulation (MOC) in the Atlantic, which transports heat northward at all latitudes, heating the north and cooling the south (Crowley, 1992). If the MOC shuts down, heat is no longer drawn to the north from the south along the axis of the Atlantic, causing rapid cooling in Greenland and gradual warming in Antarctica. In this way it is suggested that cooling during D-O stadials, plausibly caused by iceberg release also affects Antarctic temperature (Hemming, 2004). Here we consider the link between IRD deposition in the north Atlantic and the larger Antarctic climate events over the last 500 kyr.

Although millennial-scale variability has been predominantly found in high-resolution records of MIS 3, some evidence for it exists also in earlier intervals (Oppo et al., 1998; McManus et al., 1999). Larrasoaña et al. (2003) found that the occurrence of sub-Milankovitch (or sub-orbital) variability became significant from as early as ~0.95 Ma. A study of millennial events recorded in the Vostok ice core has suggested that broadly similar north-south phase relationships to those of MIS 3, as described in Stocker and Johnsen (2003), existed over the last four climatic cycles (Delmotte et al., 2004). Siddall et al. (2006, 2007) described very similar millennial-scale relationships between temperature reconstructions in the Vostok ice core, methane concentrations and a simple model of the bipolar seesaw during MIS 8, compared to MIS 3. Cave speleothem records have revealed millennial-scale variability within the last several glacial cycles, which is similar in appearance to that of MIS 3 (Bar-Matthews et al., 2003; Yuan et al., 2004).

Several authors have suggested that millennial-scale climate variability is a function of ice volume, relating periods of more intense variability to periods with increases ice volume (Oppo et al., 1998; McManus et al., 1999; Siddall et al., 2006, 2007). Other work has found millennial variability during periods with reduced ice volume (Martrat et al., 2007; Desprat et al., 2009). Most recently Ganopolski et al. (2009) have linked episodes of millennial variability to the complex phase relationships between Antarctic temperature and atmospheric  $CO_2$  concentrations during the terminations, requiring millennial variability during periods of reduced ice volume.

Ice core records from Antarctica extend down to 800 ka BP (EPICA Community Members 2004), although adequate resolution to consider millennial changes only exists for the last 500 kyr. New, multi-century resolution sea-level data from the Red Sea extend down to 500 ka BP (Rohling et al., 2009). Combined with IRD data from the north Atlantic (Oppo et al., 1998; McManus et al., 1999), these data allow a systematic analysis of millennial climate variability over the last 500 kyr (Fig. 1). Here we take a simple approach to analysing millennial variability over the last 500 kyr using these records. Although 800 kyr of data is available from the EPICA Dome C (EDC) ice core, we consider only the last 500 kyr for several reasons: (i) Red Sea sea-level and north Atlantic IRD data only extend to the last 500 kyr and we compare our results to these and; (ii) the resolution of the EDC deuterium record deteriorates with depth and millennial comparison is not possible much beyond 500 kyr ago.

#### 2 Method

Our method is simple but effective. Essentially we take a representative A-event (in this case the mean of A-events 1 to 4) from the MIS 3 section from the EDC ice core record as recorded in the deuterium temperature proxy, and look for similar events in the rest of the EDC record. We further validate our results by carrying out a series of Fast Fourier Transforms with overlapping windows (known as a spectrogram analysis) in addition. Our approach consists of five steps:

1. Removing the long-term trend from the EDC record

We first treat the last 500 kyr of EDC deuterium data on the EDC3 time scale to remove the 6 kyr trend (Fig. 1a). We achieve this by carrying out least-square linear regressions on 6 kyr windows centered on each data point in turn and removing the temperature trend. We will call the detrended, evenly sampled EDC deuterium record  $AA_{detrended}$  (Fig. 1b). The window length of 6 kyr is chosen to effectively remove the trend during the glacial terminations. Sensitivity tests varying the length of this 6 kyr window by a factor of two (between 3 ky and 12 kyr) show no impact on our results.

2. Defining the probe signal

We define an average A-event and use this event to probe the rest of the signal to look for similar events. To define an average A-event we took 4 kyr windows centered around 37.9, 46.2, 52 and 58.4 kyr ago representing A-events 1 to 4, respectively (Fig. 2). The four windows were overlain around their centres and the probe signal constructed by binning the combined data using the mean over 0.1 kyr intervals. Although, in the example given here we have used an average A-event, we could have used any of the four events individually because our result is not sensitive to which of the MIS 3 A-events we use. We will call the probe signal  $A_{mean}$ .

3. Probing  $AA_{detrended}$  with  $A_{mean}$ 

We compare  $AA_{detrended}$  with  $A_{mean}$  by calculating the standard difference between  $AA_{detrended}$  and  $A_{mean}$  for overlapping windows separated by 0.1 kyr The standard difference is then normalised to its greatest value over the last 500 kyr. We call this result  $AA_{probed}$  (Fig. 1c).

4. Calculate the standard deviation of *AA*<sub>probed</sub> over 8 kyr windows

Finally, because  $AA_{\text{probed}}$  is sensitive to both the presence of, and the phase of, millennial variability similar to that of A2, it exhibits both low (in phase) and high (out of phase) values during periods of millennial variability. Contrastingly,  $AA_{\text{probed}}$  shows little variation during periods without millennial variability. Therefore we also calculate the standard deviation from the mean of  $AA_{\text{probed}}$  over 8 kyr windows overlapped by 0.1 kyr.



**Fig. 1.** Data and results discussed in the text (see text for full details). (**A**) EPICA Dome C deuterium data (EPICA Community members 2004) on the EDC3 time scale. (**B**)  $AA_{detrended}$ , which is the EPICA Dome C deuterium data with the trend removed across 6 kyr windows. (**C**)  $AA_{probed}$ , which is the standard residual between  $AA_{detrended}$  and  $A_{mean}$  normalised to the maximum value over the last 500 kyr. Periods with alternating high and low values represent periods of millennial variability similar to  $A_{mean}$ . (**D**) The standard deviation of  $AA_{probed}$  over 8 kyr windows overlapped by 0.1 kyr. High values show periods of intense millennial variability similar to  $A_{2smooth}$ , low values show periods with little or no millennial variability. Also shown is precession forcing from Berger and Loutre (1991). (**E**) Red Sea sea-level record from Rohling et al. (2009). (**F**) IRD from ODP 980 (Oppo et al., 1998; McManus et al., 1999). Light grey bars represent periods of sustained ice sheet growth and the absence of millennial variability and dark grey bars represent periods with ice volume greater than 80 m sea-level equivalent.



**Fig. 2.** MIS 3 in the EPICA Dome C deuterium record (grey line). Also shown is the record smoothed with a Gaussian window of length 1 kyr (thick black line). Vertical grey bars centered at 37.9, 46.2, 52 and 58.4 ka representing A-events 1 to 4, respectively.

This result is normalised by the maximum standard deviation over the 500 kyr study period. The output of this calculation (Fig. 1d) is simple to interpret because where the standard deviation of  $AA_{\text{probed}}$  is high, the record contains millennial signals similar to  $A_{\text{mean}}$ .

5. Validation using a spectrogram analysis

Because of the novelty of our approach, we have also carried out a more routine spectral analysis of  $AA_{smoothed}$  using FFTs over 30 kyr windows with a 29.999 kyr overlap standard. The resulting spectrogram is shown in Fig. 3a, where the millennial band (3 to 5 kyr period) is highlighted. By taking the mean of the spectrogram within this millennial window of 3 to 5 kyr period, we produce a time series of the mean spectral power in this band. This is compared to precession forcing in Fi. 3b and the running standard deviation of  $AA_{probed}$  in Fig. 3c. We note that the two approaches in Fig. 3c give a very similar result, confirming that our result is not dependent on our novel approach.

# 3 Results

The results of our analysis are given in Fig. 1, where they are also compared to the Red Sea sea-level reconstruction and the IRD record of ODP 980 from the north Atlantic. The results are repeated in Fig. 3, where they are compared to our spectrogram results.

The most obvious result is that millennial variability similar to that of MIS 3 is repeated throughout the last 500 kyr in the EDC deuterium record (high values in Figs. 1d and 3 and Table 1). Periods of increased variability occur with striking regularity throughout the last 500 kyr. Far from being unusual events, they intermittently occupy approximately half of the record of the last 500 kyr.

In order to compare our results with reconstructions of ice volume, we plot the Red Sea sea-level curve (Fig. 1e). Clustering of millennial-scale variability is found to occur during two distinct phases within the glacial cycles, namely during periods of intermediate ice volume and extensive IRD deposition and during glacial terminations (including the early parts of interglacials) (Fig. 1e, f and Table 1). Conversely, millennial-scale variability also appears to be absent during two distinct phases within the glacial cycles, namely during periods of extensive ice-sheet growth (marked with light grey bars in Fig. 1 and referenced to periods of sea-level fall in Fig. 1e), and during glacial maxima with more than 80 m of sea-level equivalent ice volume (marked with dark grey bars in Fig. 1 and noted in Table 1).

Over the last 500 kyr, we observe twenty-four clusters of increased/decreased millennial-scale variability. These clusters are typically separated according to a "cycle" of  $\sim$ 21 kyr, a value similar to that of precession. However, the relationship between precession and millennial variability is not simple. Precession is generally in phase with the presence of millennial variability during periods of intermediate ice volume (40 to 80 m sea-level equivalent) and in antiphase during periods of reduced ice volume (40 m or less sea-level equivalent) (Figs. 1d, e, f, 3 and Table 1). Between periods when clusters of millennial variability are in and out of phase with precession, there are transition intervals when precession leads the millennial variability.

## 4 Discussion

Previous work has pointed out the links between millennialscale variability and intermediate ice volume of around 40 to 80 m ice volume equivalent (e.g. Oppo et al., 1998; Mc-Manus et al., 1999; Siddall et al., 2007). Other work has suggested that the link between millennial-scale variability and ice volume may be less important (Martrat et al., 2007; Desprat et al., 2009). Yet other work has focused on the role of millennial-scale variability in glacial terminations (Ganopolski et al., 2009).

Our signal analysis approach finds that there are two distinct regimes during which millennial-scale variability develops. One is associated with intermediate ice volume and episodes of IRD deposition (in line with the conclusions of Oppo et al., 1998; McManus et al., 1999, and Siddall et al., 2007), and the other is associated with glacial terminations (in line with the conclusions of Ganopolski et al., 2009). The



**Fig. 3.** Alternative analysis using a windowed FFT or spectrogram technique. (**A**) Spectrogram of  $AA_{detrended}$  using a 30 kyr window with a 29.999 kyr overlap. Dashed lines show precession (19 and 23 kyr period) and the millennial band (4 to 5 kyr period). (**B**) The natural log of the mean power in the spectrogram with a period between 3 and 5 kyr. Precession forcing from Berger and Loutre (1991) is shown for comparison. (**C**) The standard deviation of  $AA_{probed}$  over 8 kyr windows overlapped by 0.1 kyr and the natural log of the mean power in the spectrogram with a period between 4 and 5 kyr. Light grey bars represent periods of sustained ice sheet growth and the absence of millennial variability and dark grey bars represent periods with ice volume greater than 80 m sea-level equivalent (see Fig. 1e).

Age window (ka)	0– 20	20- 40	40- 60	60- 80	80– 100	100– 120	120– 140	140– 160	160– 180	180- 200	200– 220	220– 240	240– 260	280– 300	320- 340	340– 360	360- 380	380– 400	400- 420	420- 440	440- 460	460- 480	480– 500
Millennial variability	L H	L	Н	Н	L	L	Н	L	н	н	L	Н	H L	L H	L H	L	н	L	н	н	L	н	L
Phase Sea level	O H D	T L	I I	I F	T F	0 H	O H D	T L	I I	I F	T F	T F	O H D	T F	I F	T D L	0 I	T F	I H	I D	I L	I F	0 H
IRD	$N \\ Y$	Y	Y	Y	Ν	Ν	N Y	Y	Ν	N	Ν	Ν	N Y	Ν	Ν	Y	Y	Ν	N	Y	Y	N	Ν

**Table 1.** A qualitative comparison of millennial variability, phase with respect to precession, ice volume (sea level) and IRD. These estimates are based on Fig. 1.

Millennial variability: H=high, L=low

Phase with respect to precession: O=out of phase, T=transition, I=in phase Sea level: H=high, L=low, D=deglaciation, F=falling IRD: Y=yes, N=no

similarity between both classes of millennial-scale events in the EPICA Dome C deuterium records suggests that both classes reflect the same processes.

If millennial-scale variability exists for approximately half of the time during the last 500 kyr, then there is an equally important reverse question: "What characterises periods which have reduced millennial variability or for which millennial variability is absent?" In Fig. 1 these periods are characterized by either ice volumes greater than  $\sim 80 \,\mathrm{m}$  icevolume equivalent or periods of sustained ice sheet growth. Many authors have shown evidence for millennial-scale sealevel fluctuations associated with millennial-scale climate variability during MIS 3 (Chappell 2002; Cutler et al., 2003; Siddall et al., 2003, 2008; Rohling et al., 2008; Arz et al., 2007). We suggest that sustained ice-sheet growth requires the absence of millennial variability to allow the continuous growth of ice sheets, rather than the growth and collapse of ice sheets during periods such as MIS 3. The lack of millennial variability at the glacial maxima may be evidence that the presence of the large ice sheets issues in a relatively stable climate during those periods.

Given that only one of our classes of millennial-scale event is linked with IRD deposition, we infer that massive iceberg calving may not be the ultimate cause of millennial variability. We note that IRD events only last several centuries (less than D-O stadials, which last 1–2 kyr) and peak at the conclusion of D-O stadials (Hemming, 2004). This suggests that IRD events are caused by the D-O stadials, rather than being the cause of D-O stadials. Hence, IRD deposition (iceberg calving) does not appear to be the driving mechanism, but merely a feature of the class of millennial-scale variability that occurs during periods of intermediate ice volume.

The above implies that another causative mechanism is required to explain D-O variability. Because the recurrence interval of clusters of millennial variability is close to precession any discussion of processes explaining millennial variability needs to consider precession forcing. It has been suggested that changes in the zonality of atmospheric circulation over the North Atlantic might instead be the cause of millennial variability by switching the latitude of the polar front between different modes (Seager et al., 2002; Seager and Battisti, 2007). As noted above, the mean recurrenceperiod of clusters of millennial-scale variability in Fig. 1c,d is 21 kyr, similar to precession. This implies that changes in the zonality of atmospheric circulation over the North Atlantic may be linked to precessional cycles. In turn this implies zonal teleconnections in the Northern Hemisphere. Such links are indeed suggested by Wang et al. (2008) who found both strong precessional and millennial variability in Chinese speleothem records.

If precession forcing sets the atmospheric preconditions for millennial variability, what mechanisms might link precession to individual millennial events? Several possible links have been suggested linking the atmospheric circulation in the Northern Hemisphere to millennial changes in the MOC. Changes in sea ice extent have been suggested to generate seesaw-type temperature variations in the MOC (Kaspi et al., 2004) and are linked to variations in atmospheric circulation in the Northern Hemisphere (Li et al., 2005). Sea-ice feedbacks with the MOC might therefore be the mechanism behind individual millennial events, given the correct preconditions with respect to precession forcing. A second explanation might be given by the direct link between the magnitude of the wind stress in the North Atlantic and the stability of the MOC found by Ashkenazy and Tziperman (2007). Such an explanation would remove the requirement of seaice feedbacks to force changes in the MOC. Finally Friedrich et al. (2009) have used the LOVECLIM model to find a feedback between sea-ice anomalies and the atmospheric circulation. This feedback in turn drives atmospheric anomalies over Hudson Bay which form a freshwater pulse into the Labrador Sea, which then causes the MOC to collapse.

In order to consider the possible link between precession forcing and the occurrence of millennial variability, we compare our result with precession forcing in Fig. 1d. Clusters of millennial variability are in phase with precession during periods of intermediate ice volume but alter phase during glacial maxima (when there is little or no millennial variability). Clusters of millennial variability then remain in antiphase during the glacial termination and periods of reduced ice volume. During the transition from reduced to intermediate ice volume the phase realigns again to put the reoccurrence of millennial variability back in phase with precession forcing. Marked changes to the atmospheric circulation due to the presence of large ice sheets and extended sea ice at the Last Glacial Maximum (LGM) have been found in atmospheric models (e.g. Li and Battisti, 2008). Otto-Bliesner et al. (2006) also found that atmospheric circulation was sensitive to the detailed configuration of the north American ice sheets at the LGM.

In summary, we speculate that the existence of ice sheet topography and albedo effects modifies the influence of precession forcing on millennial variability and that this may explain the existence of two classes of millennial variability. One class of millennial variability is linked to precession forcing, the other class requires precession forcing modified by the existence of ice sheets of intermediate size. Because precession forcing affects how much insolation an entire hemisphere receives in a certain season, precession may be an important driver of changes in seasonal sea ice. We therefore suggest that sea-ice feedbacks are the most likely candidate for linking millennial variability to precession forcing.

## 5 Conclusions

It is clear from our analysis that millennial-scale variability is a much more common feature of Quaternary climate than has hitherto been realised, taking up a significant proportion of the variability during periods of intermediate icesheet growth and glacial terminations.

Millennial variability occurs during distinct periods in the glacial cycles, namely periods with ice volume equivalent to 40–80 m of sea-level lowering, and glacial terminations.

Different periods give rise to the absence of millennial variability, namely periods of extensive ice-sheet growth, and periods with ice volume equivalent to more than 80 m of sea-level lowering.

The lack of a unique relationship between ice volume and millennial-scale variability calls for a causative mechanism other than iceberg release into the north Atlantic. We suggest that changes in the zonality of atmospheric circulation over the North Atlantic may be a viable alternative (Seager et al., 2002; Seager and Battisti, 2007). Because we find the mean recurrence-period of the millennial-scale variability clusters to be  $\sim 21$  kyr, we suggest that changes in the zonality of atmospheric circulation over the North Atlantic may be linked

to precession. Because the phase relationship between precession forcing and clusters of millennial variability changes over the glacial cycle, we suggest that the existence of ice sheets modifies the influence of precession forcing on millennial variability. This suggestions is supported by Chinese speleothem records, which show strong precessional and millennial variability over the last two glacial cycles, indicating important changes in the zonality of atmospheric circulation on these time scales (Wang et al., 2008).

If millennial variability requires preconditions with respect to precessional forcing, it is unlikely to occur due to natural forcing for some thousands of years from now.

Acknowledgements. Mark Siddall is supported by an RCUK research fellowship. This work has been greatly helped by interactions taking place within the PALSEA working group, which is funded by PAGES and IMAGES. EJR acknowledges support from NERC projects NE/C003152/1, NE/E01531X/1, and NE/H004424/1.

Edited by: T. Kiefer



The publication of this article was sponsored by PAGES.

#### References

- Arz, H. W., Lamy, F., Ganopolski, A., Nowaczyk, N., and Pätzold, J.: Dominant Northern Hemisphere climate control over millennial-scale glacial sea-level variability, Quat. Sci. Rev., 26(3–4), 312–321, doi:10.1016/j.quascirev.2006.07.016, 2007.
- Ashkenazy, Y. and Tziperman, E.: A Wind-Induced Thermohaline Circulation Hysteresis and Millennial Variability Regimes, J. Phys. Oceanog., 37(10), 2446–2457, 2007.
- Berger, A. and Loutre, M. F.: Insolation values for the climate of the last 10 million years. Quat. Sci. Rev., 10, 297–317, 1991.
- Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., and Hawkesworth, C. J.: Sea-land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals, Geochim. Cosmochim. Acta, 67(17), 3181– 3199, 2003.
- Blunier, T. and Brook, E.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial period, Science, 291, 109–112, doi:10.1126/science.291.5501.109, 2001.
- Blunier, T., Chappellaz, J., Schwander, J., Dällenbach, A., Stauffer, B., Stocker, T., Raynaud, D., Jouzel, J., Clausen, H. B., Hammer, C. U., and Johnsen, S. J.: Asynchrony of Antarctica and Greenland climate during the last glacial, Nature, 394, 739–743, 1998.
- Bond, G. C. and Lotti, R.: Iceberg discharges into the North-Atlantic on millennial time scales during the last glaciation, Science, 267(5200), 1005–1010, doi:10.1126/science.267.5200.1005, 1995.

- Chappell, J.: Sea level changes forced ice breakouts in the Last Glacial Cycle: New results from coral terraces, Quat. Sci. Rev., 21(10), 1229–1240, doi:10.1016/S0277-3791(01)00141-X, 2002.
- Crowley, T. J.: North Atlantic deep water cools the Southern Hemisphere, Paleoceanography, 7, 489–497, 1992.
- Cutler, K. B., Edwards, R. L., Taylor, F. W., Cheng, H., Adkins, J., Gallup, C. D., Cutler, P. M., Burr, G. S., and Bloom, A. L.: Rapid sea-level fall and deep-ocean temperature change since the last interglacial period, Earth Planet. Sci. Lett., 206(3–4), 253–271, doi:10.1016/S0012-821X(02)01107-X, 2003.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N., Hammer, C. U., and Oeschger, H.: North Atlantic climatic oscillations revealed by deep Greenland ice cores, in: Climate Processes and Climate Sensitivity, edited by: Hansen, J. E. and Takahashi, T., Geophys. Monogr. Ser., 29, 1984.
- Dansgaard, W., Johnsen, S. J., Clausen, H. B., Dahl-Jensen, D., Gundestrup, N. S., Hammer, C. U., Hvidberg, C. S., Steffensen, J. P., Sveinbjornsdottir, A. E., Jouzel, J., and Bond, G.: Evidence for general instability of past climate from a 250-ka icecore record, Nature, 364, 218–220, 1993.
- Delmotte, M., Chappellaz, J., Brook, E., Yiou, P., Barnola, J. M., Goujon, C., Raynaud, D., and Lipenkov, V. I.: Atmospheric methane during the last four glacial-interglacial cycles: Rapid changes and their link with Antarctic temperature, J. Geophys. Res., 109, D12104, doi:10.1029/2003JD004417, 2004.
- Denton, G. H., Alley, R. B., Comer, G. C., and Broecker, W. S.: The role of seasonality in abrupt climate change, Quat. Sci. Rev., 24(10–11), 1159–1182, doi:10.1016/j.quascirev.2004.12.002, 2005.
- Desprat, S., Sánchez Goñi, M. F., McManus, J. F., Duprat, J., and Cortijo, E.: Millennial-scale climatic variability between 340 000 and 270 000 years ago in SW Europe: evidence from a NW Iberian margin pollen sequence, Clim. Past, 5, 53–72, 2009, http://www.clim-past.net/5/53/2009/.
- EPICA Members: Eight glacial cycles from an Antarctic ice core, Nature, 429, 623–628, 2004.
- EPICA Community Members: One-to-one interhemispheric coupling of polar climate variability during the last glacial, Nature, 444, 195–198, doi:10.1038/nature05301, 2006.
- Friedrich, T., Timmermann, A., Timm, O., Mouchet, A., and Roche, D. M.: Orbital modulation of millennial-scale climate variability in an earth system model of intermediate complexity, Clim. Past Discuss., 5, 2019–2051, 2009,

http://www.clim-past-discuss.net/5/2019/2009/.

- Ganopolski, A. and Roche, D.: On the nature of lead-lag relationships during glacial-interglacial climate transitions, Quat. Sci. Rev., 37/38, 3361–3378, doi:10.1016/j.quascirev.2009.09.019, 2009.
- Hemming, S. R.: Heinrich events: Massive late Pleistocene detritus layers of the North Atlantic and their global climate imprint, Rev. Geophys., 42, RG1005, doi:10.1029/2003RG000128, 2004.
- Kaspi, Y., Sayag, R., and Tziperman, E.: A "triple sea-ice state" mechanism for the abrupt warming and synchronous ice sheet collapses during Heinrich events, Paleoceanography, 19(3), PA3004, doi:10.1029/2004PA001009, 2004.
- Larrasoaña, J. C., Roberts, A. P., Rohling, E. J., Winklhofer, M., and Wehausen, R.: Three million years of monsoon variability

over the northern Sahara, Clim. Dyn., 21, 689-698, 2003.

- Li, C. and Battisti, D. S.: Reduced Atlantic stormines during the Last Glacial Maximum: Evidence from a coupled climate model, J. Climate, 21, 3561–79, 2008.
- Li, C., Battisti, D. S., Schrag, D. P. and Tziperman, E.: Abrupt climate shifts in Greenland due to displacements of the sea ice edge, Geophys. Res. Lett., 32, L19702, doi:10.1029/2005GL023492, 2005.
- Martrat, B., Grimalt, J. O., Shackleton, N. J., de Abreu, L., Hutterli, M. A. and Stocker, T. F.: Four climate cycles of recurring deep and surface water destabilizations on the Iberian Margin, Science, 317, 502–507, 2007.
- McManus, J. F., Oppo, D. W., and Cullen, J. L.: A 0.5 millionyear record of millennial-scale climate variability in the North Atlantic, Science, 283(5404), 971–975, doi:10. 1126/ science.283.5404.971, 1999.
- Oeschger, H., Beer, J., Siegenthaler, U., Stauffer, B., Dansgaard, W., and Langway, C. C.: Late glacial climate history from ice cores, in: Climate Processes and Climate Sensitivity, edited by: Hansen, J. E. and Takahashi, T., Geophysical Monograph Series, 29, AGU, Washington, DC, USA, 299–306, 1984.
- Oppo, D. W., McManus, J. F., and Cullen J. L.: Abrupt climate events 500,000 to 340 000 years ago: Evidence from subpolar north Atlantic sediments, Science, 279(5355), 1335–1338, doi:10.1126/science.279.5355.1335, 1998.
- Otto-Bliesner, B. L., Brady, E. C., Clauzet, G., Tomas, R., Levis, S., and Kothavala, Z.: Last Glacial Maximum and Holocene Climate in CCSM3, J. Climate, 19, 2526–2544, 2006.
- Rohling, E. J., Mayewski, P. A., and Challenor, P.: On the timing and mechanism of millennial-scale climate variability during the last glacial cycle, Clim. Dyn., 20, 257–267, 2003.
- Rohling, E. J., Grant, K., Hemleben, C., Kucera, M., Roberts, A. P., Schmeltzer, I., Schulz, H., Siccha, M., Siddall, M., and Trommer, G.: New constraints on the timing of sea level fluctuations during early to middle marine isotope stage 3, Paleoceanography, 23, PA3219, doi:10.1029/2008PA001617, 2008.
- Rohling, E. J., Grant, K., Bolshaw, M., Roberts, A.P., Siddall M., Hemleben C., and Kucera M.: Antarctic temperature and global sea level closely coupled over the past five glacial cycles, Nat. Geosci., 2, 500–504, 2009.
- Seager, R., Battisti, D. S., Yin, J., Naik, N., Gordon, N., Clement, A. C., and Cane, M.: Is the Gulf Stream responsible for Europe's mild winters? Q. J. R. Meteo. Soc., 128, 2563–86, 2002.
- Seager, R. and Battisti, D. S.: Challenges to our understanding of the general circulation: Abrupt climate change, in: Global Circulation of the Atmosphere, edited by: Schneider, T. and Sobel, A. H., 331–371, Princeton Univ. Press, Princeton, 2007.
- Shackleton, N. J., Hall, M. A., and Vincent, E.: Phase relationships between millennial-scale events 64 000–24 000 years ago, Paleoceanography, 15, 565–569, doi:10.1029/2000PA000513, 2000.
- Siddall, M., Rohling, E. J., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmelzer, I., and Smeed, D. A.: Sea-level fluctuations during the last glacial cycle, Nature, 423, 853–858, doi:10.1038/nature01690, 2003.
- Siddall, M., Stocker, T. F., Spahni, R., Blunier, T., McManus, J. F., and Bard, E.: Using a maximum simplicity paleoclimate model to simulate millennial variability during the last four glacial cycles, Quat. Sci. Rev., 25, 3185–3197, doi:10.1016/j.quascirev.2005.12.014, 2006.

- Siddall, M., Stocker, T. F., Blunier, T., Spahni, R., Schwander, J., Barnola, J.-M., and Chappellaz, J.: Marine Isotope Stage (MIS) 8 millennial variability stratigraphically identical to MIS 3, Paleoceanography, 22, PA1208, doi:10.1029/2006PA001345, 2007.
- Siddall, M., Rohling, E. J., Thompson, W. G., and Waelbroeck, C.: MIS 3 Sea-level fluctuations: data synthesis and new outlook, Rev. Geophys., 46, RG4003, doi:10.1029/2007RG000226, 2008.
- Stocker, T. F. and Johnsen, S. J.: A minimum thermodynamic model for the bipolar seesaw, Paleoceanography, 18(4), 1087, doi:10.1029/2003PA000920, 2003.
- Voelker, A. H. L.: Global distribution of centennial-scale records for marine isotope stage (MIS) 3: A database, Quat. Sci. Rev., 21(10), 1185–1212, doi:10.1016/S0277-3791(01)00139-1, 2002.
- Wang, Y. J., Cheng, H., Edwards, R. L., An, Z. S., Wu, J. Y., Shen, C. C., and Dorale, J. A.: A high-resolution absolute-dated Late Pleistocene monsoon record from Hulu Cave, China, Science, 294(5550), 2345–2348, doi:10.1126/science.1064618, 2001.
- Wang, Y. J., Cheng, H., Edwards, R. L., Kong, X. G., Shao, X. H., Chen, S. T., Wu, J. Y., Jiang, X. Y., Wang, X. F. and An, Z. S.: Millennial- and orbital-scale changes in the East Asian monsoon over the past 224 000 years, Nature, 451, 1090–1093, 2008.
- Yuan, D., Cheng, H., Edwards, R. L., Dykoski, C. A., Kelly, M. J., Zhang, M., Qing, J., Lin, Y., Wang, Y., Wu, J., Dorale, J.A., An, Z., and Cai, Y.: Timing, Duration, and Transitions of the Last Interglacial Asian Monsoon, Science, 304(5670), 575–578, 2004.