

Turning the Table: Plants Consume Microbes as a Source of Nutrients

Chanyarat Paungfoo-Lonhienne^{1*}, Doris Rentsch², Silke Robatzek³, Richard I. Webb⁴, Evgeny Sagulenko⁵, Torgny Näsholm⁶, Susanne Schmidt^{1,9}, Thierry G. A. Lonhienne^{1,9}

1 School of Biological Sciences, The University of Queensland, Brisbane, Queensland, Australia, **2** Institute of Plant Sciences, University of Bern, Bern, Switzerland, **3** The Sainsbury Laboratory, Norwich Research Park, Norwich, United Kingdom, **4** Centre for Microscopy and Microanalysis, The University of Queensland, Brisbane, Queensland, Australia, **5** School of Biochemistry and Molecular Biosciences, The University of Queensland, Brisbane, Queensland, Australia, **6** Department of Forest Ecology and Management, Swedish University of Agricultural Sciences, Umeå, Sweden

Abstract

Interactions between plants and microbes in soil, the final frontier of ecology, determine the availability of nutrients to plants and thereby primary production of terrestrial ecosystems. Nutrient cycling in soils is considered a battle between autotrophs and heterotrophs in which the latter usually outcompete the former, although recent studies have questioned the unconditional reign of microbes on nutrient cycles and the plants' dependence on microbes for breakdown of organic matter. Here we present evidence indicative of a more active role of plants in nutrient cycling than currently considered. Using fluorescent-labeled non-pathogenic and non-symbiotic strains of a bacterium and a fungus (*Escherichia coli* and *Saccharomyces cerevisiae*, respectively), we demonstrate that microbes enter root cells and are subsequently digested to release nitrogen that is used in shoots. Extensive modifications of root cell walls, as substantiated by cell wall outgrowth and induction of genes encoding cell wall synthesizing, loosening and degrading enzymes, may facilitate the uptake of microbes into root cells. Our study provides further evidence that the autotrophy of plants has a heterotrophic constituent which could explain the presence of root-inhabiting microbes of unknown ecological function. Our discovery has implications for soil ecology and applications including future sustainable agriculture with efficient nutrient cycles.

Citation: Paungfoo-Lonhienne C, Rentsch D, Robatzek S, Webb RI, Sagulenko E, et al. (2010) Turning the Table: Plants Consume Microbes as a Source of Nutrients. PLoS ONE 5(7): e11915. doi:10.1371/journal.pone.0011915

Editor: Juergen Kroymann, CNRS UMR 8079/Université Paris-Sud, France

Received: June 7, 2010; **Accepted:** July 7, 2010; **Published:** July 30, 2010

Copyright: © 2010 Paungfoo-Lonhienne et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was funded by the Australian Research Council (DP0986495 to SS, DR, TN, and SR) and The University of Queensland (to CPL). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Chanyarat@uq.edu.au

⁹ These authors contributed equally to this work.

Introduction

Plants and microbes have evolved detrimental and beneficial relationships. Detrimental relationships involve pathogens including fungi, bacteria and viruses [1] and the hallmark of pathogenic interactions is the suppression and interference with plant immune responses [2,3]. Beneficial relationships include symbiosis [1], diazotrophic endophytes that supply the plant with fixed nitrogen [4,5] and other endophytic associations that promote plant growth by producing phytohormones, volatiles, defence compounds, and enzymes [6,7,8,9,10]. A less well-defined beneficial relationship involves the association of plant roots with microbes in the rhizosphere. Roots attract soil microbes by exuding nutrient sources including carbohydrates, organic and amino acids [11,12,13,14] and the density of microbes in the rhizosphere is much higher than in bulk soil [15]. According to the "soil microbial loop" concept, nutrients and carbon are cycled between soil and microbial pools [16,17,18], and inorganic and organic nutrients of low molecular mass become available through microbial turnover of soil organic matter and are subsequently 'scavenged' by the plant root.

However, new concepts are emerging which point to a wider range of nutrient sources for plants [19] and question the 'soil

microbial loop' concept. We recently demonstrated that roots can incorporate large organic molecules including proteins and DNA [20,21], and this implies that plants may be less dependent on microbial activity for break-down of organic matter than currently assumed. Adding to mounting questions of plant-microbe interactions in soil is the discovery that diverse microbes without known relationships with plants exist in roots [22].

Here, we explored the possibility that plants take up and digest microbes as a source of nutrients. We discovered that *Arabidopsis* (*Arabidopsis thaliana*) and tomato (*Lycopersicon esculentum*) are able to take up non-pathogenic *E. coli* and *S. cerevisiae* into root cells, digest and use these microbes as a nutrient source. Our results show that the uptake process involves modification of the walls of root cells which is followed by active incorporation and degradation of the incorporated microbes.

Results and Discussion

Bacteria and yeast are taken up by *Arabidopsis* and tomato

To examine if plants take up microbes and use them as a nutrient source, we incubated roots of intact *Arabidopsis* and tomato plants with *E. coli* BL21 and yeast *S. cerevisiae* which express

the green fluorescent protein ($GFP^{E. coli}$ and GFP^{yeast}). To examine plants with different root specialisations, we chose *Arabidopsis* which does not form symbiotic relationships and tomato which forms symbioses with mycorrhizal fungi, but was grown here without symbionts. Plants were cultivated in non-axenic hydroponic (tomato) and axenic agar (*Arabidopsis*) culture. Microbial solution was added to growth media ensuring that roots were not disturbed or damaged. After 12 h (tomato) or 4 h (*Arabidopsis*) incubation, $GFP^{E. coli}$ and GFP^{yeast} were detected in root hairs and the rhizodermis and cortex of mature zones of the roots ($GFP^{E. coli}$, Figure 1A–D; yeast, Figure 1E–F) by confocal laser scanning microscopy (CLSM). Cytoplasmic streaming in root hairs ($GFP^{E. coli}$: movie S1; GFP^{yeast} : movie S2) and other root cells (movie S3)

was indicative of live and active plant cells. Similar results were obtained with *Arabidopsis* axenic hydroponic culture and soil-grown *Arabidopsis* or tomato. To demonstrate the specificity of the uptake process, we incubated *Arabidopsis* with 5 μm nano-silica fluorescent beads similar in size to yeast (3–5 μm) but larger than *E. coli* (<2 μm). No beads were detected in roots and few beads were attached to root surface after washing (Figure S1) suggesting that roots recognize microbes and this results in targeted incorporation.

CLSM of root cross-sections revealed microbes were present in epidermis cells, cortex cells and the apoplastic space, but absent from tissue separated by the Casparian strip (Figure 2A and B; Figure S2; movie S4 and S5). Transfer of bacteria from root




Figure 1. Roots of axenically grown *Arabidopsis* and tomato were incubated with *E. coli* or yeast expressing green fluorescent protein ($GFP^{E. coli}$ or GFP^{yeast}). $GFP^{E. coli}$ was detected at the surface of roots and root hairs (A and C), and inside roots and root hairs (B and D). GFP^{yeast} was present inside roots and root hairs (E and F). (A, D and F) and (B, C and E) correspond to tomato and *Arabidopsis* root, respectively. Fluorescent images were taken by confocal laser scanning microscopy (CLSM). doi:10.1371/journal.pone.0011915.g001




Figure 2. Root transverse sections and electron micrographs of tomato and Arabidopsis show ^{GFP} *E. coli* in the apoplast and inside root cells. *E. coli* was detected inside tomato roots (A, C and D, E and F) and Arabidopsis roots (B). (A and B) Fluorescent images of transverse sectioned roots taken by CLSM. (C and D) Images taken by a transmission electron microscope. White triangles in (C) indicate *E. coli* cell present in apoplast. (D) Roots were probed with immunogold-labeled anti-GFP revealing *E. coli* in root cortex cells. Sub-image in (D) is a detail of dash-white square box. Gold labeling is marked with white arrows. Rhizodermis cell (R) and plant cell wall (pcw) is indicated. (F) is a detail image of (E) showing plant cells containing *E. coli*, and both images were taken by SEM.
doi:10.1371/journal.pone.0011915.g002

surfaces was avoided by coating roots with agar prior to processing. Transmission Electron Microscopy (TEM) verified that cells of *E. coli* Bl21 were present in the intercellular space (Figure 2C) and inside cortex cells (Figure 2D). This finding was confirmed with Scanning Electron Microscopy (SEM) showing *E. coli* Bl21 in epidermis cells (Figure 2E and F) and demonstrates that non-pathogenic and non-symbiotic microbes enter cells of mature roots.

After uptake, microbes are confined to root cortex cells where they are degraded

We investigated the fate of microbes after incorporation into root cells. Hydroponic tomato plants were incubated overnight with ^{GFP} yeast. Since expression of GFP in yeast clone *TDH3* (YGR192C) is constitutive, monitoring of GFP fluorescence allows an assessment of yeast cells activity. Three hours after incubation, fluorescing ^{GFP} yeast cells were detected at the root surface and inside root cells (Figure 3A). After 3 days, ^{GFP} yeast was only

detected inside roots, with some yeast cells alive and fluorescing, and some non-fluorescing yeast cells displaying an altered shape (Figure 3A). Few yeast cells were fluorescing after 7 days, no GFP signal was detected after 10 days, and root cells contained only debris of yeast cells after 14 days (Figure 3A). To support microscopy findings, we quantified the TDH3:GFP fusion protein (expressed constitutively by ^{GFP} yeast) in roots harvested in parallel with CLSM-inspected plants by western blot analysis (Figure 3B). TDH3:GFP in roots strongly diminished over time (Figure 3B). No protein was detected in roots after 10 and 14 days incubation, confirming CLSM findings.

Over the time of the experiment, tomato plants retained a healthy phenotype. To confirm that *E. coli* Bl21 was not a threat to plants, we incubated Arabidopsis grown on MS medium with *E. coli* Bl21 and monitored growth for 14 days. Similarly to tomato, Arabidopsis plants grown with or without *E. coli* Bl21 had similar appearance (Figure S3).



A**B**

Figure 3. Time course experiment of yeast degradation in tomato roots. (A) The number of living yeast cells (fluorescing green) in tomato decreased over time as observed by CLSM. (B) The amount of recombinant TDH3:GFP protein present inside the roots decreased over time. Equal amounts of proteins from root extracts were separated by SDS-PAGE (B, Left) and analyzed by western blot using anti-GFP antibody to detect yeast recombinant TDH3:GFP protein (B, Right).
doi:10.1371/journal.pone.0011915.g003

CLSM analysis revealed no evidence that $GFP^{E. coli}$ or GFP^{yeast} were transported to leaves. The absence of microbes in leaves was confirmed as leaf homogenates incubated on LB media containing selective antibiotics did not produce colonies. Similarly, incubation of *Arabidopsis* roots with *Salmonella typhimurium* caused proliferation of *Salmonella* in root cells, but not leaves [23].

Taken together, our results demonstrated that upon incorporation, *E. coli* BI21 and yeast are confined to the root cells where they are degraded.

E. coli BI21 induces cellulase(s) activity in roots of *Arabidopsis*

A central question is how *E. coli* and yeast enter intact root cells. Plant cells possess walls composed of a highly integrated and structurally complex network that acts as barrier to larger molecules, particles and microbes [24]. We assumed that *E. coli* BI21 and yeast can only enter intact root cells if cell walls are degraded prior to entry. Pathogens attack cell walls by secretion of polysaccharide-degrading enzymes including polygalacturonases

and cellulases [25], and *Rhizobium* infection in the legume root symbiosis occurs through the activity of a cell-bound bacterial cellulase [26]. There are no reports of non-pathogenic and non-symbiotic microbes degrading plant cell walls.

We therefore examined whether plant-derived cell wall degrading enzymes facilitate entry of the microbes studied here. Hydroponic Arabidopsis were incubated overnight with *E. coli* BI21 and transferred to liquid MS medium containing resorufin- β -D-cellobioside (Res-CB), an artificial substrate for cellulases that emits red fluorescence upon cleavage [27]. Fluorescence increased in *E. coli* BI21-incubated roots (Figure 4A) but not in *E. coli* incubated with Res-CB or roots grown without *E. coli*. The assay does not allow localization of the origin of cellulase activity because the generated resorufin diffuses rapidly through tissues. These results indicate that the presence of *E. coli* in the medium triggers the induction of cellulase expression in Arabidopsis, and this may be linked to the uptake of *E. coli* into root cells.

E. coli BI21 induces plant cell wall-like outgrowth

Additional insights into the mechanisms of microbial uptake were obtained by TEM. Clusters of *E. coli* BI21 at the surface of root cells were systematically surrounded by a thin layer of an undetermined structure. This layer resembled the structure that was equivocally reported to be a matrix-like bacterial substance suggestive of a site from which bacterial cells may gain entry into young roots [28] or mucilaginous material secreted either by plants or bacteria for binding bacteria to the root surface [29]. Our results suggest that this structure consists of cell wall components as it was connected to the cell wall of the rhizodermis (Figure 4B). Dual gold (Au)-labeling of sections with Au-labeled cellulase (10 nm) and Au-labeled anti-GFP antibody (15 nm) showed that this structure is at least partly composed of cellulose (Figure 4C), and indicate that Arabidopsis synthesizes a cell wall-like structure that contains cellulose. Thus, a step in the process of the acquisition of microbes by roots may involve ‘corralling’ microbes at the root surface by cell wall-like outgrowth for subsequent incorporation. This sophisticated mechanism has interesting connotations with the mechanism used by *Agrobacterium tumefaciens* to adhere to the root surface of plants for infection. During the infection process, *Agrobacterium* produce cellulose fibrils via the activity of its own cellulose synthases to strengthen its adherence to the surface of the roots [30].

E. coli BI21 triggers extensive alteration of the expression of genes involved in cell wall modification

To further explore mechanisms involved in the observed plant-microbe interactions, we proceeded with genome-wide transcriptome analysis of Arabidopsis roots incubated with *E. coli* BI21 for 24 hours. Microarray data revealed that a numerous number of genes involved in cell wall modification increased in expression (Figure 5). Strongly induced were the expression of cellulases (endo-glucanases) and other cell wall degrading enzymes including pectinases and xyloglucan endotransglycosidases (Figure 5A), supporting our biochemical analysis (Figure 4A). Expression of expansins, involved in cell wall loosening, was also highly up-regulated (Figure 5B). Consistent with EM data demonstrating cell wall-like outgrowth containing cellulose (Figure 4B and C), cellulose synthases, cellulose synthase-like, and extensins were strongly up-regulated (Figure 5C–D). Cellulose synthase-like proteins (CLSs) are involved in the linkage of non-cellulosic polysaccharides [31]. The induction of cellulose synthase-like genes (*CSLs*) observed here is interesting in the view that mutation of *CSLA9* (At5G03760) in Arabidopsis leads to inhibition of *Agrobacterium*-mediated root transformation through reduced ability

of the roots to bind *A. tumefaciens* [32]. Zhu et al. (2003) did not observe any major differences in the linkage structure of the non-cellulosic polysaccharides in the *CSLA9* defective mutant and hypothesized that the defect in binding *A. tumefaciens* may arise from the altered ability of the Arabidopsis mutant to secrete particular polysaccharides necessary for bacterial recognition of the host and subsequent attachment. Our results suggest that CLSs may be involved in the recognition and attachment of *E. coli* BI21 at the root surface. Extensins are also potential candidates for such function because they have been reported to be involved in similar processes. Cannon et al. (2008) [33] have established that extensins are involved in the formation of the cross wall (cell plate) during cytokinesis and they proposed that self-assembling extensins serve as scaffolds for ordered pectin deposition in the cell plate. It is reasonable to conceive that the extensins that are over expressed in reaction of *E. coli* BI21 treatment have a role in the cell wall-like outgrowth.

In relation to microbe attachment at the surface of the roots, the over-expression of arabinogalactan-proteins (Figure 5E) involved in formation of wall ingrowths [34] is also likely to be relevant. Mutation of arabinogalactan-protein AGP17 (At2G23130) in Arabidopsis resulted in decreased efficiency of *Agrobacterium*-induced transformation due to altered binding to the root surface caused by reduced direct binding or impaired signaling pathway(s) [35]. The strong induction of nearly all arabinogalactan-genes of Arabidopsis treated with *E. coli* BI21 (Figure 5E) corroborates with arabinogalactan proteins promoting binding of microbes to the root surface.

Further, it is possible that endocytosis or a related process is involved in the incorporation of microbes into root cells. Induction of genes involved in cytoskeleton structure and re-organization (Figure S4) supports this hypothesis.

Altogether, our results indicate that uptake of microbes by roots occurs through major structural modifications of root cells controlled by the plant, including outgrowth of a cell-wall like structure capturing microbes, and degradation and/or loosening of cell walls with plant-derived enzymes. In contrast, entry of pathogenic and symbiotic microbes into root cells is controlled predominantly or partly by microbes; pathogenic fungi enter plants by secreting enzymes that degrade plant cell walls [36], and infection of legume root cells with symbiotic rhizobia requires formation of special root hairs initiated by rhizobia-secreted Nod factors [37]. The process of colonization of roots by diazotrophic endophytes is also fundamentally different from our observations in mature roots, because diazotrophic endophytes enter in elongation zones and through cracks at the point of lateral roots emergence [4,38,39]. Although root colonization by diazotrophic endophytes involves cell wall degradation processes, the source of the cell-wall degrading enzymes differs. Diazotrophs release plant cell-wall-degrading enzymes for the ingress into roots [40,41], whereas plant-derived cell-wall degrading enzymes facilitated entry of *E. coli* and yeast into mature roots. Thus, the uptake of *E. coli* and yeast involves mechanisms which have not been described by previous research, further indicating that the observed processes are hitherto un-described interactions between microbes and plants.

E. coli BI21 is a nitrogen source for plants

To determine whether microbes are a nutrient source for plants, we incubated roots of hydroponic tomato plants for 1 h with ^{15}N -labelled *E. coli* BI21 (^{15}N -*E. coli*) and analyzed new leaves for ^{15}N content. Controls included plants not incubated with *E. coli* and plants incubated with filtrate of ^{15}N -*E. coli* solution to account for possible ^{15}N release from bacteria during incubation. Plants were




Figure 4. Root produced cellulase and extended the cell wall when incubated with *E. coli* BI21. (A) Incubation of Arabidopsis roots in 31 $\mu\text{g/mL}$ resorufin cellulobioside after incubating overnight with *E. coli*. After 2 h incubation, roots were viewed by CLSM. (B) TEM image of cell wall-like structure of plant roots encompassing bacteria. (C) TEM image of cellulase-gold labeling on the root sections with double labeling with the anti-GFP antibody. The size of the gold particle on bacteria is 15 nm (Au-particle specific to ^{GFP}*E. coli*) and gold particles on the plant material are 10 nm (Au-particle specific to plant cellulose). (d), (e) and (f) are detail images of insets d, e and f.
doi:10.1371/journal.pone.0011915.g004




Figure 5. Arabidopsis genes involved in cell wall modification with differential expression at the time incubated with *E. coli* BI21 compared with control. Gene expression more than 3 fold changes were shown. (A) Glycosyl hydrolases and lyases (x, c, p are the symbol for xyloglucan endotransglycosylase, cellulase and pectinases/putative pectinases, respectively). (B) Expansins. (C) Cellulose synthases and cellulose synthase-like. (D) Extensins. (E) Arabinogalactan proteins.
doi:10.1371/journal.pone.0011915.g005

rinsed and grown hydroponically for 2 weeks. New leaves of ^{15}N -*E. coli*-incubated plants had a significantly higher concentration of ^{15}N than controls (Figure 6). Although this experiment does not provide unequivocal evidence that *E. coli* is digested inside root cells, it demonstrates that nitrogen derived from *E. coli* is assimilated by plants.

General considerations

Our experiments show that in the absence of pathogenic or symbiotic relationships, plants coordinated the entry of *E. coli* and yeast into root cells with an apparent expenditure of energy that is most likely justified by the benefit of using microbes as a nutrient source.

It appears that the pronounced plant responses to exposure to microbes, including induction of gene expression and remodeling of cell walls, is highly localized and strictly regulated to minimize the cost for the plant. It is possible that plant responses to non-pathogenic microbes are controlled at the cellular level, and evidence for this suggestion is provided by the patchiness of microbial uptake in the mature root zones (Figure 1A). We show that the presence of microbes induces the expression of plant enzymes with divergent functions, such as cellulases and cellulose synthases, and this suggests that the uptake process consist of a succession of distinct and tightly regulated processes, which would exclude the possibility of permanent induction of genes. In addition to minimizing energy expenditure, a transient and localized uptake process would also reduce opportunities for pathogens to invade the root.

Adding to the energetic costs of the uptake process, possible loss of turgor and cell contents could be associated with the entry of microbes into root cells. It is conceivable that one of the functions of the observed cell wall outgrowth limits loss of cell contents by preventing diffusion and leakage into the rhizosphere. Future research has to scrutinize the observed processes including each

step of incorporation and digestion, and how nutrients gains relate to energy expenditures and possibly loss of cell content.

Our discovery may explain the high diversity of root-inhabiting microbes of unknown ecological function [22,42] and brings a new dimension to current concepts of rhizosphere ecology. Much attention has focused on plant-growth-promoting bacteria for their potential to enhance plant growth [10,43]. Our discovery indicates the presence of a further category of plant-growth-promoting microbes which are used as a direct nutrient source. It is tempting to speculate that the microbe-enriched rhizosphere maintained by plants through exudation of photosynthates [16] is in part a 'microbe nursery' facilitating direct nutrients supply to plants.

Mixotrophy, the use of nutrients derived from photosynthesis and organic sources, is considered an exception in higher plants but characteristic of photosynthetic phytoplankton [44]. Our results indicate that mixotrophy may also occur in higher plants. This discovery has implications for carbon, nitrogen and phosphorus cycles in soils. High-production crop systems carry a strong pollution footprint which contributes to greenhouse gas emissions and pollutes ground and surface waters [45], and new approaches to supply soil-derived nutrients efficiently to plants are being sought. Exploiting the synergistic interactions between plants and microbes by harnessing soil microbes to supply crops with nutrients may be a further strategy.

Materials and Methods

Plasmid Construction

The *green fluorescent protein (GFP)* coding region was cloned as a glutathione S-transferase (GST) tagged recombinant gene. GFP was amplified from pDH51-GW-EGFP (GenBank: AM773753.1) using the following forward and reverse primers: 5'- GGC TCG AGA TGG TGA GCA AGG GCG AGG AG-3' and 5'- GGA AGC TTT CAC TTG TAC AGC TCG TCC ATG CC -3'. The PCR product was digested with XhoI and HindIII and cloned into pGEX-KG [46] designated pGSTGFP.

Preparation of *E. coli* Expressing GFP

E. coli strain BI21 (DE3) (Novagen) carrying the plasmid pGTF2 (TAKARA BIO INC) was transformed with pGSTGFP. Recombinant *E. coli* cells were selected on LB plates containing 200 $\mu\text{g ml}^{-1}$ ampicillin and 35 $\mu\text{g ml}^{-1}$ chloramphenicol. A single colony was used to inoculate a pre-culture containing 20 ml of LB supplemented with ampicillin (200 $\mu\text{g ml}^{-1}$) and chloramphenicol (35 $\mu\text{g ml}^{-1}$). The pre-culture was grown overnight at 37°C and used to inoculate 1 l of LB supplemented with ampicillin and chloramphenicol. *E. coli* was grown at 37°C to a cell density of 0.6 to 1 A_{600} units. Cells were cooled down on ice and 1 mM of Isopropyl-1-thio- β -D-galactopyranoside (IPTG) was added to induce expression of GFP. After incubation on a shaker (160 rpm) for 16–20 hours at 18°C, cells were harvested by centrifugation and washed twice with 1 l of 5 mM MES pH 5.8 (wash buffer) and resuspended in wash buffer. Cultures were used immediately.

Preparation of *Saccharomyces cerevisiae* Expressing GFP
GFP⁺ yeast clone *TDH3* (YGR192C) (Invitrogen, California, USA) expressing glyceraldehyde-3-phosphate dehydrogenase fused to




Figure 6. Incorporation of *E. coli*-derived ^{15}N by leaves of tomato plants. Roots of tomato grown in hydroponic culture were incubated with ^{15}N -*E. coli* for 1 h. After washing of the roots, plants were further grown for 2 weeks. Then 2–3 new leaves were analyzed for ^{15}N content. Control 1 are plants grown without ^{15}N -*E. coli*. Control 2 are plants incubated for 2 h with filtered ^{15}N -*E. coli* incubation solution. Results are depicted as mean \pm SD ($n=7$). Different letters indicate significant differences at $p<0.001$ (1-way ANOVA, Tukey's posthoc test).
doi:10.1371/journal.pone.0011915.g006

GFP was selected among the entire ^{GFP} yeast clone library for its high emission of green fluorescence (yeastgfp.yeastgenome.org). A single colony was used to inoculate 1 l of Yeast-extract Peptone Dextrose (YPD) liquid media and the culture was grown for 48 h at 28°C. Cells were harvested by centrifugation, washed twice with 1 l of wash buffer and re-suspended in wash buffer. Cultures were used immediately.

Plant Growth Conditions

Arabidopsis (*Arabidopsis thaliana* ecotype *Columbia* [Col-0]) plate culture: Seeds were germinated axenically on Petri dishes containing Murashige and Skoog (MS[47]) medium solidified by 3.2 g l⁻¹ of phytagel (Sigma). Plates were positioned vertically so that germinating radicals grow downward along the gel surface. Plants were grown for 2–3 weeks in a growth room with 16/8 h light/dark, 21°C, 150 μmol m⁻² s⁻¹ light intensity *Arabidopsis* axenic hydroponic culture: sterile seeds were sown in agar-filled 1.5 ml microcentrifuge tubes without cap and bottom. Microcentrifuge tubes were filled with 1.5 ml agar (0.68%) and tube bottoms cut off after agar had solidified, standing in a rack holder, and placed into sterile Combiness boxes (Microbox, Belgium) contained 300 ml half-strength MS medium. Adding 1 *Arabidopsis* seed into each tube, the boxes were incubated in a cold room for three days and then transferred to a growth cabinet (21°C, 16 h/8 h day/night, 150 μmol m⁻² s⁻¹). Plant roots grew from tubes into the solution. The boxes were aerated from day 11 after sowing by pumping air through a sterile filter (0.22 μm Millipore Filter, Ireland). Plants were grown for another 20 days and then in N-free MS medium for 3 days. Then 20 ml of ^{GFP}*E. coli* (OD_{600 nm} = 30) was added for 24 h. Plant incubated with 20 ml of wash buffer were used as a control. Plant were harvested, rinsed in deionized water, and immediately submersed in liquid N₂ and stored at -80°C.

Tomato (*Solanum lycopersicum*) vermiculite culture: seeds were germinated in soil for 10 days prior to being transferred into 200 ml pots containing vermiculite (one seedling per pot) in a growth room (16/8 h light/dark, 21°C, 150 μmol m⁻² s⁻¹). Pots were watered daily with tap water with addition of fertilizer (N-P-K: 15-15-15) once a week. Plants were grown for 2 to 3 weeks until shoot size was 10–15 cm.

Tomato hydroponic culture: 8–12 cm tall plants grown on vermiculite were carefully transferred into hydroponic culture consisting of 0.5 l water at pH 5.8 supplemented with 10 μM CaSO₄ (hydroponic solution) with 1 seedling per pot. The hydroponic cultures were continuously aerated and mixed by gentle stirring with a magnetic stirrer bar.

Uptake of *E. coli* and Yeast by Roots of *Arabidopsis* and Tomato

To assess uptake of *E. coli* and yeast by *Arabidopsis*, 5 ml of ^{GFP}*E. coli* or ^{GFP} yeast preparation (see above) at a cell density of 2 A₆₀₀ units was carefully added to roots of plants grown axenically on MS plates (see above) and incubated for 4 h horizontally at room temperature. Plants were carefully removed from the medium and roots washed with deionized water before being analyzed by confocal laser microscopy (CLSM, see details below). To assess uptake of *E. coli* and yeast by tomato in hydroponic cultures, plants were initially grown in hydroponic solution for 3 days to ensure the integrity of the roots. 20 ml of ^{GFP}*E. coli* or ^{GFP} yeast preparation at a cell density of 50 A₆₀₀ units was then added into the 500 ml hydroponic culture. After an overnight incubation at room temperature, roots were washed with deionized water and analyzed by CLSM.

For analysis of *Arabidopsis* and tomato root section by CLSM, visually assessed roots regions showing high fluorescence were excised (5–10 mm long), washed and embedded in 3% agarose. Hand-cut cross sections were transferred into curved slides, washed thoroughly with deionized water and analyzed by CLSM (see below). For analysis of tomato root sections by TEM, root regions showing high uptake by CLSM were coated with agarose before processing to ensure that bacteria external to roots were trapped in the agarose and not dislodged during cutting.

Time Course Experiment to Assess Status of ^{GFP} yeast in Tomato Roots

Ten tomato plants grown hydroponically for three days in hydroponic solution (see above) were incubated with ^{GFP} yeast overnight. Roots were carefully rinsed with deionized water and plants were placed in fresh hydroponic solution. The hydroponic solution was replaced every two days. Duplicate plants were removed from hydroponic culture at different time points and roots were treated with hydrogen peroxide (15%, 10 min) to sterilize the root surface. Roots from one plant were analyzed by CLSM and roots from the other were ground in liquid N₂ and analyzed for TDH3:GFP content by western blotting.

Western Blotting

Entire tomato roots were ground in liquid nitrogen and resuspended in 0.5 ml of 50 mM Tris (pH 7.5) supplemented with 0.1% Tween20. Non-soluble material was discarded by centrifugation at 14,000 rpm for 30 min. Total protein content of the extracts was determined as described by Bradford [48]. Equal amounts of protein sample was resolved by SDS-PAGE and characterized by western blot analysis using anti-GFP antibody (0.4 μL ml⁻¹, Roche) as primary antibody and Alexa Fluor 680 goat anti-mouse (Molecular Probes) as secondary antibody. Detection was performed with an Odyssey infrared imaging system (Li-COR, USA).

¹⁵N-Labeling of *E. coli*

¹⁵N-labeling of *E. coli* B121 cells was carried out as described by [49]. ¹⁵N-labeled *E. coli* cells (0.5 l) were harvested by centrifugation and washed four times with 0.5 l of deionized water. *E. coli* cells were then re-suspended in 1 l of water and used immediately for the incubation experiment.

Uptake of ¹⁵N-Labeled *E. coli* by Tomato

Twenty-one tomato plants (15 days old) were grown for three days in hydroponic solution (see above). Seven plants were incubated in 1 l of ¹⁵N-labeled *E. coli* solution for 1 h. After incubation, roots were gently rinsed with deionized water and plants were transferred to hydroponic solution. During this process, special care was given to avoid any contamination of the shoots by the bacterial solution. The remaining ¹⁵N-labeled *E. coli* incubation solution was centrifuged (3000 rpm, 15 min) and the supernatant sterilized by filtration (0.22 μm Millipore Filter, Ireland) to remove remaining *E. coli* cells. Seven plants were incubated for 2 h in the filtered supernatant (“control 2”). After incubation, roots were gently rinsed with sterile deionized water and plants were further grown in hydroponic solution. A further seven plants were grown in hydroponic culture without addition of *E. coli* (“control 1”). All plants were grown for a further 2 weeks with hydroponic solution changed daily. Subsequently, 2–3 new leaves of each plant were excised and dried at 60°C overnight, weighted and homogenized. The samples were analyzed for total nitrogen (N) and ¹⁵N content with continuous flow Isotope Ratio

Mass Spectrometer (IRMS, Stable Isotope Facility, University of California, Davis).

Microarray Analysis

Total RNA of Arabidopsis roots grown in hydroponic culture were extracted using NucleoSpin® Plant Kits (BD Biosciences Clontech, Japan). RNA of plants incubated with or without *E. coli* (control) were labelled with Cy3 or Cy5 fluorescent dye, mixed and used for subsequently hybridization onto 4x44K Agilent Arabidopsis GeneChip arrays (Agilent Technologies, USA). Labelling and hybridization of RNA, including scanning of the chips were performed by the Australian Genome Research Facility (AGRF, Victoria, Australia). Expression values (log10) for three biological replicates were extracted using robust multi-array analysis with perfect match correction and quantile normalization. Genes with ≥ 3 fold change were computed using one-way ANOVA ($p < 0.05$) with Partek Genomics suite.

Accession Numbers

The microarray hybridization data have been submitted to the National Center for Biotechnology Information (NCBI) Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo>) under accession number GSE22277.

Electron Microscopy

Roots of Arabidopsis and tomato incubated with *GFP* *E. coli* or *GFP* yeast were fixed in 4% paraformaldehyde in 0.1 M phosphate buffer pH 6.8 overnight at 4°C. After washing in 0.1 M phosphate buffer, roots were dehydrated through a graded ethanol series and infiltrated with LR White Resin and polymerized overnight at 50°C. Thin sections were cut with a Leica Ultracut UC6 ultramicrotome, picked up on carbon coated copper grids, stained with uranyl acetate and Reynold's lead citrate [50] and viewed in a JEOL 1010 transmission electron microscope operated at 80 kV and images were captured on a Olympus Soft Imaging Solutions Megaview III digital camera.

Gold Labeling

Thin sections were labeled using an anti-GFP antibody (Clontech, Mountain View, USA) as the primary antibody and a goat anti-mouse secondary labeled with 10 nm colloidal gold (British Biocell International, Cardiff, UK). Sections were also labeled with cellulase gold, made according to [51]. The cellulase was 1,4-(1,3;1,4)- β -D-Glucan 4-glucano-hydrolase from *Trichoderma reesei* (Sigma Aldrich, St Louis, USA). As a control, root sections were exposed to 2 mg ml⁻¹ cellulase for 16 h prior to labeling.

Cellulase Activity Analysis

26-days old hydroponically grown Arabidopsis were incubated with *E. coli* BI21 overnight. Plants not incubated with *E. coli* were used as negative control. Roots were rinsed twice in fresh medium and then transferred to fresh medium containing 31 μ g ml⁻¹ resorufin- β -D-cellobioside (Res-CB) (Marker Gene Technologies Inc., Eugene, OR, USA), a long-wavelength fluorescent substrate, which releases red fluorescent fluorophore resorufin upon cleavage. Roots were incubated for 2 h at room temperature, washed and inspected under CLSM.

References

- Baron C, Zambryski PC (1995) Notes from the underground: Highlights from plant-microbe interactions. Trends Biotechnol 13: 356–362.
- Boller T, He SY (2009) Innate immunity in plants: an arms race between pattern recognition receptors in plants and effectors in microbial pathogens. Science 324: 742–744.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ (2006) Host-microbe interactions: Shaping the evolution of the plant immune response. Cell 124: 803–814.
- Reinhold-Hurek B, Hurek T (1998) Life in grasses: diazotrophic endophytes. Trends Microbiol 6: 139–144.

Confocal Microscopy

A Zeiss LSM510 META (Carl Zeiss, Germany) confocal laser scanning microscope (CLSM) was used with 10x dry, 20x water immersion objectives, 40x and 60x oil immersion objectives. GFP and Res-CB were visualized by excitation with an argon laser at 488 nm and HeNe1 laser at 543 nm; detection with a 505–530 nm and 560–615 nm band-pass filter, respectively.

Supporting Information

Figure S1 Roots of Arabidopsis (A) and tomato (B) plant incubated with nano-silica fluorescent beads. No nano-beads were detected inside roots. Bar corresponds to 50 μ M.

Found at: doi:10.1371/journal.pone.0011915.s001 (5.72 MB TIF)

Figure S2 Roots of tomato plant incubated with yeast expressing GFP.

Found at: doi:10.1371/journal.pone.0011915.s002 (5.27 MB TIF)

Figure S3 Arabidopsis grown with or without *E. coli* BI21 incubation maintained a healthy phenotype.

Found at: doi:10.1371/journal.pone.0011915.s003 (7.36 MB TIF)

Figure S4 Arabidopsis genes involved in the cytoskeleton structure and re-organization with differential expression at the time incubated with *E. coli* BI21 compared with control.

Found at: doi:10.1371/journal.pone.0011915.s004 (9.60 MB TIF)

Movie S1 Presence of *E. coli* inside root hairs of Arabidopsis incubated with *E. coli* containing green fluorescent protein.

Found at: doi:10.1371/journal.pone.0011915.s005 (7.99 MB AVI)

Movie S2 Presence of yeast inside root hairs of Arabidopsis incubated with yeast containing green fluorescent protein.

Found at: doi:10.1371/journal.pone.0011915.s006 (7.99 MB AVI)

Movie S3 Presence of *E. coli* inside roots of tomato incubated with *E. coli* containing green fluorescent protein.

Found at: doi:10.1371/journal.pone.0011915.s007 (5.59 MB AVI)

Movie S4 Tomato root transverse sections show *GFP* *E. coli* in the apoplast and inside the root cells.

Found at: doi:10.1371/journal.pone.0011915.s008 (9.59 MB AVI)

Movie S5 Tomato root transverse sections show *GFP* yeast in the apoplast and inside the root cells.

Found at: doi:10.1371/journal.pone.0011915.s009 (6.43 MB AVI)

Acknowledgments

We thank Kelly Hanson and Yoganand Sundaravadanam (for help with microarray data analysis), Anthony Young (for helpful discussion) and Sabrina Latansio-Aidar (for technical assistance). We are grateful to the ARC Centre of Excellence for Integrative Legume Research for access to research facilities.

Author Contributions

Conceived and designed the experiments: CPL DR SR RIW ES TN SS TL. Performed the experiments: CPL RIW ES TL. Analyzed the data: CPL DR SR RIW TN SS TL. Wrote the paper: CPL DR SR RIW TN SS TL.

5. Triplett EW (1996) Diazotrophic endophytes: Progress and prospects for nitrogen fixation in monocots. *Plant Soil* 186: 29–38.
6. Ryu CM, Farag MA, Hu CH, Reddy MS, Wei HX, et al. (2003) Bacterial volatiles promote growth in Arabidopsis. *Proc Natl Acad Sci U S A* 100: 4927–4932.
7. Timmusk S, Wagner EGH (1999) The plant-growth-promoting rhizobacterium *Paenibacillus polymyxa* induces changes in Arabidopsis thaliana gene expression: A possible connection between biotic and abiotic stress responses. *Mol Plant-Microbe Interact* 12: 951–959.
8. Kuklinsky-Sobral J, Araujo WL, Mendes R, Geraldi IO, Pizzirani-Kleiner AA, et al. (2004) Isolation and characterization of soybean-associated bacteria and their potential for plant growth promotion. *Environ Microbiol* 6: 1244–1251.
9. Bajwa R, Abuarghub S, Read DJ (1985) The Biology of Mycorrhiza in the Ericaceae.10. The utilization of proteins and the production of proteolytic-enzymes by the mycorrhizal endophyte and by mycorrhizal plants. *New Phytol* 101: 469–486.
10. Lugtenberg B, Kamilova F (2009) Plant-Growth-Promoting Rhizobacteria. *Annu Rev Microbiol* 63: 541–556.
11. Bertin C, Yang X, Weston LA (2003) The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil* 256: 67–83.
12. Jones DL, Edwards AC, Donachie K, Darrah PR (1994) Role of Proteinaceous Amino-Acids Released in Root Exudates in Nutrient Acquisition from the Rhizosphere. *Plant Soil* 158: 183–192.
13. Grayston SJ, Vaughan D, Jones D (1996) Rhizosphere carbon flow in trees, in comparison with annual plants: The importance of root exudation and its impact on microbial activity and nutrient availability. *Appl Soil Ecol* 5: 29–56.
14. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM (2006) The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol* 57: 233–266.
15. Marschner H (1995) Mineral Nutrition of Higher Plants, 2nd edn. London: Academic Press, Ltd. 889 p.
16. Bonkowski M (2004) Protozoa and plant growth: the microbial loop in soil revisited. *New Phytol* 162: 617–631.
17. Morgan JA (2002) Looking beneath the surface. *Science* 298: 1903–1904.
18. Clarholm M (1985) Interactions of Bacteria, Protozoa and Plants Leading to Mineralization of Soil-Nitrogen. *Soil Biol Biochem* 17: 181–187.
19. Näsholm T, Kielland K, Ganeteg U (2009) Uptake of organic nitrogen by plants. *New Phytol* 182: 31–48.
20. Paungfoo-Lonhienne C, Lonhienne TGA, Mudge SR, Schenk PM, Christie M, et al. (2010) DNA is taken up by root hairs and pollen, and stimulates root and pollen tube growth. *Plant Physiol* 153: 799–805.
21. Paungfoo-Lonhienne C, Lonhienne TGA, Rentsch D, Robinson N, Christie M, et al. (2008) Plants can use protein as a nitrogen source without assistance from other organisms. *Proc Natl Acad Sci U S A* 105: 4524–4529.
22. Vandenkoornhuyse P, Mahe S, Ineson P, Staddon P, Ostle N, et al. (2007) Active root-inhabiting microbes identified by rapid incorporation of plant-derived carbon into RNA. *Proc Natl Acad Sci U S A* 104: 16970–16975.
23. Schikora A, Carreri A, Charpentier E, Hirt H (2008) The Dark Side of the Salad: Salmonella typhimurium Overcomes the Innate Immune Response of Arabidopsis thaliana and Shows an Endopathogenic Lifestyle. *Plos One* 3: e2279. doi:2210.1371/journal.pone.0002279.
24. Hematy K, Cherk C, Somerville S (2009) Host-pathogen warfare at the plant cell wall. *Curr Opin Plant Biol* 12: 406–413.
25. English PD, Jurale JB, Albersheim P (1971) Host-Pathogen Interactions: II. Parameters Affecting Polysaccharide-degrading Enzyme Secretion by Colletotrichum lindemuthianum Grown in Culture. *Plant Physiol* 47: 1–6.
26. Robledo M, Jimenez-Zurdo JI, Velazquez E, Trujillo ME, Zurdo-Pineiro JL, et al. (2008) Rhizobium cellulase CelC2 is essential for primary symbiotic infection of legume host roots. *Proc Natl Acad Sci U S A* 105: 7064–7069.
27. Coleman DJ, Studler MJ, Naleway JJ (2007) A long-wavelength fluorescent substrate for continuous fluorometric determination of cellulase activity: resorufin-[beta]-d-cellobioside. *Anal Biochem* 371: 146–153.
28. Bacon CW, Hinton DM (2007) Bacterial Endophytes: The Endophytic Niche, Its Occupants, and Its Utility. In: Gnanamanickam SS, ed. Plant-Associated Bacteria Netherlands: Springer. pp 155–194.
29. Puente ME, Holguin G, Glick BR, Bashan Y (1999) Root-surface colonization of black mangrove seedlings by *Azospirillum halopraeferens* and *Azospirillum brasilense* in seawater. *FEMS Microbiol Ecol* 29: 283–292.
30. Matthysse AG (1983) Role of bacterial cellulose fibrils in Agrobacterium tumefaciens infection. *J Bacteriol* 154: 906–915.
31. Richmond TA, Somerville CR (2000) The Cellulose Synthase Superfamily. *Plant Physiol* 124: 495–498.
32. Zhu Y, Nam J, Carpita NC, Matthysse AG, Gelvin SB (2003) Agrobacterium-mediated root transformation is inhibited by mutation of an Arabidopsis cellulose synthase-like gene. *Plant Physiol* 133: 1000–1010.
33. Cannon MC, Terneus K, Hall Q, Tan L, Wang Y, et al. (2008) Self-assembly of the plant cell wall requires an extensin scaffold. *Proc Natl Acad Sci U S A* 105: 2226–2231.
34. McCurdy DW, Patrick JW, Offler CE (2008) Wall ingrowth formation in transfer cells: novel examples of localized wall deposition in plant cells. *Curr Opin Plant Biol* 11: 653–661.
35. Gaspar YM, Nam J, Schultz CJ, Lee LY, Gilson PR, et al. (2004) Characterization of the Arabidopsis lysine-rich arabinogalactan-protein AtAGP17 mutant (rat1) that results in a decreased efficiency of agrobacterium transformation. *Plant Physiol* 135: 2162–2171.
36. Mendgen K, Deising H (1993) Tansley Review No 48 Infection Structures of Fungal Plant-Pathogens - a Cytological and Physiological Evaluation. *New Phytol* 124: 193–213.
37. Ferguson BJ, Indrasumunar A, Hayashi S, Lin M-H, Lin Y-H, et al. (2010) Molecular Analysis of Legume Nodule Development and Autoregulation. *J Integr Plant Biol* 52: 61–76.
38. James EK, Gyaneshwar P, Mathan N, Barraquio QL, Reddy PM, et al. (2002) Infection and colonization of rice seedlings by the plant growth-promoting bacterium Herbaspirillum seropedicae Z67. *Mol Plant-Microbe Interact* 15: 894–906.
39. Rothballer M, Eckert B, Schmid M, Fekete A, Schlöter M, et al. (2008) Endophytic root colonization of gramineous plants by Herbaspirillum frisingense. *FEMS Microbiol Ecol* 66: 85–95.
40. Kovtunovych G, Lar O, Kamalova S, Kordyum V, Kleiner D, et al. (1999) Correlation between pectate lyase activity and ability of diazotrophic Klebsiella oxytoca VN 13 to penetrate into plant tissues. *Plant Soil* 215: 1–6.
41. Reinhold-Hurek B, Krause A, Leyser B, Mische L, Hurek T (2007) The rice apoplast as a habitat for endophytic N₂-fixing bacteria. In: Sattelmacher B, Horst W, eds. The Apoplast of Higher Plants: Compartment of Storage, Transport, and Reactions. , Netherlands: Springer. pp 427–443.
42. Vandenkoornhuyse P, Baldauf SL, Leyval C, Straczek J, Young JJPW (2002) Extensive fungal diversity in plant roots. *Science* 295: 2051–2051.
43. Yang J, Kloepper JW, Ryu CM (2009) Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci* 14: 1–4.
44. Raven JA, Beardall J, Flynn KJ, Maberly SC (2009) Phagotrophy in the origins of photosynthesis in eukaryotes and as a complementary mode of nutrition in phototrophs: relation to Darwin's insectivorous plants. *J Exp Bot* 60: 3975–3987.
45. Gruber N, Galloway JN (2008) An Earth-system perspective of the global nitrogen cycle. *Nature* 451: 293–296.
46. Hakes DJ, Dixon JE (1992) New vectors for high level expression of recombinant proteins in bacteria. *Anal Biochem* 202: 293–298.
47. Murashige T, Skoog F (1962) A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiol Plant* 15: 473–497.
48. Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72: 248–254.
49. Marley J, Lu M, Bracken C (2001) A method for efficient isotopic labeling of recombinant proteins. *J Biomol NMR* 20: 71–75.
50. Reynolds ES (1963) The use of lead citrate at high pH as an electron-opaque stain for electron microscopy. *J Cell Biol* 17: 208–212.
51. Berg RH, Erdos GW, Gritzali M, Brown RD (1988) Enzyme-gold affinity labelling of cellulase. *J Electron Microsc Tech* 8: 371–379.