Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

ArticleTitle	Recent anthropogenic impact in ancient Lake Ohrid (Macedonia/Albania): a palaeolimnological approach	
Article Sub-Title		
Article CopyRight	Springer Science+Business Media Dordrecht (This will be the copyright line in the final PDF)	
Journal Name	Journal of Paleolimnology	
Corresponding Author	Family Name	Lorenschat
	Particle	
	Given Name	Julia
	Suffix	
	Division	
	Organization	
	Address	Schunterstraße 56, 38106, Brunswick, Germany
	Email	j.lorenschat@t-online.de
Author	Family Name	Zhang
	Particle	
	Given Name	Xiaosen
	Suffix	
	Division	Department of Geography, Environment and Earth Sciences
	Organization	University of Hull
	Address	Cottingham Road, HU6 7RX, Hull, UK
	Email	xiaosen.zhang@2011.hull.ac.uk
Author	Family Name	Anselmetti
	Particle	
	Given Name	Flavio S.
	Suffix	
	Division	Institute of Geological Sciences and Oeschger Centre for Climate Change Research
	Organization	University of Bern
	Address	Baltzerstrasse 1+3, 3012, Bern, Switzerland
	Email	flavio.anselmetti@geo.unibe.ch
Author	Family Name	Reed
	Particle	
	Given Name	Jane M.
	Suffix	
	Division	Department of Geography, Environment and Earth Sciences
	Organization	University of Hull
	Address	Cottingham Road, HU6 7RX, Hull, UK
	Email	j.m.reed@hull.ac.uk
Author	Family Name	Wessels
	Particle	
	Given Name	Martin

	Suffix	
	Division	Institut für Seenforschung
	Organization	LUBW
	Address	Argenweg 50/1, 88045, Langenargen, Germany
	Email	Martin.Wessels@lubw.bwl.de
Author	Family Name	Schwalb
	Particle	
	Given Name	Antje
	Suffix	
	Division	Institut für Geosysteme und Bioindikation
	Organization	Technische Universität Braunschweig
	Address	Langer Kamp 19 c, 38106, Brunswick, Germany
	Email	antje.schwalb@tu-bs.de
Schedule	Received	19 September 2013
	Revised	
	Accepted	27 June 2014
Abstract	<p>Ancient lakes, which are important centres of biodiversity and endemism, are threatened by a wide variety of human impacts. To assess environmental impact on ancient Lake Ohrid we have taken short sediment cores from two contrasting site locations, comprising a site of urban pollution and an apparently pristine area. Recent impacts on water quality and ecology were assessed using sediment, geochemical, ostracode, and diatom data derived from analysis of two ^{210}Pb-dated sediment cores spanning the period from 1918 to 2009. According to the index of geoaccumulation, sediments were often moderately contaminated with As, Fe and Ni concentrations often exceeded reported maximum limits above which harmful effects on sediment-dwelling organisms are expected. Productivity in the (pristine) south-eastern part of Lake Ohrid (Sveti Naum) is generally lower than in the north, probably due to the strong influence of spring discharge. Low ostracode and diatom concentrations, low abundance of the epilimnetic diatom <i>Cyclotella ocellata</i>, and low values of TOC and TIC indicate a lower productivity from the early 1920s to the late 1980s. Since the mid 1970s, increased relative abundance of <i>C. ocellata</i> and increasing diatom concentration indicate increasing productivity in the south-eastern part. Rising numbers of ostracode valves and higher TIC and TOC contents in both sediment cores indicate an increase in productivity during the late 1980s. A slight increase in productivity near Sveti Naum continued from the early 1990s until 2009, witnessed by rising TC, TIC, and TOC content and a generally high number of ostracode valves and ostracode diversity. The area near the City of Struga (site of urban pollution) is also characterized by rising TOC and TIC contents and, furthermore, by increasing Cu, Fe, Pb, and Zn concentrations since the early 1990s. The recent reduction in the number of ostracode valves and ostracode diversity is probably caused by a higher heavy metal load into the lake. This suggests that living conditions for the endemic species in Lake Ohrid have become less favourable in the northern part of the lake, which might threaten the unique flora and fauna of Lake Ohrid.</p>	
Keywords (separated by '-')	Lake Ohrid - Palaeolimnology - Eutrophication - Geochemistry - Ostracodes - Diatoms	
Footnote Information	Electronic supplementary material The online version of this article (doi:10.1007/s10933-014-9783-5) contains supplementary material, which is available to authorized users.	

Metadata of the article that will be visualized in OnlineAlone

Electronic supplementary
material

Below is the link to the electronic supplementary material. Supplementary material 1 (DOC 31 kb)Supplementary material 2 (DOC 35 kb)

Recent anthropogenic impact in ancient Lake Ohrid (Macedonia/Albania): a palaeolimnological approach

Julia Lorenschat · Xiaosen Zhang · Flavio S. Anselmetti ·
Jane M. Reed · Martin Wessels · Antje Schwalb

Received: 19 September 2013 / Accepted: 27 June 2014
© Springer Science+Business Media Dordrecht 2014

Abstract Ancient lakes, which are important centres of biodiversity and endemism, are threatened by a wide variety of human impacts. To assess environmental impact on ancient Lake Ohrid we have taken short sediment cores from two contrasting site locations, comprising a site of urban pollution and an apparently pristine area. Recent impacts on water quality and ecology were assessed using sediment, geochemical, ostracode, and diatom data derived from analysis of two ^{210}Pb -dated sediment cores spanning the period from 1918 to 2009. According to the index of geoaccumulation, sediments were often moderately

contaminated with As. Fe and Ni concentrations often exceeded reported maximum limits above which harmful effects on sediment-dwelling organisms are expected. Productivity in the (pristine) south-eastern part of Lake Ohrid (Sveti Naum) is generally lower than in the north, probably due to the strong influence of spring discharge. Low ostracode and diatom concentrations, low abundance of the epilimnetic diatom *Cyclotella ocellata*, and low values of TOC and TIC indicate a lower productivity from the early 1920s to the late 1980s. Since the mid 1970s, increased relative abundance of *C. ocellata* and increasing diatom concentration indicate increasing productivity in the south-eastern part. Rising numbers of ostracode valves and higher TIC and TOC contents in both sediment cores indicate an increase in productivity

Electronic supplementary material The online version of this article (doi:10.1007/s10933-014-9783-5) contains supplementary material, which is available to authorized users.

J. Lorenschat (✉)
Schunterstraße 56, 38106 Brunswick, Germany
e-mail: j.lorenschat@t-online.de

X. Zhang · J. M. Reed
Department of Geography, Environment and Earth
Sciences, University of Hull, Cottingham Road,
Hull HU6 7RX, UK
e-mail: xiaosen.zhang@2011.hull.ac.uk

J. M. Reed
e-mail: j.m.reed@hull.ac.uk

F. S. Anselmetti
Institute of Geological Sciences and Oeschger Centre for
Climate Change Research, University of Bern,
Baltzerstrasse 1+3, 3012 Bern, Switzerland
e-mail: flavio.anselmetti@geo.unibe.ch

M. Wessels
Institut für Seenforschung, LUBW, Argenweg 50/1,
88045 Langenargen, Germany
e-mail: Martin.Wessels@lubw.bwl.de

A. Schwalb
Institut für Geosysteme und Bioindikation, Technische
Universität Braunschweig, Langer Kamp 19 c,
38106 Brunswick, Germany
e-mail: antje.schwalb@tu-bs.de

during the late 1980s. A slight increase in productivity near Sveti Naum continued from the early 1990s until 2009, witnessed by rising TC, TIC, and TOC content and a generally high number of ostracode valves and ostracode diversity. The area near the City of Struga (site of urban pollution) is also characterized by rising TOC and TIC contents and, furthermore, by increasing Cu, Fe, Pb, and Zn concentrations since the early 1990s. The recent reduction in the number of ostracode valves and ostracode diversity is probably caused by a higher heavy metal load into the lake. This suggests that living conditions for the endemic species in Lake Ohrid have become less favourable in the northern part of the lake, which might threaten the unique flora and fauna of Lake Ohrid.

Keywords Lake Ohrid · Palaeolimnology · Eutrophication · Geochemistry · Ostracodes · Diatoms

Introduction

Lakes respond chemically and biologically to human impact. Commonly used proxies, such as ostracodes, diatoms, and geochemical parameters, have been used effectively to reconstruct anthropogenic influence through time on lakes from analysis of lake sediment cores (Reed et al. 2008; Pérez et al. 2010). Aquatic ecosystems such as lakes (Löffler et al. 1998; Matzinger et al. 2006a; Patceva et al. 2006) and rivers (Patceva et al. 2004; Veljanoska-Sarafiloska et al. 2004; Bilali et al. 2012) in Macedonia and Albania are under increasing human impact and this also applies to some ancient lakes in the world. Lakes Baikal, Biwa, and Tanganyika are examples. The lakes are influenced by lake level changes (mainly due to irrigation) and particularly the littoral areas are affected by sediment loading which leads to a disturbance of microhabitats and, as a result, to a drop in the number of animal and plant species (Cohen et al. 1999; Alin et al. 1999; Asaeda and Shinohara 2012; Touchart 2012). However, so far there is no evidence that a tipping point is imminent. The biodiversity hotspot of deep, ancient Lake Ohrid may equally be threatened (Matzinger 2006b). Recently, concern has been raised related to a “creeping biodiversity crisis” in Lake Ohrid (Kostoski et al. 2010), which poses a serious threat to the endemic species (Albrecht and Wilke

2008) whose extinction would cause an irreversible loss. To date the potential of palaeolimnological techniques to assess the influence of accelerated human impact on the ecology of the lake has not been explored.

The town of Ohrid is one of the oldest human settlements in Europe (UNESCO ROSTE 2004), and the shores of the adjacent lake have been inhabited since prehistoric times. Archaeological investigations have documented settlements from as early as 6,000 BC (Ministry of Environment and Physical Planning, undated). The first evidence of settled human communities and domesticated animals at about 8.5 ka BP is indicated by the presence of coprostanol, a biomarker for human and animal faeces, in a sediment core taken in Lake Ohrid (Holtvoeth et al. 2010). Wagner et al. (2009) identified the onset of human impact on catchment vegetation at about 5,000 BP and a distinct increase at 2,400 BP. After the end of World War II the population increased 5–6 times. Today, 106,000 people live in the Macedonian part of the watershed, about 61,000 residents in the Albanian part, and about 25,600 residents in the Greek part (Avramoski et al. 2003). Agriculture is one of the most important economic sectors in the region (Spirkovski et al. 2001), and run-off from cultivated land and pastures is an important source of total phosphorus (TP) input into Lake Ohrid (Spirkovski et al. 2001). Besides agriculture, households are the main anthropogenic source of phosphorus (Matzinger et al. 2004). Avramoski et al. (2003) and Matzinger et al. (2004) documented that phosphorus concentration has increased at least fourfold over the past 100 years and Matzinger et al. (2004) found an increase in sediment carbonate content over the last 50 years which is indicative of the early stages of eutrophication. To date, the TP concentration in the centre of Lake Ohrid is still low enough to consider the lake as “oligotrophic”, but there are major concerns over water quality in the littoral zone. Veljanoska-Sarafiloska et al. (2004) showed that certain areas of the shoreline are in an alarming condition, in particular where rivers enter the lake, and suggested that much of the littoral zone was mesotrophic. The River Velgoska, for example, flows through industrial zones, is exposed to sources of untreated sewage, and is classed as eutrophic. The mesotrophic River Koselska flows through rural and agricultural areas and during heavy rains sometimes receives overflow sewage water from

the sewage system. The River Sateska, diverted into Lake Ohrid in 1962, flows through agricultural and urban areas and carries a high load of sediment, drainage water, and communal wastewater which is deposited in the littoral zone. From evidence for a switch to more organic sediment character in the littoral zone, Matter et al. (2010) estimated that major impact in the shallow-water zone had persisted since ca. 1955. Other pollution sources are metal component factories in Pogradec, which discharge untreated waste into the lake, and old mines, north-west of Pogradec (Avramoski et al. 2003). The two chromium mines, three nickel–iron mines and one coal mine went out of use at the turn of the century, but many piles of waste material remain and are a permanent pollution source (Spirkovski et al. 2001). To improve the water quality of Lake Ohrid, major improvements to the sewage treatment system have been carried out recently. Since June 1988 the Regional Sewerage System for the Protection of Lake Ohrid collects wastewater from about 65 % of the Ohrid-Struga region. After treatment, the water is discharged into the River Crni Drim. Two additional construction phases should allow treatment of most of the shoreline on the Macedonian part of the lake (UNESCO ROSTE 2004), although several households in the City of Ohrid and nearby settlements are still not connected to any sewage system (Lokoska 2012). In Pogradec, three wastewater treatment plants have been opened in the last 5 years, but some unconnected areas remain (Neugebauer and Vallerien 2012).

The focus of this study is to explore past impacts on Lake Ohrid caused by anthropogenic pollution using selected proxies comprising ostracodes and diatoms, representing both water column and lake-bottom conditions, as well as geochemical parameters. To achieve the aim, we used ^{210}Pb and ^{137}Cs dated sediment cores taken from localities with contrasting degrees of human impact.

Site description

Lake Ohrid (Fig. 1) straddles the border between Macedonia and Albania and is located at 695 m a.s.l. It has a surface area of 358.2 km² (230 km² belongs to Macedonia and 128.2 km² to Albania). The length of the shoreline is 87.5 km, the maximum length of the lake is 30.8 km, and its maximum width is 14.8 km.

The lake has a maximum depth of 289 m and an average depth of 164 m. The total watershed incorporates its sister lake, Prespa, and covers an area of 2,340 km² (Dodeva 2012) extending into Greece (Watzin 2003). Lake Ohrid is directly connected with Lake Prespa via underground karstic channels and these springs contribute ~53 % to Ohrid's inflow. Only a small proportion of the inflow originates from rivers (~23 %) and direct precipitation (~23 %) (Albrecht and Wilke 2008). The main tributaries are the rivers Velgoska (mean annual inflow 0.4 m³ s⁻¹), Sateska (5.5 m³ s⁻¹), Koselska (1.3 m³ s⁻¹), and Čerava (0.2 m³ s⁻¹) (Patceva et al. 2004; Matzinger et al. 2007). The only outlet is the River Crni Drim (Dodeva 2012). Lake Ohrid is a Quaternary graben-shaped lake formed by a combination of post-Pliocene uplift and gradual subsidence (Aliaj et al. 2001). West of the lake, the landscape is characterized by the “Mokra” mountain chain, which reaches ~1,500 m a.s.l. and in the east, by the “Galičica” mountain chain (1,750 m a.s.l.) (Wagner et al. 2009). The Mokra is composed of serpentine (peridotites) overlain by Triassic limestone and the Galičica consists mainly of Triassic limestone (Stanković 1960). The catchment of Lake Ohrid is characterized by continental climate. Between 1961 and 1990, average annual air temperature was 11.1 °C in the City of Ohrid. The maximum air temperature was 31.5 °C, the minimum –5.7 °C, and the lake never freezes (Popovska and Bonacci 2007). Maximum precipitation occurs in December and March, and the late summer is dry (Salemaa 1994). Mean annual precipitation averages ~750 mm (Wagner et al. 2009).

Materials and methods

Sediment cores were collected in September 2009 from 50 m water depth in Lake Ohrid (Fig. 1). The sampling depth was chosen because Mikulić and Pljakić (1970) reported maximum candonid ostracode diversity at this depth. The northern sampling location offshore from the City of Struga (core St09) (41°09.411'N, 20°40.986'E) represented a site of high urban pollution, being the largest town on the Macedonian shoreline of Lake Ohrid (63,376 residents in 2002) (GeoHive). The south-eastern area near the springs of Sveti Naum represented a relatively pristine location (core Sv09) (40°55.760'N, 20°45.175'E),

Fig. 1 Location of Lake Ohrid (square) on the border of Macedonia/Albania (a) and a bathymetric map of Lake Ohrid showing coring locations and cities and rivers discussed in the text (b)

with low intensity tourism and scattered domestic dwellings. At each location, three parallel cores, with a diameter of 11 cm, were retrieved 36 cm apart with a gravity multicorer. One core per location was sub-sampled for ^{210}Pb and ^{137}Cs dating in the field. The top 15 cm were subsampled every 0.5 cm and below

15 cm down to the base of the core every 1 cm. The cores taken for ostracode, diatom, and geochemical analyses were sampled in the field every 1 cm throughout. Cores for sediment description and photography were split in two halves at the Institut für Seenforschung, Langenargen.

 Springer

Fig. 2 ^{137}Cs and total and unsupported ^{210}Pb (Bq kg^{-1}) concentrations plotted on log scale in sediment cores Sv09 and St09 and the respective CFCS model with 1955, 1963, and 1986 fallout maxima determined by ^{137}Cs

of the species *Cyclotella radiosa* (Grunow in Van Heurck) Lemmermann 1900, the genus name for which should revert to *Cyclotella* rather than *Puncticulata* (Houk et al. 2010). The F index of the endemic *Cyclotella fottii* Hustedt in Huber-Pestalozzi 1942 was estimated based on the ratio of pristine valves to all valves (sum of pristine and partially dissolved valves), where $F = 1$ implies valves preserved well while $F = 0$ shows valves are appreciably dissolved (Ryves et al. 2001). Biostratigraphic zone boundaries were defined using constrained incremental sum of squares cluster analysis (CONISS; Grimm 1987).

Results

Chronology

^{137}Cs peaks (1955, 1963, and 1986) were first identified independently and then compared with results from sedimentation rates based on the ^{210}Pb data so that the three marker ages could be assigned to the ^{137}Cs curve. For both cores, the differences of these ages to the averaged CFCS age line (constant sedimentation rate) are minimal (Fig. 2) so a linear age-depth model based on the ^{210}Pb data was appropriate.

The total ^{210}Pb activities in core St09 (Fig. 2) ranged between 155 Bq kg^{-1} (2.25 cm) and 26 Bq kg^{-1} (39.50 cm). Unsupported ^{210}Pb activity was highest at 10.25 cm (131 Bq kg^{-1}) and minimum activity (6 Bq kg^{-1}) was found at a depth of 27.50 cm. Using the CFCS ^{210}Pb model, an average sedimentation rate of $0.40 \text{ cm year}^{-1}$ has been determined. Maximum ^{137}Cs activities were 220 and 97 Bq kg^{-1} at 12.25 and 21.50 cm, respectively, and correspond to the Chernobyl peak from 1986 and the nuclear weapons testing ^{137}Cs maximum in 1963. The onset of ^{137}Cs activities around the year 1955 was identified at 30.5 cm. According to the CFCS model, the total age of the sediment core is ~ 80 years (~ 1928).

In core Sv09, total ^{210}Pb activity was highest at the top of core (174 Bq kg^{-1}) and declined relatively evenly down to the base of the core, with a minimum at 35.50 cm (33 Bq kg^{-1}) (Fig. 2). Unsupported ^{210}Pb activities ranged from 138 Bq kg^{-1} (0.25 cm) to 6 Bq kg^{-1} (20.50 cm). Using the CFCS ^{210}Pb model, an average sedimentation rate of $0.47 \text{ cm year}^{-1}$ was

determined. ^{137}Cs activities in core Sv09 failed to display a sharp peak that might identify the onset of ^{137}Cs production in 1955 and the maximum fallout of 1963, nevertheless, the ^{137}Cs maximum of 676 Bq kg^{-1} at 14.75 cm indicates the 1986 Chernobyl peak. According to the CFCS model, the base of Sv09 is dated to ~ 1918 .

A reason for the difference in absolute values of ^{137}Cs and ^{210}Pb activities in cores Sv09 and St09 could be the different lithologies: St09 has a higher carbonate content than Sv09, which mostly consists of siliciclastics. That could result in different affinities of the sediment to take up the radionuclides and a varying degree of reworking.

Sedimentology and geochemistry

Sediments from core St09 (Fig. 3) were relatively homogenous with a dark greyish brown colour. From the base of the core to 37.5 cm, sandy silt occurred, which was overlaid by clayey silt. Organic matter was low and fluctuated between 3.5 and 6.4 %. Carbonate content was higher from the core base to 7 cm depth with only slightly varying content (minimum of 18.5 % at 35 cm; maximum 23.2 %). Above, the content decreased to 12.8 %, rose again to 16.7 % at 2 cm. The water content was lowest (44.5 %) at the core base and increased towards the top (61.0 %). Ni and Zr decreased slightly upcore and fluctuated irregularly (Fig. 4). These fluctuations were also shown in the concentrations of As, Cu, Fe, Hg, Pb, and Zn but these elements show a slight increased upcore trend. C/N ratios increased to the core top and fluctuated to a greater or lesser extent. The maximum Hg concentration (0.08 mg kg^{-1}) was measured close to the base of the core between 48 and 49 cm. According to the Index of Geoaccumulation (I_{geo}) (Müller 1986) this corresponds to the pollution class, “moderately contaminated”. However, this sample is a single peak with a value much higher than the rest of the St09 sequence, and may be an outlier. Arsenic concentrations correspond in 17 samples to the pollution class “moderately contaminated” and in one sample (29–28 cm) to the pollution class “moderately to strongly contaminated” (34.74 mg kg^{-1}). The probable effect concentrations (PECs) of Ni (48.6 mg kg^{-1}) (MacDonald et al. 2000) were exceeded in a total of 26 samples, mostly in the upper part of the core, and As concentration exceeded the

PEC (33.0 mg kg^{-1}) (MacDonald et al. 2000) between 1957 and 1959 AD (29–28 cm) (34.74 mg kg^{-1}).

From the base of core Sv09 to 22.5 cm, the sediment consisted of silty clay with an upcore decreasing clay content (Fig. 3). Between 22.5 and 17.5 cm, a sand–silt–clay unit occurred that was overlaid by silty clay up to 12.5 cm. The uppermost 12.5 cm were characterized by clayey silt. The sediment colour was olive brown at the base of the core, dark greyish brown above 33 cm, and brown in the uppermost 12.5 cm. Organic matter and carbonate content were generally low and fluctuating. The maximum organic content (7.0 %) occurred at 10 cm depth and the minimum (1.6 %) at 21 cm. Maximum carbonate content (7.5 %) was measured at 7 cm and minimum (1 %) at 25 cm depth. Between the base of the core and 13 cm, water content fluctuated between 27.0 and 33.2 %. Above, the content increased with some fluctuations to 43.8 % at the core top. As and Hg show an increasing trend over time in Sv09 (Fig. 5), and concentrations of Cu, Fe, Ni, Zn, and Zr fluctuated irregularly throughout the core. Pb is the only element in Sv09 which shows an upcore decrease. The C/N ratios fluctuated throughout the sediment profile and vary between 9.90 and 17.70. According to the I_{geo} , As concentrations in core Sv09 correspond in nine samples to the pollution class “moderately contaminated”, mostly in the upper core sequence. Fe concentrations exceed the PEC (43.77 g kg^{-1}) (Jaagumagi 1993) in three samples [1969–1972 AD (58.04 g kg^{-1}), 1986–1989 AD (47.81 g kg^{-1}), and 2003–2005 AD cm (54.32 g kg^{-1})] and Ni concentrations exceed the PEC (48.6 mg kg^{-1}) (MacDonald et al. 2000) in all samples.

TIC and TOC contents in both cores were similar to the LOI values and show matching patterns (Figs. 4, 5). TIC content in St09 fluctuated between 4.29 % (6–5 cm) and 6.71 % (32–31 cm), TOC between 0.66 % (32–31 cm) and 2.16 % (4–3 cm). Lowest TIC (0.19 %) and TOC (0.59 %) contents in core Sv09 occurred at a depth of 22–21 cm. Highest TOC (1.74 %) and TIC values (1.36 %) occurred between 1–0 and 3–2 cm, respectively.

Ostracodes

A total of 19 ostracode species was found in core St09 (Fig. 6; ESM 1), with a relatively high number of

Fig. 3 Core photographs, organic matter, carbonate content, and siliciclastics in cores Sv09 and St09

juvenile candonids. Dominant species are *Candona media* Klie 1939 (up to 54 %) and *Cypria lacustris* Sars 1890 (up to 43 %). The Shannon index and the Evenness do not show any distinct patterns. The highest Shannon (1.96) occurred in 16–15 cm, the lowest (0.75) in 14–13 cm. Evenness ranged between 0.19 in 23–22 cm and 0.71 in 16–15 cm. The Bray–Curtis dissimilarity shows the highest value in 36–35 cm (0.63) and the lowest in 12–11 cm (0.08). The sample from 6 to 5 cm is, with a Jaccard similarity of 0.86, most similar to the core top sample. The lowest similarity occurred in 37–36 cm (0.25). Cluster analysis yielded four major zones in core St09: In Zone O-I (49–36 cm, 1922–1945 AD) 14 ostracode species and juvenile candonids occurred. The juvenile candonids show a high dominance (31–77 %), whereas the other species were relatively rare. In Zone O-II (36–23 cm, 1945–1968 AD) the number of species was 16 and in Zone O-III (23–15 cm, 1968–1982 AD) the number of species dropped down to 14. Zone O-IV (15–0 cm, 1982–2009 AD) yields

the highest number of valves (3,001 valves) in 12–11 cm depth (1988–1990 AD) throughout the core.

In core Sv09, a total of 15 ostracode taxa was identified (Fig. 7; ESM 1). Furthermore, juvenile individuals of the family Candonidae, of the genera *Cypria*, and of the species *Prionocypris zenkeri* (Chyzer and Toth 1858) as well as *Cyclocypris* sp. (juv.), were found. Mostly, *Candona trapeziformis* Klie 1939 is the dominant species in core Sv09 (up to 60 %). Only in the upper core part (3–0 cm; 2004–2009 AD) *Cypria obliqua* Klie 1939 dominates the assemblage (13–23 %). The total number of valves in core Sv09 was rather low. Highest abundance is reached in 12–11 cm (406 valves). The Shannon index increased upcore and the Evenness decreased. The Bray–Curtis dissimilarity ranged between 0.17 (23–22 cm) and 0.67 (15–14 cm). Jaccard similarity was lowest in 22–21 cm (0.09) and highest in 2–1 cm (0.58). Cluster analysis yielded five major assemblage zones: Zone O-I (34–26 cm, 1947–1962 AD) comprised six ostracode species and juvenile candonids.

Fig. 4 Summary of geochemical concentrations (black dots and lines) as well as mass accumulation rates (MAR) and C/N atomic ratios [C/N (atomic)] (grey dots and lines) measured in core St09

Fig. 5 Summary of geochemical concentrations (black dots and lines) as well as mass accumulation rates (MAR) and C/N atomic ratios [C/N (atomic)] (grey dots and lines) measured in core Sv09

527 The total number of valves was low; with a maximum
528 of 45 valves in 31–30 cm and 27–26 cm. In Zone O-II
529 (26–17 cm, 1962–1978 AD) seven species and juve-
530 nile candonids were found and in Zone O-III
531 (17–13 cm, 1978–1986 AD) seven species and

juvenile candonids occurred. The abundance
increased slightly. Zone O-IV (13–3 cm, 1986–2004
AD) revealed the highest number of valves in the core
(maximum in 12–11 cm with 406 valves). In Zone
O-V (3–0 cm, 2004–2009 AD) ostracode abundance

St09

Fig. 6 Ostracode species assemblages, Heip's index of evenness, Shannon index, Bray–Curtis dissimilarity, and Jaccard similarity coefficient in core St09

Sv09

Fig. 7 Ostracode species assemblages, Heip's index of evenness, Shannon index, Bray–Curtis dissimilarity, and Jaccard similarity coefficient in core Sv09

537 was lower compared to Zone O-IV. The maximum
538 number of valves was 193 in 3–2 cm and dropped
539 down to 76 valves in 1–0 cm. This zone included the
540 highest number of species in the entire core (13
541 species; exclusively juvenile candonids).

542 Diatoms

543 A total of 274 diatom species was identified in core
544 Sv09. The majority are only found in Lake Ohrid,
545 Sveti Naum and the hydrologically-connected Lake

Prespa, underlining the high level of biodiversity and
endemism in the lake. 24 groups and complexes
(Fig. 8; ESM 2) were established through combination
of species with similar morphological features and
apparent ecological preferences. Four main zones
(Fig. 8) can be recognised. In Zone D-I (33–24 cm,
1949–1965 AD), the endemic planktonic *Cyclotella*
fottii was dominant (20–40 %), while the planktonic
Cyclotella ocellata Pantocsek 1902 occurred at rela-
tively low abundances (5–10 %). The benthic
Amphora pediculus (Kützing) Grunow in Schmidt

Fig. 8 Summary diatom assemblages, concentration, and F index of the endemic species *C. fottii* in core Sv09

et al. 1875 was present consistently at low abundance (5 %). Zone D-II (24–18 cm, 1965–1976 AD) exhibited very low diatom concentrations. A minor peak in the planktonic *Cyclotella radiosa* occurred at 24–22 cm depth, at the expense of benthic taxa, and was followed by an increase in the relative abundance of *A. pediculus*, *Staurosirella pinnata*, and *Navicula sensu lato* species with an associated reduction in the abundance of *C. fottii*. Zone D-III (18–7 cm, 1976–1996 AD) exhibited a gradually increasing concentration, and an increase to 10–20 % throughout in *C. ocellata*. Zone D-IV (7–0 cm, 1996–2009 AD) is marked by a trend towards the increasing abundance of *C. ocellata* at the expense of *C. fottii*, and there was an abrupt increase in diatom concentration towards the top. The higher relative abundance of *A. pediculus* and *Staurosira pinnata* Ehrenberg 1843 is maintained throughout the depth of 22–0 cm. The common effect of diatom valve deformation due to high toxic metal pollution (Cattaneo et al. 2004) was not observed in core Sv09.

Discussion

The combination of geochemical and biological proxies used here provides evidence by which to assess changes in toxic metal pollution and

eutrophication over time linked to accelerated anthropogenic impact on Lake Ohrid. The exceeded PECs of Fe and Ni in cores St09 and Sv09 throughout the period, and without any notable increases over the last decades, indicate that the source is natural and derived from catchment geology. The south-west and west of Lake Ohrid consists of ultramafic extrusive rocks with associated weathering crusts containing chromium and iron-nickel ore deposits (Vogel et al. 2010). Higher concentrations of Fe and Ni in core Sv09 (Sveti Naum) could result from the closer proximity of the south-eastern part of the lake these deposits and to the piles of waste and ore dump sites of disused mines. Furthermore, the observed counterclockwise rotating surface water current in Lake Ohrid (Vogel et al. 2010) would transport these elements from the western to the eastern part of the lake. Malaj et al. (2011) found that concentrations of heavy metals in sediments are 100 times higher at sample locations in the Albanian sector of the lake, which are also closer to the mining sites than those from the Macedonian area. Many samples were also moderately contaminated (and in one case, moderately to strongly contaminated) with arsenic. The most common sources for As, for over a 100 years, are pesticides and wood preservatives (Alloway 1995), presumably derived from agricultural activity in the catchment as agriculture is one of the most important economic sectors around Lake Ohrid

(Spirkovski et al. 2001) and natural source of As do not exist around Lake Ohrid.

The generally higher abundance of ostracode valves and species diversity near Struga, in comparison to Sveti Naum correlates with slightly higher TOC and TIC values near Struga, indicating higher productivity in the northern part of the lake. C/N ratios near Sveti Naum are higher than near the City of Struga. Such higher ratios were also observed by Vogel et al. (2010) in the south-eastern part of Lake Ohrid near to the river mouth of Čerava, which passes through agricultural and populated areas. Ratios above 10 indicate that most of the organic matter comes from autochthonous production (Meyers and Ishiwatari 1993).

Near the City of Struga, the low ostracode abundance correlates with some peaks in the concentration of As, Cu, Fe, Ni, Zn, and Zr. The number of ostracode valves increases during time intervals when the heavy metal concentrations are low and vice versa. Since species composition does not shift in parallel, these fluctuations may be explained simply by changes in precipitation or amount of snow melt and a subsequently higher sediment load into the lake, rather than being a direct indicator of ecological impact.

The period between the early 1920s and the late 1980s is characterized by low ostracode abundance and low Shannon diversity in both sequences. The low numbers of valves near Sveti Naum could be explained by very low values of TOC and TIC, which indicate a low productivity near the spring discharge. This is confirmed by the low diatom concentration in Zone D–I, and a low abundance of mesotrophic *Cyclotella ocellata* indicating lower productivity in the south-eastern part of Lake Ohrid, with little nutrient input from Lake Prespa in the 1950s and early 1960s. Lake Prespa underwent a relatively high lake-level phase from 1950 to 1962 (Popovska and Bonacci 2007; Popovska 2011), which reduced nutrient enrichment in Lake Prespa. This decreased nutrient input to Lake Ohrid could have been amplified by the retention of nutrients within the karst aquifer (Matzinger et al. 2006a) and by the dilution of Lake Prespa subterranean outflow by mountain range precipitation (Popovska and Bonacci 2007). Only juvenile valves of the ostracode *P. zenkeri* were found in core Sv09. This species prefers waters connected to springs (Meisch 2000) and was probably imported from the springs of Sveti Naum into the lake.

The peak in the diatom species *Cyclotella radiosia* corresponds to a low diatom concentration, correlating with an abrupt lake-level increase in Lake Prespa in 1963 (Popovska and Bonacci 2007; Popovska 2011). This would have, increased the subterranean flow into Lake Ohrid, decreased the nutrient concentration (Matzinger et al. 2006a) and would be likely to have an impact on sediment accumulation rate. The age model does not show a clear change of sediment accumulation rate. Since, the F index of *Cyclotella fottii* does not show evidence of increased diatom dissolution, the low concentration of diatom valves supports a reduction in productivity, supported also by low ostracode abundance. While small forms of diatoms such as *Amphora pediculus* and *Staurosirella pinnata* are known to be vulnerable to sediment focusing processes on steep slopes in boreal lakes (Biskaborn et al. 2012), their abundance decreases rather than increases in this part of the record, which is dominated by planktonic taxa. Instead, the increased subterranean inflow may have resulted in small forms being less likely to settle out of the water column.

Matter et al. (2010) analyzed sediment cores, taken near the north-western shore in Lake Ohrid from ~5 to 10 m and at 53 m water depth. In the cores from shallower water they found a boundary between two distinct stratigraphic units, dated to ~1955. The sediment above this boundary was darker and characterized by lower carbonate content but higher TOC, Fe, Si, and diatom contents. Moreover, a sewage smell was noticeable during core opening. Matter et al. (2010) related this change to increasing anthropogenic impact at that time, but there was no evidence for a similar boundary in the deep water core, other than a slight increase in TOC. Our results show a similar pattern at 50 m depth, with a slight increase in TOC but no evidence for dramatic eutrophication. It appears that the shallow waters in Lake Ohrid show a faster and more drastic response to anthropogenic influences than the deeper water areas (Matter et al. 2010).

Since the mid 1970s, there has been an accelerated, zigzag lake-level decline in Lake Prespa due to the usage of water for irrigation. The most dramatic drop occurred between 1987 and 1995 with a decrease of 5–6 m (Popovska 2011). A lake-level lowering of Lake Prespa by <20 m can increase the nutrient concentration of the lake and thus lead to increased nutrient input via springs to Lake Ohrid, in spite of a decrease in underground flow. Lake Prespa was

undergoing eutrophication at the time due to intensified agriculture and associated water abstraction, fertilizer utilization, and enhanced soil erosion (Matzinger et al. 2006a), which amplified the effects of the lake-level decrease. The increase in the abundance of *Cyclotella ocellata* corresponds to the accelerated nutrient input to Lake Ohrid during this period, and may represent a response to productivity. The diatom record does not show an oscillation of nutrient input linked to the renewed lake-level rise in Lake Prespa between 1979 and 1986 and the dramatic decline between 1987 and 1995, however. An alternate explanation may be that the increase relates instead to associated warming, resulting in an increase in epilimnetic taxa with stronger summer thermal stratification, as appears to be the case in longer term transitions between glacial and interglacial phases (Reed et al. 2010). However, the ostracode data do provides evidence of this lake-level decline in Lake Prespa as the number of valves near the City of Struga and near Sveti Naum increased. This increase resulted in the highest valve concentration in the entire core St09 (maximum = 3,001 valves per 50 g wet sediment). In Sv09, high ostracode abundance (406 and 327 valves per 50 g wet sediment) was also reached during this time. In both cores, this period is characterized by low Shannon species diversity. Increasing productivity in Lake Ohrid is confirmed by high concentrations of TIC and TOC in St09 during this time span and a slight increase near Sveti Naum. It seems that in the highly oligotrophic condition of Lake Ohrid, subtle changes in nutrients have no clear effect on the endemic planktonic diatom *C. fottii* which inhabits the deep, open waters.

After ~1996 AD, the further increase in the epilimnetic diatom *C. ocellata* is mainly the result of the overall decreasing trend of the Prespa lake level (Popovska 2011) and increasing nutrient input into Lake Ohrid (Matzinger et al. 2006a). Between 1991 and 2009 AD, the area next to Sveti Naum was characterized by the highest As concentrations in the entire core. TIC and TOC increased slightly, pointing to increased productivity. This increase could be the reason for the upward increase of the total number of ostracode valves in comparison to the period between the early 1920s and the mid 1980s. Furthermore, the total number of species reached a maximum (13 species), which was the highest number in the entire core. This high biodiversity is also apparent in the coinciding high Shannon index.

The diatom record in core Sv09 does not show the clear changes for the major eutrophication, but there has been an increasing trend in nutrient concentration and productivity in south-eastern Lake Ohrid since the mid 1960s, in spite of its consistent oligotrophic condition. The measured average total phosphorus (TP) concentration in 2002–2004 was 4.6 g l^{-1} , and a simple linear model may estimate the Ohrid TP concentration increasing from $\sim 3.7 \text{ g l}^{-1}$ in the mid 1960s to $\sim 4.8 \text{ g l}^{-1}$ in the late 2000s (Matzinger et al. 2006b). The productivity in this part of Lake Ohrid is strongly influenced by the subterranean inflow and its nutrient supply, which are directly linked to the trophic status and water level of Lake Prespa (Matzinger et al. 2006a; Wagner et al. 2009). If closely connected, the shifts of diatom flora in Sv09 occur 1–2 years later than the changes of water level in Lake Prespa, maybe because the average drainage time from Lake Prespa to the springs near Lake Ohrid is 18 months (Popovska and Bonacci 2007). But a more detailed analysis of the basin-wide diatom response would be necessary to test whether the influence of Prespa has an impact on diatom ecology across the lake as a whole.

In 1988, the first sewage-water treatment system started to operate in the Ohrid-Struga region (UNESCO ROSTE 2004), and Watzin (2003) reported that after the system was completed, an improvement in the water quality in the Ohrid Bay was visible, namely the number of bacteria decreased one thousand fold. However, this positive effect is not clearly visible near the City of Struga. The concentrations of As, Cu, Fe, Hg, Ni, Pb, Zn, and Zr show a downward trend after the water-treatment plant came into operation but the concentrations fluctuated during the time and in the last years, mostly all concentrations show an increase. TIC concentrations were relatively stable and TOC shows a strong upcore increase reaching the maximum concentration between 2002 and 2004 AD. The number of ostracode valves and the total number of species decreased, which could point to the fact that the living conditions in this part of the lake became less favourable.

Conclusions

This multi-proxy approach using sediment records with a high sample resolution from Lake Ohrid provide a detailed insight into the environmental history of the lake. Geochemical analysis reveal

relatively high As concentrations in the northern and south-eastern part of the lake. In core St09 from the northern part, evenly distributed throughout the core, the concentrations correspond to the I_{geo} class “moderately contaminated” and in one sample from the late 1950s to the class “moderately to strongly contaminated”. Sediments from the upper core part (Sv09), taken in the south-eastern sector, were according to the I_{geo} “moderately contaminated” with As. These high concentrations may have been originated from pesticides and wood preservatives used in agriculture around Lake Ohrid. Furthermore, Fe and Ni concentrations often exceeded the PEC levels in both sediment cores, which could have been caused by the ultramafic extrusive rocks with associated weathering crust containing chromium and iron-nickel ore deposits in the west and south-west of Lake Ohrid (Vogel et al. 2010).

Between the early 1920s and the late 1980s, the lake shows generally a low productivity in the northern and south-eastern part, which is indicated by low numbers of ostracode valves, low abundance of the mesotrophic diatom *Cyclotella ocellata*, a general low diatom concentration, as well as low values of TOC and TIC. Furthermore, the low numbers of ostracode valves correlated near the City of Struga with some high concentrations of As, Cu, Fe, Ni, Zn, and Zr. Since the mid 1970s, the increase of *C. ocellata* and an increasing diatom concentration corresponds to rising productivity in the south-eastern lake area. A high number of ostracode valves, the highest number in both cores, indicate an increasing productivity in the late 1980s. This was also confirmed by higher concentrations of TIC, and TOC. A slight increasing productivity trend in the south-eastern part of Lake Ohrid continued from the early 1990s until 2009, which is visible in the increasing TIC and TOC values. During this time, the total number of ostracode valves and the number of ostracode species are also generally high in this area. However, since the early 1990s, the area near the City of Struga in the northern part of the lake is characterized by a decreasing trend in the number of ostracode valves and in the total number of species. This corresponds to an increase of, e.g., TIC, TOC, As, Cu, Fe, Pb, and Zn. This might be an indication that the conditions in the northern lake part became less favourable for ostracodes, which might have dramatic consequences as a loss of the endemic Ohrid ostracode species would be irrevocable.

Acknowledgments We thank Nicole Mantke (Institut für Geologie und Mineralogie, Universität zu Köln, Germany) and Melanie Leng (NERC Isotope Geosciences Laboratory, British Geological Survey, Nottingham, United Kingdom) for carrying out geochemical analyses. Furthermore, we want to thank Meike List for ostracode analysis, Janine Dannecker for quantifying major and trace elements (Institut für Geosysteme und Bioindikation, Technische Universität Braunschweig, Germany) and Burkhard Scharf (Bremen, Germany) for the support with ostracode taxonomy. We owe a debt of gratitude to Goce Kostoski, Sasho Trajanovski, and Zoran Brdarovski (Hydrobiological Institute, Ohrid, Macedonia) for the support and commitment during the field campaign. Funding was provided by the Deutsche Forschungsgemeinschaft (Schw 671/11) and the authors would like to thank the two anonymous reviewers for their constructive comments, which helped us to improve the manuscript.

References

- Albrecht C, Wilke T (2008) Ancient Lake Ohrid: biodiversity and evolution. *Hydrobiologia* 615:103–140
- Aliaj S, Baldassarre G, Shkupi D (2001) Quaternary subsidence zones in Albania: some case studies. *B Eng Geol Environ* 59:313–318
- Alin SR, Cohen AS, Bills R, Gashagaza MM, Michel E, Tiercelin JJ, Martens K, Coveliers P, Mboko SK, West K, Soreghan M, Kimbadi S, Ntakimazi G (1999) Effects of landscape disturbance on animal communities in Lake Tanganyika, East Africa. *Conserv Biol* 13:1017–1033
- Alloway BJ (1995) Heavy metals in soils. Blackie Academic and Professional, London
- Appleby PG, Oldfield F (1992) Application of lead-210 to sedimentation studies. In: Ivanovich M, Harmon RS (eds) Uranium series disequilibrium, applications to earth, marine and environmental sciences. Clarendon Press, Oxford, pp 731–778
- Asaeda T, Shinohara R (2012) Japanese lakes. In: Bengtsson L, Herschy RW, Fairbridge RW (eds) Encyclopedia of lakes and reservoirs. Springer, New York, pp 415–421
- Avramoski O, Kycyku S, Naumoski T, Panovski D, Puka V, Selfo L, Watzin M (2003) Lake Ohrid: experience and lessons learned brief (Lake Basin Management Initiative). n.d. Web 06 Oct 2012. <http://www.ilec.or.jp/eg/lbmi/index.htm>
- Balascio N, Bradley R (2012) Evaluating Holocene climate change in northern Norway using sediment records from two contrasting lake systems. *J Paleolimnol* 48:259–273
- Battarbee RW, Jones VJ, Flower RJ, Cameron NG, Bennion H (2001) Diatoms. In: Smol JP, Birks HJB, Last WM (eds) Tracking environmental change using lake sediments, vol 3., Terrestrial, algal, and siliceous indicators Kluwer, Dordrecht, pp 155–202
- Bilali I, Musai M, Shemo M (2012) Managing the river “Velgozda-Grasnica” from the different chemical pollutants fluted in the Lake Ohrid. Balwois—conference of water observation and information system for decision support, Ohrid

- Biskaborn BK, Herzschuh U, Bolshiyarov DY, Savelieva LA, Diekmann B (2012) Environmental variability in north-eastern Siberia during the last ~13300 yr inferred from lake diatoms and sediment-geochemical parameters. *Palaeogeogr Palaeoclimatol Palaeoecol* 329–330:22–36
- Bray JR, Curtis JT (1957) An ordination of the upland forest communities of Southern Wisconsin. *Ecol Monogr* 27:325–349
- California Academy of Sciences (2011) Catalogue of diatom names, on-line version. n.d. Web 27 Feb 2013. <http://researcharchive.calacademy.org/research/diatoms/names/index.asp>
- Cattaneo A, Couillard Y, Wunsam S, Courcelles M (2004) Diatom taxonomic and morphological changes as indicators of metal pollution and recovery in Lac Dufault (Québec, Canada). *J Paleolimnol* 32:163–175
- Cheburkin AK, Shotykh W (1996) An energy-dispersive miniprobe multielement analyzer (EMMA) for direct analysis of Pb and other trace elements in peats. *Fresen J Anal Chem* 354:688–691
- Cohen AS, Bills R, Cocquyt CZ, Caljon AG (1999) The impact of sediment pollution on biodiversity in Lake Tanganyika. *Conserv Biol* 7:667–677
- Dodeva S (2012) Macedonian lakes. In: Bengtsson L, Herschy RW, Fairbridge RW (eds) *Encyclopedia of lakes and reservoirs*. Springer, New York, pp 503–508
- Grimm EC (1987) CONISS: a FORTRAN 77 program for stratigraphically constrained cluster analysis by the method of incremental sum of squares. *Comput Geosci* 13:13–35
- GeoHive. n.d. Web 10 Nov 2012. <http://www.geohive.com/cntry/europe.aspx>
- Hammer Ø, Harper DA, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4:1–9
- Heiri O, Lotter AF, Lemcke G (2001) Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J Paleolimnol* 25:101–110
- Holtvoeth J, Vogel H, Wagner B, Wolff GA (2010) Lipid biomarkers in Holocene and glacial sediments from ancient Lake Ohrid (Macedonia, Albania). *Biogeosciences* 7:4607–4640
- Houk V, Klee R, Tanaka H (2010) Atlas of freshwater centric diatoms, with a brief key and descriptions. Part 3: Stephanodiscaceae A, Cyclotella, Tertiaris, Discostella. Fottea 10 Supplement. Czech Phycological Society, Benátská
- Jaagumagi R (1993) Development of the Ontario provincial sediment quality guidelines for arsenic, cadmium, chromium, copper, iron, lead, manganese, mercury, nickel, and zinc. Water resources branch, Ministry of the Environment, Ontario
- Kostoski G, Albrecht C, Trajanovski S, Wilke T (2010) A freshwater biodiversity hotspot under pressure—assessing threats and identifying conservation needs for ancient Lake Ohrid. *Biogeosciences* 7:3999–4015
- Krammer K (2002) Diatoms of Europe. Diatoms of the European inland waters and comparable habitats, volume 3: *Cymbella*. A.R.G. Gantner Verlag, Ruggell
- Krammer K, Lange-Bertalot H (1986) Süßwasserflora von Mitteleuropa. Bacillariophyceae. 1. Teil: Naviculaceae (vol. 2/1). Gustav Fischer Verlag, Stuttgart
- Krammer K, Lange-Bertalot H (1988) Süßwasserflora von Mitteleuropa. Bacillariophyceae. 2. Teil: Epithemiaceae, Bacillariaceae, Surirellaceae (vol. 2/2). Gustav Fischer Verlag, Stuttgart
- Krammer K, Lange-Bertalot H (1991a) Süßwasserflora von Mitteleuropa. Bacillariophyceae. 3. Teil: Centrales, Fragilariaceae, Eunotiaceae (vol. 2/3). Gustav Fischer Verlag, Stuttgart
- Krammer K, Lange-Bertalot H (1991b) Süßwasserflora von Mitteleuropa. Bacillariophyceae. 4. Teil: Achnantheaceae (vol. 2/4). Gustav Fischer Verlag, Stuttgart
- Krebs CJ (1989) *Ecological methodology*. Harper and Row, New York
- Lange-Bertalot H (2001) Diatoms of Europe. Diatoms of the European inland waters and comparable habitats, volume 2: *Navicula* sensu stricto, 10 genera separated from *Navicula* sensu lato, *Frustulia*. A.R.G. Gantner Verlag, Ruggell
- Levkov Z (2009) Diatoms of Europe. Diatoms of the European inland waters and comparable habitats, volume 5: *Amphora* sensu lato. A.R.G. Gantner Verlag, Ruggell
- Levkov Z, Williams DM (2011) Fifteen new diatom (Bacillariophyta) species from Lake Ohrid, Macedonia. *Phytotaxa* 30:1–41
- Levkov Z, Krstic S, Metzeltin D, Nakov T (2007) Diatoms of Lakes Prespa and Ohrid (Macedonia). *Iconographia Diatomologica* 16. A.R.G. Gantner Verlag, Ruggell
- Löffler H, Schiller E, Kusel E, Kraill H (1998) Lake Prespa, a European natural monument, endangered by irrigation and eutrophication? *Hydrobiologia* 384:69–74
- Lokoska L (2012) Microbiological investigation of the water and sediment in the north part of Lake Ohrid, Macedonia. Balwois—conference of water observation and information system for decision support, Ohrid
- MacDonald DD, Ingersoll CG, Berger TA (2000) Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 39:20–31
- Magurran AE (2004) *Measuring biological diversity*. Blackwell, Oxford
- Malaj E, Rousseau D, Du Laing G, Lens P (2011) Near-shore distribution of heavy metals in the Albanian part of Lake Ohrid. *Environ Monit Assess* 184:1–17
- Matter M, Anselmetti FS, Jordanoska B, Wagner B, Wessels M, Wüest A (2010) Carbonate sedimentation and effects of eutrophication observed at the Kališta subaquatic springs in Lake Ohrid (Macedonia). *Biogeosciences* 7:4715–4747
- Matzinger A, Veljanoska-Sarafiloska E, Jordanoski M, Naimoski T (2004) Lake Ohrid—a unique ecosystem endangered by eutrophication? Balwois—conference of water observation and information system for decision support, Ohrid
- Matzinger A, Jordanoski M, Veljanoska-Sarafiloska E, Sturm B, Müller M, Wüest A (2006a) Is Lake Prespa jeopardizing the ecosystem of ancient Lake Ohrid? *Hydrobiologia* 553:89–109
- Matzinger A, Spirkovski Z, Patceva S, Wüest A (2006b) Sensitivity of ancient Lake Ohrid to local anthropogenic impacts and global warming. *J Gt Lakes Res* 32:158–179
- Matzinger A, Schmid M, Veljanoska-Sarafiloska E, Patceva S, Guseska D, Wagner B, Müller B, Sturm M, Wüest A (2007)

- 1031 Eutrophication of ancient Lake Ohrid: global warming
1032 amplifies detrimental effects of increased nutrient inputs.
1033 *Limnol Oceanogr* 52:338–353
- 1034 Meisch C (2000) Freshwater Ostracoda of Western and Central
1035 Europe. Spektrum Akademischer, Heidelberg
- 1036 Meyers PA, Ishiwatari R (1993) Lacustrine organic geochem-
1037 istry—an overview of indicators of organic matter sources
1038 and diagenesis in lake sediments. *Org Geochem* 20:
1039 867–900
- 1040 Meyers PA, Teranes JL (2001) Sediment organic matter. In:
1041 Last WM, Smol JP (eds) Tracking environmental change
1042 using lake sediments, vol 2., Physical and geochemical
1043 methods Kluwer Academic Publishers, Dordrecht, pp 239–
1044 269
- 1045 Mikulić F, Pljakić MA (1970) Die Merkmale der kvalitativen
1046 distribution der endemischen Candonaarten im Ochridsee.
1047 *Ekologija* 5:101–115
- 1048 Ministry of environment and physical planning. Human activi-
1049 ties in the basin. n.d. Web 09 Nov 2012. <http://www.moepp.gov.mk/WBStorage/Files/SOER%203%20History,%20socioeconomics.pdf>
- 1050 Müller G (1986) Schadstoffe in sedimenten—sedimente als
1051 Schadstoffe. *Mitteilungen der Österreichischen Mineral-
1052 ogischen Gesellschaft* 79:107–126
- 1053 Neugebauer T, Vallerien D (2012) Energy-efficient protection
1054 of UNESCO natural world heritage—Lake Ohrid, Albania.
1055 *Wasser Abfall* 14:25–29
- 1056 Patceva S, Mitic V, Jordanoski M, Veljanoska-Sarafiloska E
1057 (2004) Influence of the main tributaries on the trophic state
1058 of Lake Ohrid. Balwois—conference of water observation
1059 and information system for decision support, Ohrid
- 1060 Patceva S, Mitic V, Momcula J, Matzinger A, Veljanoska-
1061 Sarafiloska E (2006) Trophic state of Lake Prespa. Balwois—
1062 conference of water observation and information
1063 system for decision support, Ohrid
- 1064 Pérez L, Bugja R, Massaferró J, Steeb P, von Geldern R, Frenzel
1065 P, Brenner M, Scharf B, Schwalb A (2010) Post-Colum-
1066 bian environmental history of Lago Petén Itzá, Guatemala.
1067 *Rev Mex Cienc Geol* 27:490–507
- 1068 Popovska C (2011) Tectonic lakes—climatic and anthropogenic
1069 impacts. EGU General Assembly, Vienna
- 1070 Popovska C, Bonacci O (2007) Basic data on the hydrology of
1071 Lakes Ohrid and Prespa. *Hydrol Process* 21:658–664
- 1072 Reed JM, Leng MJ, Ryan S, Black S, Altinsacılı S, Griffiths HI
1073 (2008) Recent habitat degradation in karstic Lake Uluabat,
1074 western Turkey: a coupled limnological-palaeolimnological
1075 approach. *Biol Conserv* 141:2765–2783
- 1076 Reed JM, Cvetkoska A, Levkov Z, Vogel H, Wagner B (2010)
1077 The last glacial-interglacial cycle in Lake Ohrid (Mace-
1078 donia/Albania): testing diatom response to climate. *Bio-
1079 geosciences* 7:3083–3094
- 1080 Ryves DB, Juggins S, Fritz SC, Battarbee RW (2001) Experi-
1081 mental diatom dissolution and the quantification of
1082 microfossil preservation in sediments. *Palaeogeogr Palaeo-
1083 climatol Palaeoecol* 172:99–113
- 1084 Salemaa H (1994) Lake Ohrid. *Arch Hydrobiol* 44:55–64
- 1085 Schwalb A, Hadorn P, Thew N, Straub F (1998) Evidence for
1086 Late Glacial and Holocene environmental changes from
1087 subfossil assemblages in sediments of Lake Neuchâtel,
1088 Switzerland. *Palaeogeogr Palaeoclimatol Palaeoecol* 140:
1089 307–323
- 1090 Spirkovski Z, Avramovski O, Kodzoman A (2001) Watershed
1091 management in the Lake Ohrid region of Albania and
1092 Macedonia. *Lakes Reserv Res Manag* 6:237–242
- 1093 Stanković S (1960) The Balkan Lake Ohrid and its living world.
1094 Uitgeverij Dr. W. Junk, Den Haag
- 1095 Touchart L (2012) Lake Baikal. In: Bengtsson L, Herschy RW,
1096 Fairbridge RW (eds) Encyclopedia of lakes and reservoirs.
1097 Springer, New York, pp 83–91
- 1098 UNESCO ROSTE (2004) Report about the Lake Ohrid water-
1099 shed region. n.d. Web 24 Oct 2012. http://portal.unesco.org/en/files/24220/11032790451ohrid_prespa_report_august_2004.pdf/ohrid+prespa+report+august+2004.pdf
- 1100 Veljanoska-Sarafiloska E, Jordanoski M, Mitic V, Patceva S
1101 (2004) Influence of River Velgoska and Koselska on the
1102 trophic state of Lake Ohrid. Balwois—conference of water
1103 observation and information system for decision support,
1104 Ohrid
- 1105 Vogel H, Wessels M, Albrecht C, Stich HB, Wagner B (2010)
1106 Spatial variability of recent sedimentation in Lake Ohrid
1107 (Albania/Macedonia)—a complex interplay of natural and
1108 anthropogenic factors and their possible impact on biodi-
1109 versity patterns. *Biogeosciences* 7:3333–3342
- 1110 Wagner B, Lotter A, Nowaczyk N, Reed J, Schwalb A, Sulpizio
1111 R, Valsecchi V, Wessels M, Zanchetta G (2009) A 40,000-
1112 year record of environmental change from ancient Lake
1113 Ohrid (Albania and Macedonia). *J Paleolimnol* 41:407–430
- 1114 Watzin MC (2003) Lake Ohrid and its watershed: our lake, our
1115 future. A state of the environment report. Lake Ohrid
1116 conservation project, Tirana, Albania and Ohrid