

Post-concussive Symptoms and Neuropsychological Performance in the Post-acute Period following Pediatric Mild Traumatic Brain Injury

Martina Studer,^{1,2} Barbara Goeggel Simonetti,^{1,2,3} Alexander Joeris,⁴ Katja Margelisch,^{1,2,5} Maja Steinlin,^{1,2} Claudia Maria Roebers,^{2,5} AND Theda Heinks^{1,2}

¹Department of Pediatric Neurology, Development & Rehabilitation, University Children's Hospital Bern, Bern

²Center for Cognition and Learning, University of Bern, Bern

³Department of Neurology, University Hospital Bern, Bern

⁴Department of Pediatric Surgery, University Children's Hospital Bern, Bern

⁵Department of Developmental Psychology, University of Bern, Bern

(RECEIVED February 18, 2014; FINAL REVISION September 6, 2014; ACCEPTED October 7, 2014; FIRST PUBLISHED ONLINE November 10, 2014)

Abstract

Objective: There is evidence that children after mild traumatic brain injuries (mTBI) suffer ongoing post-concussive symptoms (PCS). However, results concerning neuropsychological outcome after mTBI are controversial. Thus, our aim was to examine group differences regarding neuropsychological outcome and PCS. Additionally, we explored the influence of current and pre-injury everyday attention problems on neuropsychological outcome in children after mTBI. **Method:** In a prospective short-term longitudinal study, 40 children (aged 6–16 years) after mTBI and 38 children after orthopedic injury (OI) underwent neuropsychological, socio-behavioral and PCS assessments in the acute stage and at 1 week, at 4 weeks, and 4 months after the injury. **Results:** Parents of children after mTBI observed significantly more PCS compared to parents of children after OI, especially in the acute stage. Our results revealed no neuropsychological or socio-behavioral differences over time between both groups. However, in children after mTBI, we found negative correlations between elevated levels of everyday attention problems and reduced neuropsychological performance. Furthermore, there was a negative influence of pre-injury everyday attention problems on neuropsychological performance in children after mTBI. **Conclusion:** In accordance with earlier studies, parents of children after mTBI initially observed significantly more PCS compared to parents of children after OI. There were no neuropsychological or socio-behavioral group differences between children after mTBI and OI in the post-acute period. However, our exploratory findings concerning the influence of everyday attention problems on neuropsychological outcome indicate that current and pre-injury everyday attention problems were negatively associated with neuropsychological performance in children after mTBI. (*JINS*, 2014, 20, 982–993)

Keywords: Concussion, Children, Recovery, Pre-injury attention problems, Everyday attention problems, Verbal learning and memory performance

INTRODUCTION

With an annual incidence of 200–500 per 100,000 children (Kraus, 1995), traumatic brain injuries (TBI) are the leading cause of long-term disability in children and adolescents, with serious effects on the lives of patients and their families (Feigin et al., 2013). Although 80–90% of all pediatric TBI cases are of mild severity, this large population has been neglected in research for a long time, due to the assumption that mild TBI (mTBI) is a “benign” injury, without any

cognitive sequelae (Carroll et al., 2004; Menascu & MacGregor, 2007; Yeates, 2010a). Even though this assumption is supported by meta-analyses and reviews, there is a high variability within outcome studies after mTBI (Babikian & Asarnow, 2009; Carroll et al., 2004; Satz et al., 1997).

Cognitive outcome after mTBI has been controversially discussed in the literature. Various possible explanations for the inconsistencies in results have been debated, such as divergent definitions of mTBI, the use of different types of control groups (healthy or with other injuries), different types of outcome measures (performance-based testing vs. parent questionnaires) as well as the lack of adequate control of pre-injury factors that might influence outcome (Asarnow, Satz, Light, Lewis, & McCleary, 1995; Babikian, & Asarnow, 2009;

Correspondence and reprint requests to: Martina Studer, Department of Pediatric Neurology, Development & Rehabilitation, University Children's Hospital, Inselspital, 3010 Bern. E-mail: martina.studer@insel.ch

Babikian et al., 2011). These points will be outlined in the following sections.

There is compelling evidence that children after mTBI suffer from an array of *post-concussive symptoms* (PCS) that include somatic (e.g., headache, blurry vision, nausea), cognitive (e.g., attention and memory problems, reduced processing speed), behavioral (e.g., sleep difficulties), and emotional (e.g., mood lability, aggressive behavior) complaints (Mittenberg, Wittner, & Miller, 1997; Taylor et al., 2010; Yeates et al., 2009). Compared to healthy controls as well as to children with orthopedic injuries, children after mTBI exhibit more PCS following the injury that typically recede within the first months after the injury (Barlow et al., 2010; Rieger et al., 2013; Sroufe et al., 2010; Taylor et al., 2010; Yeates et al., 2009). Although most parents of children with mTBI do not report elevated PCS after the post-acute period of 3 months, roughly 10% of children after mTBI exhibit persisting PCS (Barlow et al., 2010; Taylor et al., 2010). A recent study investigating predictive factors of PCS after pediatric mTBI demonstrated that a variety of non-injury child and family factors were substantially associated with persistent PCS, while injury related factors were only significant predictors in the acute period (McNally et al., 2013). This result suggests that, depending on the time point after the injury, physiological and psychological factors might differentially influence the development of PCS.

Persisting PCS often cover cognitive problems, especially attention and memory impairments (Taylor et al., 2010). These reported cognitive impairments correspond with recent performance-based studies in children after mTBI in which, compared to healthy non-injured control children, subtle impairments in verbal memory, selective attention, working memory, or switching were identified (Anderson, Catroppa, Morse, Haritou, & Rosenfeld, 2001; Babikian et al., 2011; Catale, Marique, Closset, & Meulemans, 2009; Loher, Fatzer, & Roebbers, 2014; Scherwath et al., 2011). Selective attention as well as the executive functions working memory and switching are higher-order cognitive functions and strongly related to verbal memory capacities (Duff, Schoenberg, Scott, & Adams, 2005). It is assumed that these complex functions are slightly impaired in children after mTBI, compared to healthy children.

However, when comparing neuropsychological outcome in children after mTBI to that of children with “other” injuries, studies report comparable or even better neuropsychological performance in mTBI patients (Babikian et al., 2011; Maillard-Wermelinger et al., 2009; Rieger et al., 2013). In a comprehensive neuropsychological outcome study of Babikian et al. (2011), children after mTBI performed similarly to children with orthopedic injuries (OI), but worse compared to healthy children. The authors interpreted the comparable performance between mTBI and OI children in terms of a *general injury effect*, pointing to a potential influence of pre-injury differences. Since injuries like mTBI or OI do not often occur at random, there is some evidence which suggests that injured children are more likely to have a history of pre-existing behavioral or learning problems and

often have lower socioeconomic backgrounds (Asarnow et al., 1995; Durkin, Davidson, Kuhn, O'Connor, & Barlow, 1994; Ramsay et al., 2003). To control for such pre-existing systematic differences as well as for characteristics associated with traumatic injuries (e.g., hospitalization), it has been suggested to compare outcome of children after mTBI to children with a different injury not involving the head (Yeates, 2010b).

Besides choosing an appropriate control group, it is highly important to detect pre-injury risk factors like attention or learning problems that might aggravate cognitive problems after childhood TBI. It was shown that pre-injury risk factors lead to more pronounced impairments in cognition, daily functioning or elevated cognitive PCS after the injury (Babikian, MacArthur, & Asarnow, 2013; Bonfield, Lam, Lin, & Greene, 2013; Farmer et al., 2002; Fay et al., 2010; Yeates et al., 2005). Some of these findings were explained using the theoretical construct of *cognitive reserve capacity* that refers to the ability to optimize cognitive performance through differential recruitment of brain networks (Satz, 1993; Stern, 2002). According to this theory, possessing intact cognitive abilities is a resource, protecting against the effects of a TBI, while pre-injury problems may reduce cognitive resources after an injury and thus contribute to an impaired outcome. To date, only few studies have investigated and demonstrated the negative effect of pre-injury risk factors on outcome after pediatric mTBI (Babikian et al., 2013; Bonfield et al., 2013; Fay et al., 2010), and to the best of our knowledge, no published study examined the role of pre-existing attention problems on neuropsychological outcome after pediatric mTBI. Because pre-injury attention problems moderate outcome after more severe TBI (Yeates et al., 2005), it is important to examine whether pre-injury attention problems might influence as well cognitive outcome in children after mTBI.

Summarizing, cognitive outcome in children after mTBI is controversially discussed among researchers. Although some of the inconsistencies are explained by the type of control group, there seems to exist a correspondence between the reported attention and memory impairments in performance-based tests and observed continuing PCS like inattentiveness or forgetfulness in children after mTBI. Because memory, attention, and executive functions are not only crucial for academic achievement, but also for social life, problems in these higher-order cognitive functions might impede various aspects of daily life (Heim & Keil, 2012; Roebbers, Röthlisberger, Cimeli, Michel, & Neuenschwander, 2011; Smallwood, Fishman, & Schooler, 2007). Consequently, it is important to study the post-acute trajectory of such susceptible cognitive functions to increase the possibility of detecting problems as early as possible.

Thus, the main aim of our short-term longitudinal study presented here was to examine PCS as well as attention, executive function, and verbal memory in the post-acute period after mTBI and OI. According to the literature we hypothesized that parents of children with mTBI will observe more PCS than parents of children with OI. Furthermore, we hypothesized that children after mTBI show a similar

neuropsychological outcome as children after OI. In addition to these main hypotheses, we explored the association of pre-injury and current everyday attention problems on neuropsychological outcome in children after mTBI.

METHODS

The study was approved by the Ethics Committee of the University Children's Hospital in Bern and by the Bernese Cantonal Ethics Committee. All caregivers provided informed written consent before participation, consistent with the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Participants and Sample Attrition

German-speaking participants between 6 and 16 years of age were recruited for this study between February 2012 and September 2012 (for children after OI, we extended the recruitment period to April 2013 to have similar subsample sizes) in the Emergency Department (ED) of the University Children's Hospital in Bern, Switzerland. Inclusion criteria were (a) the diagnosis of an mTBI, confirmed by an emergency physician and defined according to the American Congress of Rehabilitation Medicine (ACRM, 1993); and (b) no pre-existing psychiatric or neurodevelopmental diseases (except for ADHD, known to be more common in children sustaining an injury, DiScala, Lescohier, Barthel, & Li, 1998). Consistent with the ACRM (1993), we defined mTBI with an initial Glasgow Coma Scale (GCS) of 13–15, as deriving from a traumatically induced physiological disruption of brain function, as manifested by at least one of the following symptoms: loss of consciousness < 30 min, duration of post-traumatic amnesia (PTA) (< 24 hr), any alteration in mental state at the time of the accident, with possible focal deficits of transient nature. Inclusion Criteria for the OI were (a) fractures of the non-dominant upper extremity or of the lower extremity below the knee or soft wound tissue wounds requiring surgical intervention, (b) no head involvement as well as no injury of the dominant upper extremity (because children were tested with speed dependent neuropsychological tasks 1 month after the injury) and (c) the absence of pre-existing psychiatric or neurodevelopmental disease (except ADHD, for the same reason as above, DiScala et al., 1998). A total of 92 children (46 children per group) were recruited for this study. Between the ED visit and the first neuropsychological testing session, six children (13%) of the mTBI group and eight children (17%) of the OI group declined further participation. Between the first and second neuropsychological assessment, three children (8%) of the mTBI group and four children (11%) of the OI group declined further participation. Overall, the dropout rate was comparable to other studies (e.g., McCullagh & Feinstein, 2003) and a retrospective analysis concerning the participation rate revealed similar participation rates (mTBI: 46%, OI: 30%) to a study by Yeates et al. (2009). All participants who completed the first neuropsychological assessment were

included in data analyses. While all children in the mTBI group were inpatients and hospitalized for GCS supervision, seven children after OI (18%) were not inpatients and were released after initial treatment in the ED had been completed. Descriptive information of both groups is shown in Table 1.

Procedure

Figure 1 illustrates the study design of our prospective short-term longitudinal study with four time-points (acute = T0, 1 week after the injury = T1, 1 month after the injury = T2, and 4 months after the injury = T3). In the ED, parents were asked to fill out questionnaires to rate acute PCS (T0 post-injury) as well as the presence and severity of these symptoms for the time before the injury (T0 pre-injury) (Gioia, Schneider, Vaughan, & Isquith, 2009). Furthermore, parents were asked to fill out the same questionnaire at all other time points. At T2 and T3, comprehensive neuropsychological assessments were performed, with a counterbalanced task order using alternate versions of the tests at T3, if available. Children were tested individually in a quiet room by a child neuropsychologist (M.S.) or by a trained psychology graduate student. Assessment duration was approximately 90 min each time. Regular breaks were offered and at the end of testing sessions, children were allowed to pick a small gift as reward for their participation.

Measures

At T2 and T3, participants performed neuropsychological tests to assess verbal learning and memory, executive functions (working memory and switching), selective attention, nonverbal intelligence, and processing speed. Raw scores of the tests were transformed into age-corrected standard scores (SS) or percentiles (PR), as requested by the respective test manual.

Verbal learning and memory performance was assessed with the German Version of the Rey Auditory Verbal Learning Test (RAVLT, Helmstaedter, Lendt, & Lux, 2001). The dependent variables were verbal learning (the sum of the recalled words over the five learning trials) and verbal memory (delayed recall). We used an alternate list of words at T3.

Working memory was assessed using the backward digit recall task of the Working Memory Test Battery (WMTB-C, Pickering & Gathercole, 2001).

Switching was assessed using the fourth condition (number-letter switching) of the Trail Making Task of the Delis-Kaplan Executive Function System (D-KEFS, Delis, Kaplan, & Kramer, 2001). This test could only be conducted with children aged 8 years and older. The dependent variable was time for completion.

Selective Attention was measured with the Conners' Continuous Performance Test (CPT, Conners, 2004). The dependent variables were the errors of omission (number of targets to which the individual did not respond) as a proxy for inattention/inhibition and the errors of commission [number of times the individual responded to a nontarget ("X")] as a proxy for impulsivity (Epstein et al., 2003).

Table 1. Demographic and injury characteristics of both groups

	mTBI ¹ group	OI group
<i>N</i>		
T0, <i>n</i> (% male)*	40 (42.5)	38 (64.9)
T1, <i>n</i> (% male)*	40 (42.5)	38 (64.9)
T2, <i>n</i> (% male)*	40 (42.5)	38 (64.9)
T3, <i>n</i> (% male)*	37 (43.6)	34 (70.6)
Age in years at injury, <i>M</i> (<i>SD</i>)	11.05 (3.1)	10.54 (2.65)
SES (highest parental education level), <i>M</i> (<i>SD</i>)	2.45 (.93)	2.37 (.75)
University degree, <i>n</i> (%)	8 (20)	5 (13.2)
High School degree, <i>n</i> (%)	6 (15)	5 (13.2)
Apprenticeship, <i>n</i> (%)	22 (55)	27 (71.1)
Obligatory school, <i>n</i> (%)	4 (10)	1 (2.6)
Therapy requirement before injury ^a , <i>n</i> (%)	10 (25)	15 (39.5)
Attention deficit hyperactivity disorder, <i>n</i> (%)	1 (2.5) ^b	1 (2.6)
Injury characteristics		
GCS at injury, <i>M</i> (<i>SD</i> , range)	14.78 (0.53, 13–15)	15
Intracranial injury ^c , <i>n</i> (%)	3 (7.5)	0
Loss of consciousness, <i>n</i> (%)	14 (35)	0
Duration of loss of consciousness, <i>M</i> (<i>SD</i> , range), min	0.45 (1.19, 0–5)	0
Retrograde Amnesia, <i>n</i> (%)	12 (30)	0
Anterograde Amnesia, <i>n</i> (%)	20 (50)	1 (2.2) ^d
Previous mTBI, <i>n</i> (%)	11 (27.5)	6 (15.8)
Cause of injury		
Fall, <i>n</i> (%)	25 (62.5)	22 (57.9)
Blow, <i>n</i> (%)	7 (17.5)	1 (2.6)
Bike accident, <i>n</i> (%)	6 (15)	5 (13.2)
Motorbike accident, <i>n</i> (%)	1 (2.5)	—
MVC pedestrian, <i>n</i> (%)	1 (2.5)	1 (2.6)
Others, like spraining, <i>n</i> (%)	—	9 (23.7)

Note. ¹mTBI: mild traumatic brain injury; * $p < .05$. OI = orthopedic injury; T0 = acute; T1 = 1 week following the injury; T2 = 1 month following the injury; T3 = 4 months following the injury; SES = socio-economic status (highest parental education level); GCS = Glasgow Coma Scale; MVC = motor vehicle collision.

^a“Therapy requirement” summarizes the following therapies: medication (Ritalin), speech therapy, psychomotor therapy, occupational therapy, any kind of math/literacy intervention and special school support.

^bThis child required stimulant medication.

^cAll intracranial injuries were extra cerebral.

^dThis child had a toe fracture, without any evidence for an additional mTBI.

Processing speed was measured using the Processing Speed Index score of the German version of the Wechsler Intelligence Scale for children (WISC-IV, Petermann, & Petermann, 2007).

Nonverbal Intelligence was assessed using the Test of Nonverbal Intelligence (TONI 3, L. Brown, Sherbenou, & Johnsen, 1997), a language and motor-free measure of intellectual functioning.

Socio-behavioral strengths and difficulties were measured with the Strength and Difficulties Questionnaire (SDQ, Goodman, 1997), rated by parents at T2 and T3. Subscale scores above three (conduct problems), four (peer problems), five (emotional symptoms), six (hyperactivity/inattention), or below five (prosocial behavior) are considered abnormal. Our dependent variables were the raw scores of all five subscales.

Post-concussive symptoms were rated by parents and children using a translated and slightly adapted version of the Post-concussion symptom inventory (PCSI, Gioia et al., 2009). It consists of 29 items that have to be rated on a Likert Scale (0–3, never/seldom/sometimes/often), covering

physical, cognitive, sleep-related, and emotional symptoms which have been derived from factor analytic approaches (Gioia, Vaughan, & Isquith, 2011). To attain more information on cognitive items, we added the following cognitive items from the Health and Behavior Inventory (HBI, Ayr, Yeates, Taylor, & Browne, 2009): “easily distracted” / “forgetful” / “difficulties to complete a task” / “difficulties to stay focused on a task”. For this study, we used the sum of all PCS items of the parent ratings similar to other studies (e.g., Rieger et al., 2013). Furthermore, we concatenated the parent-rated cognitive items covering *everyday attention problems* (concentration problems / easily distracted / difficulties to execute an order / difficulties to complete a task / difficulties to stay focused on a task), creating the variable *everyday attention problems*. Internal consistency was strong for the total scale as well as for the subscale *everyday attention problems* ($\alpha = .87-.91$). Furthermore, test–retest reliability of the *everyday attention problems* subscale (Pearson r between T2 and T3, $r_{T2,T3} = .65$, $p < .001$) was acceptable and there were hints for convergent validity (Pearson r between the




Fig. 1. Flow chart with the design and drop-out rate of the short-term longitudinal study.

parent-rated SDQ subscale hyperactivity / inattention and the everyday attention problems subscale, $r_{T2} = .57$, $p < .001$; $r_{T3} = .70$, $p < .001$) in children after mTBI. For one analysis, we dichotomized the variable everyday attention problems into *presence of pre-injury everyday attention problems* (sum of the attention items ≥ 1) and into *absence of pre-injury everyday attention problems* (sum of the attention items = 0).

Highest level of education in the family was used as a proxy for socio-economic status (SES) and was coded using the following scale: 1 = obligatory schooling, 2 = high school/on the job training, 3 = college degree, 4 = university/graduate degree.

Statistical Analyses

For the statistical analyses we used the Statistical Package for Social Sciences software for Windows, version 21 (SPSS IBM, New York). Demographic characteristics were compared using two-tailed independent samples t tests (age at injury, nonverbal IQ) and nonparametric Pearson Chi-square test (χ^2 , for gender, SES, previous mTBI, and pre-injury therapy requirements). Before our analysis, we executed exploratory analyses to identify possible confounding covariates. Due to our gender imbalance, we included gender as covariate because there is evidence that gender has a moderating effect on neuropsychological outcome and may influence PCS rating in parents (Donders & Woodward, 2003; Taylor et al., 2010). In a first step, repeated-measures analyses of covariance (ANCOVAs) (Group \times Time, controlled for gender) were performed to investigate the influence of the injury group and time on the parental rating of PCS and everyday attention problems. Because Mauchly's test of sphericity was violated in these analyses, we adapted

the degrees of freedom (df) using the Huynh-Feldt correction. To follow-up on the very recent assumption that emotional symptoms may be associated with persistent PCS (Donlon, & Jones, 2014), we additionally performed a hierarchical regression analysis to explore the contribution of demographic variables, injury group, pre-injury PCS, and emotional symptoms at T3 (SDQ subscale) on PCS at T3. In a second step, repeated-measures ANCOVAs (Group \times Time, controlled for gender) were used to test the influence of injury group and time on neuropsychological and socio-behavioral outcome. To measure associations between everyday attention problems and neuropsychological performance, cross-sectional two-sided partial Pearson correlations (controlled for gender and pre-injury everyday attention problems) were computed for T2 and T3. In a last step, independent samples t tests were computed to investigate the influence of pre-injury attention problems on neuropsychological outcome in children after mTBI. Statistical significance was defined as $p < .05$, however, in the case of multiple comparisons (correlations and independent samples t tests), we applied the Bonferroni correction with a modified p -value. Effect size is reported as Cramer V , r , or partial Eta² (η_p^2) values.

RESULTS

Descriptive Data

Descriptive and injury-related characteristics of both groups are presented in Table 1. Injury severity of our mTBI group was very mild, with a mean GCS of 14.8 and only a third of participants suffered from a short period of unconsciousness (≤ 5 min). Both groups were comparable concerning age at injury, $t(76) = .77$, $p = .44$, SES, $\chi^2(3) = 3.04$, $p = .39$,




Fig. 2. Development of post-concussive symptoms across all time points, rated by parents (*M, SD*).

nonverbal IQ, $t(76) = 1.90$, $p = .06$, previous mTBI, $\chi^2(1) = 1.89$, $p = .17$ and pre-injury therapy requirement, $\chi^2(1) = 1.87$, $p = .17$. At T2, $\chi^2(1) = 4.25$, $p = .04$, $V = .23$ as well as at T3, $\chi^2(1) = 5.38$, $p = .02$, $V = .28$, there was a gender imbalance between both groups with significantly more male participants in the OI group, compared to the mTBI group. Furthermore, three children had intracranial injuries (subarachnoid hemorrhage, epidural hematoma), confirmed by an acute CT. Although the presence of intracranial lesions is related with worse neurobehavioral outcomes (Williams, Levin, & Eisenberg, 1990), we included these children because they were eligible patients. Furthermore, our results did not change when excluding these three cases with intracranial injuries.

Change of Post-concussive Symptoms in the Post-acute Period

Changes in the sum of PCS across all assessments (T0–T3) are shown in Figure 2. Repeated-measure ANCOVA revealed no main effect of time, but a main effect of injury group, $F(1,65) = 4.72$, $p = .03$, $\eta_p^2 = .07$ (parents of children after

mTBI > parents of children after OI) and an interaction effect of time x injury group, $F(2.56, 166.49) = 2.87$, $p = .04$, $\eta_p^2 = .04$. Parents of children after mTBI observed more PCS acutely compared to 4 months after the injury. Furthermore, the covariate gender was significantly related to PCS, $F(1,65) = 5.29$, $p = .03$, $\eta_p^2 = .08$, revealing that parents of girls observed more PCS than parents of boys. Cross-sectional ANCOVAs indicated that parents of children after mTBI rated more PCS at T0 post-injury, $F(1,75) = 11.12$, $p = .001$, $\eta_p^2 = .13$ and at T1, $F(1,75) = 6.08$, $p = .02$, $\eta_p^2 = .08$. At all other time-points, there were no differences between groups.

A hierarchical regression analysis investigating the independent contribution of injury group, pre-injury PCS as well as emotional symptoms on PCS beyond the acute period (controlling for demographic variables in the first step) indicated that only pre-injury PCS ($\beta = .21$; $p < .05$) and emotional symptoms at T3 ($\beta = .68$; $p < .001$) significantly predicted PCS at T3 ($F(6,60) = 11.26$; $p < .001$; $R^2 = .53$).

Changes in everyday attention problems across all assessments (T0–T3) are illustrated in Figure 3. Repeated-measure ANCOVA revealed no effect of time or injury group on everyday attention problems.




Fig. 3. Development of everyday attention problems across all time points, rated by parents (*M, SD*).

Neuropsychological Outcome 1 Month (T2) and 4 Months (T3) following the Injury

Descriptive data of the neuropsychological assessments at T2 and T3 are reported in Table 2. Repeated-measures ANCOVA revealed a significant time effect for processing speed, $F(1,67) = 4.42, p = .04, \eta_p^2 = .06$, meaning that both groups improved their processing speed between T2 and T3 ($p < .001$). Furthermore, the covariate gender, $F(1,67) = 5.90, p = .03, \eta_p^2 = .02$ was significantly related to processing speed, revealing that girls showed a quicker processing speed than boys. Concerning all other neuropsychological functions, there were neither main effects (injury group or time) nor interaction effects found. At both time-points, the means of all participants were within the normal range.

Socio-behavioral Outcome 1 Month and 4 Months following the Injury

The socio-behavioral outcome (SDQ subscale scores), rated by parents, is shown in Table 3. Repeated-measures ANCOVA revealed no changes over time or differences between groups. At both time-points, means of all participants were within the normal range.

Relation between Everyday Attention Problems and Neuropsychological Performance

Correlation coefficients between neuropsychological performance and parent-rated everyday attention problems are shown in Tables 4 and 5. One month after the injury (T2) (Table 4), there were generally small-sized, insignificant correlations between everyday attention problems and selected neuropsychological functions in both groups.

Four months after the injury (Table 5), there were again generally small, insignificant correlations in children after OI. In children after mTBI, however, there was a significant, large-sized negative correlation between verbal learning and everyday attention problems. The medium-sized correlation between working memory and everyday attention problems ($p = .01$) fell short of the significance threshold ($p < .008$) after controlling for gender and pre-injury attention problems. The associations between the performance-based selective attention measures (inattention and impulsivity) and the ratings of everyday attention problems were small and insignificant for either testing session.

Influence of Pre-injury Everyday Attention Problems on Neuropsychological Performance in Children after mTBI

The influence of pre-injury attention problems on neuropsychological outcome in children after mTBI is shown in Table 6. On a descriptive level, children after mTBI with pre-injury attention problems showed at both time points a worse neuropsychological performance compared to children after mTBI without pre-injury attention problems. At T2, children

Table 2. Neuropsychological test performance for both groups at one (T2) and four (T3) months post-injury (M,SD)

	mTBI						OI			
	T2 (1 month post-injury)		T3 (4 months post-injury)		T2 (1 month post-injury)		T3 (4 months post-injury)		T3 (4 months post-injury)	
	Female (n = 23)	Male (n = 17)	Female (n = 21)	Male (n = 16)	Female (n = 13)	Male (n = 25)	Female (n = 10)	Male (n = 24)	Female (n = 10)	Male (n = 24)
Verbal learning (PR) ^a	54.78 (33.32)	40.00 (28.96)	56.95 (32.80)	53.80 (35.92)	57.77 (30.51)	44.28 (32.55)	73.60 (33.76)	45.46 (29.49)	73.60 (33.76)	45.46 (29.49)
Verbal memory (PR) ^a	58.17 (27.01)	49.81 (30.41)	56.24 (32.88)	45.60 (26.73)	64.08 (26.03)	54.48 (29.19)	76.40 (25.44)	61.25 (26.71)	76.40 (25.44)	61.25 (26.71)
Working memory (SS) ^a	94.35 (14.12)	98.81 (19.03)	95.81 (13.24)	100.60 (15.66)	93.69 (15.08)	93.12 (9.62)	97.80 (8.90)	94.42 (8.83)	97.80 (8.90)	94.42 (8.83)
Switching (SS) ^b	10.13 (2.92)	10.23 (3.30)	11.53 (3.04)	11.25 (3.72)	10.60 (2.46)	10.18 (3.80)	11.88 (2.23)	10.89 (2.76)	11.88 (2.23)	10.89 (2.76)
Nonverbal IQ ^c	103.91 (13.78)	108.12 (17.02)	112.71 (10.93)	110.31 (12.21)	98.85 (11.11)	100.56 (11.24)	114.50 (10.24)	104.96 (13.71)	114.50 (10.24)	104.96 (13.71)
Processing speed index (SS) [*]	109.39 (8.91)	104.06 (11.08)	48.22 (24.54)	52.37 (29.66)	54.46 (23.66)	42.11 (24.58)	53.82 (25.97)	49.47 (26.46)	53.82 (25.97)	49.47 (26.46)
Inattention (PR) ^d	51.61 (27.02)	50.68 (28.21)	64.50 (31.44)	55.37 (33.12)	65.58 (23.11)	53.40 (31.69)	71.33 (26.99)	44.65 (34.02)	71.33 (26.99)	44.65 (34.02)
Impulsivity (PR) ^d	65.76 (25.97)	49.66 (34.48)								

Note. mTBI = mild traumatic brain injury; OI = orthopedic injury; M = mean; SD = standard deviation; PR = Percentile; SS = Standard Score; IQ = Intelligence Quotient.

^aDue to language problems, one boy in the mTBI group could not be tested with the verbal learning and memory as well as working memory test at both time points.

^bThis test could only be conducted with children aged 8 years and older, leading to a smaller n in the single cells (T2: mTBI: n = 30; OI: n = 28; T3: mTBI: n = 27; OI: n = 26).

^cNonverbal IQ was only tested at T2.

^dFor both variables, inattention and impulsivity, high percentage values mean a worse attention performance.

^{*}Significant main effect of time ($p < .05$).

Table 3. Socio-behavioral strengths and weaknesses for both groups at one (T2) and four (T3) months post-injury (*M (SD)*)

	mTBI				OI			
	T2 (1 month post-injury)		T3 (4 months post-injury)		T2 (1 month post-injury)		T3 (4 months post-injury)	
	Female (<i>n</i> = 21)	Male (<i>n</i> = 17)	Female (<i>n</i> = 19)	Male (<i>n</i> = 14)	Female (<i>n</i> = 12)	Male (<i>n</i> = 22)	Female (<i>n</i> = 9)	Male (<i>n</i> = 23)
Emotional symptoms	1.71 (2.00)	1.47 (1.42)	2.26 (1.76)	1.21 (1.53)	1.25 (1.42)	1.45 (1.22)	1.89 (2.80)	1.09 (1.28)
Conduct problems	1.33 (1.39)	2.00 (1.37)	1.68 (1.64)	1.50 (1.51)	1.42 (1.51)	1.64 (1.40)	1.89 (2.03)	1.17 (1.19)
Hyperactivity/inattention problems	2.52 (1.72)	2.76 (2.54)	2.79 (1.96)	2.50 (2.35)	2.08 (1.98)	3.00 (1.95)	3.11 (2.93)	2.74 (2.18)
Peer problems	1.10 (1.45)	1.12 (1.50)	1.68 (1.95)	1.00 (1.71)	1.17 (1.03)	1.50 (1.95)	1.56 (1.81)	1.30 (2.10)
Prosocial behavior ¹	8.81 (1.08)	7.76 (1.86)	8.53 (1.61)	8.64 (1.22)	8.92 (1.00)	7.50 (1.50)	8.44 (1.81)	7.70 (1.69)

Note. No significant main effects.

¹Only for the prosocial behavior subscale: Low raw scores point to higher problems, meaning less prosocial behavior. mTBI: mild traumatic brain injury. OI: orthopaedic injury.

with pre-injury attention problems showed a significantly worse learning performance than children without pre-injury attention problems, $t(37) = 2.86, p = .007, r = .43$. At T3, children with pre-injury attention problems showed a worse verbal memory ($t(34) = 2.69; p = .01; r = .42$) and working memory ($t(34) = 2.44; p = .02; r = .39$) performance with medium effect sizes; however, after applying the Bonferroni correction, these results became insignificant.

DISCUSSION

The aim of this study was to examine neuropsychological functions as well as PCS in the post-acute period in children after mTBI in comparison to children after OI. According to earlier studies, parents of children after mTBI observed significantly more PCS, particularly in the acute stage and 1 week following the injury (Barlow et al., 2010; Sroufe et al., 2010; Taylor et al., 2010). Furthermore, our analysis revealed that gender influenced the parental reporting of PCS, replicating the previous finding that female adults or children report higher PCS after mTBI than males do (McNally et al., 2013; Ponsford et al., 2012).

The initial sharp increase of reported PCS after mTBI may reflect acute temporary neurochemical abnormalities following mTBI, including the release of excitatory neurotransmitters, changes in glucose metabolism, and impaired axonal function (Giza & Hovda, 2001). During this initial period, any physical or cognitive activity constitutes an additional neurometabolic demand on the recovering and vulnerable brain (Sady, Vaughan, & Gioia, 2011). Given that a high activity level during recovery is associated with higher symptom ratings and longer symptom duration in students, it might be essential to avoid physical and cognitive over-exertion in the initial recovery phase (Brown et al., 2014; Majerske et al., 2008; Sady et al., 2011). Furthermore, it seems crucial to prevent second head injuries during recovery because repeated head injuries are associated with prolonged recovery and complications (Eisenberg, William Meehan, & Mannix, 2013; Simma, Lütschg, & Callahan, 2013). As a consequence, physical and cognitive activity levels should be managed carefully during the first weeks after an mTBI (Kirkwood et al., 2008; Sady et al., 2011).

One and 4 months after the injury, there were no group differences in the parental rating of PCS between children

Table 4. Correlation coefficients between neuropsychological performance and parent-rated everyday attention problems one month after the injury

	Verbal learning	Verbal memory	Working memory	Switching	Inattention ³	Impulsivity ³
mTBI						
PCS attention ¹	-.33	-.19	-.23	-.15	-.01	.23
PCS attention ²	-.14	-.03	-.11	-.04	-.13	.22
OI						
PCS attention ¹	.21	.19	-.12	-.12	-.16	.08
PCS attention ²	.09	.11	-.12	-.12	-.11	.11

Note. No significant correlations ($p < .05$, Bonferroni corrected) were found.

¹Uncorrected correlations.

²Partial correlations (controlled for gender and pre-injury everyday attention problems).

³For both variables inattention and impulsivity, high percentage values mean a worse attention performance. mTBI: mild traumatic brain injury. OI: orthopedic injury.

Table 5. Correlation coefficients between neuropsychological performance and parent-rated everyday attention problems four months after the injury

	Verbal learning	Verbal memory	Working memory	Switching	Inattention ³	Impulsivity ³
mTBI						
PCS attention ¹	-.56*	-.34	-.48*	-.31	.22	.11
PCS attention ²	-.52*	-.28	-.42	-.26	.19	.10
OI						
PCS attention ¹	.34	.34	-.13	-.13	-.14	-.01
PCS attention ²	.19	.19	-.16	-.05	-.11	.03

Note. * $p < .05$ (Bonferroni corrected).

¹Uncorrected correlations.

²Partial correlations (controlled for gender and pre-injury everyday attention problems).

³For both variables inattention and impulsivity, high percentage values mean a worse attention performance. mTBI: mild traumatic brain injury. OI: orthopedic injury.

after mTBI and OI. Of interest, our data indicated that ongoing emotional symptoms and pre-injury PCS, but not injury group, were significant predictors of ongoing PCS 4 months after the injury, supporting the finding that certain non-injury child and family factors may be associated with persistent PCS, while injury related factors are only significant predictors in the acute period (McNally et al., 2013). Since the contribution of injury characteristics on long-term PCS tends to decline over time (McNally et al., 2013; Ponsford et al., 2012), the appropriateness of the term *post-concussive symptoms*, suggesting a neurological basis, is being questioned for persisting symptoms after the post-acute period (Donlon, & Jones, 2014). Hence, further research is

needed to improve the understanding of the etiology of PCS, specifically regarding the differential influence of injury and non-injury related factors on long-term PCS.

Consistent with previous studies, our results showed that socio-behavioral and neuropsychological outcome were comparable between children after mTBI and OI in the post-acute period (Asarnow et al., 1995; Babikian et al., 2011; Maillard-Wermelinger et al., 2009; Rieger et al., 2013). With the exception of an improved processing speed over time, no further group or time changes were found. Although recent studies (Anderson et al., 2001; Catale et al., 2009; Loher et al., 2014) detected subtle impairments in attention and executive functions compared to healthy control children,

Table 6. Neuropsychological performance (M , SD) at T2 and T3 of children after mTBI without and with pre-injury everyday attention problems

	Absence of pre-injury attention problems ($n = 20$)		Presence of pre-injury attention problems ($n = 20$)		p value
	M	SD	M	SD	
1 month after the injury (T2)					
Verbal learning T2 (PR) ^a	62.53	31.57	35.60	27.20	.007*
Verbal memory T2 (PR) ^a	61.37	30.67	48.45	25.15	.158
Working memory T2 (SS) ^a	97.74	19.11	94.70	13.26	.566
Switching T2 (SS) ^b	10.38	2.83	9.92	3.39	.698
Inattention T2 (PR) ^c	46.49	25.90	55.94	28.24	.290
Impulsivity T2 (PR) ^c	55.09	32.30	62.87	28.90	.439
	Absence of pre-injury attention problems ($n = 20$)		Presence of pre-injury attention problems ($n = 17$)		p value
	M	SD	M	SD	
4 months after the injury (T3)					
Verbal learning T3 (PR) ^a	63.47	30.25	46.88	35.01	.142
Verbal memory T3 (PR) ^a	63.74	26.56	38.47	29.77	.011
Working memory T3 (SS) ^a	102.95	13.68	92.06	13.01	.020
Switching T3 (SS) ^b	11.81	3.23	10.82	3.46	.452
Inattention T3 (PR) ^c	47.06	25.46	53.49	28.16	.470
Impulsivity T3 (PR) ^c	57.83	31.45	63.75	33.41	.583

Note. * $p < .05$ (Bonferroni corrected). PR = percentile; SS = Standard Score; T2 = 1 month following the injury; T3 = 4 months following the injury.

^aDue to language problems, one child of the mTBI group could not be tested with the verbal learning and memory test as well as with the working memory test at both time points.

^bThis test could only be conducted with children aged eight years and older, leading to a smaller n in the single cells (T2: no pre-injury attention problems: $n = 16$, with pre-injury attention problems: $n = 14$; T3: no pre-injury attention problems: $n = 16$, with pre-injury attention problems: $n = 11$).

^cFor inattention and impulsivity, high percentage values mean a worse attention performance.

these group differences, however, seem to disappear when comparing the outcome to another injury group. The finding of a similar cognitive performance between children after mTBI and OI was recently interpreted as *general injury effect* which is hypothesized to originate from systematic pre-injury differences and/or the experience of sustaining an injury among injured children (Babikian et al., 2011). Thus, it seems crucial to compare cognitive outcome after pediatric mTBI with a carefully selected control group. Another injury group enables to separate specific head injury effects from a general injury effect to determine the *pure* effect of the head injury. Moreover, the additional use of healthy children as a second control group would be helpful to better estimate the degree of impairment compared to non-injured children.

Despite the lack of significant group differences concerning neuropsychological outcome in the post-acute period, 4 months after the injury, we found group-specific medium-sized associations between elevated parent-rated everyday attention problems and reduced working memory as well as verbal learning performance in children after mTBI, replicating a recent finding of an adult mTBI study (Dean & Sterr, 2013). Although children after mTBI were not rated as having more everyday attention problems compared to children after OI, their ongoing everyday attention problems were related to their neuropsychological performance, while this association was less pronounced in children after OI.

Moreover, not only persistent, but also pre-injury everyday attention problems negatively influenced neuropsychological outcome in children after mTBI. Thus, our findings are similar to previous studies which have shown that pre-injury risk factors may accentuate weaknesses after pediatric mTBI (Babikian et al., 2013; Bonfield et al., 2013; Fay et al., 2010). Although our exploratory analyses indicated only few moderate-sized effects, our findings suggest that pre-injury attention problems may act as a risk factor for cognitive outcome in children after mTBI. According to the *cognitive reserve capacity* hypothesis (Stern, 2002), pre-injury attention problems might indicate a reduction in cognitive resources, putting children with pre-injury attention problems at a higher risk for cognitive difficulties after a TBI.

Thus, it is highly important to detect pre-injury risk factors like attention problems that might influence cognitive functioning after childhood mTBI. Considering our cognitive outcome, reduced neuropsychological performance was associated with pre-injury as well as with current everyday attention problems. Because pre-injury and follow-up ratings of everyday attention problems were moderately correlated, it may be assumed that most children with attention problems already have shown attention impairments before the injury. Thus, an acute rating of pre-injury attention skills might help at an early time point to select children who are likely to show ongoing attention problems and who are at risk for neuropsychological impairments after mTBI.

The following limitations of this study should be mentioned: First, our sample was relatively small and our mTBI group was very mildly injured, possibly limiting the generalization of our results. Second, our PCS questionnaire

is a translated and slightly adapted version of the PCSI (Gioia et al., 2009), without Swiss norms. However, because the PCSI questionnaire is a valid instrument, with strong psychometric properties (Sady, Vaughan, & Gioia, 2014), we do not expect significant differences when using it with Swiss children, especially when focusing on general PCS, a “simple” measure of PCS that was also investigated in earlier studies (Barlow et al., 2010; Sroufe et al., 2010; Rieger et al., 2013). Nevertheless, we are aware that our general PCS outcome may be simplified since there is evidence that a sum score of PCS covers multiple factors with different trajectories (Ayr et al., 2009; Taylor et al., 2010). Moreover, the use of a non-validated everyday attention subscale is problematic, even though acceptable basic psychometric properties were given. Third, SES data were collected at the first neuropsychological assessment. As a consequence, we could not compare included participants with patients who declined or refused participation to detect systematic demographic differences. Additionally, given that higher physical and cognitive activity during recovery is associated with higher symptom ratings, resulting in a prolonged recovery, it would have been beneficial to have inquired physical and cognitive activity levels following the injury that might have influenced recovery and outcome in our study.

To conclude, our results with a very mild mTBI sample support the assumption that mTBI is a “benign” injury, even in the post-acute period (Babikian et al., 2011; Carroll et al., 2004). Parents of children after mTBI initially observed a sharp increase in the number of PCS, which regressed over the first few weeks. Therefore, treating physicians should inform parents and teachers about typical symptoms that may occur in the first weeks following an injury to help adapting the child’s schedule and activities. Comprehensive information is not only important to prevent excessive physical and cognitive strains after the injury (Sady et al., 2011), but also to minimize parental stress and optimize coping with the incident (Ponsford et al., 2001). Although there were no post-acute neuropsychological or socio-behavioral differences between children after mTBI and OI, our findings indicate that current and pre-injury everyday attention problems were negatively associated with neuropsychological performance in children after mTBI. Thus, future research should focus on further cognitive and socio-emotional risk factors that might influence neuropsychological outcome after pediatric mTBI.

ACKNOWLEDGMENTS

This work was financially supported by following foundations: the Johanna Dürmüller-Bol Foundation, the Batzebär Foundation of the University Children’s Hospital of Bern, and the Anna Müller-Grocholski Foundation. There are no conflicts of interest. Portions of this research were presented at the Mid-Year Meeting of the International Neuropsychological Society (INS) in Amsterdam, 2013 as well as at the Annual Meeting of the INS in Seattle, 2014. We thank all parents and children for participating in this research project. In addition, we thank the Psychology graduate students Venera Gashaj, Sandra Hartmann, and Elisabeth Ruprecht for helping with data collection. Besides, the authors express gratitude to the anonymous Reviewers for their helpful suggestions and thank Thomas Mettler for support with data plotting.

REFERENCES

- Ayr, L. K., Yeates, K. O., Taylor, H. G., & Browne, M. (2009). Dimensions of postconcussive symptoms in children with mild traumatic brain injuries. *Journal of the International Neuropsychological Society, 15*, 19–30. doi:10.1017/S1355617708090188
- Anderson, V., Catroppa, C., Morse, S., Haritou, F., & Rosenfeld, J. (2001). Outcome from mild head injury in young children: A prospective study. *Journal of Clinical and Experimental Neuropsychology, 23*(6), 705–717. doi:10.1076/jcen.23.6.705.1015
- American Congress of Rehabilitation Medicine (1993). Definition of mild traumatic brain injury. *Journal of Head Trauma Rehabilitation, 8*, 86–87.
- Asarnow, R., Satz, P., Light, R., Lewis, R. S., & McCleary, C. (1995). The UCLA study of mild closed head injury in children and adolescents. In S. Broman & M. E. Michel (Eds.), *Traumatic brain injury in children* (pp 117–146). New York: Oxford University Press.
- Babikian, T., & Asarnow, R. (2009). Neurocognitive outcomes and recovery after pediatric TBI: Meta-analytic review of the literature. *Neuropsychology, 23*(3), 283–296. doi:10.1037/a0015268
- Babikian, T., McArthur, D., & Asarnow, R. F. (2013). Predictors of 1-month and 1-year neurocognitive functioning from the UCLA Longitudinal Mild, Uncomplicated, Pediatric Traumatic Brain Injury Study. *Journal of International Neuropsychological Society, 19*(2), 145–154. doi:10.1017/S135561771200104X
- Babikian, T., Satz, P., Zaucha, K., Light, R., Lewis, R. S., Asarnow, R. F. (2011). The UCLA Longitudinal Study of Neurocognitive Outcomes following mild pediatric traumatic brain injury. *Journal of International Neuropsychological Society, 17*, 886–895. doi:10.1017/S1355617711000907
- Barlow, K. M., Crawford, S., Stevenson, A., Sandhu, S. S., Belanger, F., & Dewey, D. (2010). Epidemiology of postconcussion syndrome in pediatric mild traumatic brain injury. *Pediatrics, 126*(2), e374–e381. doi:10.1542/peds.2009-0925
- Bonfield, C. M., Lam, S., Lin, Y., & Greene, S. (2013). The impact of attention deficit hyperactivity disorder on recovery from mild traumatic brain injury. *Journal of Neurosurgery: Pediatrics, 12*, 97–102. doi:10.3171/2013.5.PEDS12424
- Brown, N. J., Mannix, R. C., O'Brien, M. J., Gostine, D., Collins, M. W., & Meehan, W. P. III (2014). Effect of cognitive activity level on duration of post-concussion symptoms. *Pediatrics, 133*, e299–e304. doi:10.1542/peds.2013-2125
- Brown, L., Sherbenou, R. J., & Johnsen, S. K. (1997). *Test of nonverbal Intelligence (TONI-III). A language-free measure of cognitive ability*. Austin: Pro-ed.
- Carroll, L. J., Cassidy, D., Peloso, P. M., Borg, J., von Holst, H., Holm, L., ... Pépin, M. (2004). Prognosis for mild traumatic brain injury: Results of the WHO collaborating centre task force on mild traumatic brain injury. *Journal of Rehabilitation Medicine, 43*(Suppl), 84–105. doi:10.1080/16501960410023859
- Catale, C., Marique, P., Closset, A., & Meulemans, T. (2009). Attentional and executive functioning following mild traumatic brain injury in children using the Test for Attentional Performance (TAP) battery. *Journal of Clinical and Experimental Neuropsychology, 31*(3), 331–338. doi:10.1080/13803390802134616
- Conners, C. K. (2004). *The Conners Continuous Performance Test II (CPT-II)*. Toronto: Multi-Health Systems.
- Dean, P. J., & Sterr, A. (2013). Long-term effects of mild traumatic brain injury on cognitive performance. *Frontiers in Human Neuroscience, 7*, 1–11. doi:10.3389/fnhum.2013.00030
- Delis, D. C., Kaplan, E., & Kramer, J. (2001). *Delis-Kaplan executive function system*. San Antonio: The Psychological Corporation.
- DiScala, C., Lescohier, I., Barthel, M., & Li, G. (1998). Injuries to children with attention deficit hyperactivity disorder. *Pediatrics, 102*, 1415–1421. doi:10.1542/peds.102.6.1415
- Donders, J., & Woodward, H. R. (2003). Gender as a moderator of memory after traumatic brain injury in children. *Journal of Head Trauma Rehabilitation, 18*, 106–115.
- Donlon, K., & Jones, R. T. (2014). Applying a traumatic stress approach to understanding PCS following pediatric mild TBI. *Child Neuropsychology*. doi:10.1080/09297049.2014.944491
- Duff, K., Schoenberg, M. R., Scott, J. G., & Adams, R. L. (2005). The relationship between executive functioning and verbal and visual learning and memory. *Archives of Clinical Neuropsychology, 20*, 111–122. doi:10.1016/j.acn.2004.03.003
- Durkin, M. S., Davidson, L. L., Kuhn, L., O'Connor, P., & Barlow, B. (1994). Low-income neighborhoods and the risk of severe pediatric injury: A small-area analysis in northern Manhattan. *American Journal of Public Health, 84*(4), 587–592.
- Eisenberg, M. A., William Meehan, J. A., & Mannix, R. (2013). Time interval between concussions and symptom duration. *Pediatrics, 132*, 8–17. doi:10.1542/peds.2013-0432
- Epstein, J. N., Erkanli, A., Conners, C. K., Klaric, J., Costello, J. E., & Angold, A. (2003). Relations between continuous performance test performance measures and ADHD behaviors. *Journal of Abnormal Child Psychology, 31*(5), 543–554.
- Farmer, J. E., Kanne, S. M., Haut, J. S., Williams, J., Johnstone, B., & Kirk, K. (2002). Memory functioning following traumatic brain injury in children with premorbid learning problems. *Developmental Neuropsychology, 22*(2), 455–469. doi:10.1207/S15326942DN2202_2
- Fay, T. B., Yeates, K. O., Taylor, H. G., Bangert, B., Dietrich, A., Nuss, K. E., ... Wright, M. (2010). Cognitive reserve as a moderator of postconcussive symptoms in children with complicated and uncomplicated mild traumatic brain injury. *Journal of International Neuropsychological Society, 16*(1), 94–105. doi:10.1017/S1355617709991007
- Feigin, V., Theadom, A., Barker-Collo, S., Starkey, N. J., McPherson, K., Kahan, M., ... Ameratunga, S. (2013). Incidence of traumatic brain injury in New Zealand: A population-based study. *The Lancet, 12*, 53–64. doi:10.1016/S1474-4422(12)70262-4
- Gioia, G. A., Schneider, J. C., Vaughan, C. G., & Isquith, P. K. (2009). Which symptom assessment and approaches are uniquely appropriate for pediatric concussion? *British Journal of Sports Medicine, 43*(Suppl D), 13–22. doi:10.1136/bjmsm.2009.058255
- Gioia, G. A., Vaughan, C. G., & Isquith, P. K. (2011). *Manual for pediatric immediate post-concussion assessment and cognitive testing*. Pittsburgh, PA: ImPACT Applications.
- Giza, C. C., & Hovda, D. A. (2001). The neurometabolic cascade of concussion. *Journal of Athletic Training, 36*(3), 228–235.
- Goodman, R. (1997). The Strengths and Difficulties Questionnaire: A research note. *Journal of Child Psychology and Psychiatry, 38*, 581–586. doi:10.1111/j.1469-7610.1997.tb01545.x
- Heim, S., & Keil, A. (2012). Developmental trajectories of regulating attentional selection over time. *Frontiers in Psychology, 3*, 1–9. doi:10.3389/fpsyg.2012.00277
- Helmsaedter, C., Lendt, M., & Lux, S. (2001). *Verbaler Lern- und Merkfähigkeitstest*. Göttingen: Beltz Test GmbH.
- Kirkwood, M. W., Yeates, K. O., Taylor, H. G., Randolph, C., McCrea, M., & Anderson, V. A. (2008). Management of pediatric mild traumatic brain injury: A neuropsychological review from injury through recovery. *Clinical Neuropsychology, 22*, 769–800. doi:10.1080/13854040701543700

- Kraus, J. F. (1995). Epidemiological features of brain injury in children: Occurrence, children at risk, causes and manner of injury, severity and outcomes. In S. H. Broman & M. E. Michel (Eds.), *Traumatic head injury in children* (pp 22–39). Oxford: Oxford University Press.
- Loher, S., Fatzer, S. T., & Roebbers, C. M. (2014). Executive functions after pediatric mild traumatic brain injury: A prospective short-term longitudinal study. *Applied Neuropsychology: Child*, 3, 103–114. doi:10.1080/21622965.2012.716752
- Maillard-Wermelinger, A., Yeates, K. O., Gerry Taylor, H., Rusin, J., Bangert, B., Dietrich, A., ... Wright, M. (2009). Mild traumatic brain injury and executive functions in school-aged children. *Developmental Neurorehabilitation*, 12(5), 330–341. doi:10.3109/17518420903087251
- Majerske, C. W., Mihalik, J. P., Ren, D., Collins, M. W., Reddy, C. C., Lovell, M. R., ... Wagner, A. K. (2008). Concussion in sports: Postconcussive activity levels, symptoms, and neurocognitive performance. *Journal of Athletic Training*, 43(3), 265–274. doi:10.4085/1062-6050-43.3.265
- McCullagh, S., & Feinstein, A. (2003). Outcome after mild traumatic brain injury: An examination of recruitment bias. *Journal of Neurology, Neurosurgery, & Psychiatry*, 74, 39–43. doi:10.1136/jnnp.74.1.39
- McNally, K. A., Bangert, B., Dietrich, A., Nuss, K. E., Rusin, J., Wright, M., ... Yeates, K. O. (2013). Injury versus non-injury factors as predictors of post-concussive symptoms following mild traumatic brain injury in children. *Neuropsychology*, 27(1), 1–12. doi:10.1037/a0031370
- Menascu, S., & MacGregor, D. L. (2007). Mild closed head injury. In D. L. MacGregor, P. B. Dirks & P. Rumney (Eds.), *Head injury in children and adolescents* (pp. 41–49). London: Mac Keith Press.
- Mittenberg, W., Wittner, M. S., & Miller, L. J. (1997). Postconcussion syndrome occurs in children. *Neuropsychology*, 11(3), 447–452. doi:10.1037/0894-4105.11.3.447
- Petermann, F., & Petermann, U. (2007). *Hamburg-Wechsler-Intelligenztest für Kinder (HAWIK-IV)*. Bern: Huber.
- Pickering, S. J., & Gathercole, S. E. (2001). *Working Memory Test Battery for Children (WMTB)*. London: The Psychological Corporation.
- Ponsford, J., Cameron, P., Fitzgerald, M., Grant, M., Mikocka-Walus, A., & Schönberger, M. (2012). Predictors of postconcussive symptoms 3 months after mild traumatic brain injury. *Neuropsychology*, 26(3), 304–313. doi:10.1037/a0027888
- Ponsford, J., Willmott, C., Rothwell, A., Cameron, P., Ayton, G., Nelms, R., & Ng, K. (2001). Impact of early intervention on outcome after mild traumatic brain injury in children. *Pediatrics*, 108, 1297–1303. doi:10.1542/peds.108.6.1297
- Ramsay, L. J., Moreton, G., Gorman, D. R., Blake, E., Goh, D., Elton, R. A., & Beattie, T. F. (2003). Unintentional home injury in preschool-aged children: Looking for the key - An exploration of the inter-relationship and relative importance of potential risk factors. *Public Health*, 117(6), 404–411. doi:10.1016/S0033-3506(03)00072-6
- Rieger, B. P., Lewandowski, L. J., Callahan, J. M., Spenceley, L., Truckenmiller, A., Gathje, R., & Miller, L. A. (2013). A prospective study of symptoms and neurocognitive outcomes in youth with concussion vs orthopaedic injuries. *Brain Injury*, 27(2), 169–178. doi:10.3109/02699052.2012.729290
- Roebbers, C. M., Röthlisberger, M., Cimeli, P., Michel, E., & Neuenschwander, R. (2011). School enrolment and executive functioning: A longitudinal perspective on developmental changes, the influence of learning context, and the prediction of pre-academic skills. *European Journal of Developmental Psychology*, 8(5), 526–540. doi:10.1080/17405629.2011.571841
- Sady, M. D., Vaughan, C. G., & Gioia, G. A. (2011). School and the concussed youth - recommendations for concussion education and management. *Physical Medicine & Rehabilitation Clinics of North America*, 22(4), 701–719. doi:10.1016/j.pmr.2011.08.008
- Sady, M. D., Vaughan, C. G., & Gioia, G. A. (2014). Psychometric characteristics of the Postconcussion Symptom Inventory in children and adolescents. *Archives of Clinical Neuropsychology*, 29(4), 348–363. doi:10.1093/arclin/acu014
- Satz, P. (1993). Brain reserve capacity on symptom onset after brain injury: A formulation and review of evidence for threshold theory. *Neuropsychology*, 7, 273–295.
- Satz, P., Zaucha, K., McCleary, C., Light, R., Asarnow, R., Becker, D. (1997). Mild head injury in children and adolescents: A review of studies (1970-1995). *Psychological Bulletin*, 122, 107–131. doi:10.1037/0033-2909.122.2.107
- Scherwath, A., Sommerfeldt, D. W., Bindt, C., Nolte, A., Boiger, A., Koch, U., & Petersen-Ewert, C. (2011). Identifying children and adolescents with cognitive dysfunction following mild traumatic brain injury - Preliminary findings on abbreviated neuropsychological testing. *Brain Injury*, 25, 401–408. doi:10.3109/02699052.2011.557351
- Simma, B., Lütschg, J., & Callahan, J. M. (2013). Mild head injury in pediatrics: Algorithms for management in the ED and in young athletes. *American Journal of Emergency Medicine*, 31, 1133–1138. doi:10.1016/j.ajem.2013.04.007
- Smallwood, J., Fishman, D. J., & Schooler, J. W. (2007). Counting the cost of an absent mind: Mind wandering as an under-recognized influence on educational performance. *Psychonomic Bulletin & Review*, 14(2), 230–236.
- Sroufe, N. S., Fuller, D. S., West, B. T., Singal, B. M., Warschausky, S. A., & Maio, R. F. (2010). Postconcussive symptoms and neurocognitive function after mild traumatic brain injury in children. *Pediatrics*, 125, 1331–1339. doi:10.1542/peds.2008-2364
- Stern, Y. (2002). What is cognitive reserve? Theory and research application of the reserve concept. *Journal of the International Neuropsychological Society*, 8, 448–460. doi:10.1017/S1355617701020240
- Taylor, H. G., Dietrich, A., Nuss, K., Wright, M., Rusin, J., Bangert, B., ... Yeates, K. O. (2010). Post-concussive symptoms in children with mild traumatic brain injury. *Neuropsychology*, 24(2), 148–159. doi:10.1037/a0018112
- Williams, D. H., Levin, H. S., & Eisenberg, H. M. (1990). Mild head injury classification. *Neurosurgery*, 27(3), 422–428.
- Yeates, K. O. (2010a). Mild traumatic brain injury and postconcussive symptoms in children and adolescents. *Journal of International Neuropsychological Society*, 16, 953–960. doi:10.1017/S1355617710000986
- Yeates, K. O. (2010b). Traumatic brain injury. In K. O. Yeates, M. D. Ris, H. G. Taylor & B. F. Pennington (Eds.), *Pediatric neuropsychology. Research, theory and practice*. (Vol. 2, pp. 112–146). New York: Guilford Press.
- Yeates, K. O., Armstrong, K., Janusz, J., Taylor, H. G., Wade, S., Stancin, T., & Drotar, D. (2005). Long-term attention problems in children with traumatic brain injury. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(6), 574–584. doi:10.1097/01.chi.0000159947.50523.64
- Yeates, K. O., Taylor, H. G., Rusin, J., Bangert, B., Dietrich, A., Nuss, K., ... Jones, B. L. (2009). Longitudinal trajectories of postconcussive symptoms in children with mild traumatic brain injuries and their relationship to acute clinical status. *Pediatrics*, 123, 735–743. doi:10.1542/peds.2008-1056