

Towards the continuum limit in transport coefficient computations

A. Francis^a, O. Kaczmarek^b, M. Laine^c, M. Müller^b, T. Neuhaus^{*d}, H. Ohno^e

^a *Institute for Nuclear Physics, JGU Mainz, D-55099 Mainz, Germany*

^b *Faculty of Physics, University of Bielefeld, D-33501 Bielefeld, Germany*

^c *Institute for Theoretical Physics, AEC, University of Bern, CH-3012 Bern, Switzerland*

^d *Institute for Advanced Simulation, FZ Jülich, D-52425 Jülich, Germany*

^e *Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

*E-mail: francis@kph.uni-mainz.de, okacz@physik.uni-bielefeld.de,
laine@itp.unibe.ch, mmueller@physik.uni-bielefeld.de,
t.neuhaus@fz-juelich.de, hono@quark.phy.bnl.gov*

The analytic continuation needed for the extraction of transport coefficients necessitates in principle a continuous function of the Euclidean time variable. We report on progress towards achieving the continuum limit for 2-point correlator measurements in thermal SU(3) gauge theory, with specific attention paid to scale setting. In particular, we improve upon the determination of the critical lattice coupling and the critical temperature of pure SU(3) gauge theory, estimating $r_0 T_c \simeq 0.7470(7)$ after a continuum extrapolation. As an application the determination of the heavy quark momentum diffusion coefficient from a correlator of colour-electric fields attached to a Polyakov loop is discussed.

31st International Symposium on Lattice Field Theory LATTICE 2013

July 29 - August 3, 2013

Mainz, Germany

*Speaker.

1. Motivation

Among quantities playing a central role in the theoretical interpretation of heavy ion collision experiments at RHIC and LHC are so-called transport coefficients: shear and bulk viscosities as well as heavy and light quark diffusion coefficients. Because of strong interactions, these quantities need to be determined by numerical lattice Monte Carlo simulations. This represents a challenging problem, given that numerical simulations are carried out in Euclidean signature, whereas transport coefficients are Minkowskian quantities, necessitating an analytic continuation [1]. Nevertheless, the problem is solvable in principle [2], provided that lattice simulations reach a continuum limit and that short-distance singularities can be subtracted [3]. The purpose of this investigation is to probe the practical feasibility of these steps, by approaching the continuum limit for a particular 2-point correlator in pure SU(3) gauge theory, related to heavy-quark diffusion (cf. eq. (4.1)). An important ingredient in reaching the continuum limit is scale setting, so we start by discussing issues related to this topic in secs. 2, 3.

2. Improved determination of the SU(3) phase diagram

When SU(3) lattice gauge theory is discretized according to Wilson's classic prescription, and an infinite spatial-volume limit is taken, the theory contains two parameters: the number of lattice sites in the temporal direction, denoted by N_τ , and the lattice coupling, denoted by β . The phase diagram in the plane (N_τ, β) contains a line of first order transitions; for a given N_τ , the critical point is denoted by β_c . The continuum limit corresponds to $N_\tau \rightarrow \infty$, $\beta_c \rightarrow \infty$.

Because of computational limitations, lattice simulations were historically carried out at small values of N_τ . In fact, previous to our study, β_c had been reliably determined only for $N_\tau = 4 - 12$ [4, 5]; for instance, already at $N_\tau = 16$ the best results available came from a lattice 16×24^3 which does not correspond to the required infinite-volume limit. In addition to simulations, semi-analytic frameworks have recently been developed for estimating β_c [6, 7], however these may contain uncontrolled uncertainties.

We have carried out new simulations at $N_\tau = 10, 12, 14, 16$, in each case with at least two spatial lattice sizes, denoted by N_s , in the range $N_s \gtrsim 3N_\tau$. The critical point β_c is determined from the peak position of the susceptibility related to the Polyakov loop. A unique $\Delta\beta = h \times (N_\tau/N_s)^3$ with $h \approx 0.07$ is employed for infinite spatial-volume extrapolations. For illustration we display in fig. 1(left) our Polyakov loop susceptibility data on a 14×40^3 lattice (at $N_s/N_\tau \approx 2.9$) and their corresponding Ferrenberg-Swendsen reweighting (denoted by the curve in the figure). Using the peak position of this data (the triangle) and a similar data set for a 14×56^3 lattice (at $N_s/N_\tau = 4$) a finite-size extrapolation gives $\beta_c(N_\tau = 14) = 6.4488(59)$.

In fig. 1(right) we display our results for the critical coupling for $N_\tau = 10, 12, 14$ and 16 . Also shown are old results from refs. [4, 5] and the results of semi-analytic computations from refs. [6, 7]. Our results for $N_\tau = 10$ and 12 agree with the old ones within errors whereas for larger N_τ , β_c has now been determined relatively reliably for the first time. Even though our current estimate at $N_\tau = 16$ is preliminary ($\beta_c(N_\tau = 16) \simeq 6.5509(39)$), it can be seen that the semi-analytic calculations miss some of the structure in the data.




Figure 1: Left: Polyakov loop susceptibility $\chi_P \equiv V_{\text{spatial}}(\langle P^2 \rangle - \langle P \rangle^2)$ for a 14×40^3 box in pure SU(3) lattice gauge theory with the Wilson action. Right: Published data for β_c , from Bielefeld (BI) [4] and Lucini et al [5], compared with our new data points at $N_\tau = 10, 12, 14, 16$. Data from Langelage et al [6] and an interpolation to the results of Cheng and Tomboulis [7] represent semi-analytic studies.

3. Conversion of results to physical units

When transport coefficients are measured, then simulations are carried out at a temperature above the critical temperature (denoted by T_c); for a fixed N_τ , this corresponds to $\beta > \beta_c$. For any given β , it is possible to carry out a corresponding zero-temperature simulation, on a lattice N_s^4 , in order to measure some physical quantity. Thereby the temperature can be expressed in terms of a chosen reference scale; from the results of sec. 2, in turn, the reference scale can be determined in terms of T_c . Expressing everything in units of T_c may increase the value of the results, given that T/T_c is a quantity which allows e.g. to compare many related theories.

Various reference scales have been used in the past. The most traditional one is the root of the string tension, denoted by $\sqrt{\sigma}$. The problem is that it is determined from a fit to the large-distance asymptotics of a static potential (or its derivative, the static force), but such fits are delicate, because the correct ansatz needs to be known and also because measurements at large distances are subtle. Quite concretely, some of the numbers cited by various groups, $T_c/\sqrt{\sigma} = 0.630(5)$ [4] versus $T_c/\sqrt{\sigma} = 0.646(3)$ [5], differ by a value much larger than the statistical error.

Another possible scale is the so-called Sommer scale, denoted by r_0 [8]. In this case the scale is determined from the static force at intermediate distances, which removes the need for fitting. The price to pay is that the discreteness of the lattice hampers the measurement, and an interpolation together with tree-level improvement is probably needed for stable results [8]. A value for $r_0 T_c$ has been obtained in ref. [9]: $r_0 T_c = 0.7498(50)$.

Recently a possible new scale was introduced, denoted by $\sqrt{t_0}$ [10]. For any given β , the lattice configuration is ‘‘cooled’’ through a classical Wilson flow until a certain observable reaches a prescribed value, at time t_0 . There is no fitting involved, and no interpolation; t_0 therefore probably suffers from less systematic uncertainties than the other scale choices.




Figure 2: Left: Continuum extrapolation for $r_0 T_c$. The conversion from β_c to r_0/a is based on refs. [11, 12] and additional new simulations, together with a rational fit from ref. [13] (inset). The result $r_0 T_c = 0.7470(7)$ can be contrasted with $r_0 T_c = 0.7498(50)$ from ref. [9]. (For comparisons with perturbation theory, $r_0 \Lambda_{\overline{\text{MS}}} = 0.602(48)$ from ref. [14] yields $T_c/\Lambda_{\overline{\text{MS}}} = 1.24(10)$; ref. [15] suggests $r_0 \Lambda_{\overline{\text{MS}}} \simeq 0.637(32)$ which would yield $T_c/\Lambda_{\overline{\text{MS}}} \simeq 1.17(6)$.) Right: Preliminary results for $\sqrt{t_0} T_c$ as a function of a^2/t_0 .

Considering first the Sommer scale, the data of fig. 1(right) can be used for determining the dimensionless combination $r_0 T_c$; the results are shown in fig. 2(left). The results display the expected $O(a^2)$ behaviour and can be extrapolated to the continuum. We obtain $r_0 T_c \simeq 0.7470(7)$ which agrees with an earlier result from ref. [9] but displays a much reduced error.

We also have preliminary estimates for $\sqrt{t_0} T_c$, obtained at different lattice extensions from zero-temperature simulations at $\beta = \beta_c(N_\tau = 4, 6, 8, 12)$, and shown in fig. 2(right). Increasing the statistics and number of lattice ensembles hopefully yields accurate results for $\sqrt{t_0} T_c$.

4. Application to heavy quark diffusion

Heavy quarks carry a colour charge and, whenever there are gauge fields present, are subject to a coloured Lorentz force, which adjusts their velocities to those corresponding to kinetic equilibrium (this corresponds to the physics of diffusion). Through linear response theory the effectiveness of the adjustment can be related to a “colour-electric correlator” [16, 17],

$$G_E(\tau) \equiv -\frac{1}{3} \sum_{i=1}^3 \frac{\left\langle \text{Re Tr} \left[U\left(\frac{1}{T}; \tau\right) gE_i(\tau, \mathbf{0}) U(\tau; 0) gE_i(0, \mathbf{0}) \right] \right\rangle}{\left\langle \text{Re Tr} \left[U\left(\frac{1}{T}; 0\right) \right] \right\rangle}, \quad (4.1)$$

where gE_i denotes the colour-electric field, T the temperature, and $U(\tau_2; \tau_1)$ a Wilson line in the Euclidean time direction. A discretized version of this correlator is shown in fig. 3.

Preliminary lattice measurements of the colour-electric correlator have already been carried out [18, 19, 20]. The results look promising, hinting at a large and phenomenologically interesting non-perturbative effect in the diffusion coefficient. However, none of these results contain a systematic continuum extrapolation. The ultimate goal of our investigation is to perform one, by making use of the results of secs. 2, 3 as well as of new simulations.




Figure 3: Left: Fat links, thin links and electric fields along the time direction (cf. the text). Right: Run parameters. The values of r_0T are obtained through an interpolation/extrapolation as illustrated in fig. 2(left). With the value of r_0T_c from fig. 2(left), we have $T/T_c \approx 1.43$ for $N_\tau = 36$.

Our numerical investigations are based on standard lattice QCD techniques. We employ the Wilson gauge action on fine isotropic lattices with lattice spacings down to ~ 0.015 fm. The measurement of eq. (4.1) is performed on gauge field configurations generated using 500 sweeps between subsequent configurations, guaranteeing that the configurations are statistically independent. However the correlation function in eq. (4.1) decreases rapidly with τ , and for $\tau \sim \frac{1}{2T}$ suffers from a weak signal-to-noise ratio. In order to tackle this problem we use a multi-level update [21, 22] for the part of the operator that includes the electric field insertions, and link-integration (“PPR”) [23, 24] for the straight lines between them (the “fat links” in fig. 3). The table in fig. 3 summarizes the situation with N_{conf} labelling the number of statistically independent configurations and N_{stat} the number of additional “multilevel” updates. As demonstrated in fig. 4, these techniques suffice to yield a signal.

The lattice sizes of our simulations, corresponding to a temperature $T \sim 1.43T_c$, substantially exceed those of earlier simulations [18, 19, 20], which had $V_{\text{max}} = 24 \times 64^3$. Due to the increased number of multilevel updates we focussed on lattices with an aspect ratio $N_s/N_\tau = 4$ (cf. the table in fig. 3), which was the optimal choice given the available computational resources. However, volume-scaling checks have been performed at the smallest values of N_τ .

After tree-level improvement [8, 22] our measurements yield a correlator denoted by $G_{\text{imp}}(\tau)$, which is furthermore multiplied by a perturbative renormalization factor Z_{pert} [20]. Normalizing the resulting correlator to

$$G_{\text{norm}}(\tau T) \equiv \pi^2 T^4 \left[\frac{\cos^2(\pi\tau T)}{\sin^4(\pi\tau T)} + \frac{1}{3\sin^2(\pi\tau T)} \right], \quad (4.2)$$

the data are displayed in fig. 5. They exhibit a clear enhancement over the next-to-leading order (NLO) prediction from ref. [25]. A continuum extrapolation remains to be carried out.

5. Outlook

The Euclidean correlator of eq. (4.1) is related to a corresponding spectral function ρ_E through $G_E(\tau) = \int_0^\infty \frac{d\omega}{\pi} \rho_E(\omega) \frac{\cosh(\frac{1}{2}\tau T) \frac{\omega}{T}}{\sinh \frac{\omega}{2T}}$. Once a perturbatively determined short-distance divergence is subtracted from a continuum-extrapolated $G_E(\tau)$, the remainder may be subjected to an analytic continuation algorithm [2, 3] or a well-motivated model like in refs. [19, 20]. In particular, a “momentum diffusion coefficient”, often denoted by κ , can be obtained from $\kappa = \lim_{\omega \rightarrow 0} \frac{2T\rho_E(\omega)}{\omega}$. In




Figure 4: Left: Error reduction for $G_E(\tau = 12a)$ in a 36×144^3 box with techniques described in the text. For 77 statistically independent configurations we determine the non-improved observable (x -axis) and the fully improved one (y -axis). A dotted horizontal line and a triangle mark the average of the improved observable. Right: The statistically improved correlator $G_E(\tau)$. The inset shows the ratio of non-improved over improved statistical errors, denoted by R_1 . An error reduction by a factor $R_1 \sim \mathcal{O}(10^4)$ can be achieved.

the non-relativistic limit (i.e. for $M \gg \pi T$, where M stands for a heavy quark mass) the corresponding “diffusion coefficient” is given by $D = 2T^2/\kappa$. It will be interesting to see whether preliminary estimates of the diffusion coefficient [19, 20] can be confirmed after a continuum limit, and how well the results perform in phenomenological comparisons with LHC heavy ion data (cf. e.g. ref. [26]). Of course, taking the continuum limit plays an important role in the extraction of the light-quark diffusion coefficient and electrical conductivity as well [27, 28].

Acknowledgment

This work has been supported in part by the DFG under grant GRK881, by the SNF under grant 200021-140234, and by the European Union through HadronPhysics3 and ITN STRONGnet (grant 238353). Simulations were performed using JARA-HPC resources at the RWTH Aachen (project JARA0039), JUDGE/JUOPA at the JSC Jülich, the OCuLUS Cluster at the Paderborn Center for Parallel Computing, and a GPU cluster at Bielefeld.

References

- [1] H.B. Meyer, Eur. Phys. J. A 47 (2011) 86 [1104.3708].
- [2] G. Cuniberti *et al*, Commun. Math. Phys. 216 (2001) 59 [cond-mat/0109175].
- [3] Y. Burnier, M. Laine and L. Mether, Eur. Phys. J. C 71 (2011) 1619 [1101.5534].
- [4] B. Beinlich, F. Karsch, E. Laermann and A. Peikert, Eur. Phys. J. C 6 (1999) 133 [hep-lat/9707023].
- [5] B. Lucini, M. Teper and U. Wenger, JHEP 01 (2004) 061 [hep-lat/0307017].
- [6] J. Langelage *et al*, JHEP 02 (2011) 057 [Erratum-ibid. 07 (2011) 014] [1010.0951].




Figure 5: G_E at $T \sim 1.43T_c$. For the NLO result we have varied $T_c/\Lambda_{\overline{MS}} \in (1.11, 1.34)$, cf. fig. 2.

- [7] X. Cheng and E.T. Tomboulis, Phys. Rev. D 86 (2012) 074507 [1206.3816].
- [8] R. Sommer, Nucl. Phys. B 411 (1994) 839 [hep-lat/9310022].
- [9] S. Necco, Nucl. Phys. B 683 (2004) 137 [hep-lat/0309017].
- [10] M. Lüscher, JHEP 08 (2010) 071 [1006.4518].
- [11] M. Guagnelli *et al.* [ALPHA Collaboration], Nucl. Phys. B 535 (1998) 389 [hep-lat/9806005].
- [12] S. Necco and R. Sommer, Nucl. Phys. B 622 (2002) 328 [hep-lat/0108008].
- [13] S. Dürr, Z. Fodor, C. Hoelbling and T. Kurth, JHEP 04 (2007) 055 [hep-lat/0612021].
- [14] S. Capitani, M. Lüscher, R. Sommer and H. Wittig, Nucl. Phys. B 544 (1999) 669 [hep-lat/9810063].
- [15] N. Brambilla *et al.*, Phys. Rev. Lett. 105(2010)212001 [Erratum-ibid. 108(2012)269903] [1006.2066].
- [16] J. Casalderrey-Solana and D. Teaney, Phys. Rev. D 74 (2006) 085012 [hep-ph/0605199].
- [17] S. Caron-Huot, M. Laine and G.D. Moore, JHEP 04 (2009) 053 [0901.1195].
- [18] H.B. Meyer, New J. Phys. 13 (2011) 035008 [1012.0234].
- [19] D. Banerjee, S. Datta, R. Gavai and P. Majumdar, Phys. Rev. D 85 (2012) 014510 [1109.5738].
- [20] A. Francis, O. Kaczmarek, M. Laine and J. Langelage, PoS LATTICE 2011 (2011) 202 [1109.3941].
- [21] M. Lüscher and P. Weisz, JHEP 09 (2001) 010 [hep-lat/0108014].
- [22] H.B. Meyer, Phys. Rev. D 76 (2007) 101701 [0704.1801].
- [23] G. Parisi, R. Petronzio and F. Rapuano, Phys. Lett. B 128 (1983) 418.
- [24] P. de Forcrand and C. Roiesnel, Phys. Lett. B 151 (1985) 77.
- [25] Y. Burnier, M. Laine, J. Langelage and L. Mether, JHEP 08 (2010) 094 [1006.0867].
- [26] W.M. Alberico *et al.*, Eur. Phys. J. C 73 (2013) 2481 [1305.7421].
- [27] H.-T. Ding *et al.*, Phys. Rev. D 83 (2011) 034504 [1012.4963].
- [28] Y. Burnier and M. Laine, Eur. Phys. J. C 72 (2012) 1902 [1201.1994].