

Enantioselective Determination of Ondansetron and 8-Hydroxyondansetron in Human Plasma from Recovered Surgery Patients by Liquid Chromatography–Tandem Mass Spectrometry

F. Musshoff¹, B. Madea¹, F. Stüber^{2,3}, and U.M. Stamer^{2,3}

¹Institute of Forensic Medicine, University of Bonn, Stiftsplatz 12, 53111 Bonn, Germany; ²Department of Anesthesiology and Pain Therapy, Inselspital, University of Bern, Freiburgstr., 3010 Bern, Switzerland; and ³Department of Anesthesiology and Intensive Care Medicine, University of Bonn, Sigmund-Freud-Str. 25, 53105 Bonn, Germany

Abstract


A liquid chromatographic–mass spectrometric assay with atmospheric pressure chemical ionization for quantification of ondansetron and its main metabolite 8-hydroxyondansetron in human plasma was presented. The enantiomeric separation was achieved on a Chiralcel OD-R column containing cellulose tris-(3,5-dimethylphenylcarbamate). The validation data were within the required limits. The assay was successfully applied to authentic plasma samples. Quantitative results from postoperative patients receiving ondansetron demonstrated a great interindividual variability in postoperative plasma drug concentrations, the metabolites were not detected in their unconjugated form. A wide variation in the *S*-(+)-/*R*-(−)-ondansetron concentration ratio between 0.14 and 7.18 is indicative for a stereoselective disposition or metabolism. In further studies CYP2D6 and CYP3A4 genotype dependent metabolism of ondansetron enantiomers as well as of co-administered drugs and clinical efficacy of the medication should be tested.

Introduction


Ondansetron, (±)-9-methyl-3-[(2-methylimidazol-1-yl)methyl]-2,3-dihydro-1*H*-carbazol-4-one (Figure 1), is a highly selective and potent 5-hydroxytryptamine type 3 (5-HT₃) receptor antagonist. It is effective in the treatment of postoperative nausea and vomiting (PONV) as well as during cancer chemotherapy and radiotherapy, and it has reported anxiolytic and neuroleptic properties (1–4).

Ondansetron has a short plasma half-life of approximately 3 to 5 h (5,6). Between 70% and 76% of ondansetron is protein bound, and it is extensively metabolized in the liver by the

hydroxylation of the indole ring to 7- and mainly 8-hydroxy-ondansetron (40%) followed by glucuronide or sulfate conjugation (7); 6-hydroxylation and demethylation are minor routes of metabolism. In terms of overall ondansetron elimination, CYP3A4 plays the predominant role, but primary hydroxylation is also done via CYP2A6 and 1A2 (8–10). Approximately 5% is excreted unchanged in the urine. Although some of the non-



Ondansetron (stereocenter marked with *)



8-Hydroxyondansetron

Figure 1. Structures of ondansetron and its main metabolite 8-hydroxy-ondansetron.

conjugated metabolites have pharmacological activity, their plasma concentrations are probably too low to contribute to the biological activity of ondansetron.

In pharmacokinetic studies ondansetron concentration was measured using achiral procedures. However, ondansetron is usually be marketed as a racemic mixture of the *R*-(-)- and *S*-(+)-enantiomers. Considering the parallel rightward displacement of the 2-methyl-5-HT concentration response curve at the longitudinal smooth muscle of the Guinea-pig ileum, the potency of *R*-(-)-ondansetron was approximately eightfold higher than that of *S*-(+)-ondansetron (11). Also, administration of *R*-(-)-ondansetron avoided adverse effects including headache, constipation, and increases in transaminase levels, which are associated with the administration of racemic ondansetron (12).

Besides a number of achiral procedures for the measurement of ondansetron in plasma samples (13–17), only a few chiral methods have been reported for the determination of *R*-(-)- and *S*-(+)-ondansetron up to now. These include high performance capillary electrophoresis with heptakis-(2,6-di-O-methyl)- β -cyclodextrin as a mobile phase modifier (18) as well as high-performance liquid chromatography coupled with ultraviolet-spectroscopy (HPLC–UV) using a Chiralcel OD column (19) or a Chiralcel OD-R column (20). Recently a liquid chromatographic–tandem mass spectrometric (LC–MS–MS) procedure using an ovomucoid column was developed for the enantioselective measurement of ondansetron in human plasma samples (21).

The aim of the present work was the development of a chiral LC–MS–MS procedure for the quantitative determination of *R*-(-)- and *S*-(+)-ondansetron together with *R*-(-)- and *S*-(+)-8-hydroxyondansetron in postoperative patients receiving an antiemetic medication to proof possible stereoselective pharmacokinetic differences between both enantiomers.

Experimental

Chemicals

The (+)- and (-)-enantiomers of ondansetron and 8-hydroxyondansetron were a gift of GlaxoSmithKline (München, Germany). Racemic propanolol was purchased from LGC Promochem™ (Wesel, Germany). Triethylamine (TEA) was obtained from Fluka (Neu-Ulm, Germany), ethanol absol. and sodium hydroxide were from Merck (Darmstadt, Germany), and 95% *n*-hexane and dichloromethane were from J.T. Baker (Deventer, Netherlands).

Standard solutions and calibration samples

A methanolic stock solution containing *R*-(-)- and *S*-(+)-ondansetron and *R*-(-)- and *S*-(+)-8-hydroxyondansetron as well as a stock solution containing (\pm)-propanolol were prepared in methanol in concentrations of 10 μ g/mL. All solutions were stored at -20°C. For calibration mixed spiked plasma samples were created in following concentrations for each enantiomer: 2.5, 10, 25, 50, 75, 100, 200, and 250 ng/mL.

Sample preparation

A mixture of 0.2-mL plasma sample, internal standard solution (10 μ L of propanolol in a concentration of 10 μ g/mL methanol), and 0.2 mL of borate buffer (pH 9) was extracted with 1 mL dichloromethane. After centrifugation (4000 $\times g$ for 8 min), the organic phase was evaporated to dryness under a stream of nitrogen at 50°C. The residue was dissolved in 0.1 mL of HPLC mobile phase A, and a 10- μ L aliquot was used for chromatography.

Chromatographic and MS equipment and conditions

An LC–MS–MS system consisting of an Agilent (Waldbronn, Germany) 1100 HPLC system (binary pump, degasser, and autosampler) coupled with an Applied Biosystems (Darmstadt, Germany) API 2000 triple-quadrupole MS was used for analysis performing atmospheric pressure chemical ionization (LC–APCI–MS). The enantiomeric separation was achieved on a Chiralcel OD-R column (10 μ m, 250 \times 46 mm; Daicel Chemical Industries, Illkirch, France) containing cellulose tris-(3,5-dimethylphenylcarbamate) as chiral selector. LC separation was performed using mobile phase A (water/acetonitrile, 98:2, 5 mM ammonium formate) and mobile phase B (water/acetonitrile, 10:90, 5 mM ammonium formate; pH 3.5) in a gradient program with a total flow of 400 μ L/min (binary system): 0–15 min: 85% A; 15–16 min: 85% A \rightarrow 0% A; 16–21 min: 0% A; 21–22 min: 0% A \rightarrow 85% A; 22–30 min: 85% A. From the molecular ions ($[M+H]^+$) following transitions were detected in positive multiple reaction monitoring: ondansetron m/z 293.8 \rightarrow 170.0, 184.0; 8-hydroxyondansetron m/z 309.9 \rightarrow 185.9, 199.8; propanolol (IS) m/z 259.95 \rightarrow 116.1, 182.9 (quantitation ions underlined).

For quantification, peak-area ratios of the analytes to the internal standard were calculated as a function of the concentration of the substances.

Validation protocol

Selectivity. For evaluation of method selectivity, blank plasma samples from different sources were prepared as described, but without adding any analyte or internal standard mix ($n = 6$). Additionally, six samples were analyzed from patients receiving the same medication with metoclopramide instead of ondansetron. Furthermore, blank samples were analyzed to check the absence of analyte ions in the respective peaks of the internal standard ($n = 2$).

Linearity. For calibration, plasma samples were spiked with the *R*-(-)- and *S*-(+)-enantiomers of ondansetron and 8-hydroxyondansetron in concentrations from 2.5 to 250 ng/mL and processed as described ($n = 6$). Using mean values, calibration curves were checked for variance homogeneity (F-Test) and for linearity (Mandel-Test).

Sensitivity. The LOD was defined as the analyte concentration at which the signal-to-noise ratio was 3. The LOQ was defined as three times the LOD.

Precision data and recoveries. Additionally, spiked quality control (QC) plasma samples (high = 100 ng/mL, middle = 50 ng/mL, and low = 10 ng/mL) were prepared and analyzed in each series. Absolute recoveries were calculated comparing the peak areas of all analytes after extraction with the results

after injection of the same concentration (50 ng/mL) directly without extraction, which was defined as 100%.

Matrix effects. Matrix-related ionization effects were evaluated by comparing analyte responses of post-extraction spiked samples to those of spiked samples representing 100% recovery. Five replicates at low (10 ng/mL) and high (100 ng/mL) concentration were analyzed.

Stability studies. For the evaluation of the enantiomeric stability, blank plasma samples were spiked with 50 ng/mL (each enantiomer in single tests) and incubated at 25°C for 24 h, then extracted and analyzed as described. Furthermore long-term stability of all analytes was tested at -20°C for three months as well as the autosampler stability after extraction (ready-to-inject samples) at 20°C for 24 h. The freeze/thaw stability was evaluated after three complete freeze/thaw cycles on consecutive days. The analytes were considered stable when 85–115% of the initial concentration was found.

Application of the method

After approval by the local ethics committee and written informed consent plasma concentrations of the *R*-(-) and *S*-(+)-enantiomers of ondansetron as well as of 8-hydroxyondansetron were analyzed in patients recovering from major abdominal surgery. General anesthesia was conducted using a standardized protocol: 2 mg/kg propofol, 0.2 mg fentanyl, and *cis*-atracurium for induction and remifentanyl, isoflurane, and *cis*-atracurium for maintenance of anesthesia. About 30 min before termination of anesthesia, 4 mg ondansetron was given i.v. followed by an analgesic loading dose of dipyrone 1 g and piritramid 0.08 mg/kg.

Blood samples were obtained with ethylenediaminetetraacetic acid (EDTA) as anticoagulant between 30 and maximal 180 min after application of ondansetron. Whole blood was centrifuged and plasma was separated and frozen until analysis at -80°C.

Results and Discussion

The method's selectivity was tested by comparing the chromatograms of six different blank plasma samples with corresponding spiked plasma. Also, two blank samples were analyzed to check the absence of analyte ions in the respective peaks of the internal standard. Additionally, six samples were analyzed from patients recovering from major abdominal surgery and receiving the same medication with metoclopramide instead of ondansetron. No interferences due to matrix or other potentially interfering compounds were found.




Figure 2. Chromatogram of a blank sample (A), a standard sample containing 100 ng/mL ondansetron and 8-hydroxyondansetron enantiomers (B), and finally of an authentic plasma sample containing *S*-(+)-ondansetron (55.2 ng/mL) and *R*-(-)-ondansetron (45.3 ng/mL) (C).

A complete enantiomeric separation of the *R*-(-)- and *S*-(+)-enantiomers of ondansetron and the racemic internal standard was obtained on a cellulose tris-3,5-dimethyl-phenylcarbamate OD-R column using an ammonium formate buffer/acetonitrile as mobile phase. Figure 2 shows chromatograms of a blank sample, an extracted standard sample with the *R*-(-)- and *S*-(+)-enantiomers of ondansetron and 8-hydroxyondansetron, and at least an authentic plasma sample extract only containing racemic ondansetron. The order of elution of the enantiomers was determined by separately injecting solutions of each enantiomer. The linear regression analysis yielded following equations:

$$\begin{aligned} R(-)\text{-ondansetron:} & \quad y = 0.106x + (-1.49) \quad (r = 1.0000) \\ S(+)\text{-ondansetron:} & \quad y = 0.0555x + (-0.45) \quad (r = 0.9998) \\ R(-)\text{-8-hydroxyondansetron:} & \quad y = 0.0164x + (-0.254) \quad (r = 0.9986) \\ S(+)\text{-8-hydroxyondansetron:} & \quad y = 0.0342x + (0.148) \quad (r = 0.9991) \end{aligned}$$

The LODs of *R*-(-)- and *S*-(+)-ondansetron were 0.5 and 0.7 ng/mL with LOQs of 1.5 and 2.1 ng/mL. The LOD of *R*-(-)- and *S*-(+)-8-hydroxyondansetron was 0.6 ng/mL with an LOQ of 1.8 ng/mL. Intraassay ($n = 6$) and interassay precision ($n = 10$; two calibration curves during the study) showed acceptable values with relative standard deviations (R.S.D.) of less than 10 % for the high and middle control and less than 20% for the low control. For the ondansetron and 8-hydroxyondansetron enantiomers the recovery was between 85 and 91% ($n = 5$). Ion suppression and enhancement from plasma matrix was negligible under present conditions (84–95%). The enantiomers and metabolites were found to be stable in plasma at -20°C for 3 months, in the mobile phase at 20°C for 24 h, and during three freeze/thaw cycles. No chiral inversion was observed at 25°C for 24 h during storage, processing, and analysis. To sum up validation data were within international required limits and the assay was successfully applied to authentic plasma samples.

The analysis of 151 samples from patients routinely receiving intravenous 4 mg racemic ondansetron revealed positive results for *R*-(-)- and *S*-(+)-ondansetron in all analyzed samples. Without hydrolysis the main metabolite 8-hydroxyondansetron was not detected in any of the analyzed samples. Mean and median plasma concentrations, the concentration ranges, and

the ratios *S*-(+)-/*R*-(-)-ondansetron are described in Table I.

These present results from postoperative patients receiving ondansetron for postoperative treatment of PONV demonstrated a great interindividual variability in drug concentrations and, therefore, a wide therapeutic concentration range; the body mass index was not helpful to understand such differences in plasma concentrations. The measured concentrations of both ondansetron enantiomers (sums) are in the range described by others (13,22). According to previous reports free 8-hydroxyondansetron was not found in any case with an LOD of 0.6 ng/mL; a hydrolysis prior to analysis was not performed in the present study.

In general, large differences have been observed between individuals in the capacity and speed of drug metabolism. This variability may be due to concurrent diseases (e.g., renal failure, impaired liver function), co-medications, environmental pollutants that influence the enzymatic function of specific isoenzymes as well as other factors. Further well-recognized variables are genetic polymorphisms of metabolizing enzymes or targets of drug action which can have a significant impact on pharmacokinetic/pharmacodynamic of respective drugs. Genetic variants of the cytochrome P450 (CYP) enzymes play a major role in pharmacokinetic variability and can affect outcome in drug therapy. Individuals carrying alleles reducing enzyme activity may have a phenotype that may metabolize a given drug poorly [poor metabolizer (PM)] or at an intermediate level [intermediate metabolizer (IM)] compared to extensive metabolizers (EM) carrying two wild-type alleles. A gene duplication or multiduplication can result in largely increased enzyme activity resulting in the ultrarapid metabolizer (UM) phenotype.

Considering ondansetron, there was no difference in the AUC, C_{\max} , and $t_{1/2}$ in healthy subjects between EMs ($n = 6$) and PMs ($n = 6$) receiving a single dose (8 mg i.v.) (23). Otherwise, it was demonstrated that CYP2D6 UMs experienced more vomiting compared to EMs or PMs when given ondansetron in the treatment of chemotherapy-induced nausea and vomiting (24). But this difference was predictably more pronounced for topisetron than ondansetron because topisetron is primarily dependent on the CYP2D6 isoenzyme for metabolism. In a further study of 250 patients undergoing standardized general anesthesia UMs had increased therapeutic failure and a reduced response to ondansetron 4 mg (25). However, plasma concentrations were not determined in this study.

Data on enantiomeric disposition of ondansetron are sparse. Otherwise, stereoselective variations could be of interest in individual cases because of different pharmacodynamic properties of both enantiomers. In general a stereoselectivity can be assumed in the body distribution, the renal clearance, and also in the hepatic metabolism of drugs.

In a stereoselective pharmacokinetic study involving four healthy male Chinese, Liu et al. (21) found that the concentration of *R*-(-)-ondansetron was higher than of *S*-(+)-ondansetron at all the time points measured, implying a stereoselective disposition of ondansetron enantiomers. In contrast to this study, the present investigation enrolling a large collective of patients in a European hospital revealed interindividual differences in the concentration ratio of *S*-(+)-/*R*-(-)-

Table I. Plasma Concentrations of *S*-(+)- and *R*-(-)-Ondansetron in 151 Plasma Samples from Postoperative Patients

	<i>S</i> -(+)- Ondansetron (ng/mL)	<i>R</i> -(-)- Ondansetron (ng/mL)	Sum <i>S</i> -(+)- + <i>R</i> -(-)- Ondansetron (ng/mL)	Ratio <i>S</i> -(+)-/ <i>R</i> -(-)- Ondansetron
Mean	38.70	24.93	63.63	2.60
SD	11.59	21.21	22.57	1.52
Median	39.10	16.00	56.10	3.06
Min	8.04	4.15	33.95	0.14
Max	82.20	84.30	157.50	7.18

ondansetron between 0.14 and 7.18. These different findings could be explained by different study designs, volunteers without any co-medication versus patients receiving various drugs for anesthesia and suffering in part from diverse comorbidity, as well as varying geographic distributions and frequencies of genetic variants in populations from different regions of the world (26).

Considering these preliminary results, CYP2D6 and CYP3A genotype dependent metabolism of ondansetron enantiomers might be of specific interest, because effects on efficacy of antiemetic treatment with ondansetron might be possible. Meanwhile, CYP dependent metabolism of a large amount of frequently used drugs is well described (27) with numerous reports describing lacking efficacy, side effects, or even toxicity of individual drugs. Multiple interactions in pharmacokinetics resulting in adverse drug reactions (ADRs) and outcome in drug treatment can be assumed (28).

In general, pharmacogenetics can help to individualize pharmacotherapy and to improve care by predicting the optimal dose and avoiding side effects and toxicity in individual patients (29). Screening for variations in the expression of drug metabolizing enzymes has been suggested as a potential tool for improving therapy regimes (30,31). Further candidate genes like receptors, transporters and other targets of pharmacotherapy are currently under investigation. Aspects of genetic differences influencing efficacy, side effects and adverse outcome of pharmacotherapy will be of importance for future medication regimes.

Conclusions

A quantitative determination of ondansetron in human plasma by enantioselective LC–MS–MS was achieved on a Chiralcel OD-R column containing cellulose tris-(3,5-dimethylphenylcarbamate). The method was fully validated and successfully applied to authentic plasma samples. Quantitative results from postoperative patients receiving ondansetron demonstrated a great interindividual variability in plasma drug concentrations and, therefore, a wide therapeutic concentration range. The main metabolite 8-hydroxyondansetron was not detected in its unconjugated form. A wide variation in the *S*-(+)-/*R*-(–)-ondansetron concentration ratio is indicative for a stereoselective disposition or metabolism. In further studies, CYP-dependent metabolism of ondansetron enantiomers and clinical efficacy of the medication should be tested.

Acknowledgments

The authors extend special thanks to Dr. J. Dethling (Glaxo-SmithKline, Munich, Germany) for providing the reference substances of ondansetron enantiomers and metabolites and to Makbule Kobilay and Joerg Bayer for excellent technical assistance. The study was supported in part by a grant of the Richard-Sackler Research Foundation.

References

1. J.H. Ye, R. Ponnudurai, and R. Schaefer. Ondansetron: a selective 5-HT(3) receptor antagonist and its applications in CNS-related disorders. *CNS Drug Rev.* **7**: 199–213 (2001).
2. K.Y. Ho and T.J. Gan. Pharmacology, pharmacogenetics, and clinical efficacy of 5-hydroxytryptamine type 3 receptor antagonists for postoperative nausea and vomiting. *Curr. Opin. Anaesthesiol.* **19**: 606–611 (2006).
3. M. Aapro. 5-HT(3)-receptor antagonists in the management of nausea and vomiting in cancer and cancer treatment. *Oncology* **69**: 97–109 (2005).
4. T.J. Gan. Selective serotonin 5-HT3 receptor antagonists for postoperative nausea and vomiting: are they all the same? *CNS Drugs* **19**: 225–238 (2005).
5. GlaxoSmithKline, Zofran (ondansetron hydrochloride) injection (prescribing information), 2006.
6. K.H. Simpson and F.M. Hicks. Clinical pharmacokinetics of ondansetron. A review. *J. Pharm. Pharmacol.* **48**: 774–781 (1996).
7. D.A. Saynor and C.M. Dixon. The metabolism of ondansetron. *Eur. J. Cancer Clin. Oncol.* **25**(Suppl. 1): S75–S77 (1989).
8. V. Fischer, A.E. Vickers, F. Heitz, S. Mahadevan, J.P. Baldeck, P. Minery, and R. Tynes. The polymorphic cytochrome P-4502D6 is involved in the metabolism of both 5-hydroxytryptamine antagonists, tropisetron and ondansetron. *Drug Metab. Dispos.* **22**: 269–274 (1994).
9. P. Sanwald, M. David, and J. Dow. Characterization of the cytochrome P450 enzymes involved in the in vitro metabolism of dolasetron. Comparison with other indole-containing 5-HT3 antagonists. *Drug Metab. Dispos.* **24**: 602–609 (1996).
10. C.M. Dixon, P.V. Colthup, C.J. Serabjit-Singh, B.M. Kerr, C.C. Boehlert, G.R. Park, and M.H. Tarbit. Multiple forms of cytochrome P450 are involved in the metabolism of ondansetron in humans. *Drug Metab. Dispos.* **23**: 1225–1230 (1995).
11. A. Butler, J.M. Hill, S.J. Ireland, C.C. Jordan, and M.B. Tyers. Pharmacological properties of GR38032F, a novel antagonist at 5-HT3 receptors. *Br. J. Pharmacol.* **94**: 397–412 (1988).
12. W.J. James. U.S. patent 5470868. Methods for treating emesis and nausea using optically pure R(+)-ondansetron, 1993.
13. P.V. Colthup and J.L. Palmer. The determination in plasma and pharmacokinetics of ondansetron. *Eur. J. Cancer Clin. Oncol.* **25**(Suppl. 1): S71–S74 (1989).
14. M. Depot, S. Leroux, and G. Caille. High-resolution liquid chromatographic method using ultraviolet detection for determination of ondansetron in human plasma. *J. Chromatogr. B Biomed. Sci. Appl.* **693**: 399–406 (1997).
15. X. Xu, M.G. Bartlett, and J.T. Stewart. Determination of ondansetron and its hydroxy metabolites in human serum using solid-phase extraction and liquid chromatography/positive ion electrospray tandem mass spectrometry. *J. Mass Spectrom.* **35**: 1329–1334 (2000).
16. S. Bauer, E. Stormer, R. Kaiser, P.B. Tremblay, J. Brockmüller, and I. Roots. Simultaneous determination of ondansetron and tropisetron in human plasma using HPLC with UV detection. *Biomed. Chromatogr.* **16**: 187–190 (2002).
17. D. Chandrasekar, S. Ramakrishna, and P.V. Diwan. A rapid, sensitive and validated method for the determination of ondansetron in human plasma by reversed-phase high-pressure liquid chromatography. *Arzneimittelforschung* **54**: 655–659 (2004).
18. M. Siluveru and J.T. Stewart. Enantioselective determination of *S*-(+)- and *R*-(–)-ondansetron in human serum using derivatized cyclodextrin-modified capillary electrophoresis and solid-phase extraction. *J. Chromatogr. B Biomed. Sci. Appl.* **691**: 217–222 (1997).
19. J.W. Kelly, L. He, and J.T. Stewart. High-performance liquid chromatographic separation of ondansetron enantiomers in serum using a cellulose-derivatized stationary phase and solid-phase extraction. *J. Chromatogr.* **622**: 291–295 (1993).
20. J. Liu and J.T. Stewart. High-performance liquid chromatographic analysis of ondansetron enantiomers in human serum using a

- reversed-phase cellulose-based chiral stationary phase and solid-phase extraction. *J. Chromatogr. B Biomed. Sci. Appl.* **694**: 179–184 (1997).
21. K. Liu, X. Dai, D. Zhong, and X. Chen. Quantitative determination of ondansetron in human plasma by enantioselective liquid chromatography–tandem mass spectrometry. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.* **864**: 129–136 (2008).
 22. W.D. Figg, G.E. Dukes, J.F. Pritchard, D.J. Hermann, H.R. Lesesne, S.W. Carson, S.S. Songer, J.R. Powell, and L.J. Hak. Pharmacokinetics of ondansetron in patients with hepatic insufficiency. *J. Clin. Pharmacol.* **36**: 206–215 (1996).
 23. E.I. Ashforth, J.L. Palmer, A. Bye, and A. Bedding. The pharmacokinetics of ondansetron after intravenous injection in healthy volunteers phenotyped as poor or extensive metabolisers of debrisoquine. *Br. J. Clin. Pharmacol.* **37**: 389–391 (1994).
 24. R. Kaiser, O. Sezer, A. Papies, S. Bauer, C. Schelenz, P.B. Tremblay, K. Possinger, I. Roots, and J. Brockmoller. Patient-tailored antiemetic treatment with 5-hydroxytryptamine type 3 receptor antagonists according to cytochrome P-450 2D6 genotypes. *J. Clin. Oncol.* **20**: 2805–2811 (2002).
 25. K.A. Candiotti, D.J. Birnbach, D.A. Lubarsky, F. Nhuch, A. Kamat, W.H. Koch, M. Nikoloff, L. Wu, and D. Andrews. The impact of pharmacogenomics on postoperative nausea and vomiting: do CYP2D6 allele copy number and polymorphisms affect the success or failure of ondansetron prophylaxis? *Anesthesiology* **102**: 543–549 (2005).
 26. J. Sistonen, S. Fuselli, J.U. Palo, N. Chauhan, H. Padh, and A. Sajantila. Pharmacogenetic variation at CYP2C9, CYP2C19, and CYP2D6 at global and microgeographic scales. *Pharmacogenet. Genomics* **19**: 170–179 (2009).
 27. P 450 Drug Interaction Table: <http://medicine.iupui.edu/clin-pharm/ddis/table.asp>.
 28. U.M. Stamer and F. Stüber. The pharmacogenetics of analgesia. *Expert Opin. Pharmacother.* **8**: 2235–2245 (2007).
 29. M. Ingelman-Sundberg, S.C. Sim, A. Gomez, and C. Rodriguez-Antona. Influence of cytochrome P450 polymorphisms on drug therapies: pharmacogenetic, pharmacoeypigenetic and clinical aspects. *Pharmacol. Ther.* **116**: 496–526 (2007).
 30. M. Eichelbaum, M. Ingelman-Sundberg, and W.E. Evans. Pharmacogenomics and individualized drug therapy. *Annu. Rev. Med.* **57**: 119–137 (2006).
 31. S. Zhong and M. Romkes. Pharmacogenomics. *Methods Mol. Biol.* **520**: 231–245 (2009).

Manuscript received February 8, 2010;
revision received May 2, 2010.