



Echicetin Coated Polystyrene Beads: A Novel Tool to Investigate GPIb-Specific Platelet Activation and Aggregation

Alexey Navdaev¹✉, Hariharan Subramanian¹✉, Alexey Petunin², Kenneth J. Clemetson³, Stepan Gambaryan^{1,4*}, Ulrich Walter⁵

1 Institute of Clinical Biochemistry and Pathobiochemistry, University of Wuerzburg, Wuerzburg, Germany, **2** Institute of Biophysics, Siberian Branch of the Russian Academy of Sciences, Krasnoyarsk, Russia, **3** Theodor Kocher Institute, University of Berne, Berne, Switzerland, **4** Sechenov Institute of Evolutionary Physiology and Biochemistry, Russian Academy of Sciences, St. Petersburg, Russia, **5** Center for Thrombosis and Hemostasis (CTH), University Medical Center Mainz, Mainz, Germany

Abstract

von Willebrand factor/ristocetin (vWF/R) induces GPIb-dependent platelet agglutination and activation of α IIb β 3 integrin, which also binds vWF. These conditions make it difficult to investigate GPIb-specific signaling pathways in washed platelets. Here, we investigated the specific mechanisms of GPIb signaling using echicetin-coated polystyrene beads, which specifically activate GPIb. We compared platelet activation induced by echicetin beads to vWF/R. Human platelets were stimulated with polystyrene beads coated with increasing amounts of echicetin and platelet activation by echicetin beads was then investigated to reveal GPIb specific signaling. Echicetin beads induced α IIb β 3-dependent aggregation of washed platelets, while under the same conditions vWF/R treatment led only to α IIb β 3-independent platelet agglutination. The average distance between the echicetin molecules on the polystyrene beads must be less than 7 nm for full platelet activation, while the total amount of echicetin used for activation is not critical. Echicetin beads induced strong phosphorylation of several proteins including p38, ERK and PKB. Synergistic signaling via P2Y₁₂ and thromboxane receptor through secreted ADP and TxA₂, respectively, were important for echicetin bead triggered platelet activation. Activation of PKG by the NO/sGC/cGMP pathway inhibited echicetin bead-induced platelet aggregation. Echicetin-coated beads are powerful and reliable tools to study signaling in human platelets activated solely via GPIb and GPIb-triggered pathways.

Citation: Navdaev A, Subramanian H, Petunin A, Clemetson KJ, Gambaryan S, et al. (2014) Echicetin Coated Polystyrene Beads: A Novel Tool to Investigate GPIb-Specific Platelet Activation and Aggregation. PLoS ONE 9(4): e93569. doi:10.1371/journal.pone.0093569

Editor: Christian Schulz, King's College London School of Medicine, United Kingdom

Received: November 15, 2013; **Accepted:** March 5, 2014; **Published:** April 4, 2014

Copyright: © 2014 Navdaev et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by DFG (SFB688, TP A2, and grant GA 1561/1-1), BMBF (01EO1003) and Rudolf-Marx-stipendium 2014 funded by Gesellschaft für Thrombose- und Hämostaseforschung (GTH). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: gambaryan@klin-biochem.uni-wuerzburg.de

✉ These authors contributed equally to this work.

Introduction

In vivo, von Willebrand Factor (vWF) interacts with platelet glycoprotein Ib complex (GPIb) under high shear and triggers platelet activation, which leads to ADP and TxA₂ release and hence α IIb β 3 activation and platelet aggregation. Under normal conditions, vWF does not interact with circulating platelets [1]. Following injury under high shear conditions the globular molecule of vWF adheres to subendothelial collagen and is stretched to expose the A1 domains, the binding domain for GPIb [2]. vWF is a large multimer containing many A1 domains, which can bind GPIb molecules on the platelet surface [3]. The clustering of GPIb receptors linked to the submembranous cytoskeleton triggers platelet activation [4,5]. However, the number of A1 domains interacting with GPIb on a platelet surface as well as the distance between neighbouring A1 domains required for GPIb clustering and platelet activation was not previously determined.

The antibiotic ristocetin binds to the C-terminal part of the A1-domain of vWF as well as to GPIb, inducing the interaction

between vWF and GPIb on platelets [6]. However, vWF/R treatment strongly agglutinates stirred, washed platelets [7]. Agglutination by vWF/R is passive and does not cause intracellular signaling. It takes place even with formaldehyde-fixed platelets (often used as a clinical test). The C1 domain of vWF can also interact with α IIb β 3 integrin on the platelet surface to induce an intracellular signal independently from GPIb [8,9]. All these events caused by vWF/R treatment obscure the analysis of GPIb receptor signaling. A surface coated with recombinant dimeric A1 domain was used earlier to study GPIb specific signaling [10]. However, dimeric A1 domain cannot be used to activate platelets in suspension via their vWF receptor without cross linking the A1 domains. Therefore, we developed a new model for the study of GPIb specific signaling in human platelets based on echicetin-coated polystyrene beads (EP).

Echicetin, a snake venom C-type lectin, specifically binds to GPIb, but does not induce activation of washed platelets. Moreover, binding of echicetin to GPIb completely blocks vWF/R induced platelet agglutination [11]. However, echicetin can be cross-linked by plasma IgM κ to induce platelet agglutination and

weak aggregation [12]. A maximum of 10 echicetin molecules can bind to one IgMk molecule and this complex may not be sufficient to cluster GPIb molecules to the extent required to activate α IIB β 3 and induce full aggregation. Therefore, we used echicetin coated polystyrene beads with a determined number of GPIb-binding sites and a known average distance between neighboring echicetin molecules, and investigated platelet agglutination/aggregation. We showed that our model can be used to study signaling mechanisms in platelets specifically activated via GPIb and that for maximum platelet activation and aggregation the GPIb ligands should be spaced closer than 7 nm.

Materials and Methods

Ethics Statement

Blood was obtained from healthy volunteers, who gave written informed consent according to our institutional guidelines and the Declaration of Helsinki. Our studies with human platelets and the consent procedure were approved and reconfirmed (September 24, 2008) by the local ethics committee of the University of Wuerzburg (Studies No. 67/92 and 114/04).

Materials

Echis carinatus sochureki venom was from Latoxan (Valence, France). Ristocetin was from Biopool (Wicklow, Ireland). vWF (Haemate HS 1000, vWF activity 2200 IE) was from CSL Behring (Marburg, Germany). Polystyrene beads (0.46 μ m, aqueous suspension, 10% solids) and phospho-p38 (Thr¹⁸⁰/Tyr¹⁸²) MAPK antibody were from Sigma (Taufkirchen, Germany). R-phycoerythrin (RPE)-conjugated anti-CD62P antibody was from DAKO (Glostrup, Denmark). Anti-phosphotyrosine antibody (4 G10) was from Millipore (Schwalbach, Germany). Phospho-VASP Ser²³⁹, and phospho-ERK (Thr²⁰¹/Tyr²⁰⁴) antibodies were from Nanotools (Teningen, Germany). Antibodies against phospho-PKB (Thr³⁰⁸), VASP, and p38 were from Cell Signaling (Frankfurt, Germany).

Preparation of Echicetin-coated Polystyrene Beads

Echicetin was purified as previously described [12]. Polystyrene beads were washed and coated with echicetin in PBS overnight followed by washing and blocking with BSA. To obtain different densities, the beads were coated with 0.3 (EP0.3), 0.2 (EP0.2), 0.1 (EP0.1) and 0.05 (EP0.05) mg/ml echicetin.

Determination of the Amount of Echicetin, and the Distance between the Echicetin Molecules on the Beads

Echicetin was labeled with I¹²⁵ using Pierce Iodination Reagent (Iodogen) with a standard protocol (Thermo Scientific - Pierce, Rockford, USA). Unreacted I¹²⁵ was removed from the labeled protein by gel-filtration on Sephadex G10. Labeled echicetin (10 μ g) was mixed with 1 mg of unlabeled echicetin. This mixture was used as a stock to prepare 0.3 (EP0.3), 0.2 (EP0.2), 0.1 (EP0.1) and 0.05 mg/ml (EP0.05) echicetin solutions for adsorption on polystyrene beads using the same protocol as with unlabeled protein. The total radioactivity of the samples was measured before adsorption and the residual radioactivity on polystyrene beads was measured after adsorption using a gamma-counter. The amount of adsorbed echicetin was calculated from the ratio between the total sample radioactivity and the residual radioactivity on the polystyrene beads. We found that 28.2, 23.7, 13.2 and 7.5 μ g of echicetin (30.6 kDa) bound to EP0.3, EP0.2, EP0.1 and EP0.05, respectively. Adsorbed echicetin (A) corresponds to 5.7×10^{14} , 4.8×10^{14} , 2.7×10^{14} , and 1.5×10^{14} molecules. To find the average distance between two neighboring echicetin molecules

adsorbed on polystyrene beads the total area of all beads in the suspension was calculated as follows.

$$S_{total} = S_{bead} \times N = 0.6644 \mu m^2 \times 1.877 \times 10^{10} \\ = 1.247 \times 10^{10} \mu m^2$$

where S_{bead} is the surface area of a single bead (0.6644 μ m² for a bead with $d = 0.46 \mu$ m) and N is the total number of beads in the sample, calculated using an equation supplied by Sigma and determined to be 1.877×10^{10} .

Assuming that echicetin is evenly distributed on the polystyrene surface, the mean area on the bead occupied by each echicetin molecule was calculated by the equation: $S_{single} = S_{total}/A$, where S_{single} is the area containing a single molecule, S_{total} is the total area of polystyrene beads covered by echicetin and A is the number of adsorbed echicetin molecules. For EP0.3, EP0.2, EP0.1 and EP0.05, S_{single} was found to be 21.8, 25.7, 46.2 and 81.2 nm² respectively. The distance between two neighboring molecules (D) was calculated as $D = \sqrt{S_{single}}$ and found to be 4.7, 5.1, 6.8 and 9.0 nm for EP0.3, EP0.2, EP0.1 and EP0.05, respectively. Echicetin molecules per bead were calculated as $n = A/N$ giving 30500, 25800, 14300 and 8170 for EP0.3, EP0.2, EP0.1 and EP0.05, respectively.

Platelet Aggregation

Human platelets were prepared as previously described [13]. Platelet aggregation was measured using an Apact4004 (LabiTec, Ahrensburg, Germany) aggregometer. Washed platelets (3×10^8 /ml) were treated with echicetin beads (1% w/w suspension) or vWF/R under continuous stirring at 1000 rpm and 37°C.

Flow Cytometry

Platelets activated with echicetin beads were taken directly from the aggregation cuvette and fixed with 1% formaldehyde for 10 min. After centrifugation, platelets were resuspended in PBS/0.5% BSA, and incubated with RPE-conjugated anti-CD62P antibody. FACS analysis was performed on a Becton Dickinson FACScalibur using CELLQuest software, version 3.1f (Becton Dickinson, Heidelberg, Germany).

Platelet Adhesion and Spreading

Cover slips were coated with various concentrations of echicetin. Platelet adhesion and spreading on echicetin coated surfaces were performed as described before [14]. Adhered platelets were studied under the microscope (Zeiss, Axiovert 200) at 1000 \times magnification. Platelet images were captured using a camera (Diagnostic Instruments, SpotPursuit 23) and VisiView software.

Western Blot Analysis

For Western blot analysis, 2 \times Laemmli buffer was directly added to washed platelets stimulated with echicetin beads or vWF/R. Platelet proteins were separated by SDS-PAGE, transferred to nitrocellulose membranes, and the membranes were incubated with appropriate primary antibodies. Goat anti-rabbit or anti-mouse IgG conjugated with horseradish peroxidase were used as secondary antibodies, followed by ECL detection.

Data Analysis

All experiments were performed at least three times and the data are expressed as mean \pm SEM. Differences between groups were analyzed by ANOVA followed by Bonferroni's test and

Student t-test was used when appropriate. $P < 0.05$ was considered statistically significant.

Results

Echicetin Beads Induce α IIb β 3-dependent Platelet Aggregation

Elucidation of intracellular signaling mechanisms induced by vWF/R in washed platelet suspension is limited by strong platelet agglutination. In PRP, addition of ristocetin induces two waves of platelet activation; first one is agglutination, and the second α IIb β 3-dependent aggregation (Fig. 1A). Therefore, to examine GPIb specific signaling in washed platelets suspension, we introduced a new model of echicetin coated polystyrene beads (EP). Echicetin was shown to bind specifically to GPIb and to block platelet interactions with vWF [11,12]. EP, after a short lag-phase, induced strong platelet aggregation, which was completely blocked by preincubation with α IIb β 3 inhibitor aggrastat, or echicetin monomer (Fig. 1B). In contrast to PRP, in washed platelet suspension, agglutination of platelets by vWF/R was inhibited only by monomeric echicetin, but not by aggrastat (Fig. 1C).


Neither monomeric echicetin nor BSA coated beads (used as controls) induce platelet aggregation or agglutination (Fig. 2A). To elucidate whether the density, or the distance between neighboring molecules of adsorbed echicetin influences the ability of EP to induce platelet aggregation, polystyrene beads were coated with a defined concentration of echicetin based on the data obtained with absorption of I^{125} labeled echicetin on polystyrene beads (see methods). First, we established the range of echicetin concentrations (between 1 and 0.01 mg/ml) for coating the beads and found that maximum aggregation was induced by incubation with 0.3 mg/ml, whereas 0.05 mg/ml did not induce any significant platelet aggregation. According to our calculations (see methods) the number of echicetin molecules per bead should be 30500, 25800, 14300 and 8170 in EP0.3, EP0.2, EP0.1, and EP0.05, respectively and the average distance between the adsorbed echicetin molecules is 4.7, 5.1, 6.8 and 9.0 nm. Correspondingly, the total number of echicetin molecules added to platelet samples were 5.7×10^{14} , 4.8×10^{14} , 2.7×10^{14} , and 1.5×10^{14} for EP0.3, EP0.2, EP0.1, and EP0.05, respectively. There are about 40000 molecules of GPIb per platelet [15] and the total number of GPIb molecules in 4.6×10^7 platelets used in aggregation experiments were 1.84×10^{12} approximately, which is comparable with the total number of echicetin molecules in PE0.05.

The maximum aggregation, $94.6 \pm 5.6\%$ and $94.9 \pm 10.8\%$ was obtained when platelets were activated by EP0.3 and EP0.2, respectively. Slightly less aggregation ($86.1 \pm 8.2\%$) was observed with EP0.1 (Fig. 2B, 2C). However, EP0.05 did not induce aggregation ($3.2 \pm 1.3\%$), indicating that the amount, and/or the distance between echicetin molecules in the beads were not sufficient for GPIb mediated platelet aggregation. In order to clarify this, we added 5 times more EP0.05 (7.5×10^{14} echicetin molecules per sample) and still found no significant changes in platelet aggregation (Fig. 2C). These data suggest that the distance between the echicetin molecules is critical, but not the total number of echicetin molecules on the beads used for stimulation. The average distance between echicetin molecules should be less than 7 nm for GPIb mediated platelet aggregation.


Echicetin-coated Beads Induce P-selectin Surface Expression in Human Platelets

Platelet activation induced by echicetin beads was determined by P-selectin expression on the activated platelets. Though

A



B



C




Figure 1. Echicetin coated polystyrene beads induce α IIb β 3-dependent aggregation in washed platelets. Representative aggregation curves of platelet rich plasma stimulated with ristocetin (A), and washed human platelets stimulated with EP0.1 in the absence of vWF or fibrinogen (B) or vWF-15 μ g/ml and Ristocetin-1 mg/ml (C). Whenever indicated, platelets were preincubated with echicetin monomer (25 μ g/ml, 3 min) or aggrastat (1.25 μ g/ml, 1 min). Shown are results of three independent experiments. doi:10.1371/journal.pone.0093569.g001

thrombin and echicetin beads induce full platelet aggregation, P-selectin expression induced by echicetin beads was significantly less than in thrombin-stimulated platelets. There were no significant differences between P-selectin expression levels induced by EP0.3 and EP0.2. P-selectin expression was slightly lower with EP0.1 and strongly decreased in the case of EP0.05 (Fig. 2D). These data show that echicetin beads induce alpha-granule secretion, but less than strong agonists like thrombin.

Platelet Adhesion and Spreading on Echicetin-coated Surface

To identify whether GPIb-mediated platelet activation induces platelet adhesion and spreading, washed human platelets were incubated on cover slips coated with increasing concentrations of echicetin. We found typical platelet adhesion and spreading on echicetin coated surfaces and the number of adherent platelets increased with the density of echicetin coated on the surface (Figure 3A, B). The area covered by spread platelets was also




Figure 2. The distance between echicetin molecules is critical for inducing platelet activation and aggregation. Washed human platelets were stimulated with EP under stirring conditions (1000 rpm). BSA coated polystyrene beads were used as negative control. Platelet aggregation stimulated by EP0.3 was considered as 100%. A. Representative aggregation curves of platelets stimulated with EP0.1, BSA-coated polystyrene beads, or echicetin-monomer are shown. B. Representative aggregation curves of platelets stimulated with EP0.3, EP0.2, EP0.1, EP0.05 or 5 times concentration of EP0.05 are shown. C. The graph shows the corresponding average values of aggregation from B. D. P-selectin surface expression on platelets was measured as Mean Fluorescence Intensity (MFI) using RPE-conjugated P-selectin antibody in flow cytometry. Thrombin (0.05 U/ml) and BSA-coated beads (BSA) were used as positive and negative controls, respectively. Results are mean \pm SD from 3 independent experiments. * $p < 0.05$ versus EP0.3, and * $p < 0.05$ versus EP0.1. doi:10.1371/journal.pone.0093569.g002

dependent on the concentration of the echicetin coated on the surface (Figure 3C). These results indicated that GPIb-mediated platelet activation by echicetin induces platelet adhesion and spreading.

Echicetin Beads Induce Tyrosine Phosphorylation in Platelets


Global protein tyrosine phosphorylation is a reliable marker of activation dependent intracellular signaling. Therefore, we examined changes in protein tyrosine phosphorylation in platelets activated by EP0.1. Strongly increased tyrosine phosphorylation of several proteins was observed in platelets stimulated by EP0.1, which was partially inhibited by aggrastat (Fig. 4A). This indicates that GPIb-induced tyrosine phosphorylation is partly mediated by outside-in integrin signaling most likely via proteins released from α -granules such as fibrinogen. However, preincubation with monomeric echicetin completely abolished the increase in tyrosine phosphorylation induced by EP0.1 (Fig. 4A). In contrast, under our experimental conditions, activation of platelets with vWF/R

did not trigger any tyrosine phosphorylation (Fig. 4B), which directly corresponds to the platelet agglutination data (Fig. 1C).


Platelet Aggregation Induced by Echicetin Beads is Mediated by Src-kinases and SYK

Platelets preincubated with Src-kinase inhibitor (PP2) did not form aggregates, when stimulated with EP. PP3, an inactive analog of PP2 did not affect EP-induced platelet aggregation (Fig. 5A, C). In contrast, PP2 had no effect on vWF/R-induced platelet agglutination (Fig. 5B, D). GPIb-mediated platelet signaling was also tested using piceatannol, a SYK specific inhibitor. Piceatannol strongly inhibits aggregation induced by echicetin beads (Fig. 5A, C) and had no effect on vWF/R-induced agglutination (Fig. 5B, D). MAP kinases (p38, ERK) and PKB play essential role in vWF-stimulated platelet activation [16,17,18]. Therefore, we examined the phosphorylation of p38, ERK and PKB in EP-activated platelets. EP-induced phosphorylation of p38, ERK and PKB were completely inhibited in platelets by preincubation with PP2 and piceatannol (Fig. 5E). In contrast, in vWF/R treated platelets, PKB and MAP kinases were not

A



B



C




Figure 3. Platelet adhesion and spreading on echicetin-coated surface. Cover slips were coated with various amounts of echicetin or BSA (control). Washed human platelets (1×10^7 /ml) were incubated on the coated cover slips for 15 min. Following incubation, cover slips were washed to remove unbound platelets and the adherent platelets were fixed with 1% formaldehyde. The fixed platelets were permeabilized with Triton X-100 and stained with phalloidin-Oregon Green (actin). The stained platelets were analyzed under the microscope (Zeiss, Axiovert 200) at 1000 \times magnification. The images were captured using a camera (Diagnostic Instruments, SpotPursuit 23) and VisiView software. A. Representative images of the bright-field (upper panel) and actin staining (lower panel) of the cover slips were shown for each sample. B. The number of adhered platelets was counted in 10 independent cover slips for each sample. C. Surface area of adhered and spread platelets was measured using VisiView software in 10 independent cover slips for each sample. Results are mean \pm SEM from 3 independent experiments. doi:10.1371/journal.pone.0093569.g003

activated and Src kinase and Syk inhibitors inhibited only basal phosphorylation of these kinases (Fig. 5F).

ADP and TXA₂ Signaling are Important for Echicetin Bead Induced Platelet Activation and Aggregation

Secreted ADP and TXA₂ play significant roles in shear-induced GPIIb mediated platelet aggregation [1,19]. Mouse platelets in suspension stimulated with botrocetin/vWF also required ADP and TXA₂ for aggregation [20,21]. To assess the role of ADP and TXA₂ signaling in EP-induced platelet aggregation, platelets were pre-incubated with antagonists of P2Y₁₂ (AR-C69931), P2Y₁ (MRS-2179) or TXA₂ receptor (SQ-29548). AR-C69931 potently inhibits EP-induced platelet aggregation, whereas MRS-2179 had only a minimal effect. SQ-29548 strongly reduced EP-induced platelet aggregation (Fig. 6A, C). However, simultaneous inhibition of ADP and TXA₂ receptors did not significantly differ from the effect of AR-C69931 alone (data not shown) indicating that

other intracellular mechanisms are involved in EP-induced platelet aggregation. Pre-treatment with AR-C69931 and SQ-29548 inhibits phosphorylation of PKB, ERK and p38 (Fig. 6E) in EP-stimulated platelets. In contrast, neither ADP nor TXA₂ receptor antagonists affect vWF-induced platelet agglutination (Fig. 6B, D) and phosphorylation of PKB, ERK and p38 (Fig. 6F).

NO-dependent Activation of Soluble Guanylate Cyclase Inhibits Echicetin Beads-induced Platelet Aggregation

Previously, we [22,23] and others [24,25] found that vWF/R treatment of platelets activates the sGC/cGMP/PKG pathway and induces VASP phosphorylation. To evaluate whether in this case VASP phosphorylation is mediated by direct activation via GPIIb or by a complex reaction of vWF/R we used our established system (echicetin beads) to stimulate platelets. First, we found that PKG activation by sodium nitroprusside (NO donor) inhibits EP-induced platelet aggregation by $83 \pm 7\%$ (Fig. 7A, C). Preincuba-




Figure 4. Echicetin beads but not vWF/R induces protein tyrosine phosphorylation. Washed human platelets were stimulated with EP0.1 (A) or vWF/R (B) after preincubation with aggrastat (1.25 μ g/ml, 1 min) or echicetin monomer (25 μ g/ml, 3 min). Protein tyrosine phosphorylation was analyzed by Western blot. Shown are representative blots of three independent experiments.
doi:10.1371/journal.pone.0093569.g004

tion with sGC specific inhibitor (ODQ) significantly enhanced ($12 \pm 3\%$) EP-induced platelet aggregation (Fig. 7A, C), but had no effect on vWF/R-induced agglutination (Fig. 7B, D). Second, we tested whether VASP phosphorylation in vWF/R treated platelets is mediated by GPIb signaling. We can confirm our previous finding [22] that vWF/R treatment induces strong PKG activation and subsequent VASP phosphorylation (Fig. 7F). However, specific activation of GPIb by EP0.1 does not induce any VASP phosphorylation (Fig. 7E). These data indicate that vWF/R induced PKG activation is not directly mediated by GPIb activation but requires additional signaling pathways induced by vWF/R.

Discussion

vWF mediated signaling in platelets is complex and involves several intracellular mechanisms, which play an important role in the initiation of thrombus formation. The complexity of these mechanisms is related to the vWF molecule itself and its main domains including the D domain (the binding site for FVIII and heparin), D²-D3 domain (possible binding site for P-selectin), A3 domain (the binding site for fibrillar collagen type I and III), C1 domain (the binding site for integrin $\alpha_{IIb}\beta_3$), and A1 domain which binds GPIb [8]. In addition to vWF, other ligands like thrombospondin-1 [26] and α -thrombin [27] can bind to GPIb and induce intracellular signaling independently from vWF. Numerous approaches including genetically modified mouse models, various forms of von Willebrand disease, shear-induced

thrombus formation, specific interactions with isolated A1 domain, etc. have been used for the characterization of vWF-induced signaling [28]. In models, where platelets are used in suspension additional modulators such as ristocetin or botrocetin need to be added to initiate vWF/GPIb mediated platelet response. However, in all these models identification of specific signaling pathways mediated solely by vWF/GPIb interactions is extremely difficult for the following reasons: Apart from GPIb, vWF can bind to other receptors on the platelet surface and initiate complex signaling events; vWF/R or vWF/botrocetin, under stirring conditions (which are important to initiate aggregation of platelet suspensions), induce strong platelet agglutination, which can influence intracellular signaling mediated by vWF/GPIb interaction. The degree of agglutination probably depends on the source of vWF and the experimental conditions. Strong platelet agglutination (which is independent of integrin $\alpha_{IIb}\beta_3$) induced by vWF/R treatment has been described in several papers [7,29,30,31,32]. In our study, we used Haemate HS (vWF activity 2200IE), which is clinically used as vWF substitute in vWF deficiencies (von Willebrand diseases). We observed that vWF/R treatment induces only strong platelet agglutination, which was not affected by aggrastat (Fig. 1C), inhibitors of Src kinase and SYK (Fig. 5), ADP and TXA₂ receptors (Fig. 6).

The main aim of our study was to develop a GPIb-specific agonist to investigate the GPIb-mediated signaling events in platelets without interference from other receptor pathways. Echicetin, a C-type lectin from snake venom, seems to be applicable for investigations addressing GPIb specific signaling in washed platelet suspension. Previously, we had shown that a maximum of 10 echicetin molecules cross-linked with IgMk can induce $\alpha_{IIb}\beta_3$ -independent platelet agglutination [12]. The distance between the echicetin molecules coupled to IgMk is sufficient to cluster few GPIb molecules that lead to agglutination. However, this complex did not allow us to find out the distance between echicetin molecules required to induce $\alpha_{IIb}\beta_3$ integrin-dependent platelet aggregation. Therefore, we developed a novel tool to investigate GPIb-specific activation of $\alpha_{IIb}\beta_3$ integrin-dependent platelet aggregation. With this tool, we identified that the distance between the echicetin molecules (<7 nm), but not the amount of echicetin is critical to induce GPIb clustering and GPIb-mediated $\alpha_{IIb}\beta_3$ integrin activation (Fig. 2). Also, platelet adhesion and spreading depend on the density of echicetin on the surface used to activate platelets (Fig. 3). These data strongly support the hypothesis that clustering of GPIb molecules is an important factor for the initiation of GPIb-mediated intracellular signaling.

We then compared the aggregation/agglutination induced by EP and vWF/R and found, consistent with published data [24,33], that vWF/R induces strong agglutination in washed platelet suspension (Fig. 1C), without significant activation of tyrosine kinases (Fig. 4B). In contrast to washed platelets, in PRP addition of ristocetin induced the well characterized two waves of platelet activation; first agglutination, which is insensitive to aggrastat, and second, aggregation inhibited by aggrastat (Fig. 1A). Whereas EP, under the same experimental conditions, induces strong global tyrosine phosphorylation, which was only slightly decreased by $\alpha_{IIb}\beta_3$ inhibitor, indicating that most of the intracellular signal is mediated by activation of GPIb, but not $\alpha_{IIb}\beta_3$ outside-in signaling (Fig. 4A). It is important to note that adding echicetin monomer before stimulation by vWF/R or EP completely prevents platelet aggregation (Fig. 1B, C) and tyrosine phosphorylation (Fig. 4A). We also investigated several intracellular signaling pathways downstream of GPIb signaling, and consistent with others [16,18,24,34], we found that Src kinases, SYK, PKB,




Figure 5. Src-kinases and SYK are downstream effectors of echicetin beads induced platelet activation. Washed human platelets were preincubated with Src-kinases inhibitor (PP2, 10 μ M), an inactive analog (PP3, 10 μ M) or Syk inhibitor (piceatannol, 10 μ M) before stimulating with EP0.1 or vWF/R under stirring conditions. Representative aggregation curves of platelets stimulated with EP0.1 (A) or vWF/R (B) are shown. C, D. The graphs shows corresponding average values of aggregation from A and B. Platelets stimulated with EP0.1 (E) or vWF/R (F) were lysed at the indicated time and the phosphorylation of PKB, ERK and p38 was analyzed by Western blot. Total p38 served as loading control. Shown are representative blots of three independent experiments. * $p < 0.05$ versus EP0.1. doi:10.1371/journal.pone.0093569.g005

p38 and ERK MAP kinases are strongly activated by GPIb signalling. However, none of these kinases were activated in vWF/R agglutinated platelets (Fig. 5). We found that P2Y₁₂ and TxA₂ signaling plays a synergistic role in EP-induced platelet activation and aggregation, in line with earlier findings [1,19,20,21] (Fig. 6).

In contrast with other publications [16,21,35], we did not find significant input from P2Y₁ receptor in GPIb-induced platelet aggregation (Fig. 6). Our data clearly show that P2Y₁ receptor has only a minor role in GPIb signaling. This finding does not imply that P2Y₁ receptors are without a significant role in vWF-




Figure 6. ADP and TxA₂ play significant roles in echicetin bead induced platelet aggregation. Washed human platelets were preincubated with P2Y₁₂ antagonist (AR-C69931; 0.1 μ M, 5 min), P2Y₁ antagonist (MRS2179; 1 μ M, 5 min) or TxA₂ receptor antagonist (SQ-29548; 1 μ M, 5 min) before EP0.1 or vWF/R stimulation. Representative aggregation curves of platelets stimulated with EP0.1 (A) or vWF/R (B) are shown. C, D. The graphs shows corresponding average values of aggregation from A and B. Platelets stimulated with EP0.1 (E) or vWF/R (F) were lysed on indicated time and the phosphorylation of PKB, ERK and p38 was analyzed by Western blot. Representative blots of three independent experiments are shown. [†]p<0.05 versus EP0.1.
doi:10.1371/journal.pone.0093569.g006

mediated platelet activation. Probably activation of the P2Y₁ is connected with other pathways stimulated by vWF, but not by GPIIb-mediated signaling.

We can clearly confirm previous findings [22,23,24,25,36] that vWF/R triggers the sGC/cGMP/PKG pathway (Fig. 7F). However, this pathway is not mediated by GPIIb signaling, as we could show that EP did not increase phosphorylation of VASP (Fig. 7E), indicating that other, still unresolved mechanisms, connected with

vWF/R are responsible for activation of the sGC/cGMP/PKG pathway.

In summary, we introduced a new agonist (echicetin coated polystyrene beads, EP) that activates platelets via clustering of GPIIb without directly involving other receptors. This should allow the dissection of the complex mechanisms involved in vWF-induced platelet activation. We show that the distance (less than 7 nm) between GPIIb ligands rather than their concentration is




Figure 7. Activation of sGC inhibits platelet aggregation induced by echicetin beads. Washed human platelets were preincubated with NO donor (SNP; 10 μ M, 2 min) and/or sGC inhibitor (ODQ; 50 μ M, 5 min) before EP0.1 or vWF/R stimulation. Representative aggregation curves of platelets stimulated with EP0.1 (A) or vWF/R (B) are shown. C, D. The graphs show corresponding average values of aggregation from A and B. Platelets stimulated with EP0.1 (E) or vWF/R (F) were lysed and the phosphorylation VASP was analyzed by Western blot. Total VASP served as loading control. Representative blots of three independent experiments are shown. ⁺p<0.05 versus EP0.1. doi:10.1371/journal.pone.0093569.g007

critical for platelet activation/aggregation. We propose that echicetin-coated beads can be a powerful and reliable tool to study specific GPIb-dependent signaling pathways in platelets.

Author Contributions

Conceived and designed the experiments: AN SG UW. Performed the experiments: AN HS AP. Analyzed the data: AN HS AP KJC SG UW. Contributed reagents/materials/analysis tools: AN KJC. Wrote the paper: AN HS KJC SG UW.

References

- Jackson SP, Nesbitt WS, Kulkarni S (2003) Signaling events underlying thrombus formation. *J Thromb Haemost* 1: 1602–1612.
- Schneider SW, Nuschele S, Wixforth A, Gorzelanny C, Alexander-Katz A, et al. (2007) Shear-induced unfolding triggers adhesion of von Willebrand factor fibers. *Proc Natl Acad Sci U S A* 104: 7899–7903.

3. Springer TA (2011) Biology and physics of von Willebrand factor concatamers. *J Thromb Haemost* 9 Suppl 1: 130–143.
4. Arya M, Lopez JA, Romo GM, Cruz MA, Kasirer-Friede A, et al. (2003) Glycoprotein Ib-IX-mediated activation of integrin alpha(IIb)beta(3): effects of receptor clustering and von Willebrand factor adhesion. *J Thromb Haemost* 1: 1150–1157.
5. Wu Y, Asazuma N, Satoh K, Yatomi Y, Takafuta T, et al. (2003) Interaction between von Willebrand factor and glycoprotein Ib activates Src kinase in human platelets: role of phosphoinositide 3-kinase. *Blood* 101: 3469–3476.
6. Dong JF, Berndt MC, Schade A, McIntire LV, Andrews RK, et al. (2001) Ristocetin-dependent, but not botrocetin-dependent, binding of von Willebrand factor to the platelet glycoprotein Ib-IX-V complex correlates with shear-dependent interactions. *Blood* 97: 162–168.
7. Scott JP, Montgomery RR, Retzinger GS (1991) Dimeric ristocetin flocculates proteins, binds to platelets, and mediates von Willebrand factor-dependent agglutination of platelets. *J Biol Chem* 266: 8149–8155.
8. Reiminger AJ (2008) VWF attributes—impact on thrombus formation. *Thromb Res* 122 Suppl 4: S9–13.
9. Reiminger AJ (2008) Function of von Willebrand factor in haemostasis and thrombosis. *Haemophilia* 14 Suppl 5: 11–26.
10. Kasirer-Friede A, Cozzi MR, Mazzucato M, De Marco L, Ruggeri ZM, et al. (2004) Signaling through GP Ib-IX-V activates alpha IIb beta 3 independently of other receptors. *Blood* 103: 3403–3411.
11. Peng M, Lu W, Beviglia L, Niewiarowski S, Kirby EP (1993) Echicetin: a snake venom protein that inhibits binding of von Willebrand factor and alboataggregins to platelet glycoprotein Ib. *Blood* 81: 2321–2328.
12. Navdaev A, Dormann D, Clemetson JM, Clemetson KJ (2001) Echicetin, a GPIb-binding snake C-type lectin from *Echis carinatus*, also contains a binding site for IgMkappa responsible for platelet agglutination in plasma and inducing signal transduction. *Blood* 97: 2333–2341.
13. Gambaryan S, Kobsar A, Rukoyatkina N, Herterich S, Geiger J, et al. (2010) Thrombin and collagen induce a feedback inhibitory signaling pathway in platelets involving dissociation of the catalytic subunit of protein kinase A from an NFkappaB-IkappaB complex. *The Journal of biological chemistry* 285: 18352–18363.
14. Navdaev A, Clemetson KJ (2002) Glycoprotein Ib cross-linking/ligation on echicetin-coated surfaces or echicetin-IgMkappa in stirred suspension activates platelets by cytoskeleton modulated calcium release. *J Biol Chem* 277: 45928–45934.
15. Clemetson KJ, Lee AG (1996) Platelet membrane glycoproteins. *Biomembranes: A Multi-Volume Treatise*: JAI. 231–267.
16. Garcia A, Quinton TM, Dorsam RT, Kunapuli SP (2005) Src family kinase-mediated and Erk-mediated thromboxane A2 generation are essential for VWF/GPIb-induced fibrinogen receptor activation in human platelets. *Blood* 106: 3410–3414.
17. Yin H, Liu J, Li Z, Berndt MC, Lowell CA, et al. (2008) Src family tyrosine kinase Lyn mediates VWF/GPIb-IX-induced platelet activation via the cGMP signaling pathway. *Blood* 112: 1139–1146.
18. Yin H, Stojanovic A, Hay N, Du X (2008) The role of Akt in the signaling pathway of the glycoprotein Ib-IX induced platelet activation. *Blood* 111: 658–665.
19. Goto S, Tamura N, Eto K, Ikeda Y, Handa S (2002) Functional significance of adenosine 5'-diphosphate receptor (P2Y(12)) in platelet activation initiated by binding of von Willebrand factor to platelet GP Ibalpha induced by conditions of high shear rate. *Circulation* 105: 2531–2536.
20. Liu J, Pestina TI, Berndt MC, Jackson CW, Gartner TK (2005) Botrocetin/VWF-induced signaling through GPIb-IX-V produces TxA2 in an alphaIIb-beta3- and aggregation-independent manner. *Blood* 106: 2750–2756.
21. Liu J, Pestina TI, Berndt MC, Steward SA, Jackson CW, et al. (2004) The roles of ADP and TxA in botrocetin/VWF-induced aggregation of washed platelets. *J Thromb Haemost* 2: 2213–2222.
22. Gambaryan S, Geiger J, Schwarz UR, Butt E, Begonja A, et al. (2004) Potent inhibition of human platelets by cGMP analogs independent of cGMP-dependent protein kinase. *Blood* 103: 2593–2600.
23. Gambaryan S, Kobsar A, Hartmann S, Birschmann I, Kuhlencordt PJ, et al. (2008) NO-synthase-/NO-independent regulation of human and murine platelet soluble guanylyl cyclase activity. *J Thromb Haemost* 6: 1376–1384.
24. Marshall SJ, Senis YA, Auger JM, Feil R, Hofmann F, et al. (2004) GPIb-dependent platelet activation is dependent on Src kinases but not MAP kinase or cGMP-dependent kinase. *Blood* 103: 2601–2609.
25. Naseem KM (2008) eNOS, iNOS or no NOS, that is the question! *J Thromb Haemost* 6: 1373–1375.
26. Jurk K, Clemetson KJ, de Groot PG, Brodde MF, Steiner M, et al. (2003) Thrombospondin-1 mediates platelet adhesion at high shear via glycoprotein Ib (GPIb): an alternative/backup mechanism to von Willebrand factor. *FASEB J* 17: 1490–1492.
27. Zarpellon A, Celikel R, Roberts JR, McClintock RA, Mendolicchio GL, et al. (2011) Binding of alpha-thrombin to surface-anchored platelet glycoprotein Ib(alpha) sulfotyrosines through a two-site mechanism involving exosite I. *Proc Natl Acad Sci U S A* 108: 8628–8633.
28. Ruggeri ZM (2007) The role of von Willebrand factor in thrombus formation. *Thromb Res* 120 Suppl 1: S5–9.
29. Collier BS, Gralnick HR (1977) Studies on the mechanism of ristocetin-induced platelet agglutination. Effects of structural modification of ristocetin and vancomycin. *J Clin Invest* 60: 302–312.
30. Gangarosa EJ, Johnson TR, Ramos HS (1960) Ristocetin-induced thrombocytopenia: site and mechanism of action. *Arch Intern Med* 105: 83–89.
31. Howard MA, Firkin BG (1971) Ristocetin—a new tool in the investigation of platelet aggregation. *Thromb Diath Haemorrh* 26: 362–369.
32. Jenkins CS, Clemetson KJ, Luscher EF (1979) Studies on the mechanism of ristocetin-induced platelet agglutination: binding of ristocetin to platelets. *J Lab Clin Med* 93: 220–231.
33. Marshall SJ, Asazuma N, Best D, Wonerow P, Salmon G, et al. (2002) Glycoprotein IIb-IIIa-dependent aggregation by glycoprotein Ibalpha is reinforced by a Src family kinase inhibitor (PPI)-sensitive signalling pathway. *Biochem J* 361: 297–305.
34. Li Z, Xi X, Du X (2001) A mitogen-activated protein kinase-dependent signaling pathway in the activation of platelet integrin alpha IIb beta 3. *J Biol Chem* 276: 42226–42232.
35. Penz SM, Reiminger AJ, Toth O, Deckmyn H, Brandl R, et al. (2007) Glycoprotein Ibalpha inhibition and ADP receptor antagonists, but not aspirin, reduce platelet thrombus formation in flowing blood exposed to atherosclerotic plaques. *Thromb Haemost* 97: 435–443.
36. Li Z, Xi X, Gu M, Feil R, Ye RD, et al. (2003) A stimulatory role for cGMP-dependent protein kinase in platelet activation. *Cell* 112: 77–86.