Synthesis and Properties of 6'-Fluoro-tricyclo-DNA

Michal Medvecký, Alena Istrate and Christian J. Leumann*

Department of Chemistry and Biochemistry, University of Bern, Freiestrasse 3, CH-3012 Bern,
Switzerland

* Email: christian.leumann@dcb.unibe.ch

Table of Contents graphic:

Abstract

The synthesis of the two fluorinated tricyclic nucleosides 6'F-tc-T and 6'F-tc-5^{Me}C, as well as the corresponding building blocks for oligonucleotide assembly was accomplished. An X-ray analysis of N⁴-benzoylated 6'F-tc-5^{Me}C revealed a 2'-exo (North) conformation of the furanose ring characterizing it as an RNA mimic. In contrast to observations in the bicyclo-DNA series, no short contact between the fluorine atom and the H6 of the base, reminiscent of a non-classical F...H hydrogen bond, could be observed. *T_m* measurements of modified oligodeoxynucleotides with complementary RNA showed slightly sequence dependent duplex stabilization profiles with maximum $\Delta T_m/\text{mod}$ values of +4.5°C for 6'F-tc-5^{Me}C and +1°C for 6'F-tc-T. In comparison with parent tc-modified oligonucleotides, no relevant changes in *T_m* were detected, attributing the fluorine substituent a neutral role in RNA affinity. A structural analysis of duplexes with DNA and RNA by CD-spectroscopy revealed a shift from B- to A-type conformation induced by the 6'F-tc-nucleosides. This is not a specific 'fluorine effect' as the same is also observed for the parent tc-modifications. The two fluorinated tc-nucleosides were also incorporated into a pure tricyclo-DNA backbone and showed no discrimination in *T_m* with complementary RNA, demonstrating that 6'F substitution is also compatible within fully modified tc-oligonucleotides.

Introduction

Fluorine is widely appreciated in small molecule medicinal chemistry due its metabolic stability enhancing properties and its unique protein binding characteristics.¹⁻³ For similar reasons fluorinated DNA analogues are of interest in oligonucleotide therapeutic approaches. Amongst the first fluorinated oligonucleotides investigated were the 2'-deoxy-2'-fluoro RNA (F-RNA) and the 2'-deoxy-2'-fluoro-arabino nucleic acids (F-ANA, Figure 1). While both analogues are known for quite some time, their structural and biophysical features have only recently been characterized in detail. Compared to their 2'-hydroxy variants RNA and ANA, both the F-RNA and F-ANA analogues bind with higher affinity ($\Delta T_m = 1-2^\circ\text{C/mod}$) to complementary RNA. The origin of the higher duplex stability in the case of F-RNA was attributed to improved hydrogen-bonding and base-stacking as a consequence of the polar C2'-F bond.^{4,5} In the case of F-ANA, internucleoside F-H8 pseudo hydrogen-bonds, that are particularly strong at purine/pyrimidine sequence steps, have been invoked as stability enhancing feature.^{6,7} F-RNA and F-ANA have been shown to improve the performance of therapeutic siRNAs.^{8,9} Due to its unique RNaseH activating properties, F-ANA was also investigated in classical antisense applications.¹⁰

Recently there has been a growing interest in investigating the effect of fluorine substitution in more complex, carbohydrate modified oligonucleotide analogs such as F-HNA and its 2'-epimer Ara-F-HNA.^{11,12} While F-HNA shows increased thermal stability ($\Delta T_m = +2^\circ\text{C/mod}$) in complex with complementary RNA, the F-Ara-HNA analogue exhibits the opposite effect ($\Delta T_m = -3^\circ\text{C/mod}$). The destabilization of Ara-F-HNA was attributed to repulsive steric effects of the fluoro substituent onto the 5'-adjacent nucleotide unit.¹¹ Also fluorinated versions of CeNA¹³ and cLNA¹⁴ were investigated. In these cases fluorine substitution does not contribute significantly to duplex stability. The most recent additions to the palette of fluorinated oligonucleotide analogues

were F-NMC and Ara-F-NMC,¹⁵ both derived from the northern methanocarbacyclic nucleoside (NMC) analogues.^{16,17} Here again, F-NMC stabilized duplexes by +2.2°C/mod on average while Ara-F-NMC was destabilizing by -2.8°C/mod. The intrinsic contribution of the fluorine atom to thermal stability in the case of F-NMC was determined to be +0.6°C/mod on average.¹⁸

FIGURE 1: Chemical Structures of selected fluorinated nucleic acid analogues

In terms of biological activity it has been shown previously that F-HNA gapmers can down-regulate gene expression *in vivo* in liver tissue more potently than LNA despite lower target affinity.¹¹ Thus, the higher potency of F-HNA seems to be the consequence of either improved biostability or more efficient plasma transport or both. Other recent observations, attributing a special but yet elusive role to fluorine in antisense efficacy were reported for gapmer oligonucleotides with F-RNA or F-ANA units targeting mutant huntingtin,¹⁹ and for F-RNA antisense oligonucleotides recruiting the interleukin enhancer-binding factor complex (ILF2/3).²⁰

Given these intriguing properties of fluorinated oligonucleotides, and building on earlier work on 6'-F-bicyclo-DNA,²¹ we decided to investigate 6'-F-tc-DNA (Figure 1). In the following we present the synthesis and structural properties of the corresponding 6'-F-tc-nucleosides containing the bases thymine and 5-methyl cytosine, as well as the influence on duplex stability and conformation of these modifications if complexed with complementary RNA and DNA.

Results and Discussion

Synthesis of the phosphoramidite building blocks 11 and 18. Drawing on earlier experiences during the synthesis of 6'-fluorinated bicyclo-DNA²¹ we envisaged to introduce the fluorine atom in an early step of the synthesis via electrophilic fluorination of a metalated bicyclic sugar intermediate. Consequently we started our synthetic journey with the known bicyclic silylenol ether **1** (Scheme 1).²² Bromination of **1** with Br₂ at -78 °C gave the expected bromide **2** in 90% yield. To exclude side reactions during the following metal-halogen exchange, the free OH group in **2** was protected as a TMS ether ($\rightarrow$ **3**). Electrophilic fluorination of **3** worked best if t-BuLi was used as lithiation reagent and if NFSI was added in one portion at a temperature of -120 °C. Temperatures above -78°C, or addition of NFSI in multiple portions, lead to substantial decomposition and thus reduced yield. The treatment with an acidic ion exchange resin after quenching of the reaction quantitatively removed the TMS group, resulting in a 60% yield of fluorosilylenol ether **4**. The rationale of removing the TMS group at this stage was based on the hypothesis that the corresponding hydroxyl group could help in directing the subsequent cyclopropanation reaction to the convex side of the bicyclic ring system. Indeed, cyclopropanation of **4** with a Zn-carbene complex in homogeneous solution yielded exclusively the exo-tricyclic sugar **5** in good yield, but only if TFA was added as accelerator.²³ To prepare for

β -selective nucleosidation,^{24,25} compound **5** was converted to enolether **6** with TMSOTf, which was then reacted with in situ persilylated thymine and NIS, yielding iodo-nucleoside **7** in 85% yield in a stereospecific manner. Removal of the iodine via radical reduction with Bu₃SnH finally gave the expected O-protected 6'-fluoro tricyclothymidine **8** in excellent yields. From here the synthesis of the phosphoramidite building block was completed by standard removal of the silyl protecting groups ($\rightarrow$ **9**) followed by dimethoxytritylation ($\rightarrow$ **10**) and phosphitylation with 2-cyanoethyl diisopropylamino chlorophosphine (CEP-Cl) to give **11** in a respectable overall yield of 13.4% starting from **1**.

SCHEME 1. Synthesis of phosphoramidite **11**

Given the availability of the thymine nucleoside **8** and known procedures to interconvert pyrimidine bases on the level of nucleosides and oligonucleotides,^{26,27} we next envisioned the synthesis of the building block **18** containing the base 5-methyl cytosine. To this end compound **8**

was transformed into the triazolidine **12** with 1,2,4-triazole and POCl₃. Treatment of **12** with ammonia then afforded the 5-methyl tricyclocytidine **13** in 60% yield over two steps. N⁴-Benzoylation of **13** with Bz₂O yielded the two nucleosides **14** and **15** that differed only by the presence or absence of the 3'-O-TMS group. This was of no harm as the silyl groups in both, **14** and **15** were removed in the next step, yielding **16** in high yield. The synthesis of the phosphoramidite building block **18** was then completed via standard dimethoxytritylation (→ **17**) followed by phosphitylation as described above.

SCHEME 2. Synthesis of phosphoramidite **18**

X-ray structure of 16. To confirm the relative configuration and to determine the effect of the fluorine substituent on the conformation of the tricyclic ring system, crystals of nucleoside **16**

were grown and subjected to X-ray analysis. The molecular structure is depicted in Figure 2.

FIGURE 2. Ortep plot (50% probability ellipsoids) of the X-ray structure of nucleoside **16**: top-view (left) and side-view (right). Non relevant hydrogen atoms as well as the N⁴-benzoyl residue in **16** are omitted for clarity.

The furanose unit in **16** adopts a 2'-exo conformation with a pseudorotation phase angle P of 336° and a maximum torsion angle $\nu_{\max}$ of 31°. It thus belongs clearly to the N-type conformation, typically adopted by RNA nucleosides.²⁸ The base is oriented in the central anti-range ($\chi = -176.2^\circ$). Comparison of **16** with the structure of 6'-fluoro-bicyclo-T²¹ reveals two major differences: First, the distance F-H6 in **16** (3.194 Å) is too long for a non-classical F-H hydrogen bond while the same distance in 6'-fluoro-bc-T (2.865 Å) is indicative for such a weak interaction. Furthermore, there is no linear arrangement of F-H6-C6 in **16**, whereas this is clearly the case in 6'-fluoro-bc-T. Support for the absence of such an interaction in **16** comes also from the fact that there are no F-H6 or F-C6 couplings observable in the ¹H- and ¹³C-NMR spectra of **16**, which contrasts the findings in the case of 6'-fluoro-bc-T. Thus, compared to 6'-fluoro-bc-T, the base orienting H-F interaction of the fluorine is lost in **16**. The second structural change resides within the furanose conformation, which is 1'-exo (S-type) in the case of 6'-

fluorobicyclo-T and 2'-exo (N-type) in the case of **16**. With respect to the non-fluorinated tricyclo-T nucleoside which coexists in a 2'-endo (S-type) and a 4'-endo (E-type) conformation in the crystal,²⁹ it could well be that the fluoro atom helps to drive the furanose conformation of the tricyclic scaffold into a N-type conformation. We cannot exclude, however, that the higher propensity of N-conformation in **16** is also simply an effect of the base 5-methyl cytosine. Unfortunately, there are currently no X-ray structures for the non-fluorinated tc-C or tc-⁵MeC nucleosides available.

Synthesis of oligonucleotides and T_m -data. Oligodeoxynucleotides **ON1-10**, containing the 6' fluoro-tc-nucleosides, (Table 1) were synthesized on the 1.3 μ mol scale by standard phosphoramidite chemistry, utilizing the building blocks **11** and **18** (for details see experimental section). Crude oligonucleotides were deprotected and detached from solid support by standard ammonia treatment (33% NH_4OH , 55°C, overnight) and purified by ion exchange HPLC. The composition of all oligonucleotides was verified by ESI- mass spectrometry (Tables 1 and 3).

Table 1. Analytical data of oligodeoxynucleotides **ON1-ON7**, containing 6' fluoro-tc-T (**t**) or 6' fluoro-tc-⁵MeC (**c**) units, as well as T_m -data of duplexes with complementary DNA and RNA.

Sequence	ESI-MS	ESI-MS	T_m (°C) vs DNA ^{a,b}	T_m (°C) vs RNA ^{a,c}
	m/z calc	m/z found	(ΔT_m /mod)	(ΔT_m /mod)
ON1 d(AACTGtCACG)	3067.6	3067.5	45.5 (+2.0)	44.4 (0.0)
ON2 d(AACtGTCACG)	3067.6	3067.5	45.1 (+1.6)	45.4 (+1.0)
ON3 d(AACtGtCACG)	3123.6	3123.5	43.5 (0.0)	44.3 (0.0)
ON4 d(GCAtttttACCG)	3890.7	3890.6	43.1 (-0.6) ^d	45.6 (+0.5) ^e

ON5	d(AACTGT <u>c</u> ACG)	3081.6	3081.5	44.6 (+1.1)	46.3 (+1.9)
ON6	d(AA <u>c</u> TGTCACG)	3081.6	3081.5	44.9 (+1.4)	48.9(+4.5)
ON7	d(AA <u>c</u> TGT <u>c</u> ACG)	3151.6	3151.6	48.2 (+2.4)	51.0 (+3.3)

^a Total strand conc: 2 μ M in 10 mM NaH₂PO₄, 150 mM NaCl, pH 7.0. Estimated error in T_m = $\pm 0.5^\circ\text{C}$. ^b T_m of unmodified duplex: 43.5 $^\circ\text{C}$. ^c T_m of unmodified duplex: 44.4 $^\circ\text{C}$. ^d T_m of unmodified duplex: 46.3 $^\circ\text{C}$. ^e T_m of unmodified duplex: 42.9 $^\circ\text{C}$

Oligodeoxynucleotides containing single 6'fluoro tc-T mutations (**ON1-3**) lead to duplexes with complementary DNA and RNA with neutral to slightly enhanced stability in a slightly sequence dependent context, compared to unmodified duplexes. Interestingly, the stabilization ($\Delta T_m/\text{mod.}$) is a bit stronger in duplexes with complementary DNA as compared to RNA. Double substitutions in a non-contiguous order (**ON3**) tendentially lead to poorer duplex stabilization with RNA compared to multiple substitutions in a consecutive manner (**ON4**). This is in agreement with earlier observations in the series of tc-DNA,³⁰ and has been ascribed to an incremental energetic penalty arising from increasing numbers of structural hetrobackbone junctions. Replacing natural dC with 6'fluoro-tc-⁵MeC units (**ON5-7**) leads to a somewhat different picture. Again, in a slightly sequence dependent context, duplex stabilization is significantly higher (up to +4.5 $^\circ\text{C}/\text{mod}$) as compared to dC and more pronounced with RNA as a complement. In addition, there seems to be almost no energetic penalty as a function of the number of non-contiguous substitutions (**ON5** and **ON6** vs **ON7**). While it is known that the base thymine in the tc-DNA context is least stabilizing compared to the other three bases,^{30,31} the origin of this effect is yet unknown and awaits further structural investigations.

To elucidate the role of the fluorine atom and the methyl group in 5-methyl cytosine on RNA duplex stability we compared $\Delta T_m/\text{mod}$ data with that of oligonucleotides containing tc-T, tc-C and tc-⁵MeC residues, resp. (Table 2). From this set of data it becomes evident that in both the T- and C-series, the 6'-fluorine atom behaves neutral and does not significantly add to duplex stability. This is in agreement with the absence of any F-H5 pseudo-hydrogen bond, as found in the X-ray structure of **16**, and supports our earlier hypothesis that this interaction is responsible for the increase in stability in the bc-DNA series.²¹ At the same time it is in agreement with the properties of other 6'-modified tc-DNA derivatives for which it was shown before that this position can be chemically modified without compromising RNA affinity.²⁴ In the C-series, the 5-methyl group of cytosine brings about 0.2-1.2 °C/mod of additional thermal stability also in the context of the tricyclic nucleoside structure. As for the case of 5-methyldeoxycytidine in DNA duplexes, this is most likely the consequence of improved stacking interactions and/or improved hydrogen bonding induced by the molecular polarizability of the size extended base.³²

Table 2. Structure affinity relationship: $\Delta T_m/\text{mod}$ data for oligodeoxynucleotides containing parent or substituted tc-nucleosides in complex with complementary RNA.

X	tc-T	6'F-tc-T		tc-C	tc- ⁵ MeC	6'F-tc- ⁵ MeC
d(AACTG X CACG)	-0.1	0.0	d(AACTGT X ACG)	+2.0	+2.2	+1.9
d(AAC X GTACG)	+1.4	+1.0	d(AA X TGTACG)	+3.0	+4.2	+4.5
d(AAC X G X CACG)	+0.4	0.0	d(AA X TGT X ACG)	+2.5	+3.6	+3.3

Experimental conditions as in Table 1.

In the context of future applications as steric block or splice switching oligonucleotides we investigated also the fully modified tc-oligonucleotides **ON8-10** containing 6'-fluoro-tc-T units.

These oligonucleotides all carry a 5'-phosphate unit in order to confer chemical stability to the 5'-terminal nucleoside unit during oligonucleotide deprotection.³³ As can be seen from Table 3, duplexes with complementary DNA (non 5'-phosphorylated) are somewhat destabilized in the presence of the fluorine atom, while a slight stabilization in a sequence dependent manner occurs with RNA (non 5'-phosphorylated) as complement. Thus 6'-fluorination is fully compatible with the tc-DNA backbone and does not lead to loss of RNA affinity.

Table 3. Analytical data of tc-oligonucleotides **ON8-ON10**, containing 6'-fluoro-tc-T (**t**), and T_m -data of duplexes with complementary DNA and RNA.

Sequence ^a	ESI-MS	ESI-MS	T_m (°C) vs DNA ^{b,c}	T_m (°C) vs RNA ^{b,d}
	<i>m/z</i> calc	<i>m/z</i> found	(ΔT_m /mod)	(ΔT_m /mod)
ON8 d(pAACTG t CACG)	3490.5	3489.6	55.0 (0.0)	68.0 (+1.6)
ON9 d(pAA Ct GTCACG)	3490.5	3489.6	53.4 (-1.6)	66.7 (+0.3)
ON10 d(pAA Ct G t CACG)	3508.5	3507.6	53.2 (-1.8)	66.3 (-0.1)

^a Characters in italic denote regular tc-DNA residues, p denotes a 5'-phosphate group. ^b Total strand conc: 2 μ M in 10 mM NaH₂PO₄, 150 mM NaCl, pH 7.0. Estimated error in $T_m = \pm 0.5^\circ\text{C}$. ^c T_m of unmodified duplex: 55.0°C. ^d T_m of unmodified duplex: 66.4°C

To determine the effect of 6'-F-tc-T units on duplex conformation we measured CD spectra of duplexes of **ON4** with complementary DNA and RNA and compared them with the corresponding unmodified duplexes and with duplexes containing tc-T instead of 6'-F-tc-T units (Figure 3). The largest structural deviation occurs in the DNA/DNA duplex series where both, the 6'-F-tc-T and the tc-T units drive the duplex conformation from B to A-like. There are no

significant differences between duplexes with tc-T or 6'fluoro-tc-T, indicating that both adopt an N-type nucleoside conformation. The tendency to adopt a more A-like conformation in duplexes with tc-T or 6'F-tc-T units is also present in the DNA/RNA duplex series. Again, there are no large differences between the tc-T and 6'F-tc-T containing duplexes, perhaps with the exception that the maximum positive ellipticity around 270 nm is blue shifted by ca 10 nm in the case of the latter duplex, with a yet unknown implication on the helix structure.

FIGURE 3. CD-spectra of left: DNA/RNA duplexes and right: DNA/DNA duplexes. Black lines: unmodified duplexes, red lines duplexes with **ON4**, and blue lines duplexes with **ON4** in which 6'-fluoro-tc-T was replaced by tc-T. Experimental conditions as indicated in Table 1.

Conclusion:

We have accomplished the synthesis of the two 6'fluorinated tc-nucleoside building blocks **11** and **18** and have incorporated them into oligodeoxynucleotides and tc-oligonucleotides. We analyzed complementary DNA and RNA affinity by T_m -measurements and determined structural effects of fluorine substitution on duplex conformation by CD spectroscopy. Based on the X-ray structure of **16** as well as on ^1H - and ^{13}C -NMR coupling data on the nucleosides and derivatives we could not find any indications for short fluorine-base F-H6 contacts. This is in surprising

contrast to findings in the 6'-fluoro-bc-DNA series, where such short contacts were observed. Compared to the non-fluorinated tc-nucleosides we find that the fluorine substituent does not significantly alter the thermal melting properties of the corresponding duplexes, irrespective of the nature of the base (thymine vs 5-methylcytosine). This is in agreement with the absence of the glycosidic bond constraining nature of the F-H6 interaction that adds up to +2°C/mod in T_m in the case of the bicyclo-DNA series. The 6'-fluoro modification is also compatible with the tc-DNA backbone as no change or even a slight increase in T_m with complementary RNA was observed. The fluorine atom does also not significantly alter the duplex conformation compared to non-fluorinated tc-DNA as can be seen from the corresponding CD-spectra. Based on these encouraging biophysical data we are now planning to investigate functional efficacy, cellular uptake and *in vivo* tissue distribution of antisense tc-oligonucleotides containing these 6'-F-tc-nucleosides.

Experimental Section

General Methods. All reactions were performed under argon in oven dried glassware. Solvents were dried by filtration over activated alumina or by storage over molecular sieves (4 Å). Column chromatography (CC) was performed on silica gel 60 (230-400 mesh, neutralized with 0.1% of w/Ca). All solvents for CC were of technical grade and distilled prior to use. Thin-layer chromatography (TLC) was performed on silica gel plates. Compounds were visualized either under UV light or by staining in dip solution A: Cer^{IV}-sulfate (10.5 g), phosphormolybdic acid (21 g), conc. H₂SO₄ (60 mL), H₂O (900 mL); or B: KMnO₄ (6 g), K₂CO₃ (40 g), 15% NaOH (3 mL) in H₂O (800 mL)), followed by heating with a heat gun. NMR spectra were recorded at 300 or 400 MHz (¹H), at 75MHz or 100MHz (¹³C), at 376 MHz (¹⁹F) and at 162 MHz (³¹P). Chemical

shifts (δ) are reported relative to the undeuterated residual solvent peak (CHCl_3 : 7.24 ppm (^1H) and 77.2 ppm (^{13}C); DMSO-d_6 : 2.50 ppm (^1H) and 39.5 ppm (^{13}C)). Signal assignments are based on DEPT or APT experiments, and on ^1H , ^1H - and ^1H , ^{13}C -correlation experiments (COSY, HSQC). ^{13}C signal multiplicities include ^1H - and ^{19}F -couplings. ^1H -NMR difference-NOESY experiments were recorded at 400 MHz. Chemical shifts for ^{31}P and ^{19}F NMR (fully proton coupled) are reported relative to 85% H_3PO_4 and CFCl_3 as external standards, respectively. Electrospray ionization in the positive mode (ion trap, ESI^+) was used for high resolution mass detection. The numbering scheme for tc-nucleosides is outlined in Figure 1. For non-nucleoside derivatives the von Baeyer nomenclature has been applied.

(1*S*,3*R*,5*S*)-7-Bromo-8-*tert*-butyldimethylsilyloxy-3-methoxy-2-oxabicyclo[3.3.0]oct-7-en-5-ol (2). To a stirred solution of silylenolether **1** (10.6 g, 37.0 mmol) in dry CH_2Cl_2 (100 mL) was added dropwise a solution of bromine (2.1 mL, 40.7 mmol) in CH_2Cl_2 (200 mL) over a period of 30 min at -78°C , followed by Et_3N (7.7 mL, 55.48 mmol). The cooling bath was subsequently removed and the temperature was allowed to rise to rt. Stirring was continued for another 2 h. Then the reaction mixture was poured into H_2O and extracted with Et_2O . The combined organic layers were washed with water, dried over MgSO_4 and evaporated. The residual dark oil was purified by CC (hexane/ EtOAc 4:1) to give the title compound **2** (12.1 g, 90%) as a yellow oil.

Data for 2. R_f = 0.33 (hexane/ EtOAc 7:3); ^1H NMR (CDCl_3 , 300 MHz) δ 5.04 (d, J = 4.1 Hz, 1H, H-C(3)), 4.57 (m, 1H, H-C(1)), 3.37 (s, 3H, OMe), 3.15 (s, 1H, OH), 2.76 (d, J = 15.3 Hz, 1H, H_b -C(6)), 2.67 (dd, J = 2.2, 15.3 Hz, 1H, H_a -C(6)), 2.25 (d, J = 13.5 Hz, 1H, H_b -C(4)), 2.02 (dd, J = 4.1, 13.5 Hz, 1H, H_a -C(4)), 0.97 (s, 9H, *t*-Bu), 0.24, 0.20 (2s, 2x3H, 2x CH_3); ^{13}C NMR (CDCl_3 , 75 MHz) δ 148.4 (s, C(8)), 105.3 (d, C(3)), 96.7 (s, C(7)), 90.4 (d, C(1)), 83.4 (s, C(5)), 54.7 (q, OMe), 47.2 (t, C(6)), 44.8 (t, C(4)), 25.8 (q, *t*-Bu), 18.4 (s, *t*-Bu), -3.9, -4.2 (2xs, 2x CH_3); ESI^+ -HRMS m/z calcd for $\text{C}_{14}\text{H}_{25}\text{BrNaO}_4\text{Si}$ $[\text{M}+\text{Na}]^+$ 387.0603, 389.0583, found 387.0598, 389.0577.

(1*S*,3*R*,5*S*)-7-Bromo-8-*tert*-butyldimethylsilyloxy-3-methoxy-5-trimethylsilyloxy-2-oxabicyclo[3.3.0]oct-7-en (3). To a stirred solution of bromosilylenolether **2** (12.0 g, 32.74 mmol) in dry pyridine (170 mL) was added BSA (12 mL, 49.11 mmol) at rt and the mixture left overnight. The reaction mixture was diluted with sat aq NaHCO₃ and extracted with Et₂O. The combined organic layers were dried over MgSO₄, evaporated and the residual oil purified by CC (hexane/EtOAc 95:5, with 0.5% of Et₃N) to yield the title compound **3** (13.0 g, 91%) as a yellow oil.

Data for 3. *R*_f = 0.68 (hexane/EtOAc 9:1); ¹H NMR (CDCl₃, 300 MHz) δ 4.99 (dd, *J* = 1.3, 5.3 Hz, 1H, H-C(3)), 4.64 (t, *J* = 1.4 Hz, 1H, H-C(1)), 3.35 (s, 3H, OMe), 2.73 (d, *J* = 1.5 Hz, 2H, H-C(6)), 2.33 (dd, *J* = 1.3, 13.7 Hz, 1H, H_b-C(4)), 2.06 (dd, *J* = 5.3, 13.7 Hz, 1H, H_a-C(4)), 0.97 (s, 9H, *t*-Bu), 0.24, 0.21 (2s, 2x3H, 2xCH₃), 0.15 (s, 9H, TMS); ¹³C NMR (CDCl₃, 75 MHz) δ 148.6 (s, C(8)), 105.7 (d, C(3)), 96.4 (s, C(7)), 90.3 (d, C(1)), 84.8 (s, C(5)), 55.2 (q, OMe), 49.5 (t, C(6)), 48.4 (t, C(4)), 25.8 (q, *t*-Bu), 18.4 (s, *t*-Bu), 1.9 (q, TMS), -3.9, -4.2 (2q, 2xCH₃); ESI⁺- HRMS *m/z* calcd for C₁₇H₃₃BrNaO₄Si₂ [M+Na]⁺ 459.0998, 461.0978, found 459.1002, 461.0981.

(1*S*,3*R*,5*S*)-7-Fluoro-8-*tert*-butyldimethylsilyloxy-3-methoxy-2-oxabicyclo[3.3.0]oct-7-en-5-ol (4). To a stirred solution of bromosilylenolether **3** (6.14 g, 14.04 mmol) in dry THF (211 mL) and ether (42 mL) was added dropwise a solution of *t*-BuLi (1.7 M in pentane, 16.5 mL, 28.08 mmol) at -78°C. After stirring for 20 min the reaction mixture was further cooled to -120°C and NFSI (8.85 g, 28.08 mmol) was added at once, followed by another portion of *t*-BuLi (24.8 mL, 42.12 mmol). The reaction mixture was stirred for 2 hours and then allowed to warm up to -80°C. After

quenching with water (210 mL) the mixture was warmed up to rt and extracted with EtOAc. The combined organic phases were dried over MgSO₄ and evaporated. The residue was then dissolved in dry ether (200 mL), treated with amberlyst 15 (6.1 g) and the mixture stirred for 2 hours at rt. The amberlyst was then filtered off and SiO₂ was added to the filtrate prior to evaporation. Purification by CC (CH₂Cl₂/hexane 7:3 → CH₂Cl₂, + 1% Et₂O) gave the title compound **4** (2.56 g, 60 %) in form of a yellowish solid.

Data for 4. R_f = 0.53 (hexane/EtOAc 3:1); ¹H NMR (CDCl₃, 300 MHz) δ 5.05 (d, J = 4.1 Hz, 1H, H-C(3)), 4.58 (dt, J = 1.8, 8.0 Hz, 1H, H-C(1)), 3.38 (s, 3H, OMe), 3.11 (d, J = 2.3 Hz, 1H, OH), 2.62 (m, 2H, H-C(6)), 2.28 (d, J = 13.4 Hz, 1H, H_b-C(4)), 1.97 (dd, J = 4.1, 13.4 Hz, 1H, H_a-C(4)), 0.95 (s, 9H, *t*-Bu), 0.19, 0.17 (2s, 2x3H, 2xCH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 138.5 (d, J (C,F) = 272.2 Hz, C(7)), 128.6 (d, J (C,F) = 4.0 Hz, C(8)), 104.8 (d, C(3)), 89.9 (dd, J (C,F) = 5.3 Hz, C(1)), 79.7 (d, J (C,F) = 11.6 Hz, C(5)), 54.7 (q, OMe), 47.5 (t, C(4)), 37.4 (td, J (C,F) = 18.6 Hz, C(6)), 25.7 (q, *t*-Bu), 18.3 (s, *t*-Bu), -4.3 (qd, J = 1.7 Hz, CH₃), -4.56 (qd, J = 2.1 Hz, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -138.5 (s, br); ESI⁺-HRMS m/z calcd for C₁₄H₂₅FN₄Si [M+Na]⁺ 327.1404, found 327.1412.

(1S,2S,4S,6R,8S)-2-tert-Butyldimethylsilyloxy-4-fluoro-8-methoxy-9-

oxatricyclo[4.3.0^{1,6}.0^{2,4}]nonane-6-ol (5). To dry CH₂Cl₂ (97 mL) was added a solution of Et₂Zn (1 M in hexane, 48.2 mL 48.20 mmol). The mixture was cooled to 0°C and a solution of TFA (3.69 mL, 48.21 mmol) in CH₂Cl₂ (48 mL) was slowly added. After stirring for 20 min a solution of CH₂I₂ (7.76 mL, 96.42 mmol) in CH₂Cl₂ (48 mL) was added. After another 20 min of stirring, a solution of fluorosilylenolether **4** (2.45 g, 8.04 mmol) in CH₂Cl₂ (48 mL) was added, and the ice bath was removed. After 5 hours of stirring, the reaction mixture was quenched with sat aq NH₄Cl and the layers separated. The aqueous layer was extracted with CH₂Cl₂. The combined

organic layers were washed with sat NaHCO₃, dried over MgSO₄, concentrated, and purified by CC (hexane/EtOAc, 9:1) to yield the title compound **5** (1.79 g, 70%) as a colorless oil.

Data for 5. R_f = 0.33 (hexane/EtOAc 3:1); ¹H NMR (CDCl₃, 300 MHz) δ 5.10 (dd, J = 1.6, 5.2 Hz, 1H, H-C(8)), 3.89 (d, J = 6.0 Hz, 1H, H-C(1)), 3.37 (s, 3H, OMe), 2.49 (m, 1H, H_b-C(5)), 2.43 (dd, J = 5.3, 14.0 Hz, 1H, H_b-C(7)), 2.29 (dd, J = 1.1 Hz, J = 13.4 Hz, 1H, H_a-C(5)), 2.10 (s, br, 1H, OH), 2.06 (dd, J = 1.6, 14.0 Hz, 1H, H_a-C(7)), 1.33 (ddd, J = 2.5, 7.5, 21.5 Hz, 1H, H_b-C(3)), 1.19 (dd, J = 7.5, 8.5 Hz, 1H, H_a-C(3)), 0.91 (s, 9H, *t*-Bu), 0.17 (s, 6H, 2xCH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 105.7 (d, C(8)), 88.7 (dd, J (C,F) = 3.8 Hz, C(1)), 83.2 (d, J (C,F) = 268.2 Hz, C(4)), 82.1 (d, J (C,F) = 5.5 Hz, C(6)), 63.7 (d, J (C,F) = 8.3 Hz, C(2)), 54.9 (q, OMe), 49.9 (t, C(7)), 44.4 (td, J (C,F) = 16.5 Hz, C(5)), 25.9 (q, *t*-Bu), 21.7 (td, J (C,F) = 10.3 Hz, C(3)), 18.3 (s, *t*-Bu), -3.9 (q, CH₃), -4.0 (qd, J (C,F) = 1.8 Hz, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -194.9 (m); ESI⁺-HRMS m/z calcd for C₁₅H₂₇FN₄O₄Si [M+Na]⁺ 341.1560, found 341.1561.

(1S,2S,4S,6R)-2-*tert*-Butyldimethylsilyloxy-4-fluoro-6-trimethylsilyloxy-9-

oxatricyclo[4.3.0^{1.6.0^{2.4}]}non-7-ene (**6**). To a solution of compound **5** (1.51 g, 4.38 mmol) and 2,6-lutidine (2.80 mL, 24.20 mmol) in dry CH₂Cl₂ (10 mL) was added TMSOTf (2.14 mL, 11.84 mmol) dropwise at 0°C. After stirring for 2.5 h at rt, the reaction mixture was diluted with AcOEt, washed with saturated NaHCO₃ and the aqueous phase extracted with AcOEt. The combined organic phases were dried over MgSO₄, evaporated and the residue purified by CC (hexane/Et₂O 95:5) to give the title compound **6** (1.43 g, 91%) as light brownish oil.

Data for 6. R_f = 0.83 (hexane/EtOAc 4:1); ¹H NMR (CDCl₃, 300 MHz) δ 6.33 (d, J = 2.7 Hz, 1H, H-C(8)), 5.15 (d, J = 2.7 Hz, 1H, H-C(7)), 4.24 (d, J = 6.0 Hz, 1H, H-C(1)), 2.52 (ddd, J = 2.8, 11.6, 12.9 Hz, 1H, H_b-C(5)), 2.37 (dd, J = 1.1, 12.9 Hz, 1H, H_a-C(5)), 1.45 (ddd, J = 2.8, 7.4,

10.1 Hz, 1H, H_b-C(3)), 1.02 (dd, $J = 7.4, 8.4$ Hz, 1H, H_a-C(3)), 0.90 (s, 9H, *t*-Bu), 0.15, 0.11 (2s, 2x3H, 2xCH₃), 0.09 (s, 9H, TMS); ¹³C NMR (CDCl₃, 75 MHz) δ 146.7 (d, C(8)), 108.4 (d, C(7)), 94.1 (dd, $J(C,F) = 4.1$ Hz, C(1)), 87.1 (d, $J(C,F) = 14.2$ Hz, C(6)), 84.4 (d, $J(C,F) = 250.2$ Hz, C-(4)), 64.8 (d, $J(C,F) = 8.4$ Hz, C(2)), 48.7 (td, $J(C,F) = 14.4$ Hz, C(5)), 26.0 (q, *t*-Bu), 22.9 (td, $J(C,F) = 10.6$ Hz, C(3)), 18.4 (s, *t*-Bu), 2.0 (q, TMS), -3.8 (q, CH₃), -4.3 (qd, $J(C,F) = 2.8$ Hz, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -192.6 (m); ESI⁺-HRMS m/z calcd for C₁₇H₃₂FO₃Si₂ [M+H]⁺ 359.1874, found 359.1873.

(5'-*O*-(*tert*-Butyldimethylsilyl)- 3'-*O*-(trimethylsilyl)-2'-deoxy-3',5'-ethano-6'-fluoro-2'-iodo-5',6'-methano- β -D-ribofuranosyl)thymine (7). To a suspension of thymine (1.49 g, 11.80 mmol) and compound **6** (1.41 g, 3.93 mmol) in CH₂Cl₂ (20 mL) was added BSA (2.88 mL, 11.80 mmol) and the mixture was stirred at rt for 2 h to become a clear solution. Then *N*-iodosuccinimide (1.32 g, 5.90 mmol) was added and the mixture stirred overnight. The reaction was quenched with sat NaHCO₃ (30 mL) and a 10% aq solution of Na₂S₂O₃ (10 mL). The aqueous phase was extracted with EtOAc and the combined organic phases were dried over MgSO₄ and evaporated. CC (hexane/EtOAc 9:1) afforded nucleosides **7** (2.05 g, 85%) as a yellowish solid.

Data for 7. $R_f = 0.39$ (hexane/EtOAc 4:1); ¹H NMR (CDCl₃, 300 MHz) δ 9.10 (s, 1H, NH), 7.76 (d, $J = 0.9$ Hz, 1H, H-C(6)), 6.38 (d, $J = 2.6$ Hz, 1H, H-C(1')), 4.56 (d, $J = 2.6$ Hz, 1H, H-C(2')), 4.19 (d, $J = 5.7$ Hz, 1H, H-C(4')), 2.45 (d, $J = 14.2$ Hz, 1H, H_b-C(7')), 2.34 (m, 1H, H_a-C(7')), 1.91 (d, $J = 0.9$ Hz, 3H, CH₃), 1.50 (ddd, $J = 2.2, 7.8, 20.9$ Hz, 1H, H_b-C(8')), 1.19 (t, $J = 8.4$ Hz, 1H, H_a-C(8')), 0.94 (s, 9H, *t*-Bu), 0.23 (s, 9H, TMS), 0.22, 0.20 (2s, 2x3H, 2xCH₃); ¹³C NMR (CDCl₃, 75 MHz) δ 164.3 (s, CO), 150.4 (s, CO), 135.4 (d, C(6)), 110.9 (s, C(5)), 96.8 (d, C(1')),

91.2 (dd, $J(C,F) = 3.7$ Hz, C(4')), 83.9 (d, $J(C,F) = 11.4$ Hz, C(3')), 81.4 (d, $J(C,F) = 250.9$ Hz, C(6')), 62.7 (d, $J(C,F) = 8.2$ Hz, C(5')), 41.5 (d, C(2')), 41.3 (td, $J(C,F) = 16.3$ Hz, C(7')), 25.7 (q, *t*-Bu), 21.7 (td, $J(C,F) = 10.2$ Hz, C-(8')), 18.0 (s, *t*-Bu), 12.4 (q, CH₃), 2.1 (q, TMS), -3.6 (q, CH₃), -4.0 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -197.2 (m); ESI⁺-HRMS m/z calcd for C₂₂H₃₇FIN₂O₅Si₂ [M+H]⁺ 611.1270, found 611.1265.

(5'-*O*-(*tert*-Butyldimethylsilyl)- 3'-*O*-(trimethylsilyl)-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano- β -D-ribofuranosyl)thymine (8). To a solution of idonucleoside **7** (2.05 g, 3.35 mmol) in toluene (40 mL) were added Bu₃SnH (1.1 mL, 4.02 mmol) and azoisobutyronitril (AIBN, 165 mg, 1.00 mmol) at rt. After heating to reflux for 1 h the solvent was evaporated and the residue purified by CC (hexane/EtOAc 8:2) to give nucleoside **8** (1,54 mg, 95%) as a colorless solid.

Data for 8. $R_f = 0.52$ (hexane/EtOAc 1:1); ¹H NMR (CDCl₃, 300 MHz) δ 8.52 (s, 1H, NH), 7.82 (d, $J = 1.0$ Hz, 1H, H-C(6)), 6.03 (dd, $J = 1.4, 6.4$ Hz, 1H, H-C(1')), 4.07 (d, $J = 5.6$ Hz, 1H, H-C(4')), 2.62 (dd, $J = 6.4, 13.7$ Hz, 1H, H_b-C(2')), 2.53 (dd, $J = 1.4, 13.7$ Hz, 1H, H_a-C(2')), 2.37 (dd, $J = 1.8, 13.7$ Hz, 1H, H_b-C(7')), 2.13 (m, 1H, H_a-C(7')), 1.91 (d, $J = 1.0$ Hz, 3H, CH₃), 1.40 (ddd, $J = 2.4, 7.7, 21.0$ Hz, 1H, H_b-C(8')), 1.18 (dd, $J = 7.7, 9.1$ Hz, 1H, H_a-C(8')), 0.94 (s, 9H, *t*-Bu), 0.23, 0.19 (2s, 2x3H, 2xCH₃), 0.13 (s, 9H, TMS); ¹³C NMR (CDCl₃, 75 MHz) δ 164.2 (s, CO), 150.2 (s, CO), 136.0 (d, C(6)), 110.2 (s, C(5)), 93.0 (dd, $J(C,F) = 4.0$ Hz, C(4')), 89.6 (d, C(1')), 82.4 (d, $J(C,F) = 12.3$ Hz, C(3')), 81.43 (d, $J(C,F) = 249.7$ Hz, C(6')), 63.2 (d, $J(C,F) = 8.1$ Hz, C(5')), 47.3 (t, C(2')), 44.7 (td, $J(C,F) = 14.9$ Hz, C(7')), 25.7 (q, *t*-Bu), 20.4 (td, $J(C,F) = 10.0$ Hz, C(8')), 18.0 (s, *t*-Bu), 12.3 (q, CH₃), 2.0 (q, TMS), -3.7 (q, CH₃), -3.8 (q, CH₃); ¹⁹F

NMR (CDCl₃, 376 MHz) δ -197.9 (m); ESI⁺-HRMS m/z calcd for C₂₂H₃₈FN₂O₅Si₂ [M+H]⁺ 485.2303, found 485.2295.

(2'-Deoxy- 3',5'-ethano-6'-fluoro-5',6'-methano- β -D-ribofuranosyl)thymine (9). To

a solution of compound **8** (1.48 g, 3.05 mmol) and pyridine (6 mL) in CH₂Cl₂ (30 mL) was added HF-pyridine (1.5 mL, 60.6 mmol) at 0°C. After stirring overnight at rt, SiO₂ (7 g) was added and the mixture stirred for another 15 min. After evaporation the adsorbed product was purified by CC (hexane/EtOAc/EtOH 5:5:1) to yield the title compound **9** (797 mg, 87%) as a white foam.

Data for 9. R_f = 0.22 (EtOAc); ¹H NMR (CD₃OD, 400 MHz) δ 7.81 (d, J = 0.9 Hz, 1H, H-C(6)), 6.13 (dd, J = 4.0, 6.9 Hz, 1H, H-C(1')), 3.99 (d, J = 5.7 Hz, 1H, H-C(4')), 2.56 (dd, J = 6.9, 13.9 Hz, 1H, H_b-C(2')), 2.45 (dd, J = 4.0, 13.9 Hz, 1H, H_a-C(2')), 2.38 (m, 2H, H-C(7')), 1.93 (d, J = 0.9 Hz, 3H, CH₃), 1.43 (ddd, J = 2.4, 7.4, 20.9 Hz, 1H, H_b-(8')), 1.28 (m, 1H, H_a-C(8')); ¹³C NMR (CD₃OD, 100 MHz) δ 166.6 (s, CO), 152.1 (s, CO), 137.7 (d, C(6)), 111.0 (s, C(5)), 91.5 (dd, J (C,F) = 3.7 Hz, C(4')), 89.0 (d, C(1')), 84.4 (d, J (C,F) = 248.0 Hz, C(6')), 80.9 (d, J (C,F) = 12.1 Hz, C(3')), 63.4 (d, J (C,F) = 8.4 Hz, C(5')), 49.0 (t, C(2')), 45.1 (td, J (C,F) = 15.6 Hz, C(7')), 21.1 (td, J (C,F) = 10.4 Hz, C(8')), 12.4 (q, CH₃); ¹⁹F NMR (CD₃OD, 376 MHz) δ -200.4 (m); ESI⁺-HRMS m/z calcd for C₁₃H₁₆FN₂O₅ [M+H]⁺ 299.1038, found 299.1037.

(5'-O-((4,4'-Dimethoxytriphenyl)methyl)-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano- β -D-ribofuranosyl)thymine (10). To a solution of nucleoside **9** (428 mg, 1.44 mmol) in pyridine (20 mL) was added DMTrCl (1.46 g, 4.31 mmol) at rt and the mixture was stirred for 2 days. Then reaction was diluted with sat. NaHCO₃ and extracted with EtOAc. The combined organic phases were dried over MgSO₄, evaporated and the residue purified by CC (hexane/EtOAc 8:2

→ EtOAc, 1% Et₃N) to give the title compound **10** (751 mg, 87%) as a yellowish foam.

Data for 10. R_f = 0.38 (EtOAc); ¹H NMR (CDCl₃, 400 MHz) δ 9.62 (brs, 1H, NH), 8.04 (d, J = 0.8 Hz, 1H, H-C(6)), 7.47 (m, 2H, H-arom), 7.37 (dd, J = 8.9, 10.7 Hz, 4H, H-arom), 7.16 (m, 3H, H-arom), 6.75 (dd, J = 7.6, 8.9 Hz, 4H, H-arom), 5.76 (dd, J = 1.4, 6.5 Hz, 1H, H-C(1')), 3.70 (s, 3H, OMe), 3.69 (s, 3H, OMe), 2.33 (dd, J = 6.5, 14.1 Hz, 1H, H_b-(2')), 2.23 (m, 3H, H_a-C(2')), H-C(4'), H_b-C(7')), 2.09 (m, 1H, H_a-C(7')), 1.98 (d, J = 0.8 Hz, 3H, CH₃), 1.71 (m, 1H, H_b-C(8')), 0.84 (dd, J = 8.6, 9.7 Hz, 1H, H_a-C(8')); ¹³C NMR (CDCl₃, 100 MHz) δ 164.7 (s, CO), 158.95, 158.93 (2s, 2xC-arom), 150.5 (s, CO), 145.8, 136.5, 136.4 (3s, 3xC-arom), 136.3 (d, C-6), 131.2, 131.1, 128.8, 127.8, 127.2, 113.13, 113.08 (7d, 7xC-arom), 110.3 (s, C(5)), 90.6 (dd, J (C,F) = 3.8 Hz, C(4')), 88.82 (s), 88.81 (d, C(1')), 83.3 (d, J (C,F) = 247.8 Hz, C(6')), 80.5 (d, J (C,F) = 12.4 Hz, C(3')), 64.9 (d, J (C,F) = 7.8 Hz, C(5')), 55.4 (q, 2xOMe), 47.9 (t, C(2')), 44.0 (td, J (C,F) = 16.0 Hz, C(7')), 20.2 (td, J (C,F) = 9.5 Hz, C(8')), 12.5 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -193.4 (m); ESI⁺-HRMS m/z calcd for C₃₄H₃₃FN₂O₇Na [M+Na]⁺ 623.2164, found 623.2150.

(5'-O-(4,4'-Dimethoxytriphenyl)methyl)-3'-O-(2-cyanoethoxy)-

diisopropylaminophosphanyl-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano-β-D-

ribofuranosyl)thymine (11). To a solution of compound **10** (4.24 g, 7.06 mmol) and

diisopropylethylamine (4.67 ml, 28.24 mmol) in CH₃CN (142 mL) was added 2-cyanoethoxy-

diisopropylaminochlorophosphine (3.94 mL, 17.65 mmol) at rt. After stirring for 2 h at rt, the

mixture was diluted with EtOAc and washed with sat aq NaHCO₃. The aqueous phases were

extracted with EtOAc and the combined organic phases were dried (MgSO₄), evaporated and the

resulting crude product purified by CC (hex/EtOAc 1:1, 1% NEt₃). The purified product was

dissolved in CH₂Cl₂ (10 mL), slowly added to icecold hexane (220 mL) and the precipitate

collected. This procedure was repeated 7x to yield the pure title compound **11** (3.96 g, 70%) as

a white amorphous solid.

Data for 11. R_f = 0.55 (EtOAc); ¹H NMR (CDCl₃, 400 MHz) δ 8.72 (brs, 1H, NH), 8.01 (dd, J = 1.0, 5.6 Hz, 1H, H-C(6)), 7.48 (d, J = 8.1 Hz, H-arom, 2H), 7.38 (t, J = 9.0 Hz, 4H, H-arom), 7.18 (m, 3H, H-arom), 6.75 (dd, J = 6.9, 8.7 Hz, 4H, H-arom), 5.83 (dd, J = 1.9, 5.9 Hz, 1H, H-C(1')), 3.72 (m, 6H, 2xOMe), 3.63 (m, 1H, OCH₂), 3.50 (m, 1H, OCH₂), 3.37 (m, 2H, 2x (Me₂CH)N), 2.76 (m, 1H, H_b-C(7')), 2.62 (m, 1H, H_b-C(2')), 2.48 (t, J = 6.3 Hz, 2H, CH₂CN), 2.32 (m, 2H, H_a-C(2'), H-(4')), 2.02 (m, 1H, H_a-C(7')), 2.02 (s, 3H, CH₃), 1.72 (m, 1H, H_b-C(8')), 1.01 (m, 12H, 2x(CH₃)₂CHN), 0.84 (m, 1H, H_a-C(8')); ¹³C NMR (CDCl₃, 100 MHz) δ 164.29, 164.26 (s, CO), 159.0 (m, 2xC-arom), 150.1 (s, CO), 145.8, 145.7 (s, C-arom), 136.5 (m, C-arom), 136.0 (d, C(6)), 131.2, 131.0, 128.8, 127.8, 127.3, 127.2, (6d, 6xC-arom), 117.6, 117.5 (2s, CN), 113.14, 113.10 (2d, 2xC-arom), 110.22, 110.19 (2s, C(5)), 91.3 (d, C(4')), 89.5, 89.3 (2d, C(1')), 89.0, 88.9 (2s), 83.8, 83.7 (2d, J (C,F) = 11.6 Hz and 12.4 Hz C-(3')), 83.3, 83.20 (2d, J (C,F) = 248.0, Hz C(6')), 64.5 (m, C(5')), 58.0, 57.7 (2td, J (C,P) = 19.5 Hz, OCH₂), 55.4, 55.3 (2q, 2xOMe), 45.9, 45.4 (2td, J (C,P) = 9.7 Hz and 12.9 Hz, C(2')), 43.4, 43.3 (2dd, J (C,P)

= 6.7 Hz, Me₂CH), 42.1 (m, C(7')), 24.50, 24.45, 24.42, 24.37 (4q, Me₂CH), 20.5, 20.4 (2td, CH₂CN *J* (C,P) = 3.6 Hz, 4.1 Hz), 20.0 (m, C(8')), 12.5 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -193.8 (m), -193.6 (m); ³¹P NMR (CDCl₃, 161 MHz): 145.0, 142.9; ESI⁺-HRMS *m/z* calcd for C₄₃H₅₀FN₄O₈PNa [M+Na]⁺ 823.3243, found 823.3276.

(5'-*O*-(*tert*-Butyldimethylsilyl)-3'-*O*-(trimethylsilyl)-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano-β-D-ribofuranosyl)-4-(1*H*-1,2,4-triazol-1-yl)thymine (12). A suspension of 1,2,4-triazol (19.81 g, 0.29 mol) in CH₃CN was cooled to 0°C and treated consecutively with POCl₃ (2.97 ml, 31.87 mmol) and Et₃N (40.9 ml, 293.25 mmol). The resulting mixture was stirred for 50 min before compound **8** (6.18 g, 12.75 mmol), dissolved in CH₃CN (105 mL), was added. After completion (TLC control, 3.5 h) the reaction was quenched with sat aq NaHCO₃ (200 mL). The ice bath was removed and reduced to half of the volume in vacuo. Then EtOAc (200 mL) was added and the volume again reduced to one third. After pouring onto H₂O/sat aq NaCl 1:1, the resulting mixture was extracted with EtOAc. The combined organic phases were dried (MgSO₄) and evaporated. The crude compound **12** was used directly in the next step without further purification. For analytical data a sample was purified by CC (hexane/EtOAc 50:50).

Data for 12. *R_f* = 0.35 (hexane/EtOAc 1:1); ¹H NMR (CDCl₃, 400 MHz) δ 9.28 (s, 1H, H-C(5')), 8.53 (s, 1H, H-C(6)), 8.11 (s, 1H, H-C(3')), 6.18 (t, *J* = 3.9 Hz, 1H, H-C(1')), 4.18 (d, *J* = 5.6 Hz, 1H, H-C(4')), 2.76 (d, *J* = 3.9 Hz, 2H, H-C(2')), 2.45 (s, 3H, CH₃), 2.34 (dd, *J* = 2.3, 13.9 Hz, 1H, H_b-C(7')), 1.85 (m, 1H, H_a-C(7')), 1.44 (ddd, *J* = 2.3, 7.7, 21.0 Hz, 1H, H_b-C(8')), 1.20 (dd, *J* = 7.7, 9.1 Hz, 1H, H_a-C(8')), 0.96 (s, 9H, *t*-Bu), 0.27, 0.22 (2s, 2x3H, 2xCH₃), 0.13 (s, 9H, TMS); ¹³C NMR (CDCl₃, 100 MHz) δ 158.6 (s, C(4)), 154.0 (s, CO), 153.6 (d, C(3')), 147.6 (d, C(6)), 145.2 (d, C(5')), 106.0 (s, C(5)), 93.5 (dd, *J* (C,F) = 3.9 Hz, C(4')), 91.6 (d, C(1')), 82.3 (d, *J* (C,F) = 12.1 Hz, C(3')), 81.3 (d, *J* (C,F) = 249.3 Hz, C(6')), 63.3 (d, *J* (C,F) = 8.1 Hz, C(5')), 46.4 (t, C(2')), 45.0 (td, *J* (C,F) = 14.7 Hz, C(7')), 25.7 (q, *t*-Bu), 20.2 (td, *J* (C,F) = 10.0 Hz,

C(8')), 18.0 (s, *t*-Bu), 16.9 (q, CH₃), 2.0 (q, TMS), -3.7 (q, CH₃), -3.8 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -197.7 (m); ESI⁺-HRMS *m/z* calcd for C₂₄H₃₉FN₅O₄Si₂ [M+H]⁺ 536.2519, found 536.2503.

(5'-*O*-*tert*-Butyldimethylsilyl)-3'-*O*-(trimethylsilyl)-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano-β-D-ribofuranosyl)-5-methyl cytosine (13). To a solution of the crude compound **12** of the previous step (10 g), dissolved in 1,4-dioxane (110 mL), was added conc NH₄OH (110 mL) and the mixture was stirred for 2 h at rt. The solvent was evaporated and the residue dissolved in EtOAc and extracted with H₂O and sat NaCl. The aqueous phases were extracted with EtOAc and the combined organic layers dried (MgSO₄) and evaporated. The residue was purified by CC (EtOAc → EtOAc/EtOH 9:1) to yield the title compound **13** (3.70 g, 60%) as a white foam.

Data of 13. *R*_f = 0.48 (EtOAc/EtOH 9:1); ¹H NMR (CDCl₃, 400 MHz) δ 7.92 (d, *J* = 0.7 Hz, 1H, H-C(6)), 6.08 (m, 1H, H-C(1')), 4.12 (m, 1H, H-C(4')), 2.64 (d, *J* = 3.4 Hz, 2H, H-C(2')), 2.28 (dd, *J* = 2.0, 13.7 Hz, 1H, H_b-C(7')), 1.99 (m, 1H, H_a-C(7')), 1.94 (d, *J* = 0.7 Hz, 3H, CH₃), 1.37 (ddd, *J* = 2.4, 7.6, 21.0 Hz, 1H, H_b-C(8')), 1.17 (dd, *J* = 7.6, 9.1 Hz, 1H, H_a-C(8')), 0.94 (s, 9H, *t*-Bu), 0.24, 0.19 (2s, 2x3H, 2xCH₃), 0.10 (s, 9H, TMS); ¹³C NMR (CDCl₃, 100 MHz) δ 166.1 (s, C(4)), 156.3 (s, CO), 139.0 (d, C(6)), 101.6 (s, C(5)), 92.9 (dd, *J* (C,F) = 3.9 Hz, C(4')), 90.3 (d, C(1')), 82.3 (d, *J* (C,F) = 12.3 Hz, C(3')), 81.5 (d, *J* (C,F) = 249.0 Hz, C(6')), 63.3 (d, *J* (C,F) = 8.0 Hz, C(5')), 46.9 (t, C(2')), 44.7 (td, *J* (C,F) = 14.7 Hz, C(7')), 25.7 (q, *t*-Bu), 20.2 (td, *J* (C,F) = 9.9 Hz, C(8')), 18.0 (s, *t*-Bu), 13.0 (q, CH₃), 2.0 (q, TMS), -3.7, -3.8 (2q, 2xCH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -198.0 (m); ESI⁺-HRMS *m/z* calcd for C₂₂H₃₉FN₃O₄Si₂ [M+H]⁺ 484.2458, found 484.2450.

***N*⁴-Benzoyl-1-(5′-*O*-(*tert*-butyldimethylsilyl)-3′-*O*-(trimethylsilyl)-2′-deoxy-3′,5′-ethano-6′-fluoro-5′,6′-methano-β-D-ribofuranosyl)-5-methyl cytosine (14) and *N*⁴-benzoyl-1-(5′-*O*-(*tert*-butyldimethylsilyl)-2′-deoxy-3′,5′-ethano-6′-fluoro-5′,6′-methano-β-D-ribofuranosyl)-5-methyl cytosine (15).** To a solution of nucleoside **13** (492 mg, 1.01 mmol) and DMAP (25 mg, 0.20 mmol) in CH₃CN (20 mL) was added Bz₂O (253 mg, 1.12 mmol) and the mixture was stirred for 1.5 h at rt. Then Et₃N (0.28 mL, 2.03 mmol) was added and the mixture stirred overnight. After evaporation of the solvents the residue was dissolved in EtOAc and washed with H₂O. The aqueous phase was extracted with EtOAc and the combined organic phases dried (MgSO₄) and evaporated. The crude product was purified by CC (hexane/EtOAc 95:5 → Hex:EtOAc 50:50) to give compound **14** (39%) as a white foam and compound **15** (54%) as a white solid.

Data for 14. *R*_f = 0.62 (hexane/EtOAc 1:1); ¹H NMR (CDCl₃, 400 MHz) δ 13.46 (s, 1H, NH), 8.32 (d, *J* = 7.3 Hz, 2H, H-arom), 8.03 (m, 1H, H-C(6)), 7.52 (m, 1H, H-arom), 7.44 (m, 2H, H-arom), 6.09 (dd, *J* = 1.1, 6.5 Hz, 1H, H-C(1′)), 4.11 (d, *J* = 5.6 Hz, 1H, H-C(4′)), 2.67 (dd, *J* = 6.5, 13.7 Hz, 1H, H_b-C(2′)), 2.60 (dd, *J* = 1.1, 13.7 Hz, 1H, H_a-C(2′)), 2.38 (dd, *J* = 1.9, 13.7 Hz, 1H, H_b-C(7′)), 2.12 (d, *J* = 0.9 Hz, 3H, CH₃), 2.11 (m, 1H, H_a-C(7′)), 1.43 (ddd, *J* = 2.4, 7.7, 21.0 Hz, 1H, H_b-C(8′)), 1.20 (dd, *J* = 7.7, 9.0 Hz, 1H, H_a-C(8′)), 0.97 (s, 9H, *t*-Bu), 0.26, 0.22 (2s, 2x3H, 2xCH₃), 0.14 (s, 9H, TMS); ¹³C NMR (CDCl₃, 100 MHz) δ 179.8 (s, C(4)), 160.4 (s, CO), 147.9 (s, CO), 137.49 (s, C-arom), 137.45 (d, C(6)), 132.4, 130.0, 128.2 (3d, 3xC-arom), 111.4 (s, C(5)), 93.2 (dd, *J* (C,F) = 4.0 Hz, C(4′)), 90.1 (d, C(1′)), 82.3 (d, *J* (C,F) = 12.3 Hz, C(3′)), 81.4 (d, *J* (C,F) = 249.5 Hz, C(6′)), 63.2 (d, *J* (C,F) = 8.1 Hz, C(5′)), 47.1 (t, C(2′)), 44.8 (td, *J* (C,F) = 14.9 Hz, C(7′)), 25.7 (q, *t*-Bu), 20.3 (td, *J* (C,F) = 10.1 Hz, C(8′)), 18.0 (s, *t*-Bu), 13.4 (q, CH₃), 2.0 (q, TMS), -3.7 (q, CH₃), -3.8 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -197.9 (m);

ESI⁺-HRMS *m/z* calcd for C₂₉H₄₃FN₃O₅Si₂ [M+H]⁺ 588.2720, found 588.2714.

Data for 15. *R*_f = 0.30 (hexane/EtOAc 1:1); ¹H NMR (CDCl₃, 400 MHz) δ 8.29 (m, 2H, H-arom), 8.07 (d, *J* = 0.8 Hz, 1H, H-C(6)), 7.48 (m, 4H, H-arom, NH), 6.09 (dd, *J* = 2.3, 5.6 Hz, 1H, H-C(1')), 4.13 (d, *J* = 5.5 Hz, 1H, H-C(4')), 2.64 (m, 2H, H-C(2')), 2.35 (dd, *J* = 1.6, 13.9 Hz, 1H, H_b-C(7')), 2.20 (m, 1H, H_a-C(7')), 2.12 (d, *J* = 0.8 Hz, 3H, CH₃), 1.46 (ddd, *J* = 2.4, 7.8, 21.0 Hz, 1H, H_b-C(8')), 1.28 (m, 1H, H_a-C(8')), 0.97 (s, 9H, *t*-Bu), 0.25, 0.21 (2s, 2x3H, 2xCH₃); ¹³C NMR (CDCl₃, 100 MHz) δ 179.4 (s, C(4)), 160.5 (s, CO), 148.2 (s, CO), 137.9 (d, C(6)), 137.2 (s, C-arom), 132.6, 123.0, 128.3 (3d, 3xC-arom), 111.5 (s, C(5)), 91.8 (dd, *J* (C,*F*) = 3.8 Hz, C(4')), 90.0 (d, C(1')), 81.5 (d, *J* (C,*F*) = 249.1 Hz, C(6')), 80.7 (d, *J* (C,*F*) = 12.0 Hz, C(3')), 63.5 (d, *J* (C,*F*) = 8.2 Hz, C(5')), 47.9 (t, C(2')), 44.8 (td, *J* (C,*F*) = 15.8 Hz, C(7')), 25.7 (q, *t*-Bu), 20.5 (td, *J* (C,*F*) = 10.1 Hz, C(8')), 18.0 (s, *t*-Bu), 13.4 (q, CH₃), -3.7 (q, CH₃), -3.8 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -197.6 (m); ESI⁺-HRMS *m/z* calcd for C₂₆H₃₅FN₃O₅Si [M+H]⁺ 516.2325, found 516.2328.

***N*⁴-Benzoyl-1-(2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano-β-D-ribofuranosyl)-5-methyl cytosine (16).** To separate solutions of nucleoside **14** and **15** (1.85 g, 3.15 mmol and 1.24 g, 2.41 mmol resp.) and pyridine (9.6 mL and 6.0 mL, resp.) in CH₂Cl₂ (40 mL and 30 mL, resp.) was added HF-pyridine (3.3 mL and 2.1 mL, resp.) at 0°C. After stirring for 24 h at rt, silica gel (1g per 300 mg starting material) was added and the mixtures stirred for another 15 min. After evaporation the adsorbed products were purified by CC (EtOAc) to give the title compound **16** (1.13 g from **14**, 867 mg from **15**, 90% together) as white foams.

Data for 16. *R*_f = 0.48 (EtOAc); ¹H NMR (CD₃OD, 400 MHz) δ 8.23 (m, 2H, H-arom), 8.11 (m, 1H, H-C(6)), 7.58 (m, 1H, H-arom), 7.47 (m, 2H, H-arom), 6.15 (dd, *J* = 3.1, 7.0 Hz, 1H, H-C(1')), 4.08 (d, *J* = 5.6 Hz, 1H, H-C(4')), 2.65 (dd, *J* = 7.0, 14.0 Hz, 1H, H_b-(2')), 2.53 (dd, *J* =

3.1, 14.0 Hz, 1H, H_a-C(2')), 2.32 (m, 2H, H-C(7')), 2.14 (d, 3H, *J* = 0.9 Hz, CH₃), 1.45 (dd, *J* = 7.5, 20.9 Hz, 1H, H_b-C(8')), 1.31 (t, *J* = 7.5 Hz, 1H, H_a-C(8')); ¹³C NMR (CD₃OD, 75 MHz) δ 162.1 (s, C(4)), 148.0 (s, CO), 143.4 (s, CO), 137.7 (d, C(6)), 137.6 (s, C-arom), 133.6, 130.5, 129.3 (3d, 3xC-arom), 112.0 (s, C(5)), 91.9 (dd, *J* (C,*F*) = 3.7 Hz, C(4')), 90.2 (d, C(1')), 84.3 (d, *J* (C,*F*) = 248.1 Hz, C(6')), 80.8 (d, *J* (C,*F*) = 12.0 Hz, C(3')), 63.4 (d, *J* (C,*F*) = 8.5 Hz, C(5')), 48.2 (t, C(2')), 45.2 (td, *J* (C,*F*) = 15.5 Hz, C(7')), 21.1 (td, *J* (C,*F*) = 10.3 Hz, C(8')), 13.8 (q, CH₃); ¹⁹F NMR (CD₃OD, 376 MHz) δ -200.4 (m); ESI⁺-HRMS *m/z* calcd for C₂₀H₂₁FN₃O₅ [M+H]⁺ 402.1460, found 402.1461.

***N*⁴-Benzoyl-1-(5'-*O*-((4,4'-dimethoxytriphenyl)methyl)-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano-β-D-ribofuranosyl)-5-methyl cytosine (17).** To a stirred solution of compound **16** (606 mg, 1.51 mmol) in pyridine (20 mL) was added DMTrCl (1.54 g, 4.53 mmol) at rt. After 2 days the mixture was poured onto sat aq NaHCO₃ and extracted with EtOAc. The combined organic layers were dried (MgSO₄), evaporated and the crude material purified by CC (hexane/EtOAc 8:2 → 6:4, +0,2% Et₃N) to give the title compound **17** (843 mg, 79%) as a yellowish foam.

Data for 17. *R*_f = 0.44 (hexane/EtOAc 1:3); ¹H NMR (CDCl₃, 400 MHz) δ 13.37 (brs, 1H, NH), 8.26 (m, 2H, H-arom, H-C(6)), 7.42 (m, 9H, H-arom), 7.20 (m, 4H, H-arom), 6.77 (dd, *J* = 7.6, 9.0 Hz, 4H, H-arom), 5.85 (dd, *J* = 1.5, 6.4 Hz, 1H, H-C(1')), 3.73 (s, 3H, OMe), 3.72 (s, 3H, OMe), 2.39 (dd, *J* = 1.5, 14.1 Hz, 1H, H_b-C(2')), 2.33 (dd, *J* = 6.4, 14.1 Hz, 1H, H_a-C(2')), 2.22 (d, *J* = 0.7 Hz, 3H, CH₃), 2.21 (m, 2H, H-C(4'), H_b-C(7')), 2.11 (m, 1H, H_a-C(7')), 1.77 (ddd, *J* = 2.1, 8.3, 20.4 Hz, 1H, H_b-C(8')), 1.56 (brs, 1H, OH), 0.85 (dd, *J* = 8.3, 9.6 Hz, 1H, H_a-C(8')); ¹³C NMR (CDCl₃, 100 MHz) δ 181.3 (s, C(4)), 160.4 (s, CO), 159.08, 159.06 (2s, 2xC-arom), 147.9 (s, CO), 145.9 (s, C-arom), 137.5 (d, C(6)), 137.4, 136.44, 136.37 (3s, 3xC-arom), 132.6, 131.2,

131.1, 130.0, 128.9, 128.3, 127.9, 127.4, 113.19, 113.15 (10d, 10xC-arom), 111.1 (s, C(5)), 90.9 (dd, $J(C,F) = 4.1$ Hz, C(4')), 89.4 (d, C(1')), 89.0 (s), 83.2 (d, $J(C,F) = 248.1$ Hz, C(6')), 80.8 (d, $J(C,F) = 12.2$ Hz, C(3')), 64.9 (d, $J(C,F) = 7.9$ Hz, C(5')), 55.4 (q, 2xOMe), 47.9 (t, C(2')), 44.2 (td, $J(C,F) = 16.2$ Hz, C(7')), 20.2 (td, $J(C,F) = 9.7$ Hz, C(8')), 13.6 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -193.5 (m); ESI⁺-HRMS m/z calcd for C₄₁H₃₉FN₃O₇ [M+H]⁺ 704.2767, found 704.2771.

***N*⁴-Benzoyl-1-(5'-*O*-((4,4'-dimethoxytriphenyl)methyl)- 3'-*O*-(2-cyanoethoxy)-diisopropylaminophosphanyl-2'-deoxy-3',5'-ethano-6'-fluoro-5',6'-methano- β -D-ribofuranosyl)-5-methyl cytosine (18).** To a solution of compound **17** (2.37 g, 3.37 mmol) and diisopropylethylamine (2.23 mL, 13.47 mmol) in CH₃CN (50 mL) was added 2-cyanoethoxy-diisopropylaminochlorophosphine (1.88 mL, 8.42 mmol). After stirring for 2 h at rt, the mixture was diluted with EtOAc and washed with sat aq NaHCO₃. The aqueous phases were extracted with EtOAc and the combined organic phases dried (MgSO₄), evaporated and the crude product purified by CC (hexane/EtOAc 1:1, +1% NEt₃) to give the title compound **18** (2.78 g, 90 %) as a yellowish foam.

Data for 18. $R_f = 0.71$ (hexane/EtOAc 1:3); ¹H NMR (CDCl₃, 400 MHz) δ 13.39 (brs, 1H, NH), 8.25 (m, 3H, H-arom, H-C(6)), 7.43 (m, 9H, H-arom), 7.18 (m, 3H, H-arom), 6.76 (m, 4H, H-arom), 5.88 (d, $J = 6.8$ Hz, 1H, H-C(1')), 3.72 (m, 6H, 2xOMe), 3.62 (m, 1H, OCH₂), 3.49 (m, 1H, OCH₂), 3.36 (m, 2H, 2x (Me₂CH)N), 2.76 (m, 1H, H_b-C(7')), 2.62 (m, 2H, H-C(2')), 2.47 (m, 2H, CH₂CN), 2.36 (m, 1H, H-C(4')), 2.21 (s, 3H, CH₃), 2.01 (m, 1H, H_a-C(7')), 1.74 (m, 1H, H_b-C(8')), 1.01 (m, 12H, 2x(CH₃)₂CHN), 0.85 (m, 1H, H_a-C(8')); ¹³C NMR (CDCl₃, 100 MHz) δ 179.7 (s, C(4)), 160.4 (s, CO), 159.1, 159.04, 159.02, 159.01 (4s, 4xC-arom), 147.8 (s, CO), 145.70, 145.65 (2s, C-arom), 137.4 (d, C(6)), 136.43, 136.39 136.32 (3s, 3xC-arom), 132.4,

131.2, 131.0, 130.0, 128.8, 128.2, 127.9, 127.3, 127.2, (9d, 9xC-arom), 117.5, 117.4 (2s, CN), 113.2, 113.1 (2d, 2xC-arom), 111.31, 111.27 (2s, C(5')), 91.5 (md, C(4')), 90.0, 89.9 (2d, C(1')), 89.03, 88.98 (2s), 83.7 (m, C(3')), 83.3, 83.2 (2d, $J(C,F) = 248.6$ Hz, C(6')), 64.63, 64.55 (2d, $J(C,F) = 3.6$ Hz, C(5')), 58.0, 57.7 (2td, $J(C,P) = 19.4$ Hz, OCH₂), 55.34, 55.30 (2q, 2xOMe), 45.7, 45.2 (2td, $J(C,P) = 9.7, 12.9$ Hz, C(2')), 43.4, 43.3 (2dd, $J(C,P) = 12.6$ Hz, 2xMe₂CH), 42.2, 42.1 (2td, $J(C,F) J(C,P) = 10.5, 12.3$ Hz, C(7')), 24.53, 24.47, 24.40, 24.35 (4q, 2xMe₂CH), 20.4, 20.3 (2td $J(C,P) = 2.0, 2.7$ Hz, CH₂CN), 20.0, 19.9 (2td, $J(C,F) = 10.2$ Hz, C(8')), 13.5 (q, CH₃); ¹⁹F NMR (CDCl₃, 376 MHz) δ -193.7 (m), -193.5 (m); ³¹P NMR (CDCl₃, 161 MHz) δ 145.1, 143.0; ESI⁺-HRMS m/z calcd for C₅₀H₅₆FN₅O₈P [M+H]⁺ 904.3845, found 904.3846.

Oligonucleotide synthesis and purification. Oligonucleotides **ON1-10** were synthesized by standard solid phase phosphoramidite methodology on the 1.3 μ mol scale on a Pharmacia LKB Gene Assembler Special DNA Synthesizer using a slightly modified DNA synthesis program. Natural phosphoramidites (dT, dC⁴Bz, dA⁶Bz, dG²dmf) were coupled as a 0.1 M solution in CH₃CN, tricyclophosphoramidites as 0.15 M solutions in CH₃CN with the exception of 6'F-tc-T, 6'F-tc-⁵MeC⁴Bz and tc-A that were used as 0.15 M solutions in DCE. The coupling step was 90 s for natural phosphoramidites and 12 min for tricyclo-phosphoramidites. As coupling reagent, 5-(ethylthio)-1H-tetrazole (0.25 M in CH₃CN) was used. Capping, oxidation and detritylation were carried out using standard solutions as described in the manufacturer's protocol. Deprotection of the oligonucleotides after assembly and detachment from solid support was effected by standard ammonia treatment (33% aq NH₃, 16h, 55°C). The crude oligomers were purified by ion-exchange HPLC using a DNAPAC PA200, 4 x 250 mm analytical column (Dionex). Mobile phases A: 25 mM TRIZMA in H₂O, pH 8.0. B: 25 mM TRIZMA, 1.25 M NaCl in H₂O, pH 8.0.

or A: 10 mM NaOH in H₂O, pH 12.0. B: 10 mM NaOH, 1.5 M NaCl in H₂O, pH 12.0, flow 1ml/min detection at 260 nm. Purified oligonucleotides were desalted over Sep-Pak cartridges, quantified at 260 nm using extinction coefficients as determined previously for tricyclo-nucleosides,³¹ and analyzed by ESI- mass spectrometry. Oligonucleotides were then stored at -18°C.

UV-melting curves. Absorbances were monitored at 260 nm and the heating rate was set to 0.5°C/min. A cooling-heating-cooling cycle in the temperature range 20-80°C was applied. *T_m* values were obtained from the derivative curves using the Varian WinUV software. To avoid evaporation of the solution, the sample solutions were covered with a layer of dimethylpolysiloxane. All measurements were carried out in 150 mM NaCl, 10 mM Na-phosphate, pH 7.0 with duplex concentration of 2 µM.

CD-spectroscopy. CD spectra were recorded using the same buffer conditions and oligonucleotide concentrations as for UV melting curves. All CD spectra were collected at 20°C between 210 to 320 nm at a 50 nm/min rate and were baseline-corrected against buffer. The reported spectra correspond to the average of at least three scans.

Acknowledgments

Financial support by the Swiss National Science Foundation (grant-No.: 200020-130373) and by the Association Monégasques contre les Myopathies (AMM) is gratefully acknowledged. We thank the group of Chemical Crystallography of the University of Bern (PD Dr. P. Macchi) for the X-ray structure solution and the Swiss National Science Foundation (R'equip project 206021_128724) for co-funding the single crystal X-ray diffractometer at the department of Chemistry and Biochemistry of the University of Bern.

Supporting Information

Copies of ^1H -, ^{13}C -, ^{19}F -, and ^{31}P - NMR spectra of compounds **2–18** and X-ray structural data (CIF) of compound **16**. This information is available free of charge via the Internet at <http://pubs.acs.org>

References

- (1) Biffinger, J. C.; Kim, H. W.; DiMagno, S. G. *ChemBioChem* **2004**, *5*, 622.
- (2) Dalvit, C.; Vulpetti, A. *ChemMedChem* **2011**, *6*, 104.
- (3) Hagmann, W. K. *J. Med. Chem.* **2008**, *51*, 4359.
- (4) Patra, A.; Paolillo, M.; Charisse, K.; Manoharan, M.; Rozners, E.; Egli, M. *Angew. Chem. Int. Ed.* **2012**, *51*, 11863.
- (5) Pallan, P. S.; Greene, E. M.; Jicman, P. A.; Pandey, R. K.; Manoharan, M.; Rozners, E.; Egli, M. *Nucleic Acids Res.* **2011**, *39*, 3482.
- (6) Anzahae, M. Y.; Watts, J. K.; Alla, N. R.; Nicholson, A. W.; Damha, M. J. *J. Am. Chem. Soc.* **2011**, *133*, 728.
- (7) Watts, J. K.; Martin-Pintado, N.; Gomez-Pinto, I.; Schwartzentruber, J.; Portella, G.; Orozco, M.; Gonzalez, C.; Damha, M. J. *Nucleic Acids Res.* **2010**, *38*, 2498.
- (8) Allerson, C. R.; Sioufi, N.; Jarres, R.; Prakash, T. P.; Naik, N.; Berdeja, A.; Wanders, L.; Griffey, R. H.; Swayze, E. E.; Bhat, B. *J. Med. Chem.* **2005**, *48*, 901.

- (9) Dowler, T.; Bergeron, D.; Tedeschi, A.-L.; Paquet, L.; Ferrari, N.; Damha, M. J. *Nucleic Acids Res.* **2006**, *34*, 1669.
- (10) Kalota, A.; Karabon, L.; Swider, C. R.; Viazovkina, E.; Elzagheid, M.; Damha, M. J.; Gewirtz, A. M. *Nucleic Acids Res.* **2006**, *34*, 451.
- (11) Egli, M.; Pallan, P. S.; Allerson, C. R.; Prakash, T. P.; Berdeja, A.; Yu, J.; Lee, S.; Watt, A.; Gaus, H.; Bhat, B.; Swayze, E. E.; Seth, P. P. *J. Am. Chem. Soc.* **2011**, *133*, 16642.
- (12) Pallan, P. S.; Yu, J.; Allerson, C. R.; Swayze, E. E.; Seth, P.; Egli, M. *Biochemistry* **2011**, *51*, 7.
- (13) Seth, P. P.; Yu, J.; Jazayeri, A.; Pallan, P. S.; Allerson, C. R.; Østergaard, M. E.; Liu, F.; Herdewijn, P.; Egli, M.; Swayze, E. E. *J. Org. Chem.* **2012**, *77*, 5074.
- (14) Seth, P. P.; Pallan, P. S.; Swayze, E. E.; Egli, M. *ChemBioChem* **2013**, *14*, 58.
- (15) Jung, M. E.; Dwight, T. A.; Vigant, F.; Østergaard, M. E.; Swayze, E. E.; Seth, P. P. *Angew. Chem. Int. Ed.* **2014**, *53*, 9893.
- (16) Altmann, K. H.; Kesselring, R.; Francotte, E.; Rihs, G. *Tetrahedron Lett.* **1994**, *35*, 2331.
- (17) Marquez, V. E.; Siddiqui, M. A.; Ezzitouni, A.; Russ, P.; Wang, J.; Wagner, R. W.; Matteucci, M. D. *J. Med. Chem.* **1996**, *39*, 3739.
- (18) Østergaard, M. E.; Dwight, T.; Berdeja, A.; Swayze, E. E.; Jung, M. E.; Seth, P. P. *J. Org. Chem.* **2014**, *79*, 8877.
- (19) Østergaard, M. E.; Southwell, A. L.; Kordasiewicz, H.; Watt, A. T.; Skotte, N. H.; Doty, C. N.; Vaid, K.; Villanueva, E. B.; Swayze, E. E.; Frank Bennett, C.; Hayden, M. R.; Seth, P. P. *Nucleic Acids Res.* **2013**, *41*, 9634.
- (20) Rigo, F.; Hua, Y.; Chun, S. J.; Prakash, T. P.; Krainer, A. R.; Bennett, C. F. *Nat. Chem. Biol.* **2012**, *8*, 555.

- (21) Dugovic, B.; Leumann, C. J. *J. Org. Chem.* **2014**, 79, 1271.
- (22) Steffens, R.; Leumann, C. *Helv. Chim. Acta* **1997**, 80, 2426.
- (23) Lorenz, J. C.; Long, J.; Yang, Z.; Xue, S.; Xie, Y.; Shi, Y. *J. Org. Chem.* **2003**, 69, 327.
- (24) Lietard, J.; Leumann, C. J. *J. Org. Chem.* **2012**, 77, 4566.
- (25) Luisier, S.; Silhar, P.; Leumann, C. J. *Nucleic Acids Symp. Ser.* **2008**, 581.
- (26) Divakar, K. J.; Reese, C. B. *J. Chem. Soc. Perkin 1* **1982**, 1171.
- (27) MacMillan, A. M.; Verdine, G. L. *J. Org. Chem.* **1990**, 55, 5931.
- (28) Saenger, W. *Principles of Nucleic Acid Structure*; Springer-Verlag: New York, 1984.
- (29) Dugovic, B.; Wagner, M.; Leumann, C. J. *Beilstein J. Org. Chem.* **2014**, 10, 1840.
- (30) Ittig, D.; Gerber, A. B.; Leumann, C. J. *Nucleic Acids Res.* **2011**, 39, 373.
- (31) Renneberg, D.; Leumann, C. J. *J. Am. Chem. Soc.* **2002**, 124, 5993.
- (32) Sowers, L. C.; Shaw, B. R.; Sedwick, W. D. *Biochem. Biophys. Res. Commun.* **1987**, 148, 790.
- (33) Steffens, R.; Leumann, C. J. *J. Am. Chem. Soc.* **1999**, 121, 3249.