

Optimal confidence bands for shape-restricted curves

LUTZ DÜMBGEN

Department of Mathematical Statistics and Actuarial Science, University of Berne, Sidlerstrasse 5, CH-3012 Berne, Switzerland. E-mail: duembgen@stat.unibe.ch

Let Y be a stochastic process on $[0, 1]$ satisfying $dY(t) = n^{1/2}f(t)dt + dW(t)$, where $n \geq 1$ is a given scale parameter ('sample size'), W is standard Brownian motion and f is an unknown function. Utilizing suitable multiscale tests, we construct confidence bands for f with guaranteed given coverage probability, assuming that f is isotonic or convex. These confidence bands are computationally feasible and shown to be asymptotically sharp optimal in an appropriate sense.

Keywords: adaptivity; concave; convex; isotonic; kernel estimator; local smoothness; minimax bounds; multiscale testing

1. Introduction

Nonparametric statistical models often involve some unknown function f defined on a real interval J . For instance, f might be the probability density of some distribution or a regression function. Nonparametric point estimators for such a curve f are abundant. The available methods are based on kernels, splines, local polynomials, or orthogonal series, including wavelets; see Hart (1997) and references cited therein. In order to quantify the precision of estimation, one often wishes to replace a point estimator with a confidence band $(\hat{\ell}, \hat{u})$ for f . The latter consists of two functions, $\hat{\ell} = \hat{\ell}(\cdot, \text{data})$ and $\hat{u} = \hat{u}(\cdot, \text{data})$, on J with values in $[-\infty, \infty]$ such that, hopefully, $\hat{\ell} \leq f \leq \hat{u}$ pointwise. More precisely, one is aiming at a confidence band such that

$$P\{\hat{\ell} \leq f \leq \hat{u}\} \geq 1 - \alpha \quad (1.1)$$

for a given level $\alpha \in]0, 1[$, while $\hat{\ell}$ and \hat{u} should be as close to each other as possible.

Unfortunately, curve estimation is an ill-posed problem, and usually there are no non-trivial bands $(\hat{\ell}, \hat{u})$ satisfying (1.1) for arbitrary f ; see Donoho (1988). Therefore one has to impose some additional restrictions on f . Smoothness constraints on f are one possibility, for instance an upper bound on a certain derivative of f . Under such restrictions, (1.1) can be achieved approximately for large sample sizes; see, for example, Bickel and Rosenblatt (1973), Knafl *et al.* (1985), Hall and Titterington (1988), Härdle and Marron (1991), Eubank and Speckman (1993), Fan and Zhang (2000), and the references cited therein.

A problem with the aforementioned methods is that smoothness constraints are hard to justify in practical situations. More precisely, even if the underlying curve f is infinitely often differentiable, the actual coverage probabilities of the confidence bands mentioned

above depend on quantitative properties of certain derivatives of f which are difficult to obtain from the data.

In many applications qualitative assumptions about f such as monotonicity, unimodality or concavity/convexity are plausible. Growth curves in medicine are an example, for example where $f(x)$ is the mean body height of newborns at age x . Here isotonicity of f is a plausible assumption. So-called Engel curves in econometrics are another example, where $f(x)$ is the mean expenditure on certain consumer goods of households with annual income x . Here one expects f to be isotonic and sometimes also concave. Under such qualitative assumptions it is possible to construct $1 - \alpha$ confidence sets for f based on certain goodness-of-fit tests without relying on asymptotic arguments. Examples of such procedures can be found in Davies (1995), Hengartner and Stark (1995) and Dümbgen (1998). In particular, these papers present confidence bands $(\hat{\ell}, \hat{u})$ for f such that

$$P\{\hat{\ell} \leq f \leq \hat{u}\} \geq 1 - \alpha \quad \text{whenever } f \in \mathcal{F}. \quad (1.2)$$

Here \mathcal{F} denotes the specified class of functions. Given a suitable distance measure $D(\cdot, \cdot)$ for functions, the goal is to find a band $(\hat{\ell}, \hat{u})$ satisfying (1.2) such that either $D(\hat{u}, \hat{\ell})$ or $D(\hat{\ell}, f)$ and $D(\hat{u}, f)$ are as small as possible. The phrase ‘as small as possible’ can be interpreted in the sense of optimal rates of convergence to zero as the sample size n tends to infinity. The papers of Hengartner and Stark (1995) and Dümbgen (1998) contain such optimality results.

In the present paper we investigate the optimality of confidence bands in more detail. In addition to optimal rates of convergence, we obtain optimal constants and discuss the impact of local smoothness properties of f . Compared to the general confidence sets of Dümbgen (1998), the methods developed here are more stringent and computationally simpler. They are based on multiscale tests as developed by Dümbgen and Spokoiny (2001), who considered tests of qualitative assumptions rather than confidence bands. For further results on testing in nonparametric curve estimation, see Hart (1997), Fan *et al.* (2001), and the references cited therein.

2. Basic setting and overview

For mathematical convenience we focus on a continuous white noise model. Suppose that one observes a stochastic process Y on the unit interval $[0, 1]$, where

$$Y(t) = n^{1/2} \int_0^t f(x) dx + W(t).$$

Here f is an unknown function in $L^2[0, 1]$, $n \geq 1$ is a given scale parameter (‘sample size’) and W is standard Brownian motion. In this context the bounding functions $\hat{\ell}, \hat{u}$ are defined on $[0, 1]$, but for notational convenience the function f is tacitly assumed to be defined on the whole real line with values in $[-\infty, \infty]$. From now on we assume that

$$f \in \mathcal{G} \cap L^2[0, 1],$$

where \mathcal{G} denotes one of the following two function classes:

$$\mathcal{G}_\uparrow := \{\text{non-decreasing functions } g : \mathbb{R} \rightarrow [-\infty, \infty]\},$$

$$\mathcal{G}_{\text{conv}} := \{\text{convex functions } g : \mathbb{R} \rightarrow]-\infty, \infty]\}.$$

The paper is organized as follows. In Section 3 we treat the case $\mathcal{G} = \mathcal{G}_\uparrow$ and measure the quality of a confidence band $(\hat{\ell}, \hat{u})$ by quantities related to the Lévy distance $d_L(\hat{\ell}, \hat{u})$. Generally,

$$d_L(g, h) := \inf\{\epsilon > 0 : g \leq h(\cdot + \epsilon) + \epsilon \text{ and } h \leq g(\cdot + \epsilon) + \epsilon \text{ on } [0, 1 - \epsilon]\}$$

for isotonic functions $g, h : [0, 1] \rightarrow [-\infty, \infty]$. It turns out that a confidence band which is based on a suitable multiscale test as introduced by Dümbgen and Spokoiny (2001) is asymptotically optimal in a strong sense. Throughout this paper asymptotic statements refer to $n \rightarrow \infty$, unless stated otherwise.

In Section 4 we treat both classes \mathcal{G}_\uparrow and $\mathcal{G}_{\text{conv}}$ simultaneously. We discuss the construction of confidence bands $(\hat{\ell}, \hat{u})$ satisfying (1.2) such that $D(\hat{\ell}, f)$ and $D(f, \hat{u})$ are as small as possible whenever f satisfies some additional smoothness constraints. Here $D(g, h)$ is a distance measure of the form

$$D(g, h) := \sup_{x \in [0, 1]} w(x, f)(h(x) - g(x))$$

for some weight function $w(\cdot, f) \geq 0$ reflecting local smoothness properties of f . Again it turns out that suitable multiscale procedures yield nearly optimal procedures without additional prior information on f .

In Section 5 we present some numerical examples of the procedures in Section 4. The proofs are deferred to Sections 6, 7 and 8. In particular, Section 7 contains a new minimax bound for confidence rectangles in a Gaussian shift model, which may be of independent interest.

As for the white noise model, the results of Brown and Low (1996), Nussbaum (1996) and Grama and Nussbaum (1998) on asymptotic equivalence can be used to transfer the lower bounds of the present paper to other models. Moreover, one can mimic the confidence bands developed here in traditional regression models under minimal assumptions; see Dümbgen and Johns (2003) and Dümbgen (2001).

3. Optimality for isotonic functions in terms of Lévy-type distances

In this section we consider the class \mathcal{G}_\uparrow . For isotonic functions $g, h : [0, 1] \rightarrow [-\infty, \infty]$ and $\epsilon > 0$, let

$$D_\epsilon(g, h) := \inf\{\lambda \geq 0 : g \leq h(\cdot + \epsilon) + \lambda \text{ and } h \leq g(\cdot + \epsilon) + \lambda \text{ on } [0, 1 - \epsilon]\}.$$

Then the Lévy distance $d_L(g, h)$ is the infimum of all $\epsilon > 0$ such that $D_\epsilon(g, h) \leq \epsilon$. We use these functionals $D_\epsilon(\cdot, \cdot)$ in order to quantify differences between isotonic functions. Figure 1 depicts one such function g , and the shaded areas represent the set of all functions h with $D_{0.05}(g, h) \leq 0.1$ (Figure 1(a)) and $D_{0.05}(g, h) \leq 0.025$ (Figure 1(b)).




Figure 1. Two $D_{0.05}(\cdot, \cdot)$ -neighbourhoods of some function g .

The following theorem provides lower bounds for $D_\epsilon(\hat{\ell}, \hat{u})$, $0 < \epsilon \leq 1$. Throughout this paper the dependence of probabilities, expectations and distributions on the functional parameter f is sometimes indicated by a subscript f .

Theorem 3.1. *There exists a universal function b on $]0, 1]$ with $\lim_{\epsilon \downarrow 0} b(\epsilon) = 0$ such that*

$$\sup_{f \in \mathcal{G}_f \cap L^2[0,1]} P_f \left\{ \hat{\ell} \leq f \leq \hat{u} \text{ and } D_\epsilon(\hat{\ell}, \hat{u}) < \frac{(8 \log(e/\epsilon))^{1/2} - b(\epsilon)}{(n\epsilon)^{1/2}} \right\} \leq b(\epsilon)$$

for any confidence band $(\hat{\ell}, \hat{u})$ and arbitrary $\epsilon \in]0, 1]$.

Theorem 3.1 entails a lower bound for $d_L(\hat{\ell}, \hat{u})$. For let $\epsilon = \epsilon_n := c(\log(n)/n)^{1/3} - \delta n^{-1/3}$ with any fixed $c, \delta > 0$. Then one can show that, for sufficiently large n ,

$$\frac{(8 \log(e/\epsilon))^{1/2} - b(\epsilon)}{(n\epsilon)^{1/2}} = \left(\frac{8}{3c}\right)^{1/2} \left(\frac{\log n}{n}\right)^{1/3} + o(n^{-1/3}) \geq \epsilon,$$

provided that c equals $(8/3)^{1/3} \approx 1.387$.

Corollary 3.2. *For each $n \geq 1$, there exists a universal constant β_n such that $\beta_n \rightarrow 0$ and*

$$\sup_{f \in \mathcal{G}_f \cap L^2[0,1]} P_f \left\{ \hat{\ell} \leq f \leq \hat{u} \text{ and } d_L(\hat{\ell}, \hat{u}) < \left(\frac{8}{3}\right)^{1/3} \left(\frac{\log n}{n}\right)^{1/3} - \beta_n n^{-1/3} \right\} \leq \beta_n$$

for any confidence band $(\hat{\ell}, \hat{u})$. □

It is possible to get close to these lower bounds for $D_\epsilon(\hat{\ell}, \hat{u})$ *simultaneously* for all $\epsilon \in]0, 1]$ as long as (1.2) is satisfied. For let κ_α be a real number such that

$$P\left\{\left|\frac{W(t) - W(s)}{(t - s)^{1/2}}\right| \leq \Gamma(t - s) + \kappa_\alpha \text{ for } 0 \leq s < t \leq 1\right\} \leq \alpha,$$

where

$$\Gamma(u) := (2 \log(e/u))^{1/2} \quad \text{for } 0 < u \leq 1.$$

The existence of such a critical value κ_α follows from Dümbgen and Spokoiny (2001, Theorem 2.1). With the local averages

$$F_f(s, t) := \frac{1}{t - s} \int_s^t f(x) dx$$

of f and their natural estimators

$$\hat{F}(s, t) := \frac{Y(t) - Y(s)}{n^{1/2}(t - s)},$$

it follows that

$$P_f\left\{\left|\hat{F}(s, t) - F_f(s, t)\right| \leq \frac{\Gamma(t - s) + \kappa_\alpha}{(n(t - s))^{1/2}} \text{ for } 0 \leq s < t \leq 1\right\} \geq 1 - \alpha.$$

But for $0 \leq s < t \leq 1$,

$$f(s) \leq F_f(s, t) \leq f(t) \quad \text{whenever } f \in \mathcal{G}_\uparrow.$$

This implies the first assertion of the following theorem.

Theorem 3.3. *With the critical value κ_α above, let*

$$\begin{aligned} \hat{\ell}(x) &:= \sup_{0 \leq s < t \leq x} \left(\hat{F}(s, t) - \frac{\Gamma(t - s) + \kappa_\alpha}{\sqrt{n(t - s)}} \right), \\ \hat{u}(x) &:= \inf_{x \leq s < t \leq 1} \left(\hat{F}(s, t) + \frac{\Gamma(t - s) + \kappa_\alpha}{\sqrt{n(t - s)}} \right). \end{aligned}$$

This defines a confidence band $(\hat{\ell}, \hat{u})$ for f satisfying (1.2) with $\mathcal{F} = \mathcal{G}_\uparrow \cap L^2[0, 1]$. Moreover, provided that $\hat{\ell} \leq \hat{u}$,

$$D_\epsilon(\hat{\ell}, \hat{u}) \leq \frac{(8 \log(e/\epsilon))^{1/2} + 2\kappa_\alpha}{(n\epsilon)^{1/2}} \quad \text{for } 0 < \epsilon \leq 1,$$

$$d_L(\hat{\ell}, \hat{u}) \leq \left(\frac{8}{3}\right)^{1/3} \left(\frac{\log n}{n}\right)^{1/3} + o(n^{-1/3}).$$

Proof. The preceding upper bound for $D_\epsilon(\hat{\ell}, \hat{u})$ follows from the fact that, for any $x \in [0, 1 - \epsilon]$,

$$\begin{aligned} \hat{u}(x) - \hat{\ell}(x + \epsilon) &\leq \left(\hat{F}(x, x + \epsilon) + \frac{\Gamma(\epsilon) + \kappa_\alpha}{(n\epsilon)^{1/2}} \right) - \left(\hat{F}(x, x + \epsilon) - \frac{\Gamma(\epsilon) + \kappa_\alpha}{(n\epsilon)^{1/2}} \right) \\ &= \frac{2\Gamma(\epsilon) + 2\kappa_\alpha}{(n\epsilon)^{1/2}} \\ &= \frac{(8 \log(e/\epsilon))^{1/2} + 2\kappa_\alpha}{(n\epsilon)^{1/2}}. \end{aligned}$$

Letting $\epsilon = \epsilon_n = (8/3)^{1/3}(\log(n)/n)^{1/3}$ yields the upper bound for $d_L(\hat{\ell}, \hat{u})$. \square

4. Bands for potentially smooth functions

A possible criticism of the preceding results is the fact that the minimax bounds are attained at special step functions. On the other hand, one often expects the underlying curve f to be smooth in some vague sense. Therefore, we now aim for confidence bands satisfying (1.2) with $\mathcal{F} = \mathcal{G} \cap L^2[0, 1]$, which are as small as possible whenever f satisfies some additional smoothness conditions. As before, \mathcal{G} stands for \mathcal{G}_\uparrow or $\mathcal{G}_{\text{conv}}$.

In what follows let $\langle g, h \rangle := \int_{-\infty}^{\infty} g(x)h(x)dx$ and $\|g\| := \langle g, g \rangle^{1/2}$ for measurable functions g, h on the real line such that these integrals are defined. The confidence bands presented here can be described either in terms of kernel estimators for f or in terms of tests. Both viewpoints have their own merits.

4.1. Kernel estimators for f

Let ψ be some kernel function in $L^2(\mathbb{R})$. For technical reasons we make the following assumption on ψ .

Assumption 4.1. *The kernel function ψ satisfies the following three regularity conditions: ψ has bounded total variation; ψ is supported by $[-a, b]$, where $a, b \geq 0$; and $\langle 1, \psi \rangle > 0$.*

For any bandwidth $h > 0$ and location parameter $t \in \mathbb{R}$, let

$$\psi_{h,t}(x) := \psi\left(\frac{x-t}{h}\right).$$

Then $\langle g, \psi_{h,t} \rangle = h\langle g(t+h\cdot), \psi \rangle$ and $\|\psi_{h,t}\| = h^{1/2}\|\psi\|$. A kernel estimator for $f(t)$ with kernel function ψ and bandwidth h is given by

$$\hat{f}_h(t) := \frac{\psi Y(h, t)}{n^{1/2}h\langle 1, \psi \rangle},$$

where

$$\psi Y(h, t) := \int_0^1 \psi_{h,t}(x) dY(x).$$

From now on suppose that $ah \leq t \leq 1 - bh$. Then $\psi_{h,t}$ is supported by $[0, 1]$ and one may write

$$\begin{aligned} E\hat{f}_h(t) &= \frac{\langle f, \psi_{t,h} \rangle}{h\langle 1, \psi \rangle} = \frac{\langle f(t+h\cdot), \psi \rangle}{\langle 1, \psi \rangle}, \\ \text{var}(\hat{f}_h(t)) &= \frac{\langle \|\psi_{t,h}\|^2 \rangle}{nh^2\langle 1, \psi \rangle^2} = \frac{\|\psi\|^2}{nh\langle 1, \psi \rangle^2}. \end{aligned}$$

The random fluctuations of these kernel estimators can be bounded uniformly in $h > 0$. For that purpose we define the multiscale statistic

$$\begin{aligned} T(\pm\psi) &:= \sup_{h>0} \sup_{t \in [ah, 1-bh]} \left(\frac{\pm\psi W(h, t)}{h^{1/2}\|\psi\|} - \Gamma((a+b)h) \right) \\ &= \sup_{h>0} \sup_{t \in [ah, 1-bh]} \left(\pm \frac{\hat{f}_h(t) - E\hat{f}_h(t)}{\text{var}(\hat{f}_h(t))^{1/2}} - \Gamma((a+b)h) \right), \end{aligned}$$

similarly as in Dümbgen and Spokoiny (2001). It follows from Theorem 2.1 in the latter paper that $0 \leq T(\pm\psi) < \infty$ almost surely. In particular, $|\hat{f}_h(t) - E\hat{f}_h(t)| \leq (nh)^{-1/2} \log(e/h)^{1/2} O_p(1)$, uniformly in $h > 0$ and $ah \leq t \leq 1 - bh$.

It is well known that kernel estimators are biased in general. But our shape restrictions may be used to construct two kernel estimators whose bias is always non-positive or non-negative, respectively. Specifically, let $\psi^{(\ell)}$ and $\psi^{(u)}$ be two kernel functions satisfying Assumption 4.1 with respective supports $[-a^{(\ell)}, b^{(\ell)}]$ and $[-a^{(u)}, b^{(u)}]$. In addition, suppose that

$$\langle g, \psi^{(\ell)} \rangle \leq g(0)\langle 1, \psi^{(\ell)} \rangle \quad \text{for all } g \in \mathcal{G} \cap L^2[-a^{(\ell)}, b^{(\ell)}], \tag{4.1}$$

$$\langle g, \psi^{(u)} \rangle \geq g(0)\langle 1, \psi^{(u)} \rangle \quad \text{for all } g \in \mathcal{G} \cap L^2[-a^{(u)}, b^{(u)}]. \tag{4.2}$$

These inequalities imply that the corresponding kernel estimators satisfy the inequalities $E\hat{f}_h^{(\ell)}(t) \leq f(t) \leq E\hat{f}_h^{(u)}(t)$, and the definition of $T(\pm\psi)$ yields that

$$f(t) \geq \hat{f}_h^{(\ell)}(t) - \frac{\|\psi^{(\ell)}\|(\Gamma(d^{(\ell)}h) + T(\psi^{(\ell)}))}{\langle 1, \psi^{(\ell)} \rangle (nh)^{1/2}}, \tag{4.3}$$

$$f(t) \leq \hat{f}_h^{(u)}(t) + \frac{\|\psi^{(u)}\|(\Gamma(d^{(u)}h) + T(-\psi^{(u)}))}{\langle 1, \psi^{(u)} \rangle (nh)^{1/2}}. \tag{4.4}$$

Here $d^{(z)} := a^{(z)} + b^{(z)}$. Now let κ_α be the $1 - \alpha$ quantile of the combined statistic

$T^* := \max(T(\psi^{(\ell)}), T(-\psi^{(u)}))$, that is, the smallest real number such that $P\{T^* \leq \kappa_\alpha\} \geq 1 - \alpha$. Then

$$\hat{\ell}(t) := \sup_{h>0: t \in [a^{(\ell)}h, 1-b^{(\ell)}h]} \left(\hat{f}_h^{(\ell)}(t) - \frac{\|\psi^{(\ell)}\|(\Gamma(d^{(\ell)}h) + \kappa_\alpha)}{\langle 1, \psi^{(\ell)} \rangle (nh)^{1/2}} \right),$$

$$\hat{u}(t) := \inf_{h>0: t \in [a^{(u)}h, 1-b^{(u)}h]} \left(\hat{f}_h^{(u)}(t) + \frac{\|\psi^{(u)}\|(\Gamma(d^{(u)}h) + \kappa_\alpha)}{\langle 1, \psi^{(u)} \rangle (nh)^{1/2}} \right)$$

defines a confidence band $(\hat{\ell}, \hat{u})$ for f satisfying (1.2).

Equality holds in (1.2) if $\mathcal{G} = \mathcal{G}_\uparrow$ and f is constant, or if $\mathcal{G} = \mathcal{G}_{\text{conv}}$ and f is linear, provided that $\kappa_\alpha > 0$. For then it follows from (4.1) and (4.2) with $g(x) = \pm 1$ or $g(x) = \pm x$ that the kernel estimators are unbiased. Thus $\hat{\ell} \leq f \leq \hat{u}$ is equivalent to $T^* > \kappa_\alpha$. Moreover, using general theory for Gaussian measures on Banach spaces, one can show that the distribution of T^* is continuous on $]0, \infty[$.

Sufficient conditions for requirements (4.1) and (4.2) in general are provided by Lemma 8.1 below. The confidence band presented in Section 3 is a special case of the one derived here, if we define $\psi^{(\ell)}(x) := 1_{\{x \in [-1, 0]\}}$ and $\psi^{(u)}(x) := 1_{\{x \in [0, 1]\}}$ and apply postprocessing as described below.

4.2. Postprocessing of confidence bands

Any confidence band $(\hat{\ell}, \hat{u})$ for f can be enhanced if we replace $\hat{\ell}(x)$ and $\hat{u}(x)$ with

$$\hat{\hat{\ell}}(x) := \inf \{g(x) : g \in \mathcal{G}, \hat{\ell} \leq g \leq \hat{u}\} \quad \text{and} \quad \hat{\hat{u}}(x) := \sup \{g(x) : g \in \mathcal{G}, \hat{\ell} \leq g \leq \hat{u}\},$$

respectively. Here we tacitly assume that the set $\{g \in \mathcal{G} : \hat{\ell} \leq g \leq \hat{u}\}$ is non-empty.

If $\mathcal{G} = \mathcal{G}_\uparrow$ one can easily show that

$$\hat{\hat{\ell}}(x) = \sup_{t \in [0, x]} \hat{\ell}(t) \quad \text{and} \quad \hat{\hat{u}}(x) = \inf_{s \in [x, 1]} \hat{u}(s).$$

Note also that $\hat{\hat{\ell}}$ and $\hat{\hat{u}}$ are isotonic, whereas the raw functions $\hat{\ell}$ and \hat{u} need not be.

If $\mathcal{G} = \mathcal{G}_{\text{conv}}$ the modified upper bound $\hat{\hat{u}}$ is the greatest convex minorant of \hat{u} and can be computed (in discrete models) by means of the pool-adjacent-violators algorithm (cf. Robertson *et al.* 1988). The modified lower bound $\hat{\hat{\ell}}$ can be shown to be

$$\hat{\hat{\ell}}(x) = \max \left\{ \sup_{0 \leq s < t \leq x} \left(\hat{u}(s) + \frac{\hat{\ell}(t) - \hat{u}(s)}{t - s} (x - s) \right), \sup_{x \leq s < t \leq 1} \left(\hat{u}(t) - \frac{\hat{u}(t) - \hat{\ell}(s)}{t - s} (t - x) \right) \right\}.$$

This improved bound $\hat{\hat{\ell}}$ is not a convex function, though more regular than the raw function $\hat{\ell}$. Figure 2 depicts some hypothetical confidence band $(\hat{\ell}, \hat{u})$ for a function $f \in \mathcal{G}_{\text{conv}}$ and its improvement $(\hat{\hat{\ell}}, \hat{\hat{u}})$.




Figure 2. Improvement $(\hat{\ell}, \hat{u})$ of a band (ℓ, \hat{u}) if $\mathcal{G} = \mathcal{G}_{\text{conv}}$.

4.3. Adaptivity in terms of rates

Whenever we construct a band following the recipe above we end up with a confidence band adapting to the unknown smoothness of f in terms of rates of convergence. For $\beta, L > 0$, the Hölder smoothness class $\mathcal{H}_{\beta,L}$ is defined as follows. If $0 < \beta \leq 1$, let

$$\mathcal{H}_{\beta,L} := \{g : |g(x) - g(y)| \leq L|x - y|^\beta \text{ for all } x, y\}.$$

If $1 < \beta \leq 2$, let

$$\mathcal{H}_{\beta,L} := \{g \in \mathcal{C}^1 : g' \in \mathcal{H}_{\beta-1,L}\}.$$

Theorem 4.1. *Suppose that $f \in \mathcal{G} \cap \mathcal{H}_{\beta,L}$, where either $\mathcal{G} = \mathcal{G}_\uparrow$ and $\beta \leq 1$, or $\mathcal{G} = \mathcal{G}_{\text{conv}}$ and $1 \leq \beta \leq 2$. Let $(\hat{\ell}, \hat{u})$ be the confidence band for f based on test functions $\psi^{(\ell)}, \psi^{(u)}$ as described previously. Then there exists a constant Δ , depending only on (β, L) and $(\psi^{(\ell)}, \psi^{(u)})$, such that*

$$\sup_{t \in [\epsilon_n, 1 - \epsilon_n]} (\hat{u}(t) - \hat{\ell}(t)) \leq \Delta \rho_n \left(1 + \frac{\kappa_\alpha + T(\psi^{(u)}) + T(-\psi^{(\ell)})}{\log(en)^{1/2}} \right),$$

where $\epsilon_n := \rho_n^{1/\beta}$ and

$$\rho_n := \left(\frac{\log(en)}{n} \right)^{\beta/(2\beta+1)}.$$

Using the same arguments as Khas'minskii (1978), one can show that, for any $0 \leq r < s \leq 1$,

$$\inf_{f \in \mathcal{G} \cap \mathcal{H}_{\beta,L}} P_f \left\{ \sup_{t \in [r,s]} (\hat{u}(t) - \hat{\ell}(t)) \leq \Delta \rho_n \right\} \rightarrow 0,$$

provided that $\Delta > 0$ is sufficiently small. Thus our confidence bands adapt to the unknown smoothness of f .

4.4. Testing hypotheses about $f(t)$

In order to find suitable kernel functions $\psi^{(\ell)}$, $\psi^{(u)}$ we proceed similarly to Dümbgen and Spokoiny (2001, Section 3.2). That is to say, for the time being we consider tests of the null hypothesis

$$\mathcal{F}_0 := \{f \in \mathcal{G} \cap L^2[0, 1] : f(t) \leq r - \delta\}$$

versus the alternative hypothesis

$$\mathcal{F}_A := \{f \in \mathcal{G} \cap \mathcal{H}_{k,L} : f(t) \geq r\}.$$

Here $t \in [0, 1]$, $r \in \mathbb{R}$ and $L, \delta > 0$ are arbitrary fixed numbers, while

$$(\mathcal{G}, k) = (\mathcal{G}_\uparrow, 1) \quad \text{or} \quad (\mathcal{G}, k) = (\mathcal{G}_{\text{conv}}, 2). \tag{4.5}$$

Note that \mathcal{F}_0 and \mathcal{F}_A are closed, convex subsets of $L^2[0, 1]$. Suppose that there are functions $f_0 \in \mathcal{F}_0$ and $f_A \in \mathcal{F}_A$ such that

$$\int_0^1 (f_0 - f_A)(x)^2 dx = \min_{g_0 \in \mathcal{F}_0, g_A \in \mathcal{F}_A} \int_0^1 (g_0 - g_A)(x)^2 dx.$$

Then optimal tests of \mathcal{F}_0 versus \mathcal{F}_A are based on the linear test statistic $\int_0^1 (f_A - f_0)dY$, where critical values have to be computed under the assumption $f = f_0$. The problem of finding such functions f_0, f_A is treated in Section 8. Here we state our conclusion. Let

$$\psi^{(\ell)}(x) := \begin{cases} 1_{\{x \in [-1,0]\}}(1+x) & \text{if } \mathcal{G} = \mathcal{G}_\uparrow, \\ 1_{\{x \in [-2,2]\}}(1 - 3|x|/2 + x^2/2) & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}}. \end{cases} \tag{4.6}$$

Then the functions

$$f_A(s) := \begin{cases} r + L(s - t) & \text{if } \mathcal{G} = \mathcal{G}_\uparrow \\ r + L(s - t)^2/2 & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}} \end{cases} \tag{4.7}$$

and

$$f_0 := f_A - \delta \psi_{h,t}^{(\ell)}, \quad \text{with } h := (\delta/L)^{1/k},$$

solve our minimization problem, provided that $a^{(\ell)}h \leq t \leq 1 - b^{(\ell)}h$. Thus the optimal linear test statistic may be written as $\int_0^1 \psi_{h,t} dY = \psi Y(h, t)$. Elementary considerations show that the inequality

$$\tilde{f}_h^{(\ell)}(t) - \frac{\|\psi^{(\ell)}\|(\Gamma(d^{(\ell)}h) + \kappa_\alpha)}{\langle 1, \psi^{(\ell)} \rangle (nh)^{1/2}} \leq r_0$$

is equivalent to

$$\begin{aligned} \psi Y(h, t) &\leq n^{1/2} h r_0 \langle 1, \psi^{(\ell)} \rangle + h^{1/2} \|\psi^{(\ell)}\| (\Gamma(d^{(\ell)} h) + \kappa_\alpha) \\ &= E_{f_0}(\psi Y(h, t)) + \text{var}(\psi Y(h, t))^{1/2} (\Gamma(d^{(\ell)} h) + \kappa_\alpha). \end{aligned}$$

Thus our lower confidence bound $\hat{\ell}$ may be interpreted as a multiple test of all null hypotheses $\{f \in \mathcal{G} : f(t) \leq r_0\}$ with $t \in [0, 1]$ and $r_0 \in \mathbb{R}$.

Analogous considerations yield a candidate for $\psi^{(u)}$. Let

$$\mathcal{F}_0 := \{f \in \mathcal{G} \cap L^2[0, 1] : f(t) \geq r + \delta\}$$

and

$$\mathcal{F}_A := \{f \in \mathcal{G} \cap \mathcal{H}_{k,L} : f(t) \leq r\}.$$

Then the function f_A in (4.8) and

$$f_0 := f_A + \delta \psi_{h,t}^{(u)} \quad \text{with } h := (\delta/L)^{1/k}$$

form a least favourable pair (f_0, f_A) in $\mathcal{F}_0 \times \mathcal{F}_A$, where

$$\psi^{(u)}(x) := \begin{cases} 1_{\{x \in [0,1]\}}(1-x) & \text{if } \mathcal{G} = \mathcal{G}_\uparrow, \\ 1_{\{x \in [-2^{1/2}, 2^{1/2}]\}}(1-x^2/2) & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}}. \end{cases} \quad (4.8)$$

Figures 3 and 4 depict the functions $\psi^{(\ell)}$ in (4.6) and $\psi^{(u)}$ in (4.8).

4.5. Optimal constants and local adaptivity

We will now show that our multiscale confidence band $(\hat{\ell}, \hat{u})$, if constructed with the kernel functions in (4.6) and (4.8), is locally adaptive in a certain sense. Specifically, we consider




Figure 3. Kernel functions (a) $\psi^{(\ell)}$, (b) $\psi^{(u)}$ for \mathcal{G}_\uparrow .




Figure 4. Kernel functions (a) $\psi^{(\ell)}$, (b) $\psi^{(u)}$ for $\mathcal{G}_{\text{conv}}$.

an arbitrary fixed function $f_0 \in \mathcal{G} \cap \mathcal{C}^k$ with (\mathcal{G}, k) as specified in (4.5). We analyse quantities such as

$$\|(\hat{u} - f_0)w\|_{r,s}^+ \quad \text{and} \quad \|(f_0 - \hat{\ell})w\|_{r,s}^+$$

where w is some positive weight function on the unit interval and

$$\|g\|_{r,s}^+ := \sup_{t \in [r,s]} g(t).$$

The function w should reflect local smoothness properties of f_0 in an appropriate way. The following theorem demonstrates that the k th derivative of f_0 , denoted by $\nabla^k f_0$, plays a crucial role.

Theorem 4.2. For arbitrary fixed numbers $0 \leq r < s \leq 1$, let

$$L := \max_{t \in [r,s]} \nabla^k f_0(t).$$

Then, for any $\gamma \in]0, 1[$,

$$\inf_{(\hat{\ell}, \hat{u})} P_{f_0} \{ \|\hat{f} - \hat{\ell}\|_{r,s}^+ \geq \gamma \Delta^{(\ell)} L^{1/(2k+1)} \rho_n \} \geq 1 - \alpha + o(1),$$

$$\inf_{(\hat{\ell}, \hat{u})} P_{f_0} \{ \|\hat{u} - f\|_{r,s}^+ \geq \gamma \Delta^{(u)} L^{1/(2k+1)} \rho_n \} \geq 1 - \alpha + o(1),$$

where both infima are taken over all confidence bands $(\hat{\ell}, \hat{u})$ satisfying (1.2), and

$$\Delta^{(z)} := \left(\left(k + \frac{1}{2} \right) \|\psi^{(z)}\|^2 \right)^{-k/(2k+1)},$$

$$\rho_n := \left(\frac{\log(en)}{n} \right)^{k/(2k+1)}.$$

If $\mathcal{G} = \mathcal{G}_\uparrow$, the critical constants are $\Delta^{(\ell)} = \Delta^{(u)} = 2^{1/3} \approx 1.260$. If $\mathcal{G} = \mathcal{G}_{\text{conv}}$,

$$\Delta^{(\ell)} = (3/4)^{2/5} \approx 0.891 \quad \text{and} \quad \Delta^{(u)} = 3^{2/5}/128^{1/5} \approx 0.588.$$

This indicates that bounding a convex function from below is more difficult than finding an upper bound.

In view of Theorem 4.2, we introduce, for arbitrary fixed $\epsilon > 0$, the weight function

$$w_\epsilon := (\max(\nabla^k f_0, \epsilon))^{-1/(2k+1)}$$

reflecting the local smoothness of f_0 . The next theorem shows that our particular confidence band $(\hat{\ell}, \hat{u})$ attains the lower bounds of Theorem 4.2 pointwise. Suprema such as $\|(f_0 - \hat{\ell})w_\epsilon\|_{r,s}^+$ and $\|(\hat{u} - f_0)w_\epsilon\|_{r,s}^+$ attain their respective lower bounds $\Delta^{(\ell)}, \Delta^{(u)}$ up to a multiplicative factor $2^{k/(k+1/2)} + o_p(1)$.

Theorem 4.3. *Let $(\hat{\ell}, \hat{u})$ be the confidence band based on the kernel functions in (4.6) and (4.8). If $f = f_0$, then for arbitrary $\epsilon > 0$ and any $t \in]0, 1[$,*

$$(f_0 - \hat{\ell})(t)w_\epsilon(t) \leq (\Delta^{(\ell)} + o_p(1))\rho_n,$$

$$(\hat{u} - f_0)(t)w_\epsilon(t) \leq (\Delta^{(u)} + o_p(1))\rho_n.$$

Moreover,

$$\|(f_0 - \hat{\ell})w_\epsilon\|_{\epsilon,1-\epsilon}^+ \leq (2^{k/(k+1/2)}\Delta^{(\ell)} + o_p(1))\rho_n,$$

$$\|(\hat{u} - f_0)w_\epsilon\|_{\epsilon,1-\epsilon}^+ \leq (2^{k/(k+1/2)}\Delta^{(u)} + o_p(1))\rho_n.$$

If we used kernel functions differing from (4.7) and (4.9), then pointwise optimality would be lost, and the constants for the supremum distances would get worse.

5. Simulations and numerical examples

In this section we demonstrate the performance of the procedures in Section 4. We replace the continuous white noise model with a discrete one. Suppose that one observes a random vector $\bar{Y} \in \mathbb{R}^n$ with components

$$Y_i = f(x_i) + \epsilon_i, \tag{5.1}$$

where $x_i := (i - 1/2)/n$, and the random errors ϵ_i are independent with Gaussian distribution $\mathcal{N}(0, \sigma^2)$. Our kernel functions $\psi^{(\ell)}$ and $\psi^{(u)}$ are rescaled as follows:

$$\psi^{(\ell)}(x) := \begin{cases} 1_{\{x \in [-1,0]\}}(1+x) & \text{if } \mathcal{G} = \mathcal{G}_\uparrow, \\ 1_{\{x \in [-1,1]\}}(1-3|x|+2x^2) & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}}, \end{cases}$$

$$\psi^{(u)}(x) := \begin{cases} 1_{\{x \in [0,1]\}}(1-x) & \text{if } \mathcal{G} = \mathcal{G}_\uparrow, \\ 1_{\{x \in [-1,1]\}}(1-x^2) & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}}. \end{cases}$$

Note that now $a^{(\ell)}, a^{(u)}, b^{(\ell)}, b^{(u)} \in \{0, 1\}$. For convenience we compute kernel estimators and confidence bounds for f only on the grid $\mathcal{T}_n := \{1/n, 2/n, \dots, 1 - 1/n\}$, while the bandwidth parameter h is restricted to

$$H_n := \begin{cases} \{1/n, 2/n, \dots, 1\} & \text{if } \mathcal{G} = \mathcal{G}_\uparrow, \\ \{1/n, 2/n, \dots, \lfloor n/2 \rfloor / n\} & \text{if } \mathcal{G} = \mathcal{G}_{\text{conv}}. \end{cases}$$

Let ψ stand for $\psi^{(\ell)}$ or $\psi^{(u)}$ with support $[-a, b]$. Then for $h \in H_n$ and $t \in \mathcal{T}_n$ with $ah \leq t \leq 1 - bh$, we define

$$\psi \vec{Y}(h, t) := \sum_{i=1}^n \psi\left(\frac{x_i - t}{h}\right) Y_i = \sum_{j=1-anh}^{bnh} \psi\left(\frac{j-1/2}{nh}\right) Y_{nt+j}$$

and

$$\hat{f}_h(t) := \frac{\psi \vec{Y}(h, t)}{S_{nh}},$$

where S_d stands for $\sum_{j=1-d}^d \psi((j-1/2)/d)$. The standard deviation of $\hat{f}_h(t)$ equals $\sigma_h := \sigma R_{nh}^{1/2} / S_{nh}$, where $R_d := \sum_{j=1-d}^d \psi((j-1/2)/d)^2$. Tedious but elementary calculations show that if $\mathcal{G} = \mathcal{G}_\uparrow$,

$$S_d = \frac{d}{2} \quad \text{and} \quad R_d = \frac{d}{3} - \frac{1}{12d}.$$

If $\mathcal{G} = \mathcal{G}_{\text{conv}}$,

$$S_d^{(\ell)} = \frac{d}{3} - \frac{1}{3d} \quad \text{and} \quad R_d^{(\ell)} = \frac{4d}{15} - \frac{1}{2d} + \frac{7}{30d^3},$$

$$S_d^{(u)} = \frac{4d}{3} + \frac{1}{6d} \quad \text{and} \quad R_d^{(u)} = \frac{16d}{15} + \frac{7}{120d^3}.$$

Note that here $S_1^{(\ell)} = 0 = \psi^{(\ell)} \vec{Y}(1/n, \cdot)$, whence the bandwidth $1/n$ is excluded from any computation involving $\psi^{(\ell)}$.

As for the bias of these kernel estimators, one can deduce from Lemma 8.1 that $E \hat{f}_h^{(\ell)}(t) \leq f(t)$ and $E \hat{f}_h^{(u)}(t) \geq f(t)$ whenever $f \in \mathcal{G}$. Here is a discrete version of our multiscale test statistic: $T_n^* := \max(T_n(\psi^{(\ell)}), T_n(-\psi^{(u)}))$, where

$$T_n(\pm\psi) := \max_{h \in H_n} \max_{t \in \mathcal{T}_n \cap [ah, 1-bh]} (\pm \sigma^{-1} R_{nh}^{-1/2} \psi \vec{E}(h, t) - \Gamma((a+b)h))$$

with $\vec{E} := (\epsilon_i)_{i=1}^n$. Let $\kappa_{\alpha,n}$ be the $1 - \alpha$ quantile of T_n^* . Then

$$\hat{\ell}(t) := \max_{h \in H_n: t \in [a^{(\ell)}h, 1 - b^{(\ell)}h]} (\hat{f}_h^{(\ell)}(t) - \sigma_h^{(\ell)}(\Gamma(d^{(\ell)}h) + \kappa_{\alpha,n})),$$

$$\hat{u}(t) := \min_{h \in H_n: t \in [a^{(u)}h, 1 - b^{(u)}h]} (\hat{f}_h^{(u)}(t) + \sigma_h^{(u)}(\Gamma(d^{(u)}h) + \kappa_{\alpha,n})),$$

defines a confidence band for f such that

$$P\{\hat{\ell} \leq f \leq \hat{u} \text{ on } \mathcal{T}_n\} \geq 1 - \alpha \quad \text{whenever } f \in \mathcal{G}.$$

Equality holds if $\mathcal{G} = \mathcal{G}_\uparrow$ and f is constant, or if $\mathcal{G} = \mathcal{G}_{\text{conv}}$ and f is linear. If the noise variance σ^2 is unknown, it may be estimated as described in Dümbgen and Spokoiny (2001). Then, under moderate regularity assumptions on f , our confidence bands have *asymptotic* coverage probability at least $1 - \alpha$ as n tends to infinity.

For various values of n we estimated several quantiles $\kappa_{\alpha,n}$ in 9999 Monte Carlo simulations; see Table 1. One can easily show that the critical value $\kappa_{\alpha,n}$ converges to the corresponding quantile κ_α for the continuous white noise model as $n \rightarrow \infty$. Software for the computation of critical values as well as confidence bands may be obtained from <http://www.cx.unibe.ch/~duembgen/software.html>.

Figure 5 shows a simulated data vector \vec{Y} with $n = 500$ components together with the corresponding 95% confidence band $(\hat{\ell}, \hat{u})$ after postprocessing, where f is assumed to be isotonic. The latter function is depicted as well. Note that the band is comparatively narrow in the middle of $]0, \frac{1}{3}[$, on which f is constant. On $]1/3, 1]$ the width $\hat{u} - \hat{\ell}$ tends to increase, as does ∇f . These findings are in accordance with Theorem 4.3.

An analogous plot for a convex function f can be seen in Figure 6. Note that the deviation $f - \hat{\ell}$ is mostly greater than $\hat{u} - f$, as predicted by Theorem 4.3.

6. Proofs

Proof of Theorem 3.1. In order to prove lower bounds we construct unfavourable subfamilies of \mathcal{G}_\uparrow similarly to Khas'minskii (1978). For a given integer $m > 0$, we define $I_1 := [0, 1/m]$

Table 1. Some critical values for the discrete white noise model

n	\mathcal{G}_\uparrow			$\mathcal{G}_{\text{conv}}$		
	$\kappa_{0.5,n}$	$\kappa_{0.1,n}$	$\kappa_{0.05,n}$	$\kappa_{0.5,n}$	$\kappa_{0.1,n}$	$\kappa_{0.05,n}$
100	0.330	1.092	1.349	0.350	1.053	1.283
200	0.433	1.146	1.392	0.430	1.121	1.342
300	0.475	1.169	1.416	0.470	1.126	1.342
400	0.507	1.204	1.446	0.489	1.128	1.340
500	0.526	1.222	1.450	0.512	1.143	1.358
700	0.570	1.252	1.492	0.536	1.162	1.380
1000	0.585	1.250	1.483	0.552	1.178	1.393




Figure 5. Data \bar{Y} and 95% confidence band for $f \in \mathcal{G}_1$.

and $I_j := [(j - 1)/m, j/m]$ for $1 < j \leq m$. Then we define step functions g and h_ξ for $\xi \in \mathbb{R}^m$ via

$$g(t) := 2j - 1 \quad \text{and} \quad h_\xi(t) := \xi_j, \quad \text{for } t \in I_j, 1 \leq j \leq m.$$

For any $\delta > 0$ and $\xi \in [-\delta, \delta]^m$ the function $\delta g + h_\xi$ is isotonic on $[0, 1]$. Now we restrict our attention to the parametric submodel $\mathcal{F}_0 = \{\delta g + h_\xi : \xi \in [-\delta, \delta]^m\}$ of $\mathcal{G}_1 \cap L^2[0, 1]$. Any confidence band $(\hat{\ell}, \hat{u})$ for $f = \delta g + h_\xi$ defines a confidence set $S = S_1 \times S_2 \times \dots \times S_m$ for ξ via

$$S_j := \left[\sup_{t \in I_j} \hat{\ell}(t) - \delta(2j - 1), \inf_{t \in I_j} \hat{u}(t) - \delta(2j - 1) \right].$$

Here $\hat{\ell} \leq f \leq \hat{u}$ if and only if $\xi \in S$. Moreover,

$$D_\epsilon(\hat{\ell}, \hat{u}) \geq \max_{j=1, \dots, m} \text{length}(S_j) \quad \text{for } 1/(m + 1) \leq \epsilon < 1/m.$$

However,




Figure 6. Data \tilde{Y} and 95% confidence band for $f \in \mathcal{G}_{\text{conv}}$.

$$\begin{aligned} \log \frac{dP_{\delta g + h_\xi}}{dP_{\delta g}}(Y) &= n^{1/2} \int_0^1 h_\xi d\tilde{Y} - n \int_0^1 h_\xi(t)^2 dt / 2 \\ &= \sum_{j=1}^m ((n/m)^{1/2} \xi_j X_j - (n/m) \xi_j^2 / 2) \\ &= \log \frac{d\mathcal{N}((n/m)^{1/2} \xi, I)}{d\mathcal{N}(0, I)}(X), \end{aligned}$$

where $\tilde{Y}(t) := Y(t) - n^{1/2} \int_0^t \delta g(s) ds$ and $X := (X_j)_{j=1}^m$ with components

$$X_j := m^{1/2} (\tilde{Y}(j/m) - \tilde{Y}((j-1)/m)).$$

If $f = \delta g$ these random variables are independent and standard normal. Consequently, X is a sufficient statistic for the parametric submodel \mathcal{F}_0 with distribution $\mathcal{N}_m((n/m)^{1/2} \xi, I)$ if $f = \delta g + h_\xi$. In particular, the conditional distribution of S given X does not depend on ξ .

Hence, letting $\delta = (n/m)^{-1/2}c_m$ with $c_m := (2 \log m)^{1/2}$, it follows from Theorem 7.1(b) below that, for $1/(m+1) \leq \epsilon < 1/m$,

$$\begin{aligned} & \sup_{f \in \mathcal{G}_1 \cap L^2[0,1]} P_f \left\{ \hat{\ell} \leq f \leq \hat{u} \text{ and } D_\epsilon(\hat{\ell}, \hat{u}) \leq 2 \frac{c_m - b_m}{(n/m)^{1/2}} \right\} \\ & \leq \max_{\xi \in [-\delta, \delta]^m} P_\xi \left\{ \xi \in S \text{ and } \max_{j=1, \dots, m} \text{length}(S_j) \leq 2 \frac{c_m - b_m}{(n/m)^{1/2}} \right\} \leq b_m, \end{aligned}$$

where b_1, b_2, b_3, \dots are universal positive numbers such that $\lim_{m \rightarrow \infty} b_m = 0$. This entails the assertion of Theorem 3.1 with $\log(1/\epsilon)$ in place of $\log(e/\epsilon)$ and

$$b(\epsilon) := (2 \log(1/\epsilon))^{1/2} - (m\epsilon)^{1/2}(c_m - b_m) \quad \text{for } 1/(m+1) \leq \epsilon < 1/m.$$

Finally, note that $\log(e/\epsilon)^{1/2} = \log(1/\epsilon)^{1/2} + o(1)$ as $\epsilon \downarrow 0$. \square

Proof of Theorem 4.1. Instead of an upper bound for $\hat{u} - \hat{\ell}$ we prove an upper bound for $\hat{u} - f$, because analogous arguments apply to $f - \hat{\ell}$. In what follows let $\psi = \psi^{(u)}$ with support $[-a, b]$. For $t \in [0, 1]$ and $h > 0$ with $ah \leq t \leq 1 - bh$,

$$\begin{aligned} \hat{u}(t) - f(t) & \leq \hat{f}_h(t) - f(t) + \frac{\|\psi\|(\Gamma((a+b)h) + \kappa_\alpha)}{\langle 1, \psi \rangle (nh)^{1/2}} \\ & = \frac{\langle f(t+h\cdot) - f(t), \psi \rangle}{\langle 1, \psi \rangle} + \frac{\psi W(h, t)}{n^{1/2} h \langle 1, \psi \rangle} + \frac{\|\psi\|(\Gamma((a+b)h) + \kappa_\alpha)}{\langle 1, \psi \rangle (nh)^{1/2}} \\ & \leq \frac{\langle f(t+h\cdot) - f(t), \psi \rangle}{\langle 1, \psi \rangle} + \frac{\|\psi\|(2\Gamma((a+b)h) + \kappa_\alpha + T(\psi))}{\langle 1, \psi \rangle (nh)^{1/2}}. \end{aligned} \quad (6.1)$$

For any function $g \in \mathcal{H}_{\beta, L}$,

$$|g(x) - g(0)| \leq L|x|^\beta \quad \text{if } \beta \leq 1,$$

$$|g(x) - g(0) - g'(0)x| \leq L|x|^\beta \quad \text{if } 1 < \beta \leq 2.$$

Since $f(t+h\cdot) \in \mathcal{H}_{\beta, Lh^\beta}$ if $f \in \mathcal{H}_{\beta, L}$, this implies that

$$\frac{\langle f(t+h\cdot) - f(t), \psi \rangle}{\langle 1, \psi \rangle} \leq \frac{Lh^\beta \int_{-a}^b |x|^\beta |\psi(x)| dx}{\langle 1, \psi \rangle} \leq \Delta h^\beta.$$

Here and subsequently Δ denotes a generic constant depending only on (β, L) and ψ . Its value may vary from one line to another. If $t \in [\epsilon_n, 1 - \epsilon_n]$ and $h = \epsilon_n / \max(a, b)$, the right-hand side of (6.1) is not greater than

$$\Delta \epsilon_n^\beta + \frac{\Delta(\log(en)^{1/2} + \kappa_\alpha + T(\psi))}{(n\epsilon_n)^{1/2}} = \Delta \rho_n \left(1 + \frac{\kappa_\alpha + T(\psi)}{\log(en)^{1/2}} \right).$$

\square

Proof of Theorem 4.2. We prove only the lower bound for $f_0 - \hat{\ell}$, because $\hat{u} - f_0$ can be treated analogously. It suffices to consider the case $L > 0$ and to show that for any fixed number $\gamma \in]0, 1[$,

$$P_{f_0} \{ \|f_0 - \hat{\ell}\|_{r,s}^+ \geq \gamma \Delta^{(\ell)} L^{1/(2k+1)} \rho_n \} \geq 1 - \alpha + o(1)$$

for arbitrary confidence bands $(\hat{\ell}, \hat{u}) = (\hat{\ell}_n, \hat{u}_n)$ satisfying (1.2). Without loss of generality one may assume that

$$\nabla^k f_0 \geq L \quad \text{on } [r, s].$$

Otherwise one could increase γ and decrease L without changing $\gamma L^{1/(2k+1)}$, and replace $[r, s]$ with some non-degenerate subinterval. Let ψ stand for $\psi^{(\ell)}$ with support $[-a, b]$. For $0 < h \leq (s-r)/(a+b)$ and positive integers $j \leq m := \lfloor (s-r)/((a+b)h) \rfloor$, let

$$t_j := s + ah + (j-1)(a+b)h \quad \text{and} \quad f_j := f_0 - Lh^k \psi_{h,t_j}.$$

It follows from Lemma 8.4 that these functions f_j belong to $\mathcal{G} \cap L^2[0, 1]$. Thus (1.2) implies that the event

$$A := \{ \hat{\ell} \leq f_j \text{ for some } j \leq m \}$$

satisfies the inequality $P_{f_j}(A) \geq 1 - \alpha$ for all $j \leq m$. Since $\|f_0 - f_j\|_{r,s}^+ \geq \delta$, this entails the inequality

$$P_{f_0} \{ \|f_0 - \hat{\ell}\|_{r,s}^+ \geq Lh^k \} \geq P_{f_0}(A) \geq 1 - \alpha - \min_{j \leq m} (P_{f_j}(A) - P_{f_0}(A)).$$

Now let $h := (c\rho_n)^{1/k}$ so that $Lh^k = Lc\rho_n$, where $c > 0$ is some number to be specified later. For sufficiently large n , this bandwidth h is smaller than $(s-r)/(a+b)$. Then

$$\log \frac{dP_{f_j}}{dP_{f_0}}(Y) = n^{1/2} h^{k+1/2} L \|\psi\| X_j - nh^{2k+1} L^2 \|\psi\|^2 / 2,$$

where $X_j := h^{-1/2} \|\psi\|^{-1} \int_0^1 \psi_{h,t_j} d\tilde{Y}$ and $\tilde{Y}(t) := Y(t) - n^{1/2} \int_0^t f_0(x) dx$. Thus $X := (X_j)_{j=1}^m$ is a sufficient statistic for the restricted model $\{f_0, f_1, f_2, \dots, f_m\}$, where $\mathcal{L}_{f_0}(X)$ is a standard normal distribution on \mathbb{R}^m . Thus it follows from Theorem 7.1(a) and a standard sufficiency argument that

$$\lim_{n \rightarrow \infty} \min_{1 \leq j \leq m} (P_{f_j}(A) - P_{f_0}(A)) = 0 \quad \text{if} \quad \lim_{n \rightarrow \infty} \frac{nh^{2k+1} L^2 \|\psi\|^2}{2 \log m} < 1.$$

Since $\log m = (1 + o(1)) \log(n)/(2k+1)$, the limit on the right-hand side is equal to $c^{(2k+1)/k} L^2 \|\psi\|^2 (k+1/2)$ and smaller than 1 if $c = \gamma \Delta^{(\ell)} L^{-2k/(2k+1)}$. In that case, the lower bound $Lh^k = Lc\rho_n$ for $\|f_0 - \hat{\ell}\|_{r,s}^+$ equals $\gamma \Delta^{(\ell)} L^{1/(2k+1)} \rho_n$ as desired. \square

Proof of Theorem 4.3. Again we restrict our attention to $f_0 - \hat{\ell}$ and let $\psi := \psi^{(\ell)}$ with support $[-a, b]$. For any fixed $\epsilon > 0$ and arbitrary $t \in [0, 1]$, let $h_t > 0$ and

$$L_t := \max_{s \in [t-ah_t, t+bh_t] \cap [0, 1]} \max(\nabla^k f_0(s), \epsilon).$$

If $ah_t \leq t \leq 1 - bh_t$ the inequality $(f_0 - \hat{\ell})(t) \geq L_t h_t^k$ implies that

$$\hat{f}_{h_t}(t) - \frac{\|\psi\|(\Gamma((a+b)h_t) + \kappa_\alpha)}{(nh_t)^{1/2}\langle 1, \psi \rangle} \leq f_0(t) - L_t h_t^k.$$

Since $f = f_0$, this can be rewritten as

$$\begin{aligned} \frac{\psi W(h_t, t)}{h_t^{1/2}\|\psi\|} &\leq -\frac{(nh_t)^{1/2}}{\|\psi\|} \langle f_0(t + h_t) - f_0(t) + L_t h_t^k, \psi \rangle + \Gamma((a+b)h_t) + \kappa_\alpha \\ &\leq -n^{1/2} L_t h_t^{k+1/2} \|\psi\| + \Gamma((a+b)h_t) + \kappa_\alpha, \end{aligned}$$

where the latter inequality follows from Lemma 8.4(c). Specifically, let

$$h_t := cw_\epsilon(t)^2 \rho_n^{1/k}$$

for some positive constant c to be specified later. By continuity of $\nabla^k f_0$, the weight function w_ϵ is bounded away from zero and infinity. Hence $h_t \rightarrow 0$ and $L_t \max(\nabla^k f_0(t), \epsilon)^{-1} \rightarrow 1$, uniformly in $t \in [0, 1]$. In particular,

$$\begin{aligned} \Gamma((a+b)h_t) &\leq (k+1/2)^{-1/2} \log(en)^{1/2} \quad \text{for } n \geq n_0, \\ n^{1/2} L_t h_t^{k+1/2} \|\psi\| &\geq c^{k+1/2} \|\psi\| \log(en)^{1/2}, \\ L_t h_t^k &\leq w_\epsilon(t)^{-1} c^k (1+b_n) \rho_n, \end{aligned}$$

where n_0 and b_n are positive numbers depending only on f_0, ϵ and c such that $b_n \rightarrow 0$. Consequently, for $n \geq n_0$,

$$ah_t \leq t \leq 1 - bh_t \quad \text{and} \quad (f_0 - \hat{\ell})(t)w_\epsilon(t) \geq c^k(1+b_n)\rho_n$$

imply that

$$\frac{\psi W(h_t, t)}{h_t^{1/2}\|\psi\|} \leq -(c^{k+1/2}\|\psi\| - (k+1/2)^{-1/2}\log(en)^{1/2}) + \kappa_\alpha. \tag{6.2}$$

Whenever $c > (\Delta^{(\ell)})^{1/k}$, the right-hand side of inequality (6.2) tends to minus infinity, while the random variable on the left-hand side has mean zero and variance one. Since the limit of $c^k(1+b_n)$ can be arbitrarily close to $\Delta^{(\ell)}$, these considerations show that $(f_0 - \hat{\ell})(t)w_\epsilon(t) \leq (\Delta^{(\ell)} + o_p(1))\rho_n$ for any fixed $t \in]0, 1[$.

If n is sufficiently large, then $ah_t \leq t \leq 1 - bh_t$ and

$$\frac{\psi W(h_t, t)}{h_t^{1/2}\|\psi\|} \geq -T(-\psi) - \Gamma((a+b)h_t)$$

for all $t \in [\epsilon, 1 - \epsilon]$. Consequently,

$$\sup_{t \in [\epsilon, 1 - \epsilon]} (f_0 - \hat{\ell})(t)w_\epsilon(t) \geq c^k(1+b_n)$$

implies that

$$\begin{aligned}
T(-\psi) &\geq n^{1/2} L_t h_t^{k+1/2} \|\psi\| - 2\Gamma((a+b)h_t) - \kappa_\alpha \\
&\geq (c^{k+1/2} \|\psi\| - 2(k+1/2)^{-1/2}) \log(en)^{1/2} - \kappa_\alpha.
\end{aligned} \tag{6.3}$$

Whenever $c > 2^{1/(k+1/2)}(\Delta^{(\ell)})^{1/k}$, the right-hand side of inequality (6.3) tends to infinity. Since the limit of $c^k(1+b_n)$ can be arbitrarily close to $2^{k/(k+1/2)}\Delta^{(\ell)}$, these considerations reveal that $\|(f_0 - \hat{\ell})_{w_\epsilon}\|_{\epsilon, 1-\epsilon}^+$ is not greater than $(2^{k/(k+1/2)}\Delta^{(\ell)} + o_p(1))\rho_n$. \square

7. Some decision theory

Let $X = (X_i)_{i=1}^m$ be a random vector with distribution $\mathcal{N}_m(\theta, I)$. In what follows we consider tests $\phi : \mathbb{R}^m \rightarrow [0, 1]$ and confidence sets

$$S = S_1 \times S_2 \times \cdots \times S_m$$

for θ with random intervals $S_j \subset \mathbb{R}$. The conditional distribution of S , given X , does not depend on θ . The possibility of randomized confidence sets S , that is, confidence sets not just being a function of X , has to be included for technical reasons. Unless specified otherwise, asymptotic statements in this section refer to $m \rightarrow \infty$.

Theorem 7.1. *Let $c_m := (2 \log m)^{1/2}$. There are universal positive numbers b_m with $b_m \rightarrow 0$ such that the following two inequalities are satisfied:*

(a) *For arbitrary tests ϕ ,*

$$\min_{j=1, \dots, m} E(c_m - b_m) e_j \phi(X) - E_0 \phi(X) \leq b_m,$$

where e_1, e_2, \dots, e_m denotes the standard basis of \mathbb{R}^m .

(b) *For arbitrary confidence sets S as above,*

$$\min_{\theta \in [-c_m, c_m]^m} P_\theta \left\{ \theta \in S \text{ and } \max_{j=1, \dots, m} \text{length}(S_j) < 2(c_m - b_m) \right\} \leq b_m.$$

Proof of Theorem 7.1. Part (a) is classical and can be proved by a Bayesian argument; see, for instance, Ingster (1993a; 1993b; 1993c) or Dümbgen and Spokoiny (2001).

In order to prove part (b) we also consider a Bayesian model. Let θ have independent components each of which is uniformly distributed on the three-point set $K_m := \{-\kappa_m, 0, \kappa_m\}$, where $\kappa_m := c_m - b_m$ with constants $b_m \in [0, c_m]$ to be specified later on. Let $\mathcal{L}(X|\theta) = \mathcal{N}_m(\theta, I)$. Let $P(\cdot)$, $E(\cdot)$ denote probabilities and expectations in this Bayesian context, whereas $P_\theta(\cdot)$, $E_\theta(\cdot)$ are used in case of a fixed parameter θ . For any confidence set S ,

$$\begin{aligned} & \min_{\theta \in [-c_m, c_m]^m} P_\theta \left\{ \theta \in S \text{ and } \max_{j=1, \dots, m} \text{length}(S_j) < 2\kappa_m \right\} \\ & \leq P \left\{ \theta \in S \text{ and } \max_{j=1, \dots, m} \text{length}(S_j) < 2\kappa_m \right\} \leq P\{\theta \in \tilde{S}\}, \end{aligned}$$

where

$$\tilde{S} := \begin{cases} S & \text{if } \max_{j=1, \dots, m} \text{length}(S_j) < 2\kappa_m, \\ \{0\} \times \dots \times \{0\} & \text{otherwise.} \end{cases}$$

The conditional distribution of θ given (X, S) is also a product of m probability measures. For any $\eta \in K_m^m$

$$P(\theta = \eta | X, S) = \prod_{i=1}^m g(\eta_i | X_i) \quad \text{with } g(z|x) := \frac{\exp(-(x-z)^2/2)}{\sum_{y \in K_m} \exp(-(x-y)^2/2)}.$$

Since each factor \tilde{S}_j of \tilde{S} contains at most two points from K_m ,

$$\begin{aligned} P\{\theta \in \tilde{S}\} &= EP(\theta \in \tilde{S} | X, S) \\ &\leq E \max_{\eta \in K_m^m} P(\theta_i \neq \eta_i \text{ for } i = 1, \dots, m | X, S) \\ &= E \prod_{i=1}^m \left(1 - \min_{z \in K_m} g(z | X_i) \right) = \left(1 - E \min_{z \in K_m} g(z | X_1) \right)^m \\ &\leq \left(1 - 3^{-1} E \min_{z \in K_m} \exp(-(X_1 - z)^2/2) \right)^m. \end{aligned}$$

The latter expectation can be bounded from below as follows:

$$\begin{aligned} & 3^{-1} E \min_{z \in K_m} \exp(-(X_1 - z)^2/2) \\ & \geq 3^{-1} P\{|X_1| \leq b_m/2\} \exp(-(\kappa_m + b_m/2)^2/2) \\ & \geq 3^{-1} P\{|\theta_1| = 0, |X_1| \leq b_m/2\} \exp(-(c_m - b_m/2)^2/2) \\ & = 9^{-1} (2\pi)^{-1/2} (b_m + O(b_m^2)) \exp(c_m b_m/2 - b_m^2/8) m^{-1}. \end{aligned}$$

If $b_m := 1_{\{m>1\}} c_m^{-1/2} = o(1)$ the latter bound is easily seen to be $a_m m^{-1}$ with $a_m = a_m(b_m) \rightarrow \infty$. Thus

$$P\{\theta \in \tilde{S}\} \leq (1 - a_m m^{-1})^m \rightarrow 0.$$

Finally, one may replace b_m with $\max\{b_m, (1 - a_m m^{-1})^m\}$. This yields the assertion of part (b). \square

8. Related optimization problems

As in Section 4, let (\mathcal{G}, k) be either $(\mathcal{G}_\uparrow, 1)$ or $(\mathcal{G}_{\text{conv}}, 2)$. In view of future applications to other regression models we extend our framework slightly and consider $\langle g, h \rangle := \int gh \, d\mu$, $\|g\| := \langle g, g \rangle^{1/2}$, for some measure μ on the real line such that $\mu(C) < \infty$ for bounded intervals $C \subset \mathbb{R}$.

Let ψ be some bounded function on the real line with $\psi(x) = 0$ for $x \notin [-a, b]$ and $\langle 1, \psi \rangle \geq 0$, where $a, b \geq 0$. The next lemma provides sufficient conditions for one of the following two requirements:

$$\langle g, \psi \rangle \leq g(0) \langle 1, \psi \rangle \quad \text{whenever } g \in \mathcal{G}, 1_{[-a,b]} g \in L^1(\mu), \quad (8.1)$$

$$\langle g, \psi \rangle \geq g(0) \langle 1, \psi \rangle \quad \text{whenever } g \in \mathcal{G}, 1_{[-a,b]} g \in L^1(\mu). \quad (8.2)$$

Lemma 8.1. *Let $\mathcal{G} = \mathcal{G}_\uparrow$ and $\psi \geq 0$. Then $b = 0$ entails condition (8.1), while $a = 0$ implies condition (8.2).*

Let $\mathcal{G} = \mathcal{G}_{\text{conv}}$ and $\int_{-\infty}^{\infty} x\psi(x)\mu(dx) = 0$. Condition (8.2) is satisfied if $\psi \geq 0$. On the other hand, condition (8.1) is a consequence of the following two requirements: $\int x^\pm \psi(x)\mu(dx) = 0$ and

$$\psi \begin{cases} \geq 0 & \text{on } [c, d], \\ \leq 0 & \text{on } \mathbb{R} \setminus [c, d], \end{cases}$$

for some numbers $c < 0 < d$, where $\mu([-a, c])$, $\mu([d, b]) > 0$. (Here $y^+ := \max(y, 0)$ and $y^- := \max(-y, 0)$.)

With Lemma 8.1 at hand one can solve two minimization problems leading to the special kernels in (4.6) and (4.8). In both cases we consider two disjoint convex sets \mathcal{G}_0 , $\mathcal{G}_A \subset \mathcal{G}$ and construct functions $G_0 \in \mathcal{G}_0$, $G_A \in \mathcal{G}_A$ such that

$$\|G_0 - G_A\| = \min_{g_0 \in \mathcal{G}_0, g_A \in \mathcal{G}_A} \|g_0 - g_A\|. \quad (8.3)$$

Theorem 8.2. *Let $\mathcal{G}_0 := \{g \in \mathcal{G} : g(0) \leq -1\}$ and $\mathcal{G}_A := \{g \in \mathcal{G} \cap \mathcal{H}_{k,1} : g(0) \geq 0\}$. If $\mathcal{G} = \mathcal{G}_\uparrow$ let $G_A(x) := x$ and*

$$G_0(x) := \begin{cases} -1 & \text{if } x \in [-1, 0], \\ G_A(x) & \text{otherwise.} \end{cases}$$

If $\mathcal{G} = \mathcal{G}_{\text{conv}}$ let $G_A(x) := x^2/2$ and

$$G_0(x) := \begin{cases} -1 + (a/2 + 1/a)x^- + (b/2 + 1/b)x^+ & \text{if } x \in [-a, b], \\ G_A(x) & \text{otherwise,} \end{cases}$$

where $a, b \geq 2^{1/2}$ are chosen such that $\int x^\pm(G_A - G_0)(x)\mu(dx) = 0$. Then equation (8.3) holds in both cases. More precisely, the function $\psi := G_A - G_0$ satisfies the inequalities $\langle 1, \psi \rangle \geq \|\psi\|^2$, (8.1) and

$$\langle g, \psi \rangle \geq \|\psi\|^2 - \langle 1, \psi \rangle \quad \text{whenever } g \in \mathcal{H}_{k,1}, g(0) \geq 0. \tag{8.4}$$

If μ is Lebesgue measure, $\psi = G_A - G_0$ coincides with the function $\psi^{(\ell)}$ in (4.6), where $a = b = 2$.

Theorem 8.3. Let $\mathcal{G}_0 := \{g \in \mathcal{G} : g(0) \geq 1\}$, $\mathcal{G}_A := \{g \in \mathcal{G} \cap \mathcal{H}_{k,1} : g(0) \leq 0\}$, and define G_A as in Theorem 8.2. If $\mathcal{G} = \mathcal{G}_\uparrow$ let

$$G_0(x) := \begin{cases} 0 & \text{if } x \in [0, 1], \\ G_A(x) & \text{otherwise.} \end{cases}$$

If $\mathcal{G} = \mathcal{G}_{\text{conv}}$ suppose that $\mu(-\infty, 0] > 0$ and let

$$G_0(x) := \begin{cases} 1 + cx & \text{if } x \in [-a, b], \\ G_A(x) & \text{otherwise,} \end{cases}$$

where $a := -c + (c^2 + 2)^{1/2}$, $b := c + (c^2 + 2)^{1/2}$, and c is chosen such that $\int x(G_0 - G_A)(x)\mu(dx) = 0$. Then equation (8.3) is satisfied in both cases. More precisely, the function $\psi := G_0 - G_A$ satisfies the inequalities $\langle 1, \psi \rangle \geq \|\psi\|^2$, (8.2) and

$$\langle g, \psi \rangle \leq \langle 1, \psi \rangle - \|\psi\|^2 \quad \text{whenever } g \in \mathcal{H}_{k,1}, g(0) \geq 0. \tag{8.5}$$

If μ is Lebesgue measure, $\psi = G_0 - G_A$ coincides with the function $\psi^{(u)}$ in (4.8), where $c = 0$ and $a = b = 2^{1/2}$.

The following lemma summarizes essential properties of the optimal kernels $\psi^{(\ell)}$ and $\psi^{(u)}$.

Lemma 8.4. Let $\psi^{(\ell)}$ and $\psi^{(u)}$ be the kernel functions in (4.6) and (4.8), and let $h, L > 0$ and $t \in \mathbb{R}$.

- (a) If $\mathcal{G} = \mathcal{G}_\uparrow$, then $\langle 1, \psi^{(\ell)} \rangle = \langle 1, \psi^{(u)} \rangle = \frac{1}{2}$ and $\|\psi^{(\ell)}\|^2 = \|\psi^{(u)}\|^2 = \frac{1}{3}$. If $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $f(y) - f(x) \geq L(y - x)$ for all $x < y$, then

$$f - Lh^{-1}\psi_{h,t}^{(\ell)}, f + Lh^{-1}\psi_{h,t}^{(u)} \in \mathcal{G}_\uparrow.$$

- (b) If $\mathcal{G} = \mathcal{G}_{\text{conv}}$, then $\langle 1, \psi^{(\ell)} \rangle = \frac{2}{3}$, $\|\psi^{(\ell)}\|^2 = \frac{8}{15}$, $\langle 1, \psi^{(u)} \rangle = 2^{2.5}/3$ and $\|\psi^{(u)}\|^2 = 2^{4.5}/15$. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be absolutely continuous with derivative f' such that $f'(y) - f'(x) \geq L(y - x)$ for all $x < y$. Then

$$f - Lh^{-2}\psi_{h,t}^{(\ell)}, f + Lh^{-2}\psi_{h,t}^{(u)} \in \mathcal{G}_{\text{conv}}.$$

- (c) In general, for any function $f \in \mathcal{H}_{k,L}$,

$$\begin{aligned} \langle f(t+h\cdot) - r + Lh^k, \psi^{(\ell)} \rangle &\geq Lh^k \|\psi^{(\ell)}\|^2 && \text{if } f(t) \geq r, \\ \langle f(t+h\cdot) - r - Lh^k, \psi^{(u)} \rangle &\leq -Lh^k \|\psi^{(u)}\|^2 && \text{if } f(t) \leq r. \end{aligned}$$

Proof of Lemma 8.1. The assertions for $\mathcal{G} = \mathcal{G}_\uparrow$ are a simple consequence of $g \leq g(0)$ on $]-\infty, 0]$ and $g \geq g(0)$ on $[0, \infty[$.

Now let $\mathcal{G} = \mathcal{G}_{\text{conv}}$. If $\psi \geq 0$ and $\int x\psi(x)\mu(dx) = 0$, then condition (8.2) follows from Jensen's inequality applied to the probability measure $P(dx) = \langle 1, \psi \rangle^{-1} \psi(x)\mu(dx)$.

On the other hand, suppose that $\psi \geq 0$ on $[c, d]$ and $\psi \leq 0$ on $\mathbb{R} \setminus [c, d]$, where $c < 0 < d$ and $\mu([-a, c]), \mu([d, b]) > 0$. For $g \in \mathcal{G}_{\text{conv}}$ with $1_{[-a, b]}g \in L^1(\mu)$, both $g(c)$ and $g(d)$ have to be finite, and we define

$$\tilde{g}(x) := g(x) - \begin{cases} d^{-1}(g(d) - g(0))x & \text{if } x \geq 0, \\ c^{-1}(g(c) - g(0))x & \text{if } x \leq 0. \end{cases}$$

By convexity of g , this auxiliary function \tilde{g} satisfies $\tilde{g} \leq g(0)$ on $[c, d]$ and $\tilde{g} \geq g(0)$ on $\mathbb{R} \setminus [c, d]$. Thus $\langle \tilde{g}, \psi \rangle \leq g(0)\langle 1, \psi \rangle$. If, in addition, $\int x^\pm \psi(x)\mu(dx) = 0$, then $\langle g, \psi \rangle = \langle \tilde{g}, \psi \rangle$. \square

Proof of Theorem 8.2. One can easily deduce from Lemma 8.1 that the function $\psi = G_A - G_0$ satisfies inequality (8.1). But G_A is an extremal point of \mathcal{G}_A in the sense that

$$G_A - g \in \mathcal{G} \quad \text{for any } g \in \mathcal{H}_{k,1}.$$

For let $x < y$. If $\mathcal{G} = \mathcal{G}_\uparrow$, then

$$(G_A - g)(y) - (G_A - g)(x) = y - x - (g(y) - g(x)) \geq y - x - |y - x| = 0,$$

whence $G_A - g$ is non-decreasing. In case of $\mathcal{G} = \mathcal{G}_{\text{conv}}$ the same argument applies to the first derivative of $G_A - g$. Together with (8.1), this implies that

$$\begin{aligned} \langle g, \psi \rangle &= \langle G_A, \psi \rangle - \langle G_A - g, \psi \rangle \\ &\geq \langle G_A, \psi \rangle - (G_A - g)(0)\langle 1, \psi \rangle \\ &= \langle G_A, \psi \rangle + g(0)\langle 1, \psi \rangle \\ &= \|\psi\|^2 + \langle G_0, \psi \rangle + g(0)\langle 1, \psi \rangle \\ &= \|\psi\|^2 + (g(0) - 1)\langle 1, \psi \rangle. \end{aligned} \tag{8.6}$$

Equation (8.6) follows from $\langle G_0, \psi \rangle = \langle -1, \psi \rangle$, which is easily verified. The special case $g = 0$ yields the inequality $\langle 1, \psi \rangle \geq \|\psi\|^2$. Then inequality (8.4) becomes obvious.

It remains to be shown that in case of $\mathcal{G} = \mathcal{G}_{\text{conv}}$ there exist numbers $a, b \geq 2^{1/2}$ such that $\psi = \psi(\cdot, a, b)$ satisfies $\int x^\pm \psi(x)\mu(dx) = 0$. In fact, for any fixed x the number $\psi(x, a, b) \leq 1$ can be shown to be continuous and decreasing in a and b . To be precise, $\psi(0, a, b) = 1$ and $\lim_{a \rightarrow \infty} \psi(x, a, \cdot) = \lim_{b \rightarrow \infty} \psi(y, \cdot, b) = -\infty$ for $x < 0 < y$. Hence, the assertion is a consequence of monotone convergence. \square

Proof of Theorem 8.3. This proof is analogous to the proof of Theorem 8.2 and thus omitted. \square

Proof of Lemma 8.4. The calculations of $\langle 1, \psi \rangle$ and $\|\psi\|^2$ are elementary and thus omitted. Elementary calculations show that $g := -Lh^{-k}\psi_{t,h}^{(\ell)}$ as well as $g := Lh^{-k}\psi^{(u)}$ satisfy

$$\left. \begin{array}{l} g(y) - g(x) \\ g'(y) - g'(x) \end{array} \right\} \geq -L(y - x) \quad \text{if } \mathcal{G} = \begin{cases} \mathcal{G}_{\uparrow}, \\ \mathcal{G}_{\text{conv}}, \end{cases}$$

where $g'(x)$ denotes any number between the right- and left-sided derivative of g at x . Thus $f + g$ belongs to \mathcal{G} , whenever f satisfies the inequalities stated in parts (a) and (b).

As for part (c), for $f \in \mathcal{H}_{k,L}$ and $t \in \mathbb{R}$, $h, c > 0$, the function $cf(t + h \cdot)$ belongs to \mathcal{H}_{k,cLh^k} . If we take $c := (Lh^k)^{-1}$, inequality (8.4) implies that

$$\begin{aligned} \langle f(t + h \cdot) - r + Lh^k, \psi^{(\ell)} \rangle &= Lh^k \langle c(f(t + h \cdot) - f(t)) + 1, \psi^{(\ell)} \rangle \\ &\geq Lh^k \|\psi^{(\ell)}\|^2. \end{aligned}$$

One can deduce the lower bound for $\langle f(t + h \cdot) - r - Lh^k, \psi^{(u)} \rangle$ analogously. \square

Acknowledgements

The author is grateful to Lars Hömke for his assistance in Section 5. The constructive comments of a referee and an associate editor helped to improve the presentation. This work was supported by Deutsche Forschungsgemeinschaft, grant Du 238/5-1.

References

- Bickel, P.J. and Rosenblatt, M. (1973) On some global measures of the deviations of density function estimates. *Ann. Statist.*, **1**, 1071–1095.
- Brown, L.D. and Low, M.G. (1996) Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.*, **24**, 2384–2398.
- Davies, P.L. (1995) Data features. *Statist. Neerlandica*, **49**, 185–245.
- Donoho, D.L. (1988) One-sided inference about functionals of a density. *Ann. Statist.*, **16**, 1390–1420.
- Dümbgen, L. (1998) New goodness-of-fit tests and their application to nonparametric confidence sets. *Ann. Statist.*, **26**, 288–314.
- Dümbgen, L. (2001) Confidence bands for convex median functions using sign tests. Preprint.
- Dümbgen, L. and Johns, R.B. (2003) Confidence bands for isotonic median curves using sign tests. *J. Comput. Graph. Statist.* To appear.
- Dümbgen, L. and Spokoiny, V.G. (2001) Multiscale testing of qualitative hypotheses. *Ann. Statist.*, **29**, 124–152.
- Eubank, R.L. and Speckman, P.L. (1993) Confidence bands in nonparametric regression. *J. Amer. Statist. Assoc.*, **88**, 1287–1301.
- Fan, J. and Zhang, W. (2000) Simultaneous confidence bands and hypothesis testing in varying-coefficient models. *Scand. J. Statist.*, **27**, 715–731.

- Fan, J., Zhang, C. and Zhang, J. (2001) Generalized likelihood ratio statistics and Wilks phenomenon. *Ann. Statist.*, **29**, 153–193.
- Grama, I. and Nussbaum, M. (1998) Asymptotic equivalence for nonparametric generalized linear models. *Probab. Theory Related Fields*, **111**, 167–214.
- Hall, P. and Titterton, D.M. (1988) On confidence bands in nonparametric density estimation. *J. Multivariate Anal.*, **27**, 228–254.
- Härdle, W. and Marron, J.S. (1991) Bootstrap simultaneous error bars for nonparametric regression. *Ann. Statist.*, **19**, 778–796.
- Hart, J.D. (1997) *Nonparametric Smoothing and Lack-of-Fit Tests*. New York: Springer-Verlag.
- Hengartner, N.W. and Stark, P.B. (1995) Finite-sample confidence envelopes for shape-restricted densities. *Ann. Statist.*, **23**, 525–550.
- Ingster, Yu.I. (1993a) Asymptotically minimax hypothesis testing for nonparametric alternatives I. *Math. Methods Statist.*, **2**, 85–114.
- Ingster, Yu.I. (1993b) Asymptotically minimax hypothesis testing for nonparametric alternatives II. *Math. Methods Statist.*, **2**, 171–189.
- Ingster, Yu.I. (1993c) Asymptotically minimax hypothesis testing for nonparametric alternatives III. *Math. Methods Statist.*, **2**, 249–268.
- Khas'minskii, R.Z. (1978) A lower bound on the risks of non-parametric estimates of densities in the uniform metric. *Theory Probab. Appl.*, **23**, 794–798.
- Knafl, G., Sachs, J. and Ylvisaker, D. (1985) Confidence bands for regression functions. *J. Amer. Statist. Assoc.*, **80**, 683–691.
- Nussbaum, M. (1996) Asymptotic equivalence of density estimation and white noise. *Ann. Statist.*, **24**, 2399–2430.
- Robertson, T., Wright, F.T. and Dykstra, R.L. (1988) *Order Restricted Statistical Inference*. New York: Wiley.

Received September 2000 and revised August 2002