LITERATURE REVIEW

The 100 Most Influential Publications in Cervical Spine Research

Nicola Rügsegger,* Sufian S. Ahmad, MD,* Lorin M. Benneker, MD,* Ulrich Berlemann, MD,[†]
Marius J. B. Keel, MD,* and Sven Hoppe, MD*

Study Design. Bibliometric study of current literature.

Objective. To identify and analyze the 100 most cited publications in cervical spine research.

Summary of Background Data. The cervical spine is a dynamic field of research with many advances made within the last century. The literature has, however, never been comprehensively analyzed to identify and compare the most influential articles as measured by the number of citations.

Methods. All databases of the Thomson Reuters Web of Knowledge were utilized in a two-step approach. First, the 150 most cited cervical spine studies up to and including 2014 were identified using four keywords. Second, all keywords related to the cervical spine found in the 150 studies ($n = 38$) were used to conduct a second search of the database. The top 100 most cited articles were hereby selected for further analysis of current and past citations, authorship, geographic origin, article type, and level of evidence.

Results. Total citations for the 100 studies identified ranged from 173 to 879. They were published in the time frame 1952 to 2008 in a total of 30 different journals. Most studies ($n = 42$) were published in the decade 1991–2000. Level of evidence ranged from 1 to 5 with 39 studies in the level 4 category. A total of 13 researchers were first author more than once and nine researchers senior author more than once. The two-step approach with a secondary widening of search terms yielded an additional 27 studies, including the first ranking article.

Conclusion. This bibliometric study is likely to include some of the most important milestones in the field of cervical spine research of the last 100 years.

Key words: bibliometric study, cervical research, cervical spine, citation analysis, citation classics, orthopedic surgery, spine, surgery, top 100.

Level of Evidence: 3

Spine 2016;41:538–548

Originally described as a mean to evaluate scientific journals,¹ citation analysis has become an established tool to measure the impact of scientific articles in bibliometric studies—a subset of systematic reviews. It is based on the idea that publications with a high value to the field of study get cited more often than others. Owing to the vast growth of the scientific literature, an overview of the most relevant past publications has become more important than ever. In more recent years, bibliometric studies have therefore gained momentum and were subsequently published for fields as broad as surgery in general,² medical specialties,^{3,4} anatomic regions,⁵ surgical procedures,⁶ geographic origin of studies,^{7,8} and individual disease entities.⁹

In the field of the human spine, two bibliometric studies have been published.^{10,11} Even though the cervical portion of the spine exhibits some unique anatomic and clinical features, so far there has not been a systematic analysis of the scientific literature about cervical spine research. This study aims to close this gap and give the reader an overview over the most cited cervical spine articles of the last hundred years.

MATERIALS AND METHODS

Objective

To identify and analyze the 100 most cited cervical spine research papers published between 1900 and 2014 in any journal (medical and nonmedical).

Inclusion Criteria

In order to get considered for the top 100 list, a study had to focus on the anatomic region between the skull base and the first thoracic vertebra, and include the bony, cartilaginous, and/or ligamentous structures. If other parts of the body were also under investigation in the same study to the same or a higher degree, the study was excluded. Also, a study was

From the *Department of Orthopedic Surgery and Traumatology, Inselspital, University of Bern, Bern, Switzerland; and [†]Das Rückenzentrum, Spine Surgery, Thun, Switzerland.

Acknowledgment date: August 6, 2015. First revision date: September 10, 2015. Acceptance date: September 11, 2015.

The manuscript submitted does not contain information about medical device(s)/drug(s).

No funds were received in support of this work.

No relevant financial activities outside the submitted work.

Address correspondence and reprint requests to Sven Hoppe, MD, Department of Orthopedic Surgery and Traumatology, Inselspital, University of Bern, Freiburgstrasse, 3010 Bern, Switzerland; E-mail: svenhoppe@gmail.com

DOI: 10.1097/BRS.0000000000001261

538 www.spinejournal.com

Copyright © 2016 Wolters Kluwer Health, Inc. Unauthorized reproduction of this article is prohibited.

March 2016

Figure 1. Two-step approach.

not included if it focused on an anatomic structure unrelated to the cervical spine. For example, a study about spontaneous ruptures of the cervical portion of the vertebral artery would not be included while a study about whiplash injury would.

Material

All databases and all journals of the Thomson Reuter's Web of Knowledge were used to identify eligible studies.

Data Collection

In order to yield a high comprehensiveness, a two-step approach was chosen, as depicted in the flowchart below (Figure 1).

1. A topic search with the following Boolean query was conducted within the Web of Knowledge databases, the asterisk (*) indicating every possible ending of the corresponding word: (*spine OR vertebra**) AND (*cervic** OR *neck*).

This search yielded 86,723 results, which were then sorted in descending order with respect to the number of total citations. Within the first 632 studies, 150 cervical spine papers were identified based on title and abstract.

2. A list of keywords was created with the goal to foresee research titles involving the cervical spine. The previously identified 150 studies were systematically analyzed for possible keywords. Together with additional keywords known to be relevant to the field of cervical spine research (e.g. Klippel-Feil syndrome), a search query consisting of 38 terms was built (Table 1) and used for a second database search by title. This time, every word was delimited by an OR term, hereby further increasing the search results. The second search yielded 5,584,189 results, which were again sorted in descending order with respect to the number of total citations. Out of the 9694 first studies, 100 cervical spine studies were included based on title and abstract.

For every study included, the following information was extracted: Title, year of publication, first author, senior author, geographic origin, total citations, and citations 2014.

Analysis

Every study was assigned to a subspecialty, type of paper, type of study, and a level of evidence. For the level of evidence classification, the guidelines by *J Bone Joint Surg Am* were used to assign a number 1 (highest level of evidence) to 5 (lowest level of evidence) for every clinical study. Nonclinical (*i.e.* basic research) studies were not assigned to a level of evidence. A consensus approach

TABLE 1. Keywords Used in Second Search

Alar Ligament
Articular Process
Atla*
Axi*
C-Spine
C1-C7 (7)
Cervic*
Craniocervic*
Dens
Disc*
Disk*
Facet Joint
Hangman*
Jefferson*
Klippel Feil
Laminoplast*
Longitudinal Ligament
Myelopath*
Neck
Occipitocervic*
Odontoid
Pedic*
Radiculopath*
Spin*
Spondyl*
Tetraplegia
Transpedic*
Transverse Process
Uncinate Process
Vertebr*
Whiplash
Zygapophys*

was used in cases of disagreement between the researchers.¹² If a study had only one author, he or she was considered to be both, first and senior author.

The programming language Python (version 3.4.3) was used for the statistical analysis. All studies were ranked for their total number of citations. In cases ties between papers, a second and third degree ordering was performed for average citations per year and the number of citations in 2014, respectively.

Self-citations were not removed or analyzed. In the majority of cases, the error margin introduced by removing self-citations would have been larger than the error margin present by including self-citations.

RESULTS

The 100 studies included ranged from 879 to 173 citations and were published in a total of 30 different journals (Table 2). The top two journals (*Spine* and *J Bone Joint Surg Am*) accounted for 42 studies (Table 3). All studies fell within the time frame from 1952 to 2008. No studies from the first half of the 20th century were included even though all studies published between 1900 and 2014 were eligible for inclusion. The 1990s were the most active decade with 42 papers represented (Figure 2). There appears to be an inverse correlation between the average citations per year since publication and article age (Figure 3). A total of 14 different countries were identified as country of origin, with the USA in first position (n = 48). Nine studies were published from the UK, seven from Canada, followed by Australia and the Netherlands (n = 6), Japan and Norway (n = 5), Switzerland (n = 4), Germany (n = 3), Finland and India (n = 2) as well as China, Israel, and New Zealand with one publication each. Even though no language restrictions were imposed during the search process, all 100 studies were published in English. A total of 88 of the included studies were clinical, whereas the remaining 12 were basic research. The evidence level ranged from 1 to 5 with 4 being the most common (n = 43) (Figure 4). Average level of evidence per decade changed over time with a trend towards better evidence levels (Figure 2).

There were 13 authors with two first authorships. No authors were first author more than twice. Nine authors were identified as senior authors more than once, ranging from two to five articles each (Table 4).

Most articles were assigned to the subspecialty “degenerative”, (n = 41), followed by “trauma” (n = 28), “general c-spine” (n = 10), “rheumatology” (n = 3), and “oncology” (n = 1). A total of 17 studies addressed more than one subspecialty (“Various”) out of which most (n = 15) were assigned to the “surgical technique” type of paper. The latter was also the most common type of paper with 30 studies overall being assigned to “surgical technique”, followed by “clinical outcomes” with 11 papers (Table 5). Similarly, the majority of publications were either considered to be “therapeutic” (n = 44) or “prognostic” (n = 29) (Figure 5).

Out of the 100 publications, 73 were identified through the first four keywords search, 27 through the second 38

keywords query. During the first search, the 150 studies included fell within the 632 top ranked studies of the entire search result, leading to a hit-rate of 0.24. The second search with 38 keywords yielded a hit-rate of 0.01 (100 studies selected out of the 9694 most cited studies in the search results).

The 2012 published citation analysis of the top 100 spine studies by Murray *et al* listed 14 cervical spine papers.¹⁰ They all also appear in this study.

DISCUSSION

Out of the top three publications, two were published in 1958 and both described the anterior removal of cervical intervertebral discs—the 1st ranking one was published by Cloward and the 3rd placed by Smith and Robinson. Despite their age, both studies are still “highly active” with 31 and 46 citations in 2014, respectively (average citations in 2014 for all studies listed: 15). This demonstrates the long lasting impact the described surgical technique has had on the field of cervical spine surgery.

Second placed is the publication by Vernon and Mior from 1991 describing the Neck Disability Index, which modifies the Oswestry Low Back Pain Index for its use for the cervical spine. Interestingly, in a recently published bibliographic study of the lumbar spine, a review of the Oswestry Low Back Pain Index published in 2000 is the third most cited lumbar spine paper.^{11,13} This highlights the importance pain and disability classifications play in clinical practice, both for the lumbar and the cervical spine. The Neck Disability Index paper happened to be the most cited study in 2014, too. It was cited 88 times in 2014, more often than the first and third placed papers combined (77 citations). It is therefore possible, that in upcoming years, this paper might become the new number one in cervical spine research in terms of citations.

The most recent publication appearing on the top 100 list is the 2008 paper by Hogg-Johnson *et al* reviewing the published evidence on the burden of neck pain in the general population. It has an above average citation count in 2014 of 32 citations, indicating a possible further rise in rank in the near future. Furthermore, it has the highest average citation rate between the date of publication and 2013 with 88 citations per year (average: 12.4). This finding illustrates the inverse correlation between average citations per year since publication and article age (Figure 3). In other words: The more recent a publication is, the more often it has gotten cited per year since publication. This is, however, not surprising, since for a more recent study to accumulate enough citations to “keep up” with older publications, it must have had more citations per year since publication. And since this study only lists the most cited papers it was to be expected to see such a phenomenon for younger publications. The oldest paper was published in 1952 by Brain *et al* and described the neurological manifestations of cervical spondylosis. In 2014, it was cited only five times, thus possibly reflecting a phenomenon known as “obliteration by incorporation”: The tendency of classic papers getting cited

TABLE 2. The 100 Most Cited Papers in Cervical Spine Research

Rank	Paper	Total Citations	Citations/ Year: Publi- cation Until 2013	Citations 2014
1	Cloward RB. The anterior approach for removal of ruptured cervical disks. <i>Journal of neurosurgery</i> 1958;15:602-17.	879	15.42	31
2	Vernon H, Mior S. The neck disability index—a study of reliability and validity. <i>Journal of Manipulative and Physiological Therapeutics</i> 1991;14:409-15	877	35.86	88
3	Smith GW, Robinson RA. The treatment of certain cervical-spine disorders by anterior removal of the intervertebral disc and interbody fusion. <i>The Journal of bone and joint surgery. American volume</i> 1958;40-A:607-24.	631	10.64	46
4	Anderson LD, Dalonzo RT. Fractures of odontoid process of axis. <i>Journal of Bone and Joint Surgery-American Volume</i> 1974;A 56:1663-74.	511	12.38	28
5	Hilibrand AS, Carlson GD, Palumbo MA, et al. Radiculopathy and myelopathy at segments adjacent to the site of a previous anterior cervical arthrodesis. <i>Journal of Bone and Joint Surgery-American Volume</i> 1999;81A:519-28.	509	32.57	53
6	Spitzer WO, Skovron ML, Salmi LR, et al. Scientific monograph of the Quebec task-force on whiplash-associated disorders—redefining whiplash and its management. <i>Spine</i> 1995;20:S1-S73.	499	26.06	30
7	Harms J, Melcher RP. Posterior C1-C2 fusion with polyaxial screw and rod fixation. <i>Spine</i> 2001;26:2467-71.	462	34.00	54
8	Hoffman JR, Mower WR, Wolfson AB, et al. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. <i>New England Journal of Medicine</i> 2000;343:94-9.	413	28.77	39
9	Brooks AL, Jenkins EB. Atlanto-axial arthrodesis by wedge compression method. <i>Journal of Bone and Joint Surgery-American Volume</i> 1978;60:279-84.	403	10.94	20
10	Cote P, Cassidy JD, Carroll L. The Saskatchewan Health and Back Pain Survey—The prevalence of neck pain and related disability in Saskatchewan adults. <i>Spine</i> 1998;23:1689-98.	386	24.53	18
11	Bohlman HH. Acute fractures and dislocations of the cervical-spine—analysis of 300 hospitalized-patients and review of the literature. <i>Journal of Bone and Joint Surgery-American Volume</i> 1979;61:1119-42.	376	10.71	12
12	Nurick S. Pathogenesis of spinal-cord disorder associated with cervical spondylosis. <i>Brain</i> 1972;95:87-&.	376	8.63	22
13	Boden SD, McCowin PR, Davis DO, et al. Abnormal magnetic-resonance scans of the cervical-spine in asymptomatic subjects—a prospective investigation. <i>Journal of Bone and Joint Surgery-American Volume</i> 1990;72A:1178-84.	370	15.17	21
14	Bohlman HH, Emery SE, Goodfellow DB, et al. Robinson anterior cervical discectomy and arthrodesis for cervical radiculopathy—long-term follow-up of 100 and 22 patients. <i>Journal of Bone and Joint Surgery-American Volume</i> 1993;75A:1298-307.	359	16.70	25
15	Hirabayashi K, Miyakawa J, Satomi K, et al. Operative results and postoperative progression of ossification among patients with ossification of cervical posterior longitudinal ligament. <i>Spine</i> 1981;6:354-64.	358	9.94	40
16	Cassidy JD, Carroll LJ, Cote P, et al. Effect of eliminating compensation for pain and suffering on the outcome of insurance claims for whiplash injury. <i>New England Journal of Medicine</i> 2000;342:1179-86.	356	26.54	11
17	Silber JS, Anderson DG, Daffner SD, et al. Donor site morbidity after anterior iliac crest bone harvest for single-level anterior cervical discectomy and fusion. <i>Spine</i> 2003;28:134-9.	349	31.20	37
18	Stiell IG, Wells GA, Vandemheen KL, et al. The Canadian C-spine rule for radiography in alert and stable trauma patients. <i>Jama-Journal of the American Medical Association</i> 2001;286:1841-8.	339	25.42	34
19	Lord SM, Barnsley L, Wallis BJ, et al. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. <i>New England Journal of Medicine</i> 1996;335:1721-6.	328	17.94	23
20	Ranawat CS, O'Leary P, Pellicci P, et al. Cervical-spine fusion in rheumatoid-arthritis. <i>Journal of Bone and Joint Surgery-American Volume</i> 1979;61:1003-10.	323	9.21	10

TABLE 2 (Continued)

Rank	Paper	Total Citations	Citations/ Year: Publi- cation Until 2013	Citations 2014
21	Bovim G, Schrader H, Sand T. Neck pain in the general-population. <i>Spine</i> 1994;19:1307-9.	320	16.21	12
22	Kessel M, Balling R, Gruss P. Variations of cervical-vertebrae after expression of a hox-1.1 transgene in mice. <i>Cell</i> 1990;61:301-8.	313	13.57	1
23	Bailey RW, Badgley CE. Stabilization of the cervical spine by anterior fusion. <i>The Journal of bone and joint surgery. American volume</i> 1960;42-A:565-94.	312	5.60	15
24	Goel A, Laheri V. Plate and screw fixation for atlantoaxial subluxation. <i>Acta Neurochirurgica</i> 1994;129:47-53.	291	13.58	33
25	Fielding JW, Hawkins RJ. Atlanto-axial rotatory fixation— (fixed rotatory subluxation of atlanto-axial joint). <i>Journal of Bone and Joint Surgery-American Volume</i> 1977;59:37-44.	282	7.53	11
26	Makela M, Heliovaara M, Sievers K, et al. Prevalence, determinants, and consequences of chronic neck pain in Finland. <i>American Journal of Epidemiology</i> 1991;134:1356-67.	280	12.36	8
27	Barnsley L, Lord S, Bogduk N. Whiplash injury. <i>Pain</i> 1994;58:283-307.	277	14.26	6
28	Madawi AA, Casey ATH, Solanki GA, et al. Radiological and anatomical evaluation of the atlantoaxial transarticular screw fixation technique. <i>Journal of Neurosurgery</i> 1997;86:961-8.	274	16.06	17
29	Hirabayashi K, Watanabe K, Wakano K, et al. Expansive open-door laminoplasty for cervical spinal stenotic myelopathy. <i>Spine</i> 1983;8:693-9.	272	8.43	19
30	Payne EE, Spillane JD. The cervical spine—an anatomico-pathological study of 70 specimens (using a special technique) with particular reference to the problem of cervical spondylosis. <i>Brain</i> 1957;80:571-&.	264	4.64	4
31	Ramirezsolis R, Zheng H, Whiting J, et al. Hoxb-4 (Hox-2.6) mutant mice show homeotic transformation of a cervical vertebra and defects in the closure of the sternal rudiments. <i>Cell</i> 1993;73:279-94.	263	13.00	3
32	Robinson RA, Walker AE, Donald CF, et al. The results of anterior interbody fusion of the cervical spine. <i>Journal of Bone and Joint Surgery-American Volume</i> 1962;44 (8):1569 -1587.	259	5.00	4
33	Shields LBE, Raque GH, Glassman SD, et al. Adverse effects associated with high-dose recombinant human bone morphogenetic protein-2 use in anterior cervical spine fusion. <i>Spine</i> 2006;31:542-7.	258	30.14	47
34	Schrader H, Obelieniene D, Bovim G, et al. Natural evolution of late whiplash syndrome outside the medicolegal context. <i>Lancet</i> 1996;347:1207-11.	253	14.71	3
35	Allen BL, Ferguson RL, Lehmann TR, et al. A mechanistic classification of closed, indirect fractures and dislocations of the lower cervical-spine. <i>Spine</i> 1982;7:1-27.	249	7.45	18
36	Odom GL, Finney W, Woodhall B. Cervical disk lesions. <i>Journal of the American Medical Association</i> 1958;166:23-8.	248	4.24	15
37	Schneider RC, Cherry G, Pantek H. The syndrome of acute central cervical spinal cord injury; with special reference to the mechanisms involved in hyperextension injuries of cervical spine. <i>Journal of neurosurgery</i> 1954;11:546-77.	247	4.10	5
38	Brain WR, Northfield D, Wilkinson M. The neurological manifestations of cervical spondylosis. <i>Brain</i> 1952;75:187-225.	246	3.95	5
39	Barnsley L, Lord SM, Wallis BJ, et al. The prevalence of chronic cervical zygapophysial joint pain after whiplash. <i>Spine</i> 1995;20:20-5.	245	13.33	5
40	Gore DR, Sepic SB. Anterior cervical fusion for degenerated or protruded disks—a review of 146 patients. <i>Spine</i> 1984;9:667-71.	244	7.97	13
41	Eck JC, Humphreys SC, Lim TH, et al. Biomechanical study on the effect of cervical spine fusion on adjacent-level intradiscal pressure and segmental motion. <i>Spine</i> 2002;27:2431-4.	240	18.73	34
42	Lord SM, Barnsley L, Wallis BJ, et al. Chronic cervical zygapophysial joint pain after whiplash—A placebo-controlled prevalence study. <i>Spine</i> 1996;21:1737-44.	240	13.71	7
43	Sen CN, Sekhar LN. An extreme lateral approach to intradural lesions of the cervical-spine and foramen magnum. <i>Neurosurgery</i> 1990;27:197-204.	237	9.83	11

TABLE 2 (Continued)

Rank	Paper	Total Citations	Citations/ Year: Publication Until 2013	Citations 2014
44	Nurick S. Natural-history and results of surgical treatment of spinal-cord disorder associated with cervical spondylosis. <i>Brain</i> 1972;95:101-&.	227	5.41	5
45	Hurwitz EL, Aker PD, Adams AH, et al. Manipulation and mobilization of the cervical spine—A systematic review of the literature. <i>Spine</i> 1996;21:1746-59.	226	13.18	2
46	Scholten-Peeters GGM, Verhagen AP, Bekkering GE, et al. Prognostic factors of whiplash-associated disorders: A systematic review of prospective cohort studies. <i>Pain</i> 2003;104:303-22.	224	21.90	5
47	Radanov BP, Sturzenegger M, Distefano G. Long-term outcome after whiplash injury—a two-year follow-up considering features of injury mechanism and somatic, radiologic, and psychosocial findings. <i>Medicine</i> 1995;74:281-97.	223	12.17	4
48	Hogg-Johnson S, van der Velde G, Carroll LJ, et al. The burden and determinants of neck pain in the general population—Results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. <i>Spine</i> 2008;33:S39-S51.	222	38.00	32
49	Wright NM, Laurysen C. Vertebral artery injury in C1-2 transarticular screw fixation: Results of a survey of the American Association of Neurological Surgeons/Congress of Neurological Surgeons section on disorders of the spine and peripheral nerves. <i>Journal of Neurosurgery</i> 1998;88:634-40.	221	13.73	15
50	Borghouts JAJ, Koes BW, Bouter LM. The clinical course and prognostic factors of non-specific neck pain: a systematic review. <i>Pain</i> 1998;77:1-13.	220	14.13	8
51	Lunsford LD, Bissonette DJ, Jannetta PJ, et al. Anterior surgery for cervical disk disease .1. Treatment of lateral cervical disk herniation in 253 cases. <i>Journal of Neurosurgery</i> 1980;53:1-11.	219	6.42	7
52	Grob D, Crisco JJ, Panjabi MM, et al. Biomechanical evaluation of four different posterior atlantoaxial fixation techniques. <i>Spine</i> 1992;17:480-90.	218	9.95	9
53	Gore DR, Sepic SB, Gardner GM. Roentgenographic findings of the cervical-spine in asymptomatic people. <i>Spine</i> 1986;11:521-4.	218	7.56	14
54	Jull G, Trott P, Potter H, et al. A randomized controlled trial of exercise and manipulative therapy for cervicogenic headache. <i>Spine</i> 2002;27:1835-43.	216	18.18	16
55	Ariens GA, van Mechelen W, Bongers PM, et al. Physical risk factors for neck pain. <i>Scandinavian Journal of Work Environment & Health</i> 2000;26:7-19.	214	15.23	16
56	Radhakrishnan K, Litchy WJ, Ofallon WM, et al. Epidemiology of cervical radiculopathy—a population-based study from Rochester, Minnesota, 1976 through 1990. <i>Brain</i> 1994;117:325-35.	213	9.79	27
57	Jeanneret B, Magerl F. Primary posterior fusion C1/2 in odontoid fractures—indications, technique, and results of transarticular screw fixation. <i>Journal of Spinal Disorders</i> 1992;5:464-75.	212	9.57	11
58	Sterling M, Jull G, Vicenzino B, et al. Sensory hypersensitivity occurs soon after whiplash injury and is associated with poor recovery. <i>Pain</i> 2003;104:509-17.	211	19.30	18
59	Grob D, Jeanneret B, Aebi M, et al. Atlantoaxial fusion with transarticular screw fixation. <i>Journal of Bone and Joint Surgery-British Volume</i> 1991;73:972-6.	209	9.14	8
60	Paramore CG, Dickman CA, Sonntag VKH. The anatomical suitability of the C1-2 complex for transarticular screw fixation. <i>Journal of Neurosurgery</i> 1996;85:221-4.	207	11.18	17
61	Davis JW, Phreaner DL, Hoyt DB, et al. The etiology of missed cervical-spine injuries. <i>Journal of Trauma-Injury Infection and Critical Care</i> 1993;34:342-6.	206	10.05	5
62	Panjabi MM, Durancieu J, Goel V, et al. Cervical human vertebrae - quantitative 3-dimensional anatomy of the middle and lower regions. <i>Spine</i> 1991;16:861-9.	206	8.73	14
63	Baba H, Furusawa N, Imura S, et al. Late radiographic findings after anterior cervical fusion for spondylotic myeloradiculopathy. <i>Spine</i> 1993;18:2167-73.	205	9.40	17
64	Penning L. Normal movements of cervical-spine. <i>American Journal of Roentgenology</i> 1978;130:317-26.	205	5.57	10
65	Conlon PW, Isdale IC, Rose BS. Rheumatoid arthritis of cervical spine—an analysis of 333 cases. <i>Annals of the Rheumatic Diseases</i> 1966;25:120-&.	205	4.30	3

TABLE 2 (Continued)

Rank	Paper	Total Citations	Citations/ Year: Publication Until 2013	Citations 2014
66	Sjaastad O, Fredriksen TA, Pfaffenrath V. Cervicogenic headache—diagnostic criteria. <i>Headache</i> 1990;30:725-6.	200	8.43	6
67	Teresi LM, Lufkin RB, Reicher MA, et al. Asymptomatic degenerative disk disease and spondylosis of the cervical-spine—MR imaging. <i>Radiology</i> 1987;164:83-8.	199	7.35	8
68	Ylinen J, Takala EP, Nykanen M, et al. Active neck muscle training in the treatment of chronic neck pain in women—A randomized controlled trial. <i>Jama-Journal of the American Medical Association</i> 2003;289:2509-16.	198	17.80	20
69	Fang HSY, Ong GB. Direct anterior approach to the upper cervical spine. <i>Journal of Bone and Joint Surgery-American Volume</i> 1962;44:1588-604.	197	3.82	2
70	Jones EL, Heller JG, Silcox DH, et al. Cervical pedicle screws versus lateral mass screws—Anatomic feasibility and biomechanical comparison. <i>Spine</i> 1997;22:977-82.	194	11.25	14
71	Norris SH, Watt I. The prognosis of neck injuries resulting from rear-end vehicle collisions. <i>Journal of Bone and Joint Surgery-British Volume</i> 1983;65:608-11.	192	6.40	0
72	Youdas JW, Carey JR, Garrett TR. Reliability of measurements of cervical-spine range of motion-comparison of three methods. <i>Physical Therapy</i> 1991;71:98-104.	191	8.14	12
73	Dickman CA, Sonntag VKH. Posterior C1-C2 transarticular screw fixation for atlantoaxial arthrodesis. <i>Neurosurgery</i> 1998;43:275-80.	189	11.87	11
74	Caspar W, Barbier DD, Klara PM. Anterior cervical fusion and Caspar plate stabilization for cervical trauma. <i>Neurosurgery</i> 1989;25:491-502.	189	7.63	6
75	Rogers WA. Fractures and dislocations of the cervical spine; an end-result study. <i>The Journal of bone and joint surgery. American volume</i> 1957;39-A:341-76.	189	3.36	1
76	Buskila D, Neumann L, Vaisberg G, et al. Increased rates of fibromyalgia following cervical spine injury—A controlled study of 161 cases of traumatic injury. <i>Arthritis and Rheumatism</i> 1997;40:446-52.	188	11.38	6
77	Mair WGP, Druckman R. The pathology of spinal cord lesions and their relation to the clinical features in protrusion of cervical intervertebral discs. <i>Brain</i> 1953;76:70-91.	188	3.10	2
78	Flanders AE, Schaefer DM, Doan HT, et al. Acute cervical-spine trauma - correlation of MR imaging findings with degree of neurologic deficit. <i>Radiology</i> 1990;177:25-33.	187	7.91	5
79	Dwyer A, Aprill C, Bogduk N. Cervical zygapophyseal joint pain patterns .1. A study in normal volunteers. <i>Spine</i> 1990;15:453-7.	187	7.65	11
80	Menezes AH, Vangilder JC. Transoral-transpharyngeal approach to the anterior craniocervical junction—10-year experience with 72 patients. <i>Journal of Neurosurgery</i> 1988;69:895-903.	187	7.16	8
81	Fielding JW, Vancocochran G, Lawsing JF, et al. Tears of transverse ligament of atlas—clinical and biomechanical study. <i>Journal of Bone and Joint Surgery-American Volume</i> 1974;A 56:1683-91.	187	4.72	3
82	Emery SE, Bohlman HH, Bolesta MJ, et al. Anterior cervical decompression and arthrodesis for the treatment of cervical spondylotic myelopathy—Two to 17-year follow-up. <i>Journal of Bone and Joint Surgery-American Volume</i> 1998;80A:941-51.	186	11.53	13
83	Cattell HS, Filtzer DL. Pseudosubluxation and other normal variations in cervical spine in children. A study of 160 children. <i>Journal of Bone and Joint Surgery-American Volume</i> 1965;A 47:1295-&.	186	3.83	2
84	Wolf BS, Khilnani M, Malis L. The sagittal diameter of the bony cervical spinal canal and its significance in cervical spondylosis. <i>Jour Mt Sinai Hosp</i> 1956;23:283-92.	186	3.21	3
85	Sharp J, Purser DW. Spontaneous atlanto-axial dislocation in ankylosing spondylitis and rheumatoid arthritis. <i>Annals of the rheumatic diseases</i> 1961;20:47-77.	185	3.54	1

TABLE 2 (Continued)

Rank	Paper	Total Citations	Citations/ Year: Publi- cation Until 2013	Citations 2014
86	Obelieniene D, Schrader H, Bovim G, <i>et al.</i> Pain after whiplash: a prospective controlled inception cohort study. <i>Journal of Neurology Neurosurgery and Psychiatry</i> 1999;66:279-83.	184	12.79	5
87	Ariens GAM, van Mechelen W, Bongers PM, <i>et al.</i> Psychosocial risk factors for neck pain: A systematic review. <i>American Journal of Industrial Medicine</i> 2001;39:180-93.	183	14.33	11
88	Goel A, Desai KI, Muzumdar DP. Atlantoaxial fixation using, plate and screw method: A report of 160 treated patients. <i>Neurosurgery</i> 2002;51:1351-6.	182	15.00	17
89	Lees F, Turner JW. Natural history and prognosis of cervical spondylosis. <i>British medical journal</i> 1963;2:1607-10.	182	3.62	1
90	Stiell IG, Clement CM, McKnight RD, <i>et al.</i> The Canadian C-spine rule versus the NEXUS low-risk criteria in patients with trauma. <i>New England Journal of Medicine</i> 2003;349:2510-8.	180	15.80	22
91	Borghouts JAJ, Koes BW, Vondeling H, <i>et al.</i> Cost-of-illness of neck pain in The Netherlands in 1996. <i>Pain</i> 1999;80:629-36.	180	11.86	14
92	Henderson CM, Hennessy RG, Shuey HM, <i>et al.</i> Posterior-lateral foraminotomy as an exclusive operative technique for cervical radiculopathy—a review of 846 consecutively operated cases. <i>Neurosurgery</i> 1983;13:504-12.	180	5.70	9
93	Sjaastad O, Fredriksen TA, Pfaffenrath V, <i>et al.</i> Cervicogenic headache: Diagnostic criteria. <i>Headache</i> 1998;38:442-5.	178	11.00	13
94	Mummaneni PV, Burkus JK, Haid RW, <i>et al.</i> Clinical and radiographic analysis of cervical disc arthroplasty compared with allograft fusion: a randomized controlled clinical trial. <i>Journal of Neurosurgery-Spine</i> 2007;6:198-209.	177	24.83	28
95	Sawin PD, Traynelis VC, Menezes AH. A comparative analysis of fusion rates and donor-site morbidity for autogeneic rib and iliac crest bone grafts in posterior cervical fusions. <i>Journal of Neurosurgery</i> 1998;88:255-65.	176	11.07	10
96	Kotani Y, Cunningham BW, Abumi K, <i>et al.</i> Biomechanical analysis of cervical stabilization systems—an assessment of transpedicular screw fixation in the cervical-spine. <i>Spine</i> 1994;19:2529-39.	176	8.89	7
97	Sheehan S, Bauer RB, Meyer JS. Vertebral artery compression in cervical spondylosis—arteriographic demonstration during life of vertebral artery insufficiency due to rotation and extension of the neck. <i>Neurology</i> 1960;10:968-86.	176	3.32	0
98	Clark CR, White AA. Fractures of the dens—a multicenter study. <i>Journal of Bone and Joint Surgery-American Volume</i> 1985;67A:1340-8.	174	6.00	6
99	Satomi K, Nishu Y, Kohno T, <i>et al.</i> Long-term follow-up-studies of open-door expansive laminoplasty for cervical stenotic myelopathy. <i>Spine</i> 1994;19:507-10.	173	8.26	16
100	Zdeblick TA, Ducker TB. The use of freeze-dried allograft bone for anterior cervical fusions. <i>Spine</i> 1991;16:726-9.	173	7.22	7

MR imaging indicates magnetic resonance imaging.

less often because they are increasingly being incorporated into more recent studies.¹⁴ This same phenomenon might also account for the fact that most papers originated in the 1990s (n = 42) even though, theoretically, older publications have had more time to accumulate a high citation count.

The most cited paper accumulated 879 citations up until the end of 2014 while the 100th placed paper (a 1991 study by Zdeblick and Ducker describing the use of freeze-dried allograft bone for anterior cervical fusions) was cited a total of 173 times (average for all studies: 265 citations). While the top 10 studies spanned a range of 493 citations

(386–879 total citations), the last 10 studies spanned a range of only seven citations (173–180 total citations), thus indicating, that at the bottom there is the most “movement”: Here a single citation can decide over an appearance on the top 100 list. It is therefore likely that this study conducted again a few years later will look different, especially the bottom half.

Since during the search process no restriction was placed on the journal selection, a total of 30 different journals were identified. A total of 17 journals published, however, only one paper, while the top three journals (*Spine*, *J Bone Joint Surg Am*, and *J Neurosurg*) published half (n = 50) of all

TABLE 3. Journals With Multiple Publications

Journal	No. of Papers
<i>Spine</i>	25
<i>J Bone Joint Surg Am</i>	17
<i>J Neurosurg</i>	8
<i>Brain</i>	6
<i>Neurosurgery</i>	5
<i>Pain</i>	5
<i>N Engl J Med</i>	4
<i>JAMA</i>	3
<i>Ann Rheum Dis</i>	2
<i>Cell</i>	2
<i>Headache</i>	2
<i>J Bone Joint Surg Br</i>	2
<i>Radiology</i>	2
Total	83

studies. A similar pattern for spine research was noted before,^{10,11} indicating the degree of specialization and established leaderships among scientific journals. The top four countries of origin are all English speaking and accounted for a total of 70 studies, with the USA in first position (n = 48).

Figure 2. Publications per decade with average level of evidence for clinical studies. **Figure 3.** Correlation between article age and average citations per year since publication. **Figure 4.** Level of evidence.

As seen in bibliographic studies in different fields,⁵ evidence level 4 was the most common in this study as well with 43 papers. Only eight studies were deemed to be of level 1 by the research team. Many papers on the top 100 list provide something novel, most commonly a surgical technique and most often with relatively few patients and without a comparison group. It seems likely, that many studies conducted on the same surgical technique at a later point in time cite the classic study introducing the technique into the medical literature, thus elevating its citation count. After some time, the before mentioned phenomenon of “obliteration by incorporation” might, however, gradually offset this effect. While it is tempting to conclude that the level of

TABLE 4. First and Senior Authors With Multiple Publications

First Author	No. of Papers	Senior Author	No. of Papers
Ariens, GA	2	Bogduk, N	5
Barnsley, L	2	Sand, T	3
Bohlman, HH	2	Bohlman, HH	2
Borghouts, JAJ	2	Bouter, LM	2
Fielding, JW	2	Jones, PK	2
Goel, A	2	Nurick, S	2
Gore, DR	2	Pfaffenrath, V	2
Grob, D	2	Sonntag, VKH	2
Hirabayashi, K	2	Van der wal, G	2
Lord, SM	2		
Nurick, S	2		
Sjaastad, O	2		
Stiell, IG	2		

TABLE 5. Type of Paper

Type of Paper	No. of Papers
Surgical technique	30
Clinical outcomes	11
Anatomy	10
Natural history	10
Epidemiology	10
Clinical description	9
Classification	5
Conservative therapy	4
Clinical guidelines	3
Imaging	3
Genetics	2
Questionnaire development	1
Physical examination Technique	1
Pathogenesis	1
Total	100

evidence is not a criterion for publication success (as measured in the number of citations), this might not be true as there is a tendency towards a better level of evidence as time progresses (Figure 2). While for studies from the 1950s level of evidence averaged to 4.1, in the last decade (2001–2010) it went down to 2.4. A possible explanation might be that in earlier times a lot of studies introduced “a first”, while more recently more research has been conducted on the evaluation about previously made assumptions, requiring a much stronger statistical power and therefore a lower level of evidence. The decrease in level of evidence seen over time in this study is, however, not unique and might as well be due to a general tendency in the medical literature towards a better level of evidence.¹⁵

Because no restriction was placed on the field of study, basic research papers (n=12) appeared alongside with clinical studies (n=88). While approximately half (n=5) of the basic research was conducted on the anatomy of the cervical spine, two studies also came from the field of genetics demonstrating the effect the expression of two genes (Hox-1.1 and Hoxb-4) have on the development of cervical vertebrae in mice. Both studies (ranked 22nd and 31st) were published in the 90s and both have, despite their relatively young age, below average citations in 2014 (one time and three times, respectively). It has been noted before that basic research tends to get cited faster but not more often than clinical research.⁵

“Degenerative” is the most common type with 41 studies, followed by Trauma (n=28). A total of 11 papers within the latter category dealt with some aspect of whiplash injury, possibly reflecting the medical and political controversies this disease entity has been attracting. Only few studies fell into the categories “Rheumatology” (n=3) and “Oncology” (n=1). The 2012 bibliometric study on the entire spine also listed a table with the number of articles by subspecialty. The distribution of studies was very similar, showing that the research focus for the cervical

Figure 5. Study type.

spine is in general not much different from the entire spine.¹⁰

Even though citation analysis has its limitations,^{16,17} it is the current de facto gold standard for measuring the impact of scientific literature.

It is important to note that citation analysis can reflect the situation of only one point in time. Every day new publications appear citing previous papers. Thus, a citation analysis is as dynamic as the analyzed research field as a whole. Important progress is made constantly and a bibliographic study will most likely look different already months later.

The chosen two-step approach to identify eligible publications demonstrates the impact a widening of search terms can have: A total of 27 additional studies were identified, including the first ranking one. This improvement in comprehensiveness came, however, at the cost of a drastically decreased hit rate (studies meeting all inclusion criteria per studies manually sorted) from 0.24 with the initial four keywords down to 0.01 with 38 keywords. Even though this approach therefore consumes more of the research team's resources, because of the much higher likelihood to identify all publications relevant to the investigation, we believe it to be a valid approach and recommend it to be considered for future bibliometric studies. Often, the lack of higher-level keywords in published research prevents it from getting identified by the obvious search terms. For example, one study in our list had no keywords pointing to the spine. The sole indicator for it being a cervical spine study was the term “atlantoaxial” in the title. This demonstrates the importance of the attempt to foresee as many relevant titles as possible. As our experience goes, even in a rather narrow field such as the cervical spine it can, however, prove difficult to identify all the required terms. A preliminary search to identify keywords can help in such a situation.

Unfortunately, no matter what the approach chosen, it is impossible to know whether every eligible study really has been identified without manually sorting every paper ever published. With the two-step approach chosen, we however believe it to be at least highly likely that the “true” list of the 100 most cited cervical spine publications would not look much different than the list published here.

➤ Key Points

- ❑ The most cited article identified with a total of 879 citations was the publication in *J Neurosurg* from 1958 by the single author Cloward describing the anterior approach for the removal of ruptured cervical discs.
- ❑ Second placed with a total citation count of 877 was the publication from 1991 by Vernon and Mior creating the Neck Disability Index by modification of the established Oswestry Low Back Pain Index.
- ❑ The third most cited paper was the 1958 published study by Smith and Robinson, also describing the anterior approach for the removal of intervertebral discs and interbody fusion.
- ❑ Most papers either fell into the subspecialty of degeneration ($n = 41$) or trauma ($n = 28$) and were most commonly published in the 1990s ($n = 42$).
- ❑ Most studies were published in *Spine* ($n = 25$) and originated in the USA ($n = 48$).

References

1. Garfield E. Citation analysis as a tool in journal evaluation - journals can be ranked by frequency and impact of citations for science policy studies. *Science* 1972;178:471-9.
2. Heller AR, Lepore M, Bhangu A, et al. Citation classics in surgery: an analysis of the 100 most cited academic articles in surgical journals. *Br J Surg* 2015;102:59.
3. Lefaivre KA, Shadgan B, O'Brien PJ. 100 most cited articles in orthopaedic surgery. *Clin Orthop Relat Res* 2011;469:1487-97.
4. Tas F. An analysis of the most-cited research papers on oncology: which journals have they been published in?. *Tumor Biol* 2014;35:4645-9.
5. Ahmad SS, Evangelopoulos DS, Abbasian M, et al. The hundred most-cited publications in orthopaedic knee research. *J Bone Joint Surg* 2014;96:e190.
6. Cassar-Gheiti AJ, Downey RE, Byrne DP, et al. The 25 most cited articles in arthroscopic orthopaedic surgery. *Arthroscopy* 2012;28:548-64.
7. Jia ZW, Ding F, Wu YH, et al. The 50 most-cited articles in orthopaedic surgery from Mainland China. *Clin Orthop Relat Res* 2015;473:2423-30.
8. Gurbuz Y, Sugun TS, Ozaksar K. A bibliometric analysis of orthopedic publications originating from Turkey. *Acta Orthop Traumatol* 2015;49:57-66.
9. Cao F, Li J, Li A, et al. Citation classics in acute pancreatitis. *Pancreatol* 2012;12:325-30.
10. Murray MR, Wang TY, Schroeder GD, et al. The 100 most cited spine articles. *Eur Spine J* 2012;21:2059-69.
11. Steinberger J, Skovrlj B, Caridi JM, et al. The top 100 classic papers in lumbar spine surgery. *Spine (Phila Pa 1976)* 2015;40:740-7.
12. Bhandari M, Swiontkowski MF, Einhorn TA, et al. Interobserver agreement in the application of levels of evidence to scientific papers in the American volume of the Journal of Bone and Joint Surgery. *J Bone Joint Surg Am* 2004;86-A:1717-20.
13. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976)* 2000;25:2940-52; discussion 52.
14. Garfield E. 100 citation-classics from the Journal-of-the-American-Medical-Association. *JAMA* 1987;257:52-9.
15. Grant HM, Tjoumakaris FP, Maltenfort MG, et al. Levels of evidence in the clinical sports medicine literature: are we getting better over time?. *Am J Sports Med* 2014;42:1738-42.
16. Boudoulas H. Essays on citation classics in cardiology. *Cardiology* 2014;127:263-4.
17. Ioannidis JPA, Boyack KW, Small H, et al. Bibliometrics: is your most cited work your best?. *Nature* 2014;514:561-2.