

Characterization of a Novel Composite Staphylococcal Cassette Chromosome *mec* in Methicillin-Resistant *Staphylococcus pseudintermedius* from Thailand

Patrrarat Chanchaithong,^{a,b} Nuvee Prapasarakul,^b Vincent Perreten,^a Sybille Schwendener^a

Institute of Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Bern, Switzerland^a; Department of Veterinary Microbiology, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand^b

A novel staphylococcal cassette chromosome *mec* (SCC*mec*) composite island (SCC*mec*_{AI16}-SCC*czzr*_{AI16}-CI) was identified in *Staphylococcus pseudintermedius*. Four integration site sequences for SCC subdivided the 60,734-bp island into 41,232-bp SCC*mec*_{AI16}, 19,400-bp SCC*czzr*_{AI16}, and 102-bp SCC-like_{AI16} elements. SCC*mec*_{AI16} represents a new combination of *ccrA1B3* genes with a class A *mec* complex. SCC*czzr*_{AI16} contains *ccrA1B6* and genes related to restriction modification and heavy metal resistance. SCC*mec*_{AI16}-SCC*czzr*_{AI16}-CI was found in methicillin-resistant *S. pseudintermedius* sequence type 112 (ST112) and ST111 isolated from dogs and veterinarians in Thailand.

Methicillin-resistant *Staphylococcus pseudintermedius* (MRSP) has been associated with clinical manifestations in companion animals and occasionally causes diseases in humans (1, 2). MRSP contains the staphylococcal cassette chromosome *mec* (SCC*mec*) element, which is defined by the presence of the *mec* gene that mediates β-lactam resistance and by the cassette chromosome recombinase (*ccr*) gene(s) responsible for site-specific integration/excision of the element (3). Composite SCC structures containing two *ccr* gene complexes likely resulting from multiple element integrations have been increasingly reported in *Staphylococcus aureus* and in coagulase-negative staphylococci where such structures frequently carry heavy metal resistance gene clusters (3–9). So far, only five SCC*mec* elements have been completely characterized in MRSP, including SCC*mec* II-III, SCC*mec* VII-241, SCC*mec* V, ΨSCC*mec*₅₇₃₉₅, and SCC*mec* IV, and are mainly associated with predominant clones of the following sequence types (ST): ST71, ST93, ST68, ST233, ST45, and ST261 (10–14). Recently, MRSP strains of ST112 and ST111 (a single locus variant of ST112) isolated from dogs and veterinarians in Thailand were found to contain an unusual combination of the class A *mec* gene complex with the type I *ccr* gene complex, suggesting a new SCC*mec* type (15). We therefore aimed to characterize this novel SCC*mec* element and to determine whether it is conserved within other MRSP isolates of the same clonal lineage.

Identification of the SCC*mec*_{AI16}-SCC*czzr*_{AI16} composite element (CI). MRSP strains of ST111 and ST112 were obtained from a previous study and are listed in Table 1 (15). Phenotypic and genotypic characteristics, including those determined by antimicrobial susceptibility testing and antibiotic resistance gene detection, pulsed-field gel electrophoresis (PFGE), and multilocus sequence typing (MLST), were determined previously (15). In this study, the strains were additionally tested for *mec*-associated direct repeat unit (*dru*) and *spa* types (16, 17), for the presence of the novel SCC*mec* element, and for resistance to heavy metals. The genomic DNA of all strains was extracted using the peqGOLD bacterial DNA kit (Peqlab Biotechnologie GmbH, Jena, Germany), and that of strain AI16 was sequenced using Ion Torrent semiconductor (Life Technologies, Carlsbad, CA) and Illumina MiSeq (Illumina, San Diego, CA) technologies. Sequence reads

from the Ion Torrent were assembled *de novo* using MIRA v3.4.1.1, generating 64 contigs with an N_{50} (length weighted median) of 121,469 bp and a contig sum of 2,829,899 bp (mean read length, 265 bp; average coverage, >100-fold). The sequences were corrected by read mapping with MiSeq reads using Geneious version R8 (Biomatters, Auckland, New Zealand) (18), and unresolved discrepancies were verified by Sanger sequencing (ABI Prism 3100 genetic analyzer; Applied Biosystems, Foster City, CA). Nucleotide analysis searching for SCC-associated components (*orfX* gene, *mec* gene complexes, *ccr* gene complexes, and integration site sequence [ISS] for SCC [19]) using BLAST (<http://blast.ncbi.nlm.nih.gov/>) enabled the identification of a 60,734-bp SCC composite element within a 100,011-bp contig. Open reading frames (ORF) were defined using Prodigal software for locating genes in prokaryotes (34) and were annotated manually using BLASTn and BLASTp algorithms. The element was integrated at the 3' end of the chromosomal *orfX* gene and contained a *mec* gene complex, two *ccr* gene complexes, and four ISSs. The ISSs contained direct repeats (DRs) that flanked the composite SCC element and that also divided it into three subunits, including SCC*mec*_{AI16}, SCC*czzr*_{AI16}, and SCC-like_{AI16} according to the International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements (IWG-SCC) (4) (Fig. 1). ISSs were also linked to imperfect inverted repeats (IRs) at all

Received 17 September 2015 Returned for modification 7 October 2015

Accepted 19 November 2015

Accepted manuscript posted online 7 December 2015

Citation Chanchaithong P, Prapasarakul N, Perreten V, Schwendener S. 2016. Characterization of a novel composite staphylococcal cassette chromosome *mec* in methicillin-resistant *Staphylococcus pseudintermedius* from Thailand. Antimicrob Agents Chemother 60:1153–1157. doi:10.1128/AAC.02268-15.

Address correspondence to Vincent Perreten, vincent.perreten@vetsuisse.unibe.ch.

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/AAC.02268-15>.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.

TABLE 1 Origin and molecular characteristics of the six methicillin-resistant *Staphylococcus pseudintermedius* strains used in this study

Strain	Source	Site	Sequence type ^a	PFGE result ^a	SCCmec ^b	<i>spa</i> type ^c	<i>dru</i> type ^d	Antimicrobial resistance profile [resistance genes] ^e
AI16	Dog	Groin with crusty exudate	ST112	A	SCCmec _{AI16} -SCCczr _{AI16} -CI	t05	11y	OXA-PEN-TET-GEN-KAN-STR-ERY-iCLI-SMX-TMP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (M), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (B), <i>sat4</i> and <i>dfrG</i>]
AJ1	Dog	Perineal carriage	ST112	A	SCCmec _{AI16} -SCCczr _{AI16} -CI with 1,868-bp insertion	t05	11cv	OXA-PEN-TET-GEN-KAN-STR-ERY-CLISMX-TMP-MUP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (K), <i>tet</i> (M), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (A), <i>erm</i> (B), <i>sat4</i> , <i>dfrG</i> and <i>mupA</i>]
AK5	Dog	Nasal carriage	ST112	A	SCCmec _{AI16} -SCCczr _{AI16} -CI	t05	11y	OXA-PEN-TET-GEN-KAN-STR-ERY-iCLISMX-TMP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (K), <i>tet</i> (M), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (B), <i>sat4</i> and <i>dfrG</i>]
AM33	Dog	Perineal carriage	ST111	C	SCCmec _{AI16} -SCCczr _{AI16} -CI	Negative	11y	OXA-PEN-TET-GEN-KAN-STR-ERY-iCLISMX-TMP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (K), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (B), <i>sat4</i> and <i>dfrG</i>]
VA26	Human	Nasal carriage	ST112	A	SCCmec _{AI16} -SCCczr _{AI16} -CI with 1,868-bp insertion	t06	11y	OXA-PEN-TET-GEN-KAN-STR-ERY-iCLISMX-TMP-CIP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (M), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (B), <i>sat4</i> and <i>dfrG</i>]
VB16	Human	Nasal carriage	ST112	A	SCCmec _{AI16} -SCCczr _{AI16} -CI	t06	11y	OXA-PEN-TET-GEN-KAN-STR-ERY-CLISMX-TMP-MUP-CIP [<i>mecA</i> , <i>blaZ</i> , <i>tet</i> (M), <i>aac</i> (6')-Ie, <i>ant</i> (6')-Ia, <i>aph</i> (2')-Ia, <i>aph</i> (3')-III, <i>erm</i> (B), <i>sat4</i> , <i>dfrG</i> and <i>mupA</i>]

^a MLST and PFGE analyses were performed as a part of the previous study (15).

^b SCCmec_{AI16}-SCCczr_{AI16}-CI was identified in all strains by long-range PCR and restriction analysis. AJ1 and VA26 contained a 1,868-bp insertion.

^c The *spa* typing resulted in two *spa* types, t05 (r01-r02-r03-r03-r03-r06-r05) and t06 (r01-r02-r03-r03-r06-r05), and one negative strain with no *spa* amplification by PCR.

^d The *dru* types presented *dru* repeats as follows: 11y, 5a-2d-4a-1b-2d-5b-3a-2g-3b-4e-3e; 11cv, 5a-2d-4a-1b-2d-6f-3a-2g-3b-4e-3e.

^e Antimicrobial resistance phenotypes and detection of resistance genes were determined in the previous study using broth dilution methods and microarray (15). Abbreviation of antimicrobials are as follows: OXA, oxacillin; PEN, penicillin; TET, tetracycline; GEN, gentamicin; KAN, kanamycin; STR, streptomycin; ERY, erythromycin; CLI, clindamycin; iCLI, inducible resistance to clindamycin; SMX, sulfamethoxazole; TMP, trimethoprim; MUP, mupirocin; CIP, ciprofloxacin. Antibiotic resistance genes and functions are as follows: *mecA*, penicillin-binding protein 2a; *blaZ*, β -lactamase; *tet*(K), tetracycline efflux protein; *tet*(M), ribosomal protective protein; *aac*(6')-Ie, aminoglycoside acetyltransferase; *ant*(6')-Ia, aminoglycoside nucleotidyltransferase; *aph*(2')-Ia and *aph*(3')-III, aminoglycoside phosphotransferases; *erm*(B) and *erm*(C), erythromycin resistance methylase; *sat4*, streptothricin acetyltransferase; *dfrG*, dihydrofolate reductase; *mupA*, isoleucyl-tRNA synthetase.

subunit boundaries, forming parts of the SCC attachment sites (*att*) (19) (Fig. 1).

SCCmec_{AI16}. SCCmec_{AI16} consisted of a classical SCCmec element, as it contained the *mec* gene as well as the *ccr* genes and was flanked by two DRs (DR1 and DR2). It was located downstream of *orfX*, had a size of 41,232 bp, and contained 44 ORFs. The class A *mec* gene complex was located directly downstream of *orfX* and was only separated by a small 108-bp noncoding joining region (Fig. 1). The second joining region of SCCmec_{AI16} spanning the region between the *mec* gene complex and the *ccr* gene complex contained several genes coding for hypothetical protein, cadmium resistance (*cadCAD*), transposase B and C of transposon Tn554, and a putative cyclopentanol dehydrogenase (*cpnA*). The entire 5' region of SCCmec_{AI16}, including the *mec* gene complex up to *cpnA*, was highly identical (99%) to the nucleotide sequence of SCCmec VII-241 (without *cpnA*) and to a hybrid SCCmec-*mecC* region of *Staphylococcus sciuri* GVG52 (including *cpnA*) (10, 21). However, SCCmec_{AI16} differed from those two elements by the presence of a different type of *ccr* gene complex and a different 3' end-joining region that encoded various hypothetical proteins, some of them already identified in other SCCmec elements (see Table S1 in the supplemental material). The *ccr* complex of SCCmec_{AI16} contained *ccrA1* and *ccrB3* genes and shared the closest identity to those of *S. aureus* JCSC6945 and *S. sciuri* MCS24 with 92% and

89% nucleotide identity, respectively (20, 22) (see Fig. S1 in the supplemental material). *ccrA1B3* was assigned as a type 8 *ccr* complex, which was originally found in SCCmec XI together with a class E *mec* complex in *S. aureus* M10/0061 (23). However, the *ccrA1* of SCCmec_{AI16} shared only 81% nucleotide identity with the *ccrA1* of SCCmec XI but shared up to 90% sequence similarity with the *ccrA1* in SCCmec IX and X in livestock-associated methicillin-resistant *S. aureus* (LA-MRSA) in Thailand (20, 24). The combination of the type 8 *ccr* complex and the class A *mec* complex of SCCmec_{AI16} has not been previously reported and represents a new SCCmec type in *S. pseudintermedius*.

SCCczr_{AI16} and SCC-like_{AI16}. The SCCczr_{AI16} element contained recombinase genes but no *mec* gene. It was flanked by DR2 and DR3, arranged in tandem with the SCCmec_{AI16}, and followed by a 102-bp noncoding SCC-like_{AI16} fragment demarcated by DR3 and DR4. SCCczr_{AI16} was 19 kb in length, contained 18 ORFs (see Table S1 in the supplemental material), and consisted of three main segments displaying different functions. The first segment contained genes coding for proteins (HsdR, HsdM, and HsdS) associated with a type I restriction-modification system. HsdR and HsdM showed 99% and 98% amino acid (aa) identity with those encoded on SCCfusC, respectively (25), and were combined with a novel HsdS that shared 61% aa identity to the closest HsdS (see Table S1), indicating a new restriction specificity in this




FIG 1 Genetic organization of the novel composite $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ element of *Staphylococcus pseudintermedius* strain AI16 and formation of circular intermediates. The single units of the $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ are flanked by direct repeats (DR) containing the integration site sequence (ISS) (red) and by inverted repeats (IR) (harpoon arrows) forming part of the attachment sites (*att*). White arrows represent sequences encoding hypothetical proteins, and other genes are color coded as follows: *orfX*, gray; *IS431*, pale green; *mec* operon, sky blue; transposases of Tn554, light gray; cyclopentanol dehydrogenase, brick red; type I restriction-modification system, orange; and *czr* operon, brown. The *ccrA1*, *ccrB3*, and *ccrB6* recombinase genes are shown in blue, green, and cyan, respectively. Small blue arrows indicate the primers used for PCR, identifying circularization and excision of the $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ subunits (see Tables S2 and S3 in the supplemental material). The genetic map of $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ was drawn using the Easyfig software (32).

MRSP lineage (26). The second segment contained a type 7 *ccr* complex, with *ccrA1* and *ccrB6* sharing 99% and 89% identity to those of SCC_{mec} X from LA-MRSA ST398 (JCSC6945) isolated from a human in Thailand (20) (see Fig. S1 in the supplemental material). This *ccr* complex was preceded by three ORFs that also displayed high similarity (96%, 100%, and 99%) to those found upstream of the *ccr* genes of SCC_{mec} X. The third segment situated at the 3' region of $SCC_{czr_{AI16}}$ showed 99% DNA identity to the 3' region of SCC_{SH32} identified in *Staphylococcus haemolyticus* SH32 isolated from a Chinese patient (27) and to a fragment of the SCC_{mec} V(5C2&5)c found in *S. aureus* (28) and *S. haemolyticus* (9). It carried six ORFs, including *czrC*, which has been shown to confer resistance to zinc and cadmium in *S. aureus* (29). While resistance to cadmium increased up to 4-fold in *S. pseudintermedius* strains containing *cadA* compared to that in strains which lack *cadA* or carry only *czrC*, we did not observe reduced susceptibility to zinc in any of the tested strains, including those containing *czrC*, using broth microdilution assays after 20-h and 48-h (data not shown) incubations at 37°C (30) (Table 2). The same MIC of zinc for strains with and without *czrC* indicates that this gene does not confer measurable zinc resistance in *S. pseudintermedius*. Additionally, decreased susceptibility to other heavy metals was not observed (Table 2). The presence of *cadA* and *czrC* genes was confirmed by PCR using the primers listed in Table S2 in the supplemental material.

Excision and circularization of $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ subunits. Ccr recombinases recognize the *att* sites of SCC and catalyze DNA cleavage, strand exchange, and recombination to integrate an element or to release a nonreplicative circular element. Sequencing PCR products obtained using primer pairs reading inwards and outwards from the *att* sites identified circular forms containing one copy of ISS as a joining region as well as chromosomal segments remaining after element excision. Such circular forms were detected for the individual $SCC_{mec_{AI16}}$ and $SCC_{czr_{AI16}}$ and for the composites $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}$ and $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ (Fig. 1; see also Table S3 in the supplemental material). This indicated CcrAB activity in the strain AI16 and possible mobilization of the individual $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ circular subunits. Notably, ISS3 was preferred over ISS4 for recombination, which may be caused by the presence of a divergent IR in the ISS4-containing *att* site (Fig. 1; see also Table S3).

Presence of $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ in additional MRSP isolates. Long-range PCR amplification followed by restriction analysis (see Tables S2 and S4 in the supplemental material) confirmed the overall structure of $SCC_{mec_{AI16}}-SCC_{czr_{AI16}}-CI$ in MRSP strain AI16 and allowed detection of similar composite elements in five additional clonally related MRSP strains of ST111 and ST112 isolated from dogs and humans (Table 1). Two strains, AJ1 (ST112) and VA26 (ST112), showed a minor variation in the restriction analysis caused by a 1,868-bp insertion between *czrC*

TABLE 2 Heavy metal susceptibility of *Staphylococcus aureus* and *Staphylococcus pseudintermedius* strains as determined using broth microdilution test

Strain	Metal resistance gene(s)	MIC (mM) of metal						Source
		CuSO ₄	CdCl ₂	ZnSO ₄	AsNaO ₂	Pb(CH ₃ COO) ₂	HgCl ₂	
<i>S. aureus</i> RN4220		4	≤0.015	0.5	0.03	≤0.125	≤0.015	33
<i>S. pseudintermedius</i> CCUG 49543 (=LMG 22219)		16	0.25	1	2	1	≤0.015	Culture Collection, University of Göteborg, Sweden
Methicillin-resistant <i>S. pseudintermedius</i> E120		16	0.25	1	2	1	≤0.015	1
KM241	<i>cadA</i>	16	1	1	2	1	≤0.015	10
196511	<i>czrC</i>	16	0.25	1	2	1	≤0.015	D. Elad, Kimron Veterinary Institute, Bet Dagan, Israel
AI16	<i>cadA</i> , <i>czrC</i>	16	1	1	2	1	≤0.015	This study, 15

and ISS3 as confirmed by primer walking and Sanger sequencing. In addition, the *SCCmec* element of AJ1 displayed an alternative hypervariable repeat region (*dru* 11cv) that was different from that of the other strains (*dru* 11y), suggesting different evolutionary processes within SCC of the same clonal lineage.

Characterization of the novel *SCCmec*_{AI16}-*SCCczr*_{AI16}-CI element in *S. pseudintermedius* revealed regions with homology to other *SCCmec* elements. The 5' region of *SCCmec*_{AI16} obviously shared a common ancestral origin with *SCCmec* elements found in *S. pseudintermedius* KM241 and in *S. sciuri* GVGS2 but displayed a substituted 3' region resulting in a new combination of the class A *mec* complex with *ccrA1B3* genes. The presence of *SCCczr*_{AI16} harboring a second *ccrAB* complex illustrates the challenge for PCR-based *SCCmec* typing. Furthermore, the location of two related *ccrAB* gene complexes aligned in proximity offer a source for homologous recombination-mediated deletion, a mechanism observed previously with *ccrC* sequences (28, 31). The presence of this composite element in other strains of the same lineage rather supported a spread of this cassette with a clone as opposed to an autonomous transfer between strains. However, the ability of the *SCCmec*_{AI16}-*SCCczr*_{AI16}-CI subunits to circularize independently may play a role in the mobilization and in the further evolutionary diversification of this composite *SCCmec* structure.

Nucleotide sequence accession numbers. The nucleotide sequences of the *SCCmec*_{AI16}-*SCCczr*_{AI16}-CI of *S. pseudintermedius* AI16 and the 1,868-bp insertion in *S. pseudintermedius* VA26 have been deposited in GenBank under accession numbers LN864705 and LN874217, respectively.

ACKNOWLEDGMENTS

We thank Matt Riley (University of Tennessee, Knoxville, TN) for advice with sequence analysis, Arshnee Moodley (University of Copenhagen, Copenhagen, Denmark) for help with *spa* typing, Richard V. Goering (Creighton University School of Medicine, Omaha, NE) for help with *dru* typing, Daniel Elad (Kimron Veterinary Institute, Bet Dagan, Israel) for providing strain 196511, and Alexandra Collaud (Institute of Veterinary Bacteriology, University of Bern) for technical assistance.

FUNDING INFORMATION

Institute of Veterinary Bacteriology, University of Bern provided funding to Vincent Perreten and Sybille Schwendener under grant number 35-539. Ratchadaphiseksomphot Endowment Fund, Chulalongkorn University provided funding to Patrrarat Chanchaithong under grant number GDNS 57-045-31-006.

REFERENCES

- Perreten V, Kadlec K, Schwarz S, Grönlund Andersson U, Finn M, Greko C, Moodley A, Kania SA, Frank LA, Bemis DA, Franco A, Iurescia M, Battisti A, Duim B, Wagenaar JA, van Duijkeren E, Weese JS, Fitzgerald JR, Rossano A, Guardabassi L. 2010. Clonal spread of methicillin-resistant *Staphylococcus pseudintermedius* in Europe and North America: an international multicentre study. *J Antimicrob Chemother* 65:1145–1154. <http://dx.doi.org/10.1093/jac/dkq078>.
- Stegmann R, Burnens A, Maranta CA, Perreten V. 2010. Human infection associated with methicillin-resistant *Staphylococcus pseudintermedius* ST71. *J Antimicrob Chemother* 65:2047–2048. <http://dx.doi.org/10.1093/jac/dkq241>.
- Shore AC, Coleman DC. 2013. Staphylococcal cassette chromosome *mec*: recent advances and new insights. *Int J Med Microbiol* 303:350–359. <http://dx.doi.org/10.1016/j.ijmm.2013.02.002>.
- International Working Group on the Classification of Staphylococcal Cassette Chromosome Elements (IWG-SCC). 2009. Classification of staphylococcal cassette chromosome *mec* (*SCCmec*): guidelines for reporting novel *SCCmec* elements. *Antimicrob Agents Chemother* 53:4961–4967. <http://dx.doi.org/10.1128/AAC.00579-09>.
- Kinnevey PM, Shore AC, Brennan GI, Sullivan DJ, Ehrlich R, Monecke S, Slickers P, Coleman DC. 2013. Emergence of sequence type 779 methicillin-resistant *Staphylococcus aureus* harboring a novel pseudo staphylococcal cassette chromosome *mec* (*SCCmec*)-SCC-*SCC*_{CRISPR} composite element in Irish hospitals. *Antimicrob Agents Chemother* 57:524–531. <http://dx.doi.org/10.1128/AAC.01689-12>.
- Balakuntla J, Prabhakara S, Arakere G. 2014. Novel rearrangements in the staphylococcal cassette chromosome *mec* type V elements of Indian ST772 and ST672 methicillin resistant *Staphylococcus aureus* strains. *PLoS One* 9(4):e94293. <http://dx.doi.org/10.1371/journal.pone.0094293>.
- Hill-Cawthorne GA, Hudson LO, El Ghany MF, Piepenburg O, Nair M, Dodgson A, Forrest MS, Clark TG, Pain, A. 2014. Recombinations in staphylococcal cassette chromosome *mec* elements compromise the molecular detection of methicillin resistance in *Staphylococcus aureus*. *PLoS One* 9(6):e101419. <http://dx.doi.org/10.1371/journal.pone.0101419>.
- Martins Simoes P, Rasigade JP, Lemriss H, Butin M, Genevra C, Lemriss S, Goering RV, Ibrahim A, Picaud JC, El Kabbaj S, Vandenesch F, Laurent F. 2013. Characterization of a novel composite staphylococcal cassette chromosome *mec* (*SCCmec*-*SCCcad/ars/cop*) in the neonatal sepsis-associated *Staphylococcus capitis* pulstotype NRCS-A.

- Antimicrob Agents Chemother 57:6354–6357. <http://dx.doi.org/10.1128/AAC.01576-13>.
9. Xue H, Wu Z, Li L, Li F, Wang Y, Zhao X. 2015. Coexistence of heavy metal and antibiotic resistance within a novel composite staphylococcal cassette chromosome in a *Staphylococcus haemolyticus* isolate from bovine mastitis milk. Antimicrob Agents Chemother 59:5788–5792. <http://dx.doi.org/10.1128/AAC.04831-14>.
 10. Descloux S, Rossano A, Perreten V. 2008. Characterization of new staphylococcal cassette chromosome *mec* (SCCmec) and topoisomerase genes in fluoroquinolone- and methicillin-resistant *Staphylococcus pseudintermedius*. J Clin Microbiol 46:1818–1823. <http://dx.doi.org/10.1128/JCM.02255-07>.
 11. Perreten V, Chanchaithong P, Prapasarakul N, Rossano A, Blum SE, Elad D, Schwendener S. 2013. Novel pseudo-staphylococcal cassette chromosome *mec* element (Ψ SCCmec₅₇₃₉₅) in methicillin-resistant *Staphylococcus pseudintermedius* CC45. Antimicrob Agents Chemother 57:5509–5515. <http://dx.doi.org/10.1128/AAC.00738-13>.
 12. Youn JH, Moodley A, Park YH, Sugimoto C. 2013. Genome sequence of methicillin-resistant *Staphylococcus pseudintermedius* sequence type 233 (ST233) strain K7, of human origin. Genome Announc 1:e00310-13.
 13. McCarthy AJ, Harrison EM, Stanczak-Mrozek K, Leggett B, Waller A, Holmes MA, Lloyd DH, Lindsay JA, Loeffler A. 2015. Genomic insights into the rapid emergence and evolution of MDR in *Staphylococcus pseudintermedius*. J Antimicrob Chemother 70:997–1007. <http://dx.doi.org/10.1093/jac/dku496>.
 14. Black CC, Solyman SM, Eberlein LC, Bemis DA, Woron AM, Kania SA. 2009. Identification of a predominant multilocus sequence type, pulsed-field gel electrophoresis cluster, and novel staphylococcal chromosomal cassette in clinical isolates of *mecA*-containing, methicillin-resistant *Staphylococcus pseudintermedius*. Vet Microbiol 139:333–338. <http://dx.doi.org/10.1016/j.vetmic.2009.06.029>.
 15. Chanchaithong P, Perreten V, Schwendener S, Tribuddharat C, Chongthaleong A, Niyomtham W, Prapasarakul N. 2014. Strain typing and antimicrobial susceptibility of methicillin-resistant coagulase-positive staphylococcal species in dogs and people associated with dogs in Thailand. J Appl Microbiol 117:572–586. <http://dx.doi.org/10.1111/jam.12545>.
 16. Moodley A, Stegger M, Ben Zakour NL, Fitzgerald JR, Guardabassi L. 2009. Tandem repeat sequence analysis of staphylococcal protein A (*spa*) gene in methicillin-resistant *Staphylococcus pseudintermedius*. Vet Microbiol 135:320–326. <http://dx.doi.org/10.1016/j.vetmic.2008.09.070>.
 17. Goering RV, Morrison D, Al Doori Z, Edwards GF, Gemmell CG. 2008. Usefulness of *mec*-associated direct repeat unit (*dru*) typing in the epidemiological analysis of highly clonal methicillin-resistant *Staphylococcus aureus* in Scotland. Clin Microbiol Infect 14:964–969. <http://dx.doi.org/10.1111/j.1469-0691.2008.02073.x>.
 18. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics 28:1647–1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>.
 19. Ito T, Ma XX, Takeuchi F, Okuma K, Yuzawa H, Hiramatsu K. 2004. Novel type V staphylococcal cassette chromosome *mec* driven by a novel cassette chromosome recombinase, *ccrC*. Antimicrob Agents Chemother 48:2637–2651. <http://dx.doi.org/10.1128/AAC.48.7.2637-2651.2004>.
 20. Li S, Skov RL, Han X, Larsen AR, Larsen J, Sørum M, Wulf M, Voss A, Hiramatsu K, Ito T. 2011. Novel types of staphylococcal cassette chromosome *mec* elements identified in clonal complex 398 methicillin-resistant *Staphylococcus aureus* strains. Antimicrob Agents Chemother 55:3046–3050. <http://dx.doi.org/10.1128/AAC.01475-10>.
 21. Harrison EM, Paterson GK, Holden MT, Ba X, Rolo J, Morgan FJ, Pichon B, Kearns A, Zadoks RN, Peacock SJ, Parkhill J, Holmes MA. 2014. A novel hybrid SCCmec-*mecC* region in *Staphylococcus sciuri*. J Antimicrob Chemother 69:911–918. <http://dx.doi.org/10.1093/jac/dkt452>.
 22. Urushibara N, Paul SK, Hossain MA, Kawaguchiya M, Kobayashi N. 2011. Analysis of staphylococcal cassette chromosome *mec* in *Staphylococcus haemolyticus* and *Staphylococcus sciuri*: identification of a novel *ccr* gene complex with a newly identified *ccrA* allotype (*ccrA7*). Microb Drug Resist 17:291–297. <http://dx.doi.org/10.1089/mdr.2010.0144>.
 23. Shore AC, Deasy EC, Slickers P, Brennan G, O'Connell B, Monecke S, Ehrlich R, Coleman DC. 2011. Detection of staphylococcal cassette chromosome *mec* type XI carrying highly divergent *mecA*, *mecI*, *mecR1*, *blaZ*, and *ccr* genes in human clinical isolates of clonal complex 130 methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 55:3765–3773. <http://dx.doi.org/10.1128/AAC.00187-11>.
 24. Lulitanond A, Ito T, Li S, Han X, Ma XX, Engchanil C, Chanawong A, Wilailuckana C, Jiwakanon N, Hiramatsu K. 2013. ST9 MRSA strains carrying a variant of type IX SCCmec identified in the Thai community. BMC Infect Dis 13:214. <http://dx.doi.org/10.1186/1471-2334-13-214>.
 25. Lin YT, Tsai JC, Chen HJ, Hung WC, Hsueh PR, Teng LJ. 2014. A novel staphylococcal cassette chromosomal element, SCCfusC, carrying *fusC* and *speG* in fusidic acid-resistant methicillin-resistant *Staphylococcus aureus*. Antimicrob Agents Chemother 58:1224–1227. <http://dx.doi.org/10.1128/AAC.01772-13>.
 26. Lindsay JA. 2010. Genomic variation and evolution of *Staphylococcus aureus*. Int J Med Microbiol 300:98–103. <http://dx.doi.org/10.1016/j.ijmm.2009.08.013>.
 27. Yu D, Pi B, Chen Y, Wang Y, Ruan Z, Otto M, Yu Y. 2014. Characterization of the staphylococcal cassette chromosome composite island of *Staphylococcus haemolyticus* SH32, a methicillin-resistant clinical isolate from China. PLoS One 9(1):e87346. <http://dx.doi.org/10.1371/journal.pone.0087346>.
 28. Vandendriessche S, Vanderhaeghen W, Larsen J, de Mendonca R, Hallin M, Butaye P, Hermans K, Haesebrouck F, Denis O. 2014. High genetic diversity of methicillin-susceptible *Staphylococcus aureus* (MSSA) from humans and animals on livestock farms and presence of SCCmec remnant DNA in MSSA CC398. J Antimicrob Chemother 69:355–362. <http://dx.doi.org/10.1093/jac/dkt366>.
 29. Cavaco LM, Hasman H, Stegger M, Andersen PS, Skov R, Fluit AC, Ito T, Aarestrup FM. 2010. Cloning and occurrence of *czrC*, a gene conferring cadmium and zinc resistance in methicillin-resistant *Staphylococcus aureus* CC398 isolates. Antimicrob Agents Chemother 54:3605–3608. <http://dx.doi.org/10.1128/AAC.00058-10>.
 30. Clinical and Laboratory Standards Institute. 2008. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; approved standard—3rd ed. CLSI document M31-A3. Clinical and Laboratory Standards Institute, Wayne, PA.
 31. Chlebowicz MA, Nganou K, Kozytska S, Arends JP, Engelmann S, Grundmann H, Ohlsen K, van Dijk JM, Buist G. 2010. Recombination between *ccrC* genes in a type V (5C2&5) staphylococcal cassette chromosome *mec* (SCCmec) of *Staphylococcus aureus* ST398 leads to conversion from methicillin resistance to methicillin susceptibility *in vivo*. Antimicrob Agents Chemother 54:783–791. <http://dx.doi.org/10.1128/AAC.00696-09>.
 32. Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer. Bioinformatics 27:1009–1010. <http://dx.doi.org/10.1093/bioinformatics/btr039>.
 33. Kreiswirth BN, Löfdahl S, Betley MJ, O'Reilly M, Schlievert PM, Bergdoll MS, Novick RP. 1983. The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. Nature 305:709–712. <http://dx.doi.org/10.1038/305709a0>.
 34. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.