

ERRATUM

Open Access



# Erratum to: 'Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial'

Lukas M. Löffel<sup>1</sup>, Bettina Kleeb<sup>1</sup>, Fiona C. Burkhard<sup>2</sup> and Patrick Y. Wuethrich<sup>1\*</sup>

Unfortunately, the original version of this article [1] contained a few errors in Table 1. The correct data for Table 1 has now been included below.

#### Author details

<sup>1</sup>Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland. <sup>2</sup>Department of Urology, Inselspital, Bern University Hospital, Bern, Switzerland.

Published online: 18 February 2016

#### Reference

1. Löffel LM, Bettina K, Burkhard FC, Wuethrich PY. Perioperative use of crystalloids in patients undergoing open radical cystectomy: balanced Ringer's maleate versus a glucose 5 %/potassium-based balanced solution: study protocol for a randomized controlled trial. *Trials*. 2014;15:276.

\* Correspondence: [patrick.wuethrich@insel.ch](mailto:patrick.wuethrich@insel.ch)

<sup>1</sup>Department of Anesthesiology and Pain Medicine, Inselspital, Bern University Hospital, Berne CH-3010, Switzerland



**Table 1** Electrolyte composition of the 2 different crystalloid solutions

	Ringerfundin®	G5-K Solution®
Natrium	145.0 mmol/l	50.0 mmol/l
Kalium	4.0 mmol/l	30.0 mmol/l
Magnesium	1.0 mmol/l	2.0 mmol/l
Calcium	2.5 mmol/l	0 mmol/l
Chlorid	127.0 mmol/l	65 mmol/l
Acetat	24.0 mmol/l	0 mmol/l
Maleat acid	5.0 mmol/l	0 mmol/l
Lactate	0 mmol/l	18 mmol/l
HPO <sub>4</sub>	0 mmol/l	8.0 mmol/l
Glucose	0 g/l	50 g/l