

N-terminal fusion of a toll-like receptor 2-ligand to a *Neospora caninum* chimeric antigen efficiently modifies the properties of the specific immune response

ADRIANA AGUADO-MARTÍNEZ^{1*}, AFONSO P. BASTO², JOACHIM MÜLLER¹, VRENI BALMER¹, VERA MANSER¹, ALEXANDRE LEITÃO² and ANDREW HEMPHILL¹

¹Institute for Parasitology, Vetsuisse Faculty, University of Berne, Länggass-Strasse 122, CH-3012 Bern, Switzerland

²CIISA, Faculdade de Medicina Veterinária, ULisboa, Avenida da Universidade Técnica, 1300-477 Lisboa, Portugal

(Received 13 October 2015; revised 24 December 2015; accepted 29 December 2015)

SUMMARY

Immunoprophylactic products against neosporosis during pregnancy should induce an appropriately balanced immune response. In this respect, OprI, a bacterial lipoprotein targeting toll like receptor (TLR)2, provides promising adjuvant properties. We report on the manipulation of the innate and the T-cell immune response through the fusion of OprI with the *Neospora caninum* chimeric protein Mic3-1-R. In contrast to Mic3-1-R, OprI-MIC3-1-R significantly activated bone-marrow dendritic cells from naïve mice. Mice immunized with OprI-Mic3-1-R induced an immune response with mixed T helper (Th)1 and Th2 properties (high levels of both immunoglobulin (Ig)G1 and IgG2a and of interleukin (IL)-10, IL-12(p70) and interferon- γ responses) whereas Mic3-1-R+saponin induced a clear Th2-biased response (low IgG2a and high IL-4 and IL-10). After mating and challenge with *N. caninum*, increased expression of interferon- γ was only found in placentas from OprI-Mic3-1-R immunized dams. However, no protection against vertical transmission and neonatal mortality was observed in either of the two groups. These results indicated that more exhaustive studies must be done to elucidate the immune mechanisms associated with transplacental transmission. Antigen linkage to TLR2-ligands, such as OprI, is a useful tool to investigate this enigma by reorienting the innate and adaptive immune responses against other candidate antigens in future studies.

Key words: Neosporosis, innate and adaptive immune response, outer membrane lipoprotein I, toll-like receptor 2, chimeric antigen.

INTRODUCTION

Neospora caninum is an obligate intracellular apicomplexan parasite closely related to *Toxoplasma gondii* and represents an important causative agent of abortions in cattle, worldwide (Reichel *et al.* 2014). The efficient transplacental transmission of *N. caninum* and the apparent incapability of the host immune response to prevent this process still remain the most intriguing aspects of this pathogen despite many attempts by different immunization approaches in both cattle and the pregnant mouse model.

In pregnant animals, physiological changes lead to immunomodulation especially at the maternal-fetal interface, characterized by a downregulated T helper (Th) 1-biased response through decreasing interferon gamma (IFN- γ), tumour necrosis factor (TNF) and interleukin (IL)-12 production and increasing IL-4 and IL-10 expression (Roberts *et al.* 2001; Beagley and Gockel, 2003; Kano *et al.* 2007) and an induction of a regulatory phenotype in both

antigen presenting cells (APCs) and T cells (Fiorentino *et al.* 1991; Rowe *et al.* 2011). This profile prevents fetal rejection and helps maintaining the pregnancy but, at the same time, favours parasite reactivation, multiplication and subsequent transmission to the fetus (Roberts *et al.* 2001; Quinn *et al.* 2004). On the other hand, if dams mount a Th1-biased immune response to control parasite proliferation, the pro-inflammatory cytokines could damage the placenta and compromise the pregnancy. Thus, an appropriately balanced Th1/Th2 immune response appears to be crucial in pregnant cattle (Innes *et al.* 2007) and in experimentally infected mice (Monney and Hemphill, 2014). However, the achievement of this specific immune balance in *N. caninum* infected dams remains a big challenge. This is reflected by the fact that the vast majority of antigen formulations shown to confer protection in the non-pregnant neosporosis mouse model were rendered non-protective when assessed in pregnant mice immunized prior to mating (Monney and Hemphill, 2014). A good example is the *N. caninum* chimeric antigen Mic3-1-R which conferred excellent degree of protection in non-pregnant mice but failed to protect against vertical transmission and offspring death in the pregnant

* Corresponding author: Institute of Parasitology, University of Bern, Länggass-Strasse 122, 3012 Bern, Switzerland. Tel: +41 31 631 2593. Fax: +41 31 631 2477. E-mail: adriana.aguado@vetsuisse.unibe.ch

mouse model of neosporosis (Debache *et al.* 2010; Monney *et al.* 2011, 2012; Debache and Hemphill, 2013).

The OprI molecule is a lipoprotein from the outer membrane of *Pseudomonas aeruginosa*, which targets toll-like receptor 2 (TLR2) and therefore exhibits adjuvant properties (Basto *et al.* 2012). TLR2 activation has been associated with stimulation of Th1, Th2 and T regulatory (Treg) type immune responses. Moreover, TLR2 activation has been implicated in the induction of cross-presentation by APCs (Basto and Leitao, 2014; Basto *et al.* 2015). In addition, *N. caninum* recognition by TLR2 in mice has been shown to participate in the generation of an effective immune response (Mineo *et al.* 2010). The capacity of OprI to modulate adaptive immune responses *in vitro* and *in vivo* has been demonstrated in virus infections such as African swine fever, bacterial infections such as tuberculosis, and parasite infections such as *Leishmania major* (Leitao *et al.* 2000; Cote-Sierra *et al.* 2002; Gartner *et al.* 2007). Thus, OprI directly linked to the antigen emerges as an interesting adjuvant candidate for a sub-unit vaccine against neosporosis in pregnant mice, especially since a well-adjusted Th1/Th2 balance is required avoiding excessive inflammation.

In the present study, OprI was fused to the chimeric *N. caninum* antigen Mic3-1-R employing a previously developed cloning and production system (Basto *et al.* 2012). The potential of this new immunogenic formulation to activate the innate immunity and modulate the humoral and T cell immune response and the influence on congenital neosporosis in the pregnant mouse model was analysed and discussed.

MATERIALS AND METHODS

Neospora caninum culture, inoculum and crude extract preparation

Neospora caninum of the Nc-Spain7 isolate (Regidor-Cerrillo *et al.* 2008) was grown *in vitro* by continuous passages in Vero cell cultures. Infected cultures were maintained in RPMI 1640 medium supplemented with 2% heat-inactivated fetal calf serum (FCS), 2% L-glutamine, 1% antibiotic-antimycotic (100 U of penicillin, 100 µg of streptomycin and 0.25 µg of amphotericin B mL⁻¹) at 37 °C and 5% CO₂. The mouse inoculum was prepared as described by Arranz-Solís *et al.* (2015). Infected monolayers with more than 90% of tachyzoites being intracellular were removed from the surface with a cell scraper, passed through a 21-gauge needle to break the vacuoles, and viable tachyzoites (diluted 1:10 in Trypan blue) were counted in a Neubauer chamber. The tachyzoite suspension was diluted with culture medium to achieve a final

concentration of 5 × 10⁵ viable tachyzoites mL⁻¹. Parasites were maintained in ice and mice were infected by subcutaneous injection of 200 µL suspension (corresponding to 10⁵ tachyzoites) within the following 30 min. To prepare *N. caninum* crude extract for enzyme-linked immunosorbent assay (ELISA), infected Vero cells were scraped, passed through a 21-gauge needle and the tachyzoites purified on Sephadex G25 PD-10 columns (GE Healthcare, Buckinghamshire, England). Tachyzoites were then pelleted by centrifugation at 300 × g for 15 min at 4 °C, subjected to three freeze-thaw cycles, sonicated for 6 × 20 s at 57 W in a Branson Sonifier Cell Disruptor (Skan AG, Allschwill, Switzerland) and filtered through a 20 µm sterile filter. Final protein concentration was determined using the Protein Assay Dye Reagent Concentrate (Bio-Rad, Hercules, CA, USA).

Production of OprI-Mic3-1-R and Mic3-1-R antigens

The complete coding sequence of the chimeric *N. caninum* antigen Mic3-1-R (Monney *et al.* 2011) was cloned in the pOLP plasmid at the multiple cloning site-2 (MCS2) downstream of OprI sequence (Basto *et al.* 2012). OprI-MIC3-1-R was expressed in Rosetta(DE3)pLysS *Escherichia coli* strain (Novagen) and subsequently extracted from the outer membranes in order to exclusively purify the lipidated mature form. The entire procedure was done according to Basto *et al.* (2014); Basto *et al.* (2012). Briefly, after isopropyl-β-D-thiogalactopyranoside (IPTG) induction, the bacterial outer membranes were isolated by differential solubilization in sarkosyl, and lipopolysaccharides (LPS) were extracted by hot phenol/water treatment. The resulting outer membrane proteins contained in the phenol phase were precipitated, washed and solubilized in a binding buffer (20 mM NaH₂PO₄, 0.5 M NaCl, 20 mM imidazole, 6 M guanidine hydrochloride, 2% (v/v) Triton X-100, pH 7.4) supplemented with 20 mM β-mercaptoethanol and 1 × protease inhibitor cocktail (for use in purification of His6-tagged proteins, dimethyl sulfoxide solution, Sigma), and were purified by nickel-affinity chromatography (Basto *et al.* 2014). The non-OprI Mic3-1-R antigen was produced as previously described by Monney *et al.* (2011) with some modifications. The pET151-Mic3-1-R construct was transformed into BL21 (DE3) competent cells. After IPTG induction, bacteria were re-suspended in the binding buffer supplemented with 20 mM β-mercaptoethanol and 1 × of protease inhibitor cocktail and ultra-sonicated (2 × 10 min at 57 W). The lysate was gently shaken over night at 4 °C, clarified by centrifugation (3000 × g, 30 min) and submitted to hot phenol/water treatment. The proteins contained in the phenol phase were then precipitated, washed and solubilized in binding buffer and submitted to nickel-affinity chromatography as

above (Basto *et al.* 2014). The eluates of both purified recombinant proteins were dialyzed against phosphate-buffered saline (PBS). The protein contents were analysed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and quantified by the bicinchoninic acid (BCA) Protein Assay Kit (Thermo Scientific Pierce). Endotoxin levels were quantified by the Limulus Amebocyte Lysate (LAL) Kinetic-QCL (Lonza) following manufacturer's instructions. The final LPS content was below 0.13 EU μg^{-1} in purified OprI-Mic3-1-R and 0.03 EU μg^{-1} in Mic3-1-R.

Murine bone marrow-derived dendritic cells (BM-DCs) stimulation by OprI-Mic3-1-R and Mic3-1-R

All mouse procedures were conducted in compliance with the Swiss and European Union (Directive no. 86/609/EEC) legislation for the use of animals for experimental purposes and approved by the veterinary office of the Canton of Bern (license number 115/14). The authors certified the use of the minimum number of animals to produce statistically reproducible results.

BM-DCs were generated *in vitro* from bone marrow cells obtained from naïve BALB/c mice as previously described (Basto *et al.* 2015). After 7 days differentiating with granulocyte macrophage-colony stimulating factor (GM-CSF)-containing DC medium (RPMI 1640 medium containing 2 mM L-glutamine, 100 IU mL^{-1} penicillin, 100 $\mu\text{g mL}^{-1}$ streptomycin, 10% FCS, 20 mM HEPES, 1 mM sodium pyruvate, 50 μM β -mercaptoethanol, 20 ng mL^{-1} GM-CSF), non-adherent and loosely adherent cells were cultured and stimulated for 24 h with OprI-Mic3-1-R or Mic3-1-R (1 and 5 $\mu\text{g mL}^{-1}$), ultrapure LPS at 0.1 $\mu\text{g mL}^{-1}$ (Invivogen) or DC medium. Culture supernatants were collected and tested by ELISA for TNF- α according to manufacturer's instructions (eBioscience).

Immunization schedule, and immune response analysis

Six weeks-old BALB/c mice were purchased from Charles River Laboratories (Sulzfeld, Germany) and maintained under conventional day/night cycle housing conditions with water and food *ad libitum*. Euthanasia was performed in a CO₂ chamber. The mice were used for the experiment after 2 weeks of acclimatization.

Groups of 25 females mice were immunized three times, at 2-week intervals, by intraperitoneal injection with either 10 μg OprI-MIC3-1-R in 100 μL PBS (OprI-M group) or 10 μg MIC3-1-R plus 5 μg of saponin emulsified in 100 μL PBS (M+S group) (Monney *et al.* 2011). Saponin was chosen as a classic adjuvant because it was previously employed together with Mic3-1-R for immunization protocols allowing a comparable group between this

and previous studies (Monney *et al.* 2011, 2012). A third group (PBS) received PBS only. Six days after the third inoculation, five mice per group were euthanized. Blood was collected by intra-cardiac puncture and serum recovered for immunoglobulin G (IgG) analysis. The spleen was aseptically removed for splenic lymphocytes re-stimulation *in vitro*.

Challenge with N. caninum, clinical monitoring and sample collection

Two weeks after the third inoculation, the remaining 20 immunized mice per group were submitted to oestrus synchronization by the Whitten effect (Whitten, 1957) and mated with males (2 females and 1 male per cage) during three nights. At mid-gestation (7–9 days of pregnancy), mice were subcutaneously challenged with 10⁵ tachyzoites of Nc-Spain7 isolate. Other two groups of non-immunized animals received either the same dose of tachyzoites or PBS (PBS/INF and PBS/PBS, respectively). Two pregnant mice per group were euthanized on day 18 of pregnancy and placentas were snap-frozen in liquid nitrogen and stored at –80 °C for RNA isolation. Surviving non-pregnant mice were euthanized at day 30 post-infection (pi). Dams and their pups were maintained together until day 30 post-partum (pp) (41–44 pi) when all surviving animals were euthanized.

Data on pregnancy rate (percentage of pregnant mice), litter size (average of number of pups born per dam), neonatal mortality (stillborn pups or pups that died within the first two days pp) and post-natal mortality (pups dead between day 3 and 30 pp) were recorded. All animals were examined daily for clinical signs (pups twice per day) until the end of the experiment. All animals exhibiting nervous signs and/or a significant loss of body condition were immediately euthanized. Upon euthanasia, blood was extracted by cardiac puncture and sera stored at –80 °C. Brains from non-pregnant mice, dams and surviving pups were also collected and immediately frozen at –20 °C.

Cerebral parasite burden

Brains from non-pregnant mice, dams and surviving pups were analysed by *N. caninum*-specific real-time polymerase chain reaction (PCR) on Nc5 region of *N. caninum* DNA as previously described (Muller *et al.* 2002). DNA extraction from brain tissue was performed using the DNeasy[®] Blood & Tissue kit (Qiagen) according to the manufacturer recommendations. The DNA concentration in all samples was determined using the QuantiFluor dsDNA System (Promega, Madison, Wi.) according to the manufacturer protocol and adjusted to 5 ng μL^{-1} with sterile DNase free water. Quantitative real-time PCR was performed using the Rotor-Gene

6000 real-time PCR machine (Corbett Research, Qiagen). The parasite load was calculated by interpolation from a standard curve with DNA equivalents from 1000, 100 and 10 *N. caninum* tachyzoites included in each run.

Splenic lymphocytes re-stimulation in vitro

Freshly recovered spleens of mice sacrificed 6 days after the last immunization were aseptically disaggregated in full RPMI medium (RPMI 1640 medium containing 1% antibiotic-antimycotic-100 U of penicillin, 100 µg of streptomycin and 0.25 µg of amphotericin B mL⁻¹, 2% L-glutamine, 10% FCS and 55 µM β-mercaptoethanol) and by passing through a sieve. Red blood cells were lysed (1X RBC Lysis Buffer, ebiosciences) and the resulting cell suspensions were washed twice with full RPMI media, seeded in a 100-mm Petri dish and were incubated for 2 h at 37 °C with 5% CO₂ to remove adherent cells. Non-adherent cells were cultured at 1 × 10⁶ cells mL⁻¹ and re-stimulated with concanavalin A (ConA; 2 µg mL⁻¹; Sigma), Mic3-1-R (4 µg mL⁻¹) or remained non-stimulated (negative control). Supernatants were collected after 72 h and tested for cytokines by multiplex bead immunoassay.

Assessment of levels of cytokines by multiplex bead immunoassay

Levels of IFN-γ, IL-4, IL-10, IL-12(p70) and IL-17 were measured in the supernatants of re-stimulated splenic lymphocytes using the MILLIPLEX[®] MAP Mouse Cytokine/chemokine kit (Millipore Corp., Billerica, MA, USA) according to the manufacturer's protocol. Microtiter filter plates were run on Luminex instruments (BioPlex[™] 200 system). Calibration curves from the provided recombinant standards were calculated with Bio-Plex Manager software using a five parametric logistic curve fitting. When the cytokine concentration was below the detection limit, an arbitrary value corresponding to the detection limit of undiluted samples (provided by the kit manufacturer) was used for statistical analysis.

RNA levels of IFN-γ in placental tissues

IFN-γ transcript expression levels in placental tissues were measured by real-time quantitative reverse transcription-PCR (RT-PCR) in relation to the expression of β-actin transcripts. Frozen placentas were directly disrupted and homogenized in lysis buffer RLT containing β-mercaptoethanol (10 µL mL⁻¹ of buffer RLT) by rapid agitation in the presence of 1.4 mm ceramic beads (Lysing Matrix D, MP Biomedicals, Switzerland) in a FastPrep[®]-24 Instrument (MP Biomedicals,

Switzerland). Homogenates corresponding to no more than 30 mg of tissue were used for subsequent RNA isolation using the RNeasy[®] Mini kit (Qiagen) according to the manufacturer recommendations. An in-column DNA digestion step with DNase I (Qiagen) was included. RNA concentration was measured by spectrophotometry at 260 nm and purity was estimated by measuring the ratio 260/280 nm. Synthesis of complementary DNA (cDNA) was performed by Omniscript RT kit (Qiagen) using random primers and RNasin (Promega). The primers sequences for cDNA amplification were as follows: for IFN-γ forward 5'-GGAAGTGGCAAAGGATGG-3' and reverse 5'-CTGTGGGTTGTTGACCTCA-3'; and for β-actin forward 5'-ACACCCGCCACCAGTTTCG-3' and reverse 5'-CCATTCCCACCATCACACC-3'. Real-time PCR was carried out with FastStart Essential DNA Green Master (Roche) using SYBR green reagent in a Rotor-Gene 6000 real-time PCR machine (Corbett Research, Qiagen). For both genes a standard curve of Log of input cDNA vs cycle threshold (Ct) was built using a positive control cDNA and the slope calculated to know the PCR efficiencies. The amplification efficiencies of IFN-γ and β-actin were employed to calculate the mRNA levels of each gene for each sample as 10^{Ct/slope}. The final result was expressed in arbitrary units (AU) after normalizing mRNA levels of IFN-γ with mRNA levels of β-actin (Muller *et al.* 2007). All samples were analysed in triplicate.

Analysis of serum Ig by ELISA

Levels of *N. caninum*-specific or Mic3-1-R-specific IgG1 and IgG2a were measured by ELISA as previously described (Monney *et al.* 2011). Briefly, 96-well plates were coated over night at 4 °C with 200 ng of *N. caninum* crude extract or 100 ng of Mic3-1-R in 100 µL of coating buffer per well. After three washes, plates were blocked with 1% bovine serum albumin in wash buffer (PBS with 0.05% Tween-20 (v/v)) and serum samples incubated at 1:100. After three washes, plates were incubated with either goat anti-mouse IgG1 or IgG2a conjugated with alkaline phosphatase (AP) (SouthernBiotech, Birmingham, USA). Reaction was developed with AP substrate and absorbance values as optical density (OD) at 405 nm read in a tunable microplate reader (EnSpire[™] 2300 Multilabel Reader, Switzerland). In order to compare OD values between samples analysed in different plates, the same positive and negative serum controls were added in each plate and OD values for each sample were converted into a relative index per cent (RIPC) using the following formula RIPC = (OD₄₀₅ sample – OD₄₀₅ negative control)/(OD₄₀₅ positive control – OD₄₀₅ negative control) × 100.

Statistical analysis

Cytokine levels, symptoms score and parasite burden in brain between groups were compared by the non-parametric Kruskal–Wallis test; when statistical differences were detected a Dunn's multiple comparison post-test was applied to compare pair-by-pair. RIPC values of IgG1 and IgG2a between groups were compared by one-way analysis of variance (ANOVA) followed by a Tukey's Multiple Comparison test to compare pair-by-pair. Ratios IgG1:IgG2a were compared in each group between non-pregnant mice and dams by Mann–Whitney *U* test. To compare the mortality along time of dams and pups, survival proportions at each time-point were plotted and survival curves were compared by Log-rank test. Statistical significance for all analyses was established at $P < 0.05$. All statistical analyses were carried out using GraphPad Prism 6 (v.6.01) software.

RESULTS

OprI fused to Mic3-1-R induced a strong activation of naïve BM-DCs

In order to investigate whether the fusion of Mic3-1-R with OprI leads to the activation of BM-DCs, we stimulated BM-DCs from naïve mice with either OprI-Mic3-1-R or Mic3-1-R and measured TNF- α released in the supernatant after 24 h. While Mic3-1-R only marginally activated BM-DCs, OprI-Mic3-1-R induced a potent BM-DCs activation. At $1 \mu\text{g mL}^{-1}$ of OprI-Mic3-1-R, TNF- α levels were already higher than those induced by Mic3-1-R at $5 \mu\text{g mL}^{-1}$. At $5 \mu\text{g mL}^{-1}$ of OprI-Mic3-1-R, TNF- α levels were almost as high as in LPS-activation (Fig. 1).

OprI-Mic3-1-R induced an immune response with Th1/Th2 properties while Mic3-1-R + saponin a Th2-biased response

*OprI-Mic3-1-R induced significantly higher *N. caninum*-specific IgG2a than Mic3-1-R + saponin.* Specific Ig against *N. caninum* crude extract induced by the immunogenic formulations were analysed in sera from immunized mice (Fig. 2). Mice from the OprI-Mic3-1-R group (OprI-M) and Mic3-1-R + saponin group (M + S), but not from the PBS-treated group, developed *N. caninum*-specific IgGs. Both groups developed high and similar levels of IgG1 ($P > 0.05$) but mice immunized with OprI-M developed higher IgG2a levels than mice immunized with M + S ($P < 0.05$) (Fig. 2). Both OprI-M and M + S immunized mice developed equal and high levels of both IgG1 and IgG2a against the recombinant Mic3-1-R (data not shown).

OprI-Mic3-1-R induced a cytokine profile with mixed Th1/Th2/Th17 properties whereas Mic3-1-R induced a Th2-biased response. In order to address the




Fig. 1. *In vitro* naïve BM-DCs stimulation with OprI-Mic3-1-R, Mic3-1-R recombinant proteins (1 and $5 \mu\text{g mL}^{-1}$), LPS (100 ng mL^{-1}) or DC-full medium (Medium). Bars represent the TNF- α mean value of triplicates and the error bars the standard error of the mean (S.E.M.). Abbreviations: BM-DCs, bone marrow-derived dendritic cells; LPS, lipopolysaccharides; TNF, tumour necrosis factor.




Fig. 2. Specific IgG1 and IgG2a response against *Neospora caninum* crude extract measured by ELISA of mice immunized either with the OprI-Mic3-1-R recombinant protein (OprI-M) or with Mic3-1-R emulsified in saponin (M + S). Naïve mice (PBS) were included as a negative control. Sera were collected 6 days after the third immunization. Results are expressed as the mean of RIPC values (relative index per cent) and error bars represent the standard error of the mean (S.E.M.) in each group. * indicates significantly lower than OprI-M (* is $P < 0.05$; *** is $P < 0.001$). Abbreviations: ELISA, enzyme-linked immunosorbent assay; IgG, immunoglobulin G; PBS, phosphate-buffered saline; RIPC, relative index per cent.

polarization of cellular immune response induced by the immunogenic formulations, splenocytes were re-stimulated *in vitro* with Mic3-1-R antigen and cytokines quantified in the supernatants. Splenocytes of both the OprI-M group and M + S group exhibited a clearly distinct Th profile (Fig. 3). A significant increase of IFN- γ , IL-4, IL-




Fig. 3. Cytokine responses in immunized mice upon splenocyte restimulation with Mic3-1-R *in vitro*. (A) IFN- γ , (B) IL-4, (C) IL-10, (D) IL-12p70 and (E) IL-17. Splenocytes were obtained from mice 6 days after the third immunization either with OprI-Mic3-1-R recombinant protein (OprI-M), Mic3-1-R together with saponin (M + S) or with PBS (PBS). Boxes represent median, first and third quartiles and whiskers the 10th and 90th percentiles values. * indicates significantly higher values compared with the PBS group for each cytokine. (* is $P < 0.05$; ** is $P < 0.01$; *** is $P < 0.001$). Abbreviations: IFN- γ , interferon gamma; IL, interleukin; PBS, phosphate-buffered saline.

10, IL-12(p70) and IL-17 levels was detected in supernatants of splenocytes of OprI-M-immunized animals ($P < 0.05$) (Fig. 3). On the other hand, no significant levels of IFN- γ , IL-12(p70) and IL-17 were detected in cultures of M + S-immunized mice, but instead a significant increase of IL-4 and IL-10 levels was found compared with the PBS group ($P < 0.05$). In addition, IL-4 levels were significantly higher compared with splenocyte cultures of OprI-M mice ($P < 0.05$) (Fig. 3).

Placental IFN- γ was upregulated in OprI-Mic3-1-R mice but not in Mic3-1-R after challenge with N. caninum at mid-gestation. Two pregnant mice per group were euthanized 2 days prior to birth, and placentas were collected to quantify RNA expression levels of IFN- γ by RT-PCR. IFN- γ transcript expression was increased in placentas from both dams immunized with OprI-M compared with M + S-immunized dams (Fig. 4).

After challenge with N. caninum, non-pregnant mice immunized with OprI-Mic3-1-R, but not those immunized with Mic3-1-R, elicited lower IgG1:IgG2a ratios than dams. IgG1 and IgG2a responses against Mic3-1-R recombinant protein were analysed by ELISA (Fig. 5A). In non-pregnant mice, IgG1 levels from OprI-M and M + S groups were statistically higher than those from non-vaccinated infected mice (PBS/INF group) ($P < 0.05$). In case of IgG2a,




Fig. 4. mRNA expression levels of IFN- γ in placenta from two dams per group euthanized at day 18 of pregnancy. Mice were immunized either with OprI-Mic3-1-R recombinant protein (OprI-M), Mic3-1-R together with saponin (M + S) or with PBS (PBS/INF) and were challenged at day 7–9 of pregnancy. A non-immunized control group remained unchallenged (PBS/PBS). Relative expression of IFN- γ is expressed in AU after normalizing with mRNA expression levels of β -actin. Each dam was analysed by triplicate (dot) and bars represent the median value. Abbreviations: AU, arbitrary units; IFN- γ , interferon gamma; PBS, phosphate-buffered saline.

only OprI-M elicited significantly higher levels than PBS/INF group ($P < 0.05$). In dams, both IgG1 and IgG2a levels from OprI-M and M + S were statistically higher than those from PBS/INF ($P < 0.0001$). When IgG1:IgG2a ratios (Fig. 5B) were compared




Fig. 5. Humoral immune response against Mic3-1-R recombinant protein measured by ELISA of non-pregnant mice and dams after challenge with *Neospora caninum*, on days 30 and 41–44 pi, respectively. (A) represents the mean of RIPC values (relative index percent) of IgG1 and IgG2a levels. (B) shows the IgG1:IgG2a ratios for each group of mice. OprI-M: mice immunized with OprI-Mic3-1-R; M + S: mice immunized with Mic3-1-R with saponin; PBS/INF: mice non-immunized. These three groups were challenged with *N. caninum* at day 7–9 of pregnancy. PBS/PBS: non-immunized, non-challenged. Error bars represent the standard error of the mean (S.E.M.) in each group. (** is $P < 0.01$ between IgG1:IgG2a ratios from non-pregnant mice and dams). Abbreviations: ELISA, enzyme-linked immunosorbent assay; INF- γ , interferon gamma; IgG, immunoglobulin G; IL, interleukin; PBS, phosphate-buffered saline; pi, post-infection; RIPC, relative index per cent.

Table 1. Reproductive performance and outcome of *Neospora caninum* infection in dams

Group	Pregnancy rate (%) ^a	No. dams ^b	Litter size ^c	Clinical signs (%) ^d	Mortality (%) ^e	Nc DNA in brain (%) ^f
OprI-M	15/20 (75)	13	7.1	11/13 (84.6)	4/13 (30.7)	13/13 (100)
M + S	16/20 (80)	14	6.1	10/14 (71.4)	5/14 (35.7)	14/14 (100)
PBS/INF	15/20 (75)	13	6.1	10/13 (76.9)	2/13 (15.4)	13/13 (100)
PBS/PBS	14/20 (70)	12	6.2	0/12 (0)	0/12 (0)	0/12 (0)

INF, interferon; PBS, phosphate-buffered saline.

^a Number of pregnant mice/number of mice in the group (%).

^b Excluding two dams per group that were euthanized before giving birth in order to obtain placentas.

^c Number of delivered pups per dam.

^d Proportion of dams presenting at least one clinical sign (%).

^e Proportion of dams euthanized upon severe clinical signs (%).

^f Proportion of dams tested positive for *N. caninum* DNA in the brain (%).

between non-pregnant mice and dams, the OprI-M group showed higher ratios in dams compared with non-pregnant mice ($P < 0.001$) whereas this difference was not detected in either the M + S group or the PBS/INF group ($P > 0.05$). These results suggest that after challenge only mice immunized with OprI-Mic3-1-R that did not become pregnant were able to maintain a low ratio IgG1:IgG2a, but this tendency was overridden after pregnancy.

Neither OprI-Mic3-1-R nor Mic3-1-R immunization led to reduced cerebral infection in adults or vertical transmission to offspring

The effects of immunization of mice with OprI-Mic3-1-R and Mic3-1-R were assessed in challenge

experiments. With regard to pregnancy rate and litter size, no differences among groups were detected ($P > 0.05$) (Table 1). The antigen formulations did not induce protection against neosporosis neither in adults nor in offspring. Between 71.4 and 84.6% of dams from the three infected groups showed clinical signs (Table 1). Mortality rates were between 35.7% (5 in 14 dams) in the M + S group and 15.4% (2 in 13 dams) in the PBS/INF group, nevertheless the differences in the survival curves were not statistically significant ($P > 0.05$) (Table 1, Fig. 6A). *N. caninum* DNA was detected in the brains of all infected dams and no statistically significant differences in parasite burden were detected among immunized groups and non-immunized infected group (Fig. 6B). All non-pregnant




Fig. 6. Outcome of *Neospora caninum* infection in dams. (A) Kaplan-Meier survival curves of dams. (B) Cerebral parasite burden measured by real-time PCR in the dams. Boxes represent median, first and third quartiles; whiskers 10th and 90th percentiles; and circles outlier values. OprI-M: mice immunized with OprI-Mic3-1-R; M + S: mice immunized with Mic3-1-R with saponin; PBS/INF: mice non-immunized. These three groups were challenged with *N. caninum* at day 7–9 of pregnancy. PBS/PBS: non-immunized, non-challenged. Abbreviations: INF, interferon; PBS, phosphate-buffered saline; PCR, polymerase chain reaction.

Table 2. Outcome of *Neospora caninum* infection in pups

Group	No. pups ^a	Neonatal mortality (%) ^b	Postnatal mortality (%) ^c	Vertical transmission (%) ^d
OprI-M	91	13/91 (14.3)	77/78 (98.7)	78/78 (100)
M + S	79	11/79 (13.9)	67/68 (98.5)	68/68 (100)
PBS/INF	70	6/70 (8.6)	63/64 (98.4)	63/64 (98.4)
PBS/PBS	71	3/71 (4.2)	0/68 (0)	0/68

INF, interferon; PBS, phosphate-buffered saline.

^a Total number of pups born (alive or dead) from each group.

^b Proportion of pups born dead or that died within the first 2 days post-partum (%).

^c Proportion of pups dead from day 3 to 30 pp (%).

^d Proportion of dead pups from day 3 to 30 pp plus *N. caninum*-polymerase chain reaction (PCR) positive surviving pups (%). Dead pups are considered positive for *N. caninum* PCR as shown previously (Dellarupe *et al.* 2014).

animals were also positive by *N. caninum*-PCR and no differences in parasite burden in brain were detected (data not shown). Offspring mortality in the first 30 days pp was almost 100% in the three challenged groups without differences in the survival curves ($P > 0.05$) (data not shown). Vertical transmission could not be prevented; only the survivor from the PBS/INF group was found *N. caninum*-PCR negative (Table 2).

DISCUSSION

The cross-talk between innate and acquired immunity is a key aspect in the development of efficient immune responses against pathogens (Basto and Leitao, 2014). The activation of APCs by pathogen-associated molecular patterns (PAMPs) through TLRs leads to an efficient activation of T cells (Akira, 2009) but this aspect has not been considered frequently when potentially protective antigens have been investigated in anti-*N. caninum* immunization studies. In this work we investigated the effect on the innate immunity and the subsequent acquired immune response of the TLR2-

ligand OprI fused to the *N. caninum* Mic3-1-R, a chimeric antigen composed by predicted epitopes of NcMIC3, NcMIC1 and NcROP2 secretory proteins (Monney *et al.* 2011).

We have successfully exploited the recently developed cloning system, lipoprotein expression and depyrogenation methods (Basto *et al.* 2012, 2014) for N-terminal fusion of the OprI lipoprotein to *N. caninum* Mic3-1-R. The recombinant lipoprotein was exclusively purified from the outer membrane of the bacteria which guaranteed the presence of only the lipidated mature form, which is the TLR2-binding active form (Cote-Sierra *et al.* 2002), in the final purified elution (Basto *et al.* 2012).

Among the functions that have been described for TLR2 activation are the promotion of both Th1 and Th2 responses together with Treg activation (Basto and Leitao, 2014). The combination of these apparently antagonistic effects of TLR2 activation attracted our attention for *N. caninum* vaccination especially in the context of pregnancy. TLR2-ligands induce DC activation and the secretion of pro-inflammatory cytokines. In addition, enhancement of direct- and cross-presentation of exogenous antigens by APCs

and both CD4+ and cytotoxic CD8+ T cell activation have been attributed to TLR2 engagement (Basto and Leitao, 2014; Basto *et al.* 2015). Although these mechanisms seem to be essential to eliminate intracellular pathogens such as *N. caninum*, the damage caused by direct killing of infected host cells and the associated tissue inflammation could be the cause of the protection failure of Th1-inducing immunogenic formulations in mice in the past, especially when applied in the pregnant mouse model (Debache and Hemphill, 2013; Monney *et al.* 2013). However, TLR2 activation has also been associated with Th2 cell activation via IL-12 suppression and IL-10 production as well as anti-inflammatory effects via Treg cell activation (Basto and Leitao, 2014). We hypothesized that the fusion of the OprI molecule to *N. caninum* antigens could be used as a tool to modulate the innate immunity and the T-cell response towards a well-adjusted Th1/Th2/Treg phenotype which could be essential for an eventual protective antigen formulation during pregnancy by diminishing tissue damage, whereas still killing the parasite or at least limiting its proliferation.

First, we could confirm that OprI after fusion with Mic3-1-R stimulates BM-DCs, since high levels of TNF- α were secreted by BM-DCs upon incubation with the recombinant protein. Although Mic3-1-R also induced higher levels of TNF- α than only medium, those levels were four times lower than LPS and OprI-Mic3-1-R. Residual levels of LPS in the Mic3-1-R final product (even after hot water/phenol extraction) could explain those levels of TNF- α upon Mic3-1-R incubation with BM-DCs. In contrast, the TNF- α levels induced by OprI-Mic3-1-R were similar to those secreted upon LPS stimulation demonstrating the specific capacity of OprI as APC-activation inductor and indeed its adjuvanticity when linked to Mic3-1-R. The APC activation by OprI has been previously shown for OprI lipoprotein alone (Basto *et al.* 2012) but also for other OprI-fused antigens including the model antigen ovalbumin (OVA) (Basto *et al.* 2015) and the gp63 protein of *Leishmania major* (Cote-Sierra *et al.* 2002).

In a second step, we immunized BALB/c mice with either OprI-Mic3-1-R or Mic3-1-R emulsified in saponin, and the *in vivo* antibody profile and cytokine profile after lymphocyte recall was studied. In accordance with what has been previously described (Monney *et al.* 2011, 2012), Mic3-1-R emulsified with saponin induced a strong Th2-biased response, characterized by high ratio IgG1:IgG2a and enhancement of IL-4 and IL-10 but not IFN- γ or IL-12(p70). In contrast, OprI-Mic3-1-R elicited an immune response with mixed Th1/Th2 properties. Levels of IL-4 were lower than those obtained from splenocytes derived from mice immunized with Mic3-1-R+saponin but levels of the anti-

inflammatory cytokine IL-10 were similar, or even higher, and accompanied by higher levels of IFN- γ and IL-12(p70). Interestingly, splenocytes from the OprI-Mic3-1-R immunized group but not from the Mic3-1-R + Saponin also produced IL-17, a cytokine typical of Th17 polarization and related with inflammatory processes. In addition, the IgG1:IgG2a ratio of *N. caninum*-specific IgG from OprI-Mic3-1-R mice was significantly lower than in Mic3-1-R + saponin mice. Basto *et al.* (2015) also described an immunomodulation towards a profile with mixed characteristics by OprI-OVA, however in that study IFN- γ production was absent in recall lymphocytes. These differences highlighted that although OprI is a potent molecule to re-orientate the immune response, this should be specifically analysed for each particular antigen.

After challenge with *N. caninum*, none of the formulations could prevent vertical transmission and congenital neosporosis, however, differences in the immune response could be detected between OprI and non-OprI immunized mice. IFN- γ transcripts were upregulated in placental tissues in OprI-Mic3-1-R immunized dams but not in Mic3-1-R + saponin dams. This highlights again the capability of OprI to modify the expected Th2-biased response associated with saponin-emulsified Mic3-1-R. It is possible that a non-antigen dependent effect of the OprI have contributed for this modulation. This and other possible effects of OprI TLR2/1 stimulation should be addressed in the future. The fact that in both cases vertical transmission took place points out the necessity to increase our understanding of the immune mechanisms which should be induced during pregnancy. Previously, a Th1-biased immune response maintained after challenge was associated with protection against *L. major* infection in mice immunized with OprI fused to the Gp63 antigen (Cote-Sierra *et al.* 2002). However, in congenital neosporosis a Th1 polarized immune response with a storm of inflammatory cytokines, especially in placenta tissues, could be detrimental in terms of vertical transmission and offspring survival (Monney *et al.* 2013; Canton *et al.* 2014). Nevertheless, a predominant Th2-biased response in pregnant mice has also been correlated with susceptibility to neosporosis (Quinn *et al.* 2004; Kano *et al.* 2005). Whether the induction of specific Treg cells localized in the fetal-maternal interface could help in preventing the deleterious effects of effector immune mechanisms on pregnancy is not yet studied. Interestingly, TLR2 activation has been previously shown to induce both the development of regulatory responses (Dillon *et al.* 2006; Manicassamy *et al.* 2009) and the homing of lymphocytes to mucosae (Wang *et al.* 2011). Thus, the OprI-based system can be a relevant tool to address these issues in neosporosis pregnant models.

The analysis of the humoral immune response after challenge revealed that the IgG1:IgG2a ratio of anti-Mic3-1-R antibodies was near to 1 in OprI-Mic3-1-R immunized non-pregnant mice but higher in dams from the same group. This difference was not observed in M+S and PBS/INF infected groups. This switch in the antibody profile between non-pregnant mice and dams exclusively observed in OprI-Mic3-1-R immunized mice suggested again that a balanced Th1/Th2 response induced by OprI-Mic3-1-R in non-pregnant mice was overridden at pregnancy. This apparent loss of the OprI-inducing immunomodulation during pregnancy could explain the lack of protection against vertical transmission and subsequent neosporosis in offspring.

In conclusion, we have demonstrated that the lipoprotein OprI is a potent tool to activate the innate immunity and to modulate the adaptive immune response when it is fused to a *N. caninum* antigen. However, the fusion of OprI with the chimeric Mic3-1-R antigen did not improve protection against transplacental transmission. Attempts to accurately predict target antigens and the appropriate polarization of the cellular immune response associated with protection have failed in most studies on *N. caninum* vaccines, especially in pregnant models. Thus, further studies should focus (i) on how pregnancy affects the immune response against vaccines that are applied prior to pregnancy, and (ii) on the identification of novel antigens which generate protective T cell responses (Yang *et al.* 1991; Rocchi *et al.* 2011), both of which could contribute to the development of an efficient vaccine. The use of OprI fusion will be helpful to fine-tune the innate and adaptive immune responses induced by these new antigen candidates, and to adapt them to the pregnancy-mediated immunomodulation.

ACKNOWLEDGEMENTS

We thank Dr. Luis Miguel Ortega-Mora for kindly providing the Nc-Spain7 isolate of *N. caninum* and Dr. Denis Grandgirard from the Institute for Infection Diseases of the University of Bern for the analysis of cytokines levels by multiplex bead immunoassay.

FINANCIAL SUPPORT

This work was supported by the Swiss National Science Foundation (A. Hemphill, grant No. 310030 146162) and by Fundação para a Ciência e a Tecnologia, Portugal (A.P. Basto and A. Leitão, grant No. PTDC/CVT/113889/2009).

REFERENCES

Akira, S. (2009). Pathogen recognition by innate immunity and its signaling. *Proceedings of the Japan Academy Series B Physical and Biological Sciences* **85**, 143–156.
Arranz-Solis, D., Aguado-Martínez, A., Müller, J., Regidor-Cerrillo, J., Ortega-Mora, L.M., Hemphill, A. (2015). Dose-dependent effects of

experimental infection with the virulent *Neospora caninum* Nc-Spain7 isolate in a pregnant mouse model. *Veterinary Parasitology* **211**, 133–140.
Basto, A. P. and Leitao, A. (2014). Targeting TLR2 for vaccine development. *Journal of Immunology Research* 2014, 619410. doi: 10.1155/2014/619410.
Basto, A. P., Badenes, M., Almeida, S. C., Martins, C., Duarte, A., Santos, D.M. and Leitao, A. (2015). Immune response profile elicited by the model antigen ovalbumin expressed in fusion with the bacterial OprI lipoprotein. *Molecular Immunology* **64**, 36–45.
Basto, A. P., Morais, J., Marcelino, E., Leitao, A. and Santos, D.M. (2014). An efficient depyrogenation method for recombinant bacterial outer membrane lipoproteins. *Protein Expression and Purification* **98**, 10–17.
Basto, A.P., Piedade, J., Ramalho, R., Alves, S., Soares, H., Cornelis, P., Martins, C. and Leitao, A. (2012). A new cloning system based on the OprI lipoprotein for the production of recombinant bacterial cell wall-derived immunogenic formulations. *Journal of Biotechnology* **157**, 50–63.
Beagley, K. W. and Gockel, C. M. (2003). Regulation of innate and adaptive immunity by the female sex hormones oestradiol and progesterone. *FEMS Immunology and Medical Microbiology* **38**, 13–22.
Canton, G.J., Konrad, J.L., Moore, D.P., Caspe, S.G., Palarea-Albaladejo, J., Campero, C.M. and Chianini, F. (2014). Characterization of immune cell infiltration in the placentome of water buffaloes (*Bubalus bubalis*) infected with *Neospora caninum* during pregnancy. *Journal of Comparative Pathology* **150**, 463–468.
Cote-Sierra, J., Bredan, A., Toldos, C.M., Stijlemans, B., Brys, L., Cornelis, P., Segovia, M., de Baetselier, P. and Revets, H. (2002). Bacterial lipoprotein-based vaccines induce tumor necrosis factor-dependent type 1 protective immunity against *Leishmania major*. *Infection and Immunity* **70**, 240–248.
Debache, K. and Hemphill, A. (2013). Differential effects of intranasal vaccination with recombinant NcPDI in different mouse models of *Neospora caninum* infection. *Parasite Immunology* **35**, 11–20.
Debache, K., Guionaud, C., Alaeddine, F. and Hemphill, A. (2010). Intraperitoneal and intra-nasal vaccination of mice with three distinct recombinant *Neospora caninum* antigens results in differential effects with regard to protection against experimental challenge with *Neospora caninum* tachyzoites. *Parasitology* **137**, 229–240.
Dellarupe, A., Regidor-Cerrillo, J., Jimenez-Ruiz, E., Schares, G., Unzaga, J.M., Venturini, M.C. and Ortega-Mora, L.M. (2014). Comparison of host cell invasion and proliferation among *Neospora caninum* isolates obtained from oocysts and from clinical cases of naturally infected dogs. *Experimental Parasitology* **145**, 22–28.
Dillon, S., Agrawal, S., Banerjee, K., Letterio, J., Denning, T.L., Oswald-Richter, K., Kasproicz, D.J., Kellar, K., Pare, J., van Dyke, T., Ziegler, S., Unutmaz, D. and Pulendran, B. (2006). Yeast zymosan, a stimulus for TLR2 and dectin-1, induces regulatory antigen-presenting cells and immunological tolerance. *Journal of Clinical Investigation* **116**, 916–928.
Fiorentino, D.F., Zlotnik, A., Vieira, P., Mosmann, T.R., Howard, M., Moore, K.W. and O'Garra, A. (1991). IL-10 acts on the antigen-presenting cell to inhibit cytokine production by Th1 cells. *Journal of Immunology* **146**, 3444–3451.
Gartner, T., Baeten, M., Otieno, S., Revets, H., De Baetselier, P. and Huygen, K. (2007). Mucosal prime-boost vaccination for tuberculosis based on TLR triggering OprI lipoprotein from *Pseudomonas aeruginosa* fused to mycolyl-transferase Ag85A. *Immunology Letters* **111**, 26–35.
Innes, E.A., Bartley, P.M., Maley, S.W., Wright, S.E. and Buxton, D. (2007). Comparative host-parasite relationships in ovine toxoplasmosis and bovine neosporosis and strategies for vaccination. *Vaccine* **25**, 5495–5503.
Kano, R., Masukata, Y., Omata, Y., Kobayashi, Y., Maeda, R. and Saito, A. (2005). Relationship between type 1/type 2 immune responses and occurrence of vertical transmission in BALB/c mice infected with *Neospora caninum*. *Veterinary Parasitology* **129**, 159–164.
Kano, R., Kudo, A., Kamiya, H., Kobayashi, Y., Maeda, R. and Omata, Y. (2007). C57BL/6 mice infected with *Neospora caninum* during administration of progesterone show bias toward type 2 immune response. *The Journal of Veterinary Medical Science* **69**, 1095–1097.
Leitao, A., Malur, A., Cartaxeiro, C., Vasco, G., Cruz, B., Cornelis, P. and Martins, C.L. (2000). Bacterial lipoprotein based expression vectors as tools for the characterisation of African swine fever virus (ASFV) antigens. *Archives of Virology* **145**, 1639–1657.
Manicassamy, S., Ravindran, R., Deng, J., Oluoch, H., Denning, T.L., Kasturi, S.P., Rosenthal, K.M., Evavold, B.D. and Pulendran, B. (2009). Toll-like receptor 2-dependent induction of vitamin A-metabolizing enzymes in dendritic cells promotes T regulatory responses and inhibits autoimmunity. *Nature Medicine* **15**, 401–409.

- Mineo, T. W., Oliveira, C. J., Gutierrez, F. R. and Silva, J. S. (2010). Recognition by Toll-like receptor 2 induces antigen-presenting cell activation and Th1 programming during infection by *Neospora caninum*. *Immunology and Cell Biology* **88**, 825–833.
- Monney, T. and Hemphill, A. (2014). Vaccines against neosporosis: what can we learn from the past studies? *Experimental Parasitology* **140**, 52–70.
- Monney, T., Debache, K., Grandgirard, D., Leib, S. L. and Hemphill, A. (2012). Vaccination with the recombinant chimeric antigen recNcMIC3-1-R induces a non-protective Th2-type immune response in the pregnant mouse model for *N. caninum* infection. *Vaccine* **30**, 6588–6594.
- Monney, T., Grandgirard, D., Leib, S. L. and Hemphill, A. (2013). Use of a Th1 stimulator adjuvant for vaccination against *Neospora caninum* infection in the pregnant mouse model. *Pathogens* **2**, 193–208.
- Monney, T., Rutti, D., Schorer, M., Debache, K., Grandgirard, D., Leib, S. L. and Hemphill, A. (2011). RecNcMIC3-1-R is a microneme- and rhoptry-based chimeric antigen that protects against acute neosporosis and limits cerebral parasite load in the mouse model for *Neospora caninum* infection. *Vaccine* **29**, 6967–6975.
- Muller, J., Sterk, M., Hemphill, A. and Muller, N. (2007). Characterization of *Giardia lamblia* WB C6 clones resistant to nitazoxanide and to metronidazole. *The Journal of Antimicrobial Chemotherapy* **60**, 280–287.
- Muller, N., Vonlaufen, N., Gianinazzi, C., Leib, S. L. and Hemphill, A. (2002). Application of real-time fluorescent PCR for quantitative assessment of *Neospora caninum* infections in organotypic slice cultures of rat central nervous system tissue. *Journal of Clinical Microbiology* **40**, 252–255.
- Quinn, H. E., Miller, C. M. and Ellis, J. T. (2004). The cell-mediated immune response to *Neospora caninum* during pregnancy in the mouse is associated with a bias towards production of interleukin-4. *International Journal for Parasitology* **34**, 723–732.
- Regidor-Cerrillo, J., Gomez-Bautista, M., Pereira-Bueno, J., Aduriz, G., Navarro-Lozano, V., Risco-Castillo, V., Fernandez-Garcia, A., Pedraza-Diaz, S. and Ortega-Mora, L. M. (2008). Isolation and genetic characterization of *Neospora caninum* from asymptomatic calves in Spain. *Parasitology* **135**, 1651–1659.
- Reichel, M. P., McAllister, M. M., Pomroy, W. E., Campero, C., Ortega-Mora, L. M. and Ellis, J. T. (2014). Control options for *Neospora caninum*—is there anything new or are we going backwards? *Parasitology* **141**, 1455–1470.
- Roberts, C. W., Walker, W. and Alexander, J. (2001). Sex-associated hormones and immunity to protozoan parasites. *Clinical Microbiology Reviews* **14**, 476–488.
- Rocchi, M. S., Bartley, P. M., Inglis, N. F., Collantes-Fernandez, E., Entrican, G., Katzer, F. and Innes, E. A. (2011). Selection of *Neospora caninum* antigens stimulating bovine CD4+ve T cell responses through immuno-potency screening and proteomic approaches. *Veterinary Research* **42**, 91.
- Rowe, J. H., Ertelt, J. M., Aguilera, M. N., Farrar, M. A. and Way, S. S. (2011). Foxp3(+) regulatory T cell expansion required for sustaining pregnancy compromises host defense against prenatal bacterial pathogens. *Cell Host & Microbe* **10**, 54–64.
- Wang, S., Villablanca, E. J., De Calisto, J., Gomes, D. C., Nguyen, D. D., Mizoguchi, E., Kagan, J. C., Reinecker, H. C., Hachon, N., Nagler, C., Xavier, R. J., Rossi-Bergmann, B., Chen, Y. B., Blomhoff, R., Snapper, S. B. and Mora, J. R. (2011). MyD88-dependent TLR1/2 signals educate dendritic cells with gut-specific imprinting properties. *Journal of Immunology* **187**, 141–150.
- Whitten, M. K. (1957). Effect of exteroceptive factors on the oestrous cycle of mice. *Nature* **180**, 1436.
- Yang, D. M., Rogers, M. V. and Liew, F. Y. (1991). Identification and characterization of host-protective T-cell epitopes of a major surface glycoprotein (gp63) from *Leishmania major*. *Immunology* **72**, 3–9.